COMP90043 Cryptography and security: Special Session on Week 10

(1) Division Algorithm

Refer to (4.1) of the textbook.

You could use the identities:

 $((x \mod m) + (y \mod m)) \mod m = (x + y) \mod m;$ $((x \mod m) * (y \mod m)) \mod m = (x * y) \mod m;$ Perform the following operations:

- (a) $(7-8) \mod 11 = -1 = 10$
- (b) $(2*4+7) \mod 5 = 0$
- (c) $(10 * 10 3) \mod 11 = 9$
- (d) $(1+2+3+4....9+10) \mod 11=0$
- (e) $(1+2+4+8+16) \mod 31 = -1 = 30$
- (2) Prove the following.
 - (a) Let a be represented as decimal number. Prove that (a mod 10) is simply ones-place in the decimal notation of a. Prove that (a mod 100) is simply the two digit number made up of the tens and ones place digits in the decimal notation of a.

Represent the number in decimal system and apply the identities given above. $10^k \mod 10$ is 0.

- (b) Prove that for all positive integers n, $2*(1+2+3+...+n) \mod (n+1)=0$. Use the identity: $1+2+\cdots n=n(n+1)/2$.
- (c) Prove that for all positive integer m, $(1+2+4+8+...+2^m) \mod (2^{(m+1)} = -1.$

Use the identity: $2^0 + 2^1 + \cdots + 2^m = 2^m + 1 - 1$.

- (3) Inverse (modulo n).
 - (a) By trial and error find multiplicative inverse of 43 mod 100. (How much time do you take it?). Easy exerciselet the inverse be y.
 - (b) Now find the inverse of 57 mod 100. 57 is (100-43), i.e -43, so the inverse of 57 is -ys
 - (c) Draw multiplication table showing a*b mod 7 for all a,b from 1 to 6.

Easy exercise, the purpose is to get a concrete example for a multiplicative group.

- (d) Use the gcd algorithm to solve the following:
 - (i) Find gcd(23,77).
 - (ii) gcd(11,100).
 - (iii) gcd(2023,3212).
 - (iv) gcd(7,31).
 - (v) gcd(101,17).

Easy exercise.

- (4) Extended GCD algorithm. Read the material about it on Page 137-139 of the textbook.
 - (a) Apply extended gcd algorithms for questions in 2.
 - (b) State extended gcd algorithm.
 - (c) Try out xgcd algorithm on magma http://magma.maths.usyd.edu.au/magma/
 - (d) Write a magma function for inverse modulo n using XGCD algorithm.
 - (e) Try out the following exercises from Stallings textbook:

- (f) Challenging probs: try out the following exercises from Stallings textbook:Q. 4.8, Q. 4.9 and Q. 4.11.
- (5) Factors and Divisibility:
 - (a) Find the factors of the following numbers: $31, 63, 2^{1}1 1$. You may need to use magma or simple calculations.
 - (b) Prove the Fermats theorem: For any a , <math>p a prime number,

$$a^(p-1) = 1 \bmod p$$

We worked out in a workshop-Also in the text-book.

- (c) Eulers Totient Function: Let $\phi(n)$ = number of integers less than n but relatively prime to n.
 - (i) Find $\phi(7) = 6$.
 - (ii) Find $\phi(35) = 6 \times 4 = 24$

- (iii) Find $\phi(p)$ for any prime p=p-1.
- (iv) Find $\phi(pq)$ for any prime p and q = (p-1)(q-1)
- (v) Prove that

$$a^{(\phi(n))} = 1 \bmod n$$

Eulers Theorem: We worked out in a workshop-Also in the textbook.

(6) Complexity of long integer arithmetic.

What are the complexities in big O notation for the following operations?

- (a) Addition of two k-bit integers. O(k).
- (b) Subtraction of two k-bit integers O(k).
- (c) Multiplication of two k-bit integers $O(k^2)$.
- (d) Division of a k-bit integer by another k-bit integer $O(k^2)$.
- (e) Greatest common divisor of two k-bit integers $O(k^2)$
- (f) Exponentiation $a^e \mod n$, where a and n are k bit integers and e a m bit integerw.O(m) k bit operations
- (7) Give examples of polynomails over finite fields and illustrate multiplication and division.
- (8) Use the irreducible polynomial $1+x^2+x^3$ in the the finite field GF(8) tab

i	Elements: x^i	As Polynomials	As Vectors
$-\infty$	0	0	[0, 0, 0]
0	1	1	[1, 0, 0]
1	x	x	[0, 1, 0]
2	x^2	x^2	[0, 0, 1]
3	x^3	$1 + x^2$	[1, 0, 1]
4	x^4	$1 + x + x^2$	[1,1,1]
5	x^5	1+x	[1, 1, 0]
6	x^6	$x + x^2$	[0, 1, 1]
7	x^7	1	[1, 0, 0]

Table 1. Elements of $GF(2^3)$ as powers of x

- (a) Solve $y * x^6 = 1$. y = x
- (b) Solve $y * x^4 = x^2 . y = x^5$

- (c) Compute $(x+x^2)*(x^2+x)$ **Answer** = $x^{12} = x^5 = 1+x$
- (9) Consider the finite field GF(9) as discussed in class last week:

i	Elements: x^i	As Polynomials	As Vectors
$-\infty$	0	0	[0, 0]
0	1	1	[1, 0]
1	x	x	[0, 1]
2	x^2	1 + 2 * x	[1, 2]
3	x^3	2+2x	[2, 2]
4	$x^4 \\ x^5 \\ x^6$	2	[2, 0]
5	x^5	2x	[0, 2]
6	x^6	2+x	[2, 1]
7	x^7	1+x	[1,1]
8	x^8	1	[1, 0]

Table 2. Elements of $GF(3^2)$ as powers of x

- (a) Solve $y * x^2 = 1$.
- (b) Solve $y * x^3 = x^2$.
- (c) Compute (x + 1) * (x + 2)

Try these yourselves similar to the method in the previous question.