

Sofia Chaoui, Laith Qushair, Tianji Sun

Professor Alshaykh

EC 601 A2 Product Design in Electrical and Computer Engineering

September 30, 2024

## Sprint 1: Multiple Sclerosis

### **Introduction:**

Medical error is the third leading cause of death in the United States. Misdiagnosis is a large contributor to this as it can prevent sick patients from receiving timely treatment, while exposing other patients to unnecessary and excessive treatment that often weakens their immune system and can lead to early death. When it comes to Multiple Sclerosis (MS), which affects around 900,000 people in the U.S. alone, it is estimated that the frequency of misdiagnosis is about 50% of the time. The discovery of a patient's false diagnosis often takes years, with 50% of patients carrying a misdiagnosis for at least 3 years, 33% for ten years or longer, and more than 5% for over 20 years. Due to the fact that doctors often begin MS disease-modifying therapy (DMT) as soon as they suspect MS, 31% of patients suffer from "unnecessary morbidity as a direct result of misdiagnosis" (Solomon et. al) of MS.

Multiple Sclerosis (MS) is a chronic neurological disorder. It is an autoimmune disorder which means the immune system for MS patients would mistakenly attack healthy cells instead of pathogens such as bacteria and viruses. Multiple Sclerosis affects people differently, some will have mild symptoms and others will have worsening symptoms that lead to increased disability over time.

Usually manifest from a relatively young age, MS symptoms usually begin between the ages of 20 and 40. MS patients typically have a normal life expectancy. Pathologically speaking, the MS immune system attacks myelin and the nerve cell bodies of the central nervous system. Myelin is a mixture of protein and fatty acids that makes the myelin sheath which covers and protects nerve fibers (axons). The individual symptoms of MS patients would vary depending on the severity of the attacks and the size of the plaques (patches of scar-like tissues that have sizes

from the head of a pin to a golf ball). MS has a wide range of symptoms including Muscle weakness, vision problems, mental or physical fatigue, mood, and cognitive changes.

Currently, there are no single tests to diagnose MS, different tests are used to confirm or rule out MS. Besides medical history examination, physical and neurological examination, MRI scan, and spinal cord are often used to look for MS characteristic lesions. Additionally, optic coherence tomography (OCT), cerebrospinal fluid tests, and evoked potential tests are commonly used for MS diagnosis. There is no cure for MS, but treatments can reduce the number and severity of relapses and delay long-term disease progressions. Clinically different symptoms will be managed by various treatments. Currently, much research on MS is being conducted, topics including finding new biomarkers for MS that can help diagnose and monitor MS, genetic and environmental risk factors, gut microbiome and diet, MS risk factors such as race and gender, etc. (National Institute of Health (NIH))

Diagnosing Multiple Sclerosis is a complex task due to the nature of the disease; overlapping symptoms with other diseases, a lack of specific diagnostic tests, variable presentation, and nonspecific symptoms are some of the many ongoing obstacles to proper diagnosis.

In 2017, the McDonald criteria utilized by clinicians and providers to aid in the differential of MS were revised. These revisions were only meant to be applicable to a specific subset of patients, however, several clinicians have unknowingly inappropriately applied McDonald criteria to “patients for whom they were not intended” (Solomon MD et. al). These revisions were based on a study performed on primarily white patients under the age of 50 with low co-illness rates, indicating that “application of the criteria should be approached cautiously” (Solomon MD et al.) for patients who do not fall within this demographic. Similarly, a study of 110 patients across the United States revealed that 85% of misdiagnoses were female patients and 15% were male patients, indicating that gender may play a role in misdiagnosis as well.

A study found that 89% of patients who were referred to an MS subspecialty center based primarily on an abnormal MRI were *not* later diagnosed with MS, but may have already started treatment. The McDonald criteria, amongst other factors such as overlapping symptoms of

related diseases, has consequently led to “misinterpretation and overreliance on MRI abnormalities of nonspecific neurologic symptoms” (Solomon, MD et al). Although MRI’s and other imaging technologies are helpful in aiding healthcare systems in providing an accurate diagnosis, identifying the cause of cortical lesions is extremely challenging as there are several common diseases that may present similarly on an MRI scan. As of 2024, there is still no specific biomarker for Multiple Sclerosis that has been identified exclusively of other diseases. It is a huge responsibility on a clinician to be able to differentiate subtle and near-invisible differences in lesions caused by MS and others caused by vascular disease, diabetes mellitus, tobacco use, and small vessel ischemic disease, which may all present in similar ways. There are suggestions that tiny hair-like veins on lesions, known as the Central Vein Sign (CVS), are as close to a biomarker for MS as we have gotten. However, AI has proven to be a lot better at identifying these tiny veins in imaging, and it may prove useful to clinicians to use AI tools in order to catch this. This opens up the possibility of reduced time of diagnosis and a personalized treatment plan for MS patients.

### **Artificial Intelligence in Medical Space**

The medical space has benefited tremendously from the introduction of artificial intelligence. These applications can take a variety of forms ranging from disease outbreak prediction models to image enhancement techniques for noisy medical imaging data such as Magnetic Resonance Imaging (MRI) data. Some of the main image artifacts that cause distortions in MR images include motion corruption and the Gibbs ringing artifact. Motion corruption is caused by slight movements made by the patient while the MRI scan is underway. The Gibbs ringing artifact is caused by the truncation of data that requires a higher sampling frequency to observe. Research has shown that the effect of the Gibbs ringing artifact can be reduced in post-processed utilizing deep learning-based networks (Chen). Currently, a smoothing filter is applied to MR images to mitigate the effects of the Gibbs artifact, but the technique fails to reproduce the features lost from the artifact (Muckley).

Image enhancement is just one way that artificial intelligence is revolutionizing the medical space. Another significant utilization of AI is the employment of neural network models to recognize patterns amongst visual-based medical data. Studies have shown the convolutional

neural networks (CNNs) have shown notable promise in their ability to process this data and detect subtle patterns indicative of diseases (Pinto). These models tend to have better diagnostic results on diseases whose symptoms typically manifest in a uniform manner. Given that multiple sclerosis (MS) tends to present somewhat uniquely for every patient, larger databases are required to train models that have the capability to detect MS in its multitude of presentations. Another complication that MS poses is that the treatment plan is often burdensome on the patient. The subsequent effect is that the cost of false positives associated with MS is higher than other diseases. Since the probability of detection ( $P_D$ ) and the probability of false alarms ( $P_{FA}$ ) have a direct relationship, lowering  $P_D$  causes  $P_{FA}$  to increase. Despite this, progress has been made in leveraging AI to help diagnose MS. Researchers work around MS's numerous presentations by creating multiple databases that separate the data collected from common early symptoms such as movement impairment, cognitive decline etc. (Cartwright).

The high misdiagnosis rates of Multiple Sclerosis, coupled with the complexities involved in accurately diagnosing the disease, highlight the need for new solutions to support healthcare professionals. This project aims to investigate the ongoing challenges and limitations of current diagnostic methods for MS, including the exploration of the latest research on potential biomarkers. Additionally, it seeks to combine the capabilities of artificial intelligence in detecting subtle patterns in medical imaging and data, which can assist clinicians in reducing misdiagnosis rates. Exploring AI's potential to enhance diagnostic accuracy and offer effective and personalized treatment plans for patients would no doubt be helpful in advancing the integration of new technologies into clinical practice.

### **User Stories: Medical Professional's Perspective**

As a neurologist, Dr. Emily had a patient named Sarah, a 32-year-old woman who had been experiencing numbness, fatigue, and trouble with coordination. Sarah had seen various specialists but had yet to receive a definitive diagnosis. Given the complexity of Sarah's symptoms, Dr. Emily decided to use an artificial intelligence (AI) diagnostic tool to assist her in identifying the underlying cause. The tool indicated that MS was a likely diagnosis, and after further review from Dr. Emily and her colleagues, they agreed with this diagnosis.

Thanks to the AI tool, Dr. Emily was able to provide a faster, more accurate diagnosis, allowing Sarah to begin treatment sooner and improve her quality of life. The AI enhanced Dr. Emily's ability to detect the nuances in Sarah's case, offering insights that would have been challenging to identify through traditional methods alone.

**User Story: Patient's Perspective**

As a patient exhibiting unexplained neurological symptoms, I want to have the most accurate diagnosis possible because this is a life and death situation for me. I want my healthcare provider to have the most up-to-date technology to aid in the diagnosis process. I want the technology to be reliable and have the ability to see what the naked human eye may not be able to. I would like to have my diagnosis in a timely manner and have my provider present it to me with confidence prior to starting me on a treatment plan. I want to walk out of my doctor's office feeling confident that every possible step was taken in ensuring the accuracy of accepting or rejecting Multiple Sclerosis as the reason behind my symptoms.

For our product, the minimum viable product should be a model that can assist healthcare professionals in identifying patterns in MRI images that indicate Multiple Sclerosis, the AI tool would also provide clinicians with decision support to ensure they can confidently differentiate MS from other conditions with similar symptoms thereby reducing high misdiagnosis rate.

## Work Cited

Chen, Zhaolin et al. "Deep Learning for Image Enhancement and Correction in Magnetic Resonance Imaging-State-of-the-Art and Challenges." *Journal of digital imaging* vol. 36,1 (2023): 204-230. doi:10.1007/s10278-022-00721-9

National Institute of Health. "Multiple Sclerosis." *National Institute of Neurological Disorders and Stroke*, 2024, [www.ninds.nih.gov/health-information/disorders/multiple-sclerosis#:~:text=Multiple%20sclerosis%20\(MS\)%20is%20a%20chronic%20neurological%20disorder.%20It%20is#:~:text=Multiple%20sclerosis%20\(MS\)%20is%20a%20chronic%20neurological%20disorder.%20It%20is](http://www.ninds.nih.gov/health-information/disorders/multiple-sclerosis#:~:text=Multiple%20sclerosis%20(MS)%20is%20a%20chronic%20neurological%20disorder.%20It%20is#:~:text=Multiple%20sclerosis%20(MS)%20is%20a%20chronic%20neurological%20disorder.%20It%20is). Accessed 30 Sept. 2024.

Muckley, Matthew J et al. "Training a neural network for Gibbs and noise removal in diffusion MRI." *Magnetic resonance in medicine* vol. 85,1 (2021): 413-428. doi:10.1002/mrm.28395

Pinto-Coelho, Luís. "How Artificial Intelligence Is Shaping Medical Imaging Technology: A Survey of Innovations and Applications." *Bioengineering (Basel, Switzerland)* vol. 10,12 1435. 18 Dec. 2023, doi:10.3390/bioengineering10121435

Cartwright, Jacob et al. "Innovations in Multiple Sclerosis Care: The Impact of Artificial Intelligence via Machine Learning on Clinical Research and Decision-Making." *International journal of MS care* vol. 25,5 (2023): 233-241. doi:10.7224/1537-2073.2022-076

Solomon, Michael D., et al. "Misdiagnosis of Multiple Sclerosis: Impact of the 2017 McDonald Criteria on Clinical Practice." National Library of Medicine, PubMed Central (PMC), 1 Jan. 2019, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6336166/>.

Siva, Aksel. "Common Clinical and Imaging Conditions Misdiagnosed as Multiple Sclerosis: A Current Approach to the Differential Diagnosis of Multiple Sclerosis." *Neurologic Clinics*, vol. 36, no. 1, Feb. 2018, <https://www.sciencedirect.com/science/article/abs/pii/S0733861917301019>.

Hemond, Cheryl, and David S. Reich. "New Imaging Markers in Multiple Sclerosis and Related Disorders: Smoldering Inflammation and the Central Vein Sign." *Neuroimaging Clinics*, vol. 34, no. 3, 2024, [https://www.neuroimaging.theclinics.com/article/S1052-5149\(24\)00024-8/fulltext](https://www.neuroimaging.theclinics.com/article/S1052-5149(24)00024-8/fulltext).