

Lecture 2: Lebesgue integration

Tianpei Xie

Jul. 22nd., 2015

Contents

1	From Riemann integration to Lebesgue integration	2
1.1	Riemann Integration	2
1.2	Lebesgue Integration of Unsigned Simple Functions	4
1.3	Unsigned Lebesgue Measurable Function	8
1.4	Unsigned Lebesgue Integrals	12
1.5	Absolute Integrable	17
2	Development of Lebesgue Integration	20
3	Examples	21
3.1	Unsigned Measureable Functions	21

1 From Riemann integration to Lebesgue integration

1.1 Riemann Integration

- **Definition** Let $[a, b]$ be an interval of positive length, and $f : [a, b] \rightarrow \mathbb{R}$ be a function. A **tagged partition** $P \equiv ((x_0, x_1, \dots, x_n); (x'_1, \dots, x'_n))$ of $[a, b]$ is a finite sequence of real numbers $a = x_0 < x_1 < \dots < x_n = b$, together with additional numbers $x_{i-1} \leq x'_i \leq x_i$ for each $1 \leq i \leq n$. We abbreviate $(x_i - x_{i-1})$ as Δx_i . The quantity $\Delta(P) \equiv \sup_{1 \leq i \leq n} \Delta x_i$ will be called the **norm** of the tagged partition. The **Riemann sum** $\mathcal{R}(f; P)$ of f with respect to the tagged partition P is denoted as

$$\mathcal{R}(f; P) \equiv \sum_{i=1}^n f(x'_i) \Delta x_i$$

Then f is said to be **Riemann integrable** if there exist a number $\int_a^b f(x)dx$ such as for any $\epsilon > 0$, there exists $\delta > 0$, for all P such that $\Delta P < \delta$,

$$\left| \mathcal{R}(f; P) - \int_a^b f(x)dx \right| \leq \epsilon$$

holds.

- **Remark** Note that for f **piecewise constant** in a partition of intervals $[a, b] = \bigcup_k^n I_k$, where $f(x) = c_k$ if $x \in I_k$ interval,

$$\int_a^b f(x)dx = \sum_{k=1}^n c_k |I_k|.$$

- **Definition** Given any tagged partition $P = ((x_0, x_1, \dots, x_n); (x'_1, \dots, x'_n))$, we can define $g_i \equiv \min_{x \in [x_{i-1}, x_i]} f(x) \leq f(x'_i) \leq \max_{x \in [x_{i-1}, x_i]} f(x) \equiv h_i$, for all $1 \leq i \leq n$. Define the piecewise constant function g, h on interval $\bigcup_i [x_{i-1}, x_i]$ so that $g(x) = g_i$ if $x \in [x_{i-1}, x_i]$, and $h(x) = h_i$ if $x \in [x_{i-1}, x_i]$. Clearly, $g \leq f \leq h$ on $[0, 1]$. Denote the **upper and lower Darboux sums**

$$U(f, P) \equiv \sum_{i=1}^n h_i \Delta x_i$$

$$L(f, P) \equiv \sum_{i=1}^n g_i \Delta x_i$$

- **Proposition 1.1** We have the following properties

1. $L(f, P) \leq \mathcal{R}(f; P) \leq U(f, P)$ for all P .
2. $\overline{\int_a^b} f(x)dx = \inf_P U(f, P)$ and $\underline{\int_a^b} f(x)dx = \sup_P L(f, P)$;
3. For the **refinement of partition** $P^* \supseteq P$,

$$L(f, P^*) \geq L(f, P)$$

$$U(f, P^*) \leq U(f, P)$$

• **Definition** (*Upper and lower integral*).

Let $[a, b]$ be an interval, and $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. The **lower Riemann integral** $\int_a^b f(x)dx$ of f on $[a, b]$ is defined as

$$\int_a^b f(x)dx \equiv \sup_{\substack{g \leq f, \\ g \text{ piecewise constant}}} \int_a^b g(x)dx = \sup_P L(f, P),$$

where g ranges over all piecewise constant functions that are pointwise bounded above by f .

Similarly, define the **upper Riemann integral** $\int_a^b f(x)dx$ of f on $[a, b]$ as

$$\int_a^b f(x)dx \equiv \inf_{\substack{h \geq f, \\ h \text{ piecewise constant}}} \int_a^b h(x)dx = \inf_P U(f, P),$$

where h ranges over all piecewise constant functions that are pointwise bounded below by f .

If $\int_a^b f(x)dx = \int_a^b f(x)dx$, we say that f is **Riemann integrable**, and refer to this quantity as *the Darboux integral* of f on $[a, b]$.

• **Theorem 1.2** For any $\epsilon > 0$, there exists a partition P such that $|U(f, P) - L(f, P)| \leq \epsilon$, **if and only if** f is Riemann integrable (Darboux integrable).

• **Exercise 1.3** Show that for bounded function $f : [a, b] \rightarrow \mathbb{R}$ on $[a, b]$, f is Riemann integrable if and only if f is Darboux integrable.

Proof: Given any tagged partition $P = ((x_0, x_1, \dots, x_n); (x'_1, \dots, x'_n))$, we can define $g_i \equiv \min_{x \in [x_{i-1}, x_i]} f(x) \leq f(x'_i) \leq \max_{x \in [x_{i-1}, x_i]} f(x) \equiv h_i$, for all $1 \leq i \leq n$. Define the piecewise constant function g, h on interval $\bigcup_i [x_{i-1}, x_i]$ so that $g(x) = g_i$ if $x \in [x_{i-1}, x_i]$, and $h(x) = h_i$ if $x \in [x_{i-1}, x_i]$. Clearly, $g \leq f \leq h$ on $[0, 1]$. Denote

$$U(f, P) \equiv \sum_{i=1}^n h_i \Delta x_i$$

$$L(f, P) \equiv \sum_{i=1}^n g_i \Delta x_i$$

Thus $L(f, P) \leq \mathcal{R}(f; P) \leq U(f, P)$ and also $L(f, P) \leq \int_a^b f(x)dx$ and $U(f, P) \geq \int_a^b f(x)dx$ for any P .

Finally, for the refinement of partition $P_1 \cup P_2$, $L(f, P_1 \cup P_2) \geq L(f, P_i)$, $U(f, P_1 \cup P_2) \leq U(f, P_i)$.

– \Rightarrow Since f is Riemann integrable, for any $\epsilon > 0$, there must exist tagged partitions P_1, P_2 such that

$$\mathcal{R}(f; P_1) \leq U(f, P_1) \leq \int_a^b f(x)dx + \epsilon/2$$

$$\mathcal{R}(f; P_2) \geq L(f, P_2) \geq \int_a^b f(x)dx - \epsilon/2$$

Furthermore,

$$\begin{aligned} U(f, P_1 \cup P_2) &\leq U(f, P_1) \leq \int_a^b f(x)dx + \epsilon/2 \\ L(f, P_1 \cup P_2) &\geq L(f, P_2) \geq \int_a^b f(x)dx - \epsilon/2 \end{aligned}$$

Clearly, we have

$$\begin{aligned} \left| \overline{\int_a^b f(x)dx} - \underline{\int_a^b f(x)dx} \right| &\leq |U(f, P_1 \cup P_2) - L(f, P_1 \cup P_2)| \\ &\leq \left| U(f, P_1 \cup P_2) - \int_a^b f(x)dx \right| + \left| \int_a^b f(x)dx - L(f, P_1 \cup P_2) \right| \leq \epsilon. \end{aligned}$$

– \Leftarrow See that for any partition P , $\underline{\int_a^b f(x)dx} \leq \mathcal{R}(f; P) \leq \overline{\int_a^b f(x)dx}$. Since $\underline{\int_a^b f(x)dx} - \overline{\int_a^b f(x)dx} = \int_a^b f(x)dx$, therefore, for any $\epsilon > 0$, for all P ,

$$\begin{aligned} 0 &\leq \mathcal{R}(f; P) - \int_a^b f(x)dx \leq \overline{\int_a^b f(x)dx} - \underline{\int_a^b f(x)dx} \\ \Rightarrow \left| \mathcal{R}(f; P) - \int_a^b f(x)dx \right| &\leq \epsilon, \end{aligned}$$

which completes our proof. \blacksquare

1.2 Lebesgue Integration of Unsigned Simple Functions

- **Definition** A ***complex-valued simple function*** $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is *piecewise constant* in **a finite collection of Lebesgue measurable sets** E_1, \dots, E_n such that

$$f(x) = \sum_{k=1}^n c_k \mathbb{1}_{\{x \in E_k\}},$$

where $c_k \in \mathbb{C}$ are complex numbers. If $c_k \in [0, \infty)$, the piecewise constant function $f : \mathbb{R}^d \rightarrow [0, \infty)$ as above is called an ***unsigned simple function***.

- **Remark** For the space $\text{Simp}(\mathbb{R}^d)$
 1. The space $\text{Simp}(\mathbb{R}^d)$ of complex valued simple functions forms **a complex vector space**; also,
 2. $\text{Simp}(\mathbb{R}^d)$ is also **closed** under *pointwise product* $f, g \mapsto fg$ and *complex conjugation* $f \mapsto \bar{f}$. In short, $\text{Simp}(\mathbb{R}^d)$ is a **commutative -algebra**.
- **Remark** Note for $E_k, 1 \leq k \leq n$ disjoint, we can further partition them into disjoint sets so that f is constant on each disjoint subset.

- **Definition** (*integral of unsigned simple function*)

If $f = \sum_{k=1}^n c_k \mathbb{1}_{\{x \in E_k\}}$ is an unsigned simple function, then its *integral* is defined as

$$\text{simp} \int_{\mathbb{R}^d} f(x) dx \equiv \sum_{k=1}^n c_k m(E_k),$$

where $\int_{\mathbb{R}^d} \mathbb{1}_{\{x \in E_k\}} dx = m(E_k)$.

- **Lemma 1.4** (*Well-definedness of simple integral*). [Tao, 2011]

Let $k, k' \geq 0$ be natural numbers, $c_1, \dots, c_k, c'_1, \dots, c'_{k'} \in [0, +\infty]$, and let $E_1, \dots, E_k, E'_1, \dots, E'_{k'} \subseteq \mathbb{R}^d$ be Lebesgue measurable sets such that the identity

$$c_1 \mathbb{1}_{E_1} + \dots + c_k \mathbb{1}_{E_k} = c'_1 \mathbb{1}_{E'_1} + \dots + c'_{k'} \mathbb{1}_{E'_{k'}}$$

holds identically on \mathbb{R}^d . Then one has

$$c_1 \mu(E_1) + \dots + c_k \mu(E_k) = c'_1 \mu(E'_1) + \dots + c'_{k'} \mu(E'_{k'})$$

- **Remark** The simple integral could also be defined on finitely additive measure spaces, rather than countably additive ones, and all the above properties would still apply. However, on a finitely additive measure space one would have difficulty extending the integral beyond simple functions.

- **Definition** (*Almost everywhere and support*).

A property $P(x)$ of a point $x \in \mathbb{R}^d$ is said to hold *(Lebesgue) almost everywhere* in \mathbb{R}^d , or for *(Lebesgue) almost every point* $x \in \mathbb{R}^d$, if the set of $x \in \mathbb{R}^d$ for which $P(x)$ *fails* has *Lebesgue measure zero* (i.e. P is true *outside* of a *null set*). We usually omit the prefix Lebesgue, and often abbreviate “*almost everywhere*” or “*almost every*” as *a.e.*

The *support* of a function $f : \mathbb{R}^d \rightarrow \mathbb{C}$ or $f : \mathbb{R}^d \rightarrow [0, \infty]$ is defined to be the set $\{x \in \mathbb{R}^d : f(x) \neq 0\}$ where f is *non-zero*.

- **Remark** The followings are helpful to understand *the almost everywhere* concept:

1. Two functions $f, g : \mathbb{R}^d \rightarrow Z$ into an arbitrary range Z are said to *agree almost everywhere* if one has $f(x) = g(x)$ for almost every $x \in \mathbb{R}^d$.
2. If $P_1(x), P_2(x), \dots$ are an *at most countable* family of *properties*, each of which individually holds for *almost every* x , then they will *simultaneously* be true for *almost every* x , because *the countable union of null sets is still a null set*.
3. if $P(x)$ holds for *almost every* x , and $P(x)$ implies $Q(x)$, then $Q(x)$ holds for *almost every* x .

Because of these properties, one can (as a rule of thumb) treat the *almost universal quantifier* “for almost every” as *if* it was the truly *universal quantifier* “for every”, as long as one is only concatenating *at most countably many* properties together, and as long as one *never specialises* the free variable x to a *null set*.

- **Proposition 1.5** (*Basic properties of the simple unsigned integral*).

Let $f, g : \mathbb{R}^d \rightarrow [0, +\infty]$ be simple unsigned functions.

1. (**Unsigned linearity**) We have

$$\text{simp} \int_{\mathbb{R}^d} (f(x) + g(x)) dx = \text{simp} \int_{\mathbb{R}^d} f(x) dx + \text{simp} \int_{\mathbb{R}^d} g(x) dx$$

and

$$\text{simp} \int_{\mathbb{R}^d} c f(x) dx = c \text{simp} \int_{\mathbb{R}^d} f(x) dx$$

for all $c \in [0, +\infty]$.

2. (**Finiteness**) We have $\text{simp} \int_{\mathbb{R}^d} f(x) dx < \infty$ **if and only if** f is **finite almost everywhere**, and its **support** has **finite measure**.

3. (**Vanishing**) We have $\text{simp} \int_{\mathbb{R}^d} f(x) dx = 0$ **if and only if** f is **zero almost everywhere**.

4. (**Equivalence**) If f and g **agree almost everywhere**, then

$$\text{simp} \int_{\mathbb{R}^d} f(x) dx = \text{simp} \int_{\mathbb{R}^d} g(x) dx$$

5. (**Monotonicity**) If $f(x) \leq g(x)$ for **almost every** $x \in \mathbb{R}^d$, then

$$\text{simp} \int_{\mathbb{R}^d} f(x) dx \leq \text{simp} \int_{\mathbb{R}^d} g(x) dx$$

6. (**Compatibility with Lebesgue measure**) For any Lebesgue measurable E , one has

$$\text{simp} \int_{\mathbb{R}^d} \mathbb{1}_E(x) dx = m(E).$$

Furthermore, the simple unsigned integral $f \mapsto \text{simp} \int_{\mathbb{R}^d} f(x) dx$ is the **only map** from the space $\text{Simp}_+(\mathbb{R}^d)$ of unsigned simple functions to $[0, +\infty]$ that **obeys all of the above properties**.

Proof: – Proof of 3. *Vanishing*

$$\text{simp} \int_{\mathbb{R}^d} f(x) dx = \sum_{i=1}^k c_i m(E_i) = 0$$

for a finite collection of Lebesgue measurable set E_1, \dots, E_k .

Since Lebesgue measure is nonnegative, and $c_i \geq 0$ for all $1 \leq i \leq k$, the above holds if and only if $c_i m(E_i) = 0$ for $1 \leq i \leq k$. Then except for the null set that $m(E_i) = 0$, the coefficient $c_i = 0$ for all i . It means that $f = 0$, *a.e.*.

– Proof of 4. *Equivalence* Let $f(x) = \sum_{i=1}^k c_i \mathbb{1}_{\{x \in E_i\}}$ and $g(x) = \sum_{j=1}^m d_j \mathbb{1}_{\{x \in F_j\}}$.

Without loss of generality, assume $E_i \neq \emptyset$ for all i and $F_j \neq \emptyset$.

$$\begin{aligned} & \text{simp} \int_{\mathbb{R}^d} f(x) dx - \text{simp} \int_{\mathbb{R}^d} g(x) dx \\ &= \text{simp} \int_{\mathbb{R}^d} (f(x) - g(x)) dx \\ &= \sum_{i=1}^k \sum_{j=1}^m (c_i - d_j) m(E_i \cap F_j) + \sum_{i=1}^k c_i m \left(E_i / \bigcup_{j=1}^m F_j \right) - \sum_{j=1}^m d_j m \left(F_j / \bigcup_{i=1}^k E_i \right) \end{aligned}$$

If $f = g$, *a.e.*, or $f - g = 0$, *a.e.*. In other word, for $E_i \cap F_j \neq \emptyset$, either $m(E_i \cap F_j) = 0$ or $c_i = d_j$. For $m(E_i / \bigcup_{j=1}^m F_j) \neq 0$, $c_i = 0$ and $m(F_j / \bigcup_{i=1}^k E_i) \neq 0$, $d_j = 0$. In indicates that the above integral is zero. ■

• **Definition (*Absolutely convergent simple integral*).**

A complex-valued simple function $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is said to be **absolutely integrable** of $\text{simp} \int_{\mathbb{R}^d} |f(x)| dx < \infty$. If f is absolutely integrable, the **integral** $\text{simp} \int_{\mathbb{R}^d} f(x) dx$ is defined for **real** signed f by the formula

$$\text{simp} \int_{\mathbb{R}^d} f(x) dx = \text{simp} \int_{\mathbb{R}^d} f_+(x) dx - \text{simp} \int_{\mathbb{R}^d} f_-(x) dx$$

where $f_+(x) = \max\{f(x), 0\}$ and $f_-(x) = \max\{-f(x), 0\}$. (note that these are unsigned simple functions that are pointwise dominated by $|f|$ and thus have finite integral), and for complex-valued f by the formula

$$\text{simp} \int_{\mathbb{R}^d} f(x) dx = \text{simp} \int_{\mathbb{R}^d} \Re f(x) dx + j \text{simp} \int_{\mathbb{R}^d} \Im f(x) dx$$

• **Proposition 1.6 (*Basic properties of the complex-valued simple integral*).**

Let $f, g : \mathbb{R}^d \rightarrow \mathbb{C}$ be absolutely integrable simple functions.

1. (***-linearity**) We have

$$\text{simp} \int_{\mathbb{R}^d} (f(x) + g(x)) dx = \text{simp} \int_{\mathbb{R}^d} f(x) dx + \text{simp} \int_{\mathbb{R}^d} g(x) dx$$

and

$$\text{simp} \int_{\mathbb{R}^d} cf(x) dx = c \text{simp} \int_{\mathbb{R}^d} f(x) dx$$

for all $c \in \mathbb{C}$. Also we have

$$\text{simp} \int_{\mathbb{R}^d} \overline{f(x)} dx = \overline{\text{simp} \int_{\mathbb{R}^d} f(x) dx}$$

2. (**Equivalence**) If f and g agree almost everywhere, then

$$\text{simp} \int_{\mathbb{R}^d} f(x) dx = \text{simp} \int_{\mathbb{R}^d} g(x) dx$$

3. (*Compatibility with Lebesgue measure*) For any Lebesgue measurable E , one has

$$\text{simp} \int_{\mathbb{R}^d} \mathbb{1}_E(x) dx = m(E).$$

- **Remark** (Simple) functions that agree almost everywhere, have the same integral. We can view this as an assertion that **integration** is a **noise-tolerant operation**: one can have “noise” or “errors” in a function $f(x)$ on a **null set**, and this will not affect the final value of the integral. One can even integrate functions f that are merely *defined almost everywhere* on \mathbb{R}^d , *except for a null set*. And *the extension on null set is arbitrary*. For example $\sin(x)/x$ is not defined everywhere but almost everywhere in \mathbb{R} .

In **functional analysis**, it is convenient to abstract the notion of an almost everywhere defined function somewhat, by replacing any such function f with **the equivalence class of almost everywhere defined functions that are equal to f almost everywhere**. Such classes are then no longer functions in the standard set-theoretic sense (called the *distribution*). [Tao, 2011]

- **Remark** The “**Lebesgue philosophy**” that one is **willing to lose control** on **sets of measure zero** is a perspective that distinguishes *Lebesgue-type analysis* from other types of analysis.

1.3 Unsigned Lebesgue Measurable Function

- **Remark** Much as the piecewise constant integral can be completed to the Riemann integral, the unsigned simple integral can be completed to the unsigned Lebesgue integral, by extending the class of *unsigned simple functions* to the larger class of *unsigned Lebesgue measurable functions*.

- **Definition** (*Unsigned measurable function*).

An unsigned function $f : \mathbb{R}^d \rightarrow [0, +\infty]$ is **unsigned Lebesgue measurable**, or **measurable** for short, if it is the *pointwise limit of unsigned simple functions*, i.e. if there exists a sequence $f_1, f_2, f_3, \dots : \mathbb{R}^d \rightarrow [0, +\infty]$ of unsigned simple functions such that $f_n(x) \rightarrow f(x)$ for every $x \in \mathbb{R}^d$.

- The followings are equivalent definitions for *the unsigned measurable function*:

Lemma 1.7 (*Equivalent notions of measurability*) [Tao, 2011]

Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be an unsigned function. Then the following are equivalent:

1. f is **unsigned Lebesgue measurable**.
2. f is the **pointwise limit of unsigned simple functions** f_n (thus the limit $\lim_{n \rightarrow \infty} f_n(x)$ exists and is equal to $f(x)$ for all $x \in \mathbb{R}^d$).
3. f is the **pointwise almost everywhere limit of unsigned simple functions** f_n (thus the limit $\lim_{n \rightarrow \infty} f_n(x)$ exists and is equal to $f(x)$ for almost every $x \in \mathbb{R}^d$).
4. f is the **supremum** $f(x) = \sup_n f_n(x)$ of an **increasing** sequence $0 \leq f_1 \leq f_2 \leq \dots$ of **unsigned simple functions** f_n , each of which are **bounded** with **finite** measure support.

5. For every $\lambda \in [0, +\infty]$, the set $\{\mathbf{x} \in \mathbb{R}^d : f(\mathbf{x}) > \lambda\}$ is **Lebesgue measurable**.
6. For every $\lambda \in [0, +\infty]$, the set $\{\mathbf{x} \in \mathbb{R}^d : f(\mathbf{x}) \geq \lambda\}$ is **Lebesgue measurable**.
7. For every $\lambda \in [0, +\infty]$, the set $\{\mathbf{x} \in \mathbb{R}^d : f(\mathbf{x}) < \lambda\}$ is **Lebesgue measurable**.
8. For every $\lambda \in [0, +\infty]$, the set $\{\mathbf{x} \in \mathbb{R}^d : f(\mathbf{x}) \leq \lambda\}$ is **Lebesgue measurable**.
9. For every interval $I \subset [0, +\infty)$, the set $f^{-1}(I) \equiv \{\mathbf{x} \in \mathbb{R}^d : f(\mathbf{x}) \in I\}$ is **Lebesgue measurable**.
10. For every (relatively) **open** set $U \subset [0, +\infty)$, the set $f^{-1}(U) \equiv \{\mathbf{x} \in \mathbb{R}^d : f(\mathbf{x}) \in U\}$ is **Lebesgue measurable**.
11. For every (relatively) **closed** set $K \subset [0, +\infty)$, the set $f^{-1}(K) \equiv \{\mathbf{x} \in \mathbb{R}^d : f(\mathbf{x}) \in K\}$ is **Lebesgue measurable**.

Proof: 1. 1) to 2) is the definition. 3) is a specification of 2).

2. 3) \Rightarrow 4) Let a sequence of unsigned simple function $f_1, \dots, f_n \rightarrow f$ for almost every $\mathbf{x} \in \mathbb{R}^d$, then there exist a monotone subsequence $0 \leq f_{i_1} \leq f_{i_2} \leq \dots$, that is convergent to $f = \sup_n f_{i_n}$ for almost every $\mathbf{x} \in \mathbb{R}^d$. Then we can find a monotone subsequence $0 \leq f_{i_1} \mathbf{1}_{\{\|\mathbf{x}\| < 1\}} \leq f_{i_2} \mathbf{1}_{\{\|\mathbf{x}\| < 2\}} \leq \dots$, in which $f_{i_n}(\mathbf{x}) \mathbf{1}_{\{\|\mathbf{x}\| < n\}}$ is bounded has finite measure support. This is because f_{i_n} only takes finite values and its support is bounded by $m(\{\mathbf{x} | \|\mathbf{x}\| < n\}) < \infty$. To show that $f_{i_n}(\mathbf{x}) \mathbf{1}_{\{\|\mathbf{x}\| < n\}} \rightarrow f$ for almost every $\mathbf{x} \in \mathbb{R}^d$, we see that $f_{i_n}(\mathbf{x}) \mathbf{1}_{\{\|\mathbf{x}\| < n\}} \leq f_{i_n}$, and $f_{i_n} \rightarrow f, a.e. \mathbf{x} \in \mathbb{R}^d$, so by dominated convergence, it is true.
3. 4) \Rightarrow 5) Consider a sequence of monotone increasing unsigned simple functions $f_n \leq f_{n+1}, n \geq 1$, which is bounded with finite measure support. Let $f = \sup_n f_n$. Note that for any $\lambda \in [0, \infty]$, $(\lambda, \infty]$ is Lebesgue measurable and $f_n^{-1}((\lambda, \infty]) = \bigcup_{k=1}^s E_{n,k}$ for $m(E_{n,k}) < \infty$ (it could be $E_{n,k} = \emptyset$ for some k) for all $n \geq 1$. Since $\{f_n\}$ is monotone, $f_n^{-1}((\lambda, \infty]) \subseteq f_{n+1}^{-1}((\lambda, \infty]), n \geq 1$. Hence $f_n^{-1}((\lambda, \infty])$ are measurable, since E_k are all measurable. Therefore, we have a monotone nondecreasing sequence of nested sets, and $f^{-1}((\lambda, \infty]) = \bigcup_{n=1}^{\infty} f_n^{-1}((\lambda, \infty])$ so that $f_n^{-1}((\lambda, \infty]) \uparrow f^{-1}((\lambda, \infty])$. Therefore, $f^{-1}((\lambda, \infty])$ is Lebesgue measurable, since each $f_n^{-1}((\lambda, \infty])$ is Lebesgue measurable.
4. 5) \Rightarrow 6) See that $f^{-1}([\lambda, \infty]) = f^{-1}(\bigcap_{m=1}^{\infty} (\lambda - \frac{1}{2^m}, \infty]) = \bigcap_{m=1}^{\infty} f^{-1}((\lambda - \frac{1}{2^m}, \infty])$, which is measurable.
5. 6) \Leftrightarrow 7) The pre-image is closed under complement operation. Similarly, 7) \Rightarrow 8) follow the same argument as 5) \Leftrightarrow 6), with reversed direction.
6. 9) Any interval I is just a countable union of finite of intersection of the interval above. And for 10), any open set in \mathbb{R} is a countable union of intervals. Conversely, an open interval naturally works, while others are countable intersection of open intervals. 11) is true if and only if 10) is true, since 11) is the complement of 10).
7. 11) \Rightarrow 2) For $f \geq 0$, we should construct the simple function f_n by subdividing the values of f that fall in $[0, k]$ by partitioning $[0, k]$ into subintervals $[(j-1)2^{-k}, j2^{-k}], j = 1, \dots, k2^k$. By assumption, $E_{j,k} \equiv f^{-1}([(j-1)2^{-k}, j2^{-k}])$ are all measurable for $j = 1, \dots, k2^k$. Finally, the complement $\mathbb{R}/f^{-1}([0, k])$ is measurable.

Let

$$f_k(\mathbf{x}) = \begin{cases} \frac{j-1}{2^k} & \mathbf{x} \in E_{j,k}, j = 1, \dots, k2^k \\ k & \mathbf{x} \in \mathbb{R}^d / f^{-1}([0, k]) \end{cases}$$

Each f_k is a simple function defined everywhere in the domain of f . Clearly, $f_k \leq f_{k+1}$ since in passing from f_k to f_{k+1} , each subinterval $[(j-1)2^{-k}, j2^{-k}]$ is divided in half. Then we have constructed a simple function as

$$f_k(\mathbf{x}) = \sum_{j=1}^{k2^k} \frac{j-1}{2^k} \mathbb{1}_{\{E_{j,k}\}} + k \mathbb{1}_{\{\mathbb{R}^d / f^{-1}([0, k])\}}$$

And $f_k \uparrow f$ for every $\mathbf{x} \in \mathbb{R}^d$. Note that this example could be used to prove 3). It then completes our proof. ■.

- With these equivalent formulations, we can now generate plenty of measurable functions:
- **Exercise 1.8**
 1. Show that every **continuous** function $f : \mathbb{R}^d \rightarrow [0, \infty]$ is **measurable**;
 2. Show that every **unsigned simple function** is **measurable**
 3. Show that the **supremum, infimum, limit superior, or limit inferior** of unsigned measurable functions is unsigned measurable.
 4. Show that an **unsigned function** that is **equal almost everywhere** to an unsigned measurable function, is itself measurable.
 5. Show that if a sequence f_n of unsigned measurable functions **converges** pointwise almost everywhere to an **unsigned limit** f , then f is also measurable;
 6. If $f : \mathbb{R}^d \rightarrow [0, \infty]$ is **measurable** and $\phi : [0, +\infty] \rightarrow [0, +\infty]$ is **continuous**, show that $\phi \circ f : \mathbb{R}^d \rightarrow [0, +\infty]$ is measurable.
 7. If f, g are unsigned measurable functions, show that $f + g$ and fg are measurable.

Proof: 1. Note that for any $\lambda \in [0, \infty]$, $f^{-1}((\lambda, \infty])$ is open, thus Lebesgue measurable. So f is unsigned measurable.

2. Just by definition.

3. For any $\lambda \in [0, \infty]$, $(\inf_n f_n)^{-1}((\lambda, \infty]) = \bigcap_n f_n^{-1}((\lambda, \infty])$ so measurable. Similarly, $(\sup_n f_n)^{-1}([0, \lambda)) = \bigcap_n f_n^{-1}([0, \lambda))$ measurable. Also

$$\begin{aligned} \left(\liminf_{n \rightarrow \infty} f_n \right)^{-1}((\lambda, \infty]) &= \left(\sup_{k \geq 1} \inf_{n \geq k} f_n \right)^{-1}((\lambda, \infty]) \\ &= \bigcup_m \bigcup_{k \geq 1} \bigcap_{n \geq k} f_n^{-1}((\lambda + \frac{1}{m}, \infty]) \end{aligned}$$

is measurable and

$$\begin{aligned} \left(\limsup_{n \rightarrow \infty} f_n \right)^{-1}([0, \lambda)) &= \left(\inf_{k \geq 1} \sup_{n \geq k} f_n \right)^{-1}([0, \lambda)) \\ &= \bigcup_m \bigcup_{k \geq 1} \bigcap_{n \geq k} f_n^{-1}([0, \lambda - \frac{1}{m})) \end{aligned}$$

is measurable.

4. Assume that $g = f, a.e.$ where f is unsigned measurable, g is unsigned function. Then $f = \limsup_{n \rightarrow \infty} f_n$ for a sequence of unsigned simple function. For any $\lambda \in [0, \infty]$, $g^{-1}((\lambda, \infty])$ is equal to $f^{-1}((\lambda, \infty])$ outside a null set. Then $g^{-1}((\lambda, \infty])$ is equal to

$$\bigcup_m \bigcap_{k \geq 1} \bigcup_{n \geq k} f_n^{-1}((\lambda + \frac{1}{m}, \infty])$$

outside a null set. As each f_n is an unsigned simple function, $f_n^{-1}((\lambda + \frac{1}{m}, \infty])$ is Lebesgue measurable, so as their countable union and intersection. Modifying a Lebesgue measurable set on a null set produces another Lebesgue measurable set, so $g^{-1}((\lambda, \infty])$ is measurable.

5. Each $f_n = \sup_k f_{n,k}$ for some monotone unsigned simple function $f_{n,k} \leq f_{n,k+1}$. For every convergent sequence of unsigned measurable function f_n , we can find a monotone increasing subsequence that is convergent to f . Therefore there exists a sequence of some monotone unsigned simple function $f_{1,1}, \dots, f_{1,k}, \dots, f_{n,k}, f_{n,k+1}, \dots$ so that $f = \sup_{n \geq 1} \sup_{k \geq n} f_{n,k}$, so f is unsigned measurable.
6. Trivial, since $(\phi \circ f)^{-1}(\lambda, \infty] = f^{-1}(\phi^{-1}((\lambda, \infty]))$, which is measurable. ■

- **Exercise 1.9** Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$. Show that f is a **bounded unsigned measurable function** if and only if f is the **uniform limit** of bounded simple functions.

Proof: \Rightarrow Given $f : \mathbb{R}^d \rightarrow [0, +\infty)$ is bounded unsigned measurable function, suppose $f(\mathbf{x}) \in [0, M]$ for all \mathbf{x} , where $M = \inf_{s \in \mathbb{N}} \{s \geq \sup_{\mathbf{x}} \{f(\mathbf{x})\}\}$. For each k , we subdivided it into $M2^k$ interval $E_{j,k} \equiv [(j-1)2^{-k}, j2^{-k}]$. Then the monotone increasing bounded unsigned simple function f_k is constructed as

$$f_k(\mathbf{x}) = \sum_{j=1}^{M2^k} \frac{j-1}{2^k} \mathbb{1}_{\{\mathbf{x} \in E_{j,k}\}}.$$

Clearly $f = \sup_k f_k$ for any \mathbf{x} and $f_k \leq f_{k+1}$.

Then for any $\epsilon > 0$, there exists k , for all $n \geq k$

$$\sup_{\mathbf{x}} |f(\mathbf{x}) - f_n(\mathbf{x})| \leq \frac{1}{2^n} < \epsilon.$$

The sequence $\{f_n\}$ converges to f uniformly.

\Leftarrow Given that the unsigned function $f = \lim_{n \rightarrow \infty} f_n$ is the uniform limit of bounded simple functions $\{f_n\}$, we see that there exist $k \geq 1$, such that for all $n \geq k$, $f_n(\mathbf{x}) \geq 0$ for all \mathbf{x} . Then we can find $(i_k, i_{k+1}, \dots) \subset (k, k+1, \dots)$ such that $f_{i_k}, f_{i_{k+1}}, \dots$, is a monotone increasing sequence of bounded *unsigned* simple function so that $f = \sup_k f_{i_k}$. Therefore f is unsigned measurable and f is bounded since f_n is bounded for all n . ■

- **Exercise 1.10** Show that an **unsigned function** $f : \mathbb{R}^d \rightarrow [0, +\infty]$ is a **simple function** if and only if it is **measurable** and takes on **at most finitely many values**.
- **Exercise 1.11** Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be an unsigned measurable function. Show that the region $\{(x, t) \in \mathbb{R}^d \times \mathbb{R} : 0 \leq t \leq f(x)\}$ is a **measurable subset** of \mathbb{R}^{d+1} .

- **Definition** (*Complex measurability*).

An almost everywhere defined **complex-valued function** $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is **Lebesgue measurable**, or measurable for short, if it is the pointwise almost everywhere limit of complex-valued simple functions.

1.4 Unsigned Lebesgue Integrals

- **Definition** (*Lower unsigned Lebesgue integral*).

Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be an unsigned function (*not necessarily measurable*). We define the **lower unsigned Lebesgue integral** $\int_{\mathbb{R}^d} f(x) dx$ to be the quantity

$$\int_{\mathbb{R}^d} f(x) dx \equiv \sup_{0 \leq g \leq f, g \text{ simple}} \text{simp} \int_{\mathbb{R}^d} g(x) dx,$$

where g ranges over all unsigned simple functions $g : \mathbb{R}^d \rightarrow [0, +\infty]$ that are pointwise bounded by f .

- **Remark** One can also define the **upper unsigned Lebesgue integral**

$$\overline{\int_{\mathbb{R}^d} f(x) dx} \equiv \inf_{h \geq f, h \text{ simple}} \text{simp} \int_{\mathbb{R}^d} h(x) dx,$$

but we will use this integral much more rarely. Note that both integrals take values in $[0, +\infty]$, and that the upper Lebesgue integral is always at least as large as the lower Lebesgue integral.

- **Remark** (*Compatibility with the simple integral*)

If f is *simple*, then

$$\int_{\mathbb{R}^d} f(x) dx = \overline{\int_{\mathbb{R}^d} f(x) dx} = \text{simp} \int_{\mathbb{R}^d} f(x) dx.$$

- **Proposition 1.12** (*Basic properties of the lower Lebesgue integral*).

Let $f, g : \mathbb{R}^d \rightarrow [0, +\infty]$ be unsigned functions (*not necessarily measurable*). Then the following is true:

1. (**Equivalence**) If f, g agree almost everywhere, then $\int_{\mathbb{R}^d} f(x) dx = \int_{\mathbb{R}^d} g(x) dx$.
2. (**Monotonicity**) If $f \leq g$ almost everywhere, then $\int_{\mathbb{R}^d} f(x) dx \leq \int_{\mathbb{R}^d} g(x) dx$ and $\overline{\int_{\mathbb{R}^d} f(x) dx} \leq \overline{\int_{\mathbb{R}^d} g(x) dx}$.
3. (**Superadditivity**) $\int_{\mathbb{R}^d} (f(x) + g(x)) dx \geq \int_{\mathbb{R}^d} f(x) dx + \int_{\mathbb{R}^d} g(x) dx$.
4. (**Divisibility**) For any measurable set E , one has $\int_{\mathbb{R}^d} f(x) dx = \int_{\mathbb{R}^d} \mathbf{1}_{\{x \in E\}} f(x) dx + \int_{\mathbb{R}^d} \mathbf{1}_{\{x \in \mathbb{R}^d/E\}} f(x) dx$.
5. (**Horizontal truncation**) As $n \rightarrow \infty$, $\int_{\mathbb{R}^d} \min(f(x), n) dx$ converges to $\int_{\mathbb{R}^d} f(x) dx$.
6. (**Vertical truncation**) As $n \rightarrow \infty$, $\int_{\mathbb{R}^d} f(x) \mathbf{1}_{\{|x| \leq n\}} dx$ converges to $\int_{\mathbb{R}^d} f(x) dx$.
Hint: show that $m(E \cap \{x : |x| \leq n\}) \rightarrow m(E)$ for any measurable set E .

7. (**Reflection**) If $f + g$ is a simple function that is bounded with finite measure support (i.e. it is absolutely integrable), then $\text{simp} \int_{\mathbb{R}^d} (f(x) + g(x)) dx = \underline{\int_{\mathbb{R}^d}} f(x) dx + \overline{\int_{\mathbb{R}^d}} f(x) dx$.

8. (**Subadditivity of upper integral**) $\overline{\int_{\mathbb{R}^d}} (f(x) + g(x)) dx \leq \overline{\int_{\mathbb{R}^d}} f(x) dx + \overline{\int_{\mathbb{R}^d}} g(x) dx$.

Proof: 1. By assumption, $f = g$ outside a set of measure zero. Then for simple unsigned function $0 \leq h \leq f$, $h \leq g$ outside a set of measure zero. Then

$$\begin{aligned} \underline{\int_{\mathbb{R}^d}} f(x) dx &= \sup_{0 \leq h \leq f, h \text{ simple}} \text{simp} \int_{\mathbb{R}^d} h(x) dx \\ &= \sup_{0 \leq h \leq g, a.e., h \text{ simple}} \text{simp} \int_{\mathbb{R}^d} h(x) dx \\ &\leq \sup_{0 \leq g' \leq g, g' \text{ simple}} \text{simp} \int_{\mathbb{R}^d} g'(x) dx \\ &= \underline{\int_{\mathbb{R}^d}} g(x) dx. \end{aligned}$$

By symmetry, $\underline{\int_{\mathbb{R}^d}} f(x) dx \geq \underline{\int_{\mathbb{R}^d}} g(x) dx$. So the equality holds.

2. Trivial, since any unsigned simple function $0 \leq h \leq f \Rightarrow 0 \leq h \leq g$, thus the $\sup_{0 \leq h \leq f} I(h) \leq \sup_{0 \leq h \leq g} I(h)$.

3. Note that since $\underline{\int_{\mathbb{R}^d}} g(x) dx = \sup_{0 \leq h_g \leq g, h_g \text{ simple}} \text{simp} \int_{\mathbb{R}^d} h_g(x) dx$, then $\forall \epsilon > 0$, $\exists h_{g,f} : 0 \leq h_{g,f} \leq g$ simple, s.t. $\underline{\int_{\mathbb{R}^d}} g(x) dx - \epsilon \leq \text{simp} \int_{\mathbb{R}^d} h_{g,f}(x) dx$. So

$$\begin{aligned} \underline{\int_{\mathbb{R}^d}} f(x) dx + \underline{\int_{\mathbb{R}^d}} g(x) dx &= \sup_{0 \leq h_f \leq f, h_f \text{ simple}} \text{simp} \int_{\mathbb{R}^d} h_f(x) dx + \underline{\int_{\mathbb{R}^d}} g(x) dx \\ &= \sup_{0 \leq h_f \leq f, h_f \text{ simple}} \text{simp} \int_{\mathbb{R}^d} (h_f(x) + h_{g,f}(x)) dx \\ &\quad + \left(\underline{\int_{\mathbb{R}^d}} g(x) dx - \text{simp} \int_{\mathbb{R}^d} h_{g,f}(x) dx \right) \\ &\leq \sup_{0 \leq h_f \leq f, h_f \text{ simple}} \text{simp} \int_{\mathbb{R}^d} (h_f(x) + h_{g,f}(x)) dx + \epsilon \\ &\leq \sup_{\substack{0 \leq h_{f+g} \leq f+g, \\ h_{f+g} \text{ simple}}} \text{simp} \int_{\mathbb{R}^d} h_{f+g}(x) dx + \epsilon \quad (\text{since } h_f(x) + h_{g,f}(x) \leq f + g) \\ &= \underline{\int_{\mathbb{R}^d}} (f(x) + g(x)) dx + \epsilon, \quad \forall \epsilon > 0. \end{aligned}$$

4. Trivial, since each simple integration is divideable.

5. Note that $\min(f(x), n) \leq \min(f(x), n+1), \forall x \in \mathbb{R}^d$. By monotonicity,

$$\underline{\int_{\mathbb{R}^d}} \min(f(x), n) dx \leq \underline{\int_{\mathbb{R}^d}} \min(f(x), n+1) dx \leq \dots$$

If f is bounded above, then let $m_n = \min \left\{ n, \left\lceil \sup_{x \in \mathbb{R}^d, a.e.} f(x) \right\rceil \right\} \leq n$ and $\lceil \sup_x f(x) \rceil$ is

the smallest integer above $\sup_{x \in \mathbb{R}^d, a.e.} f(x)$; otherwise let $m_n = n$. We subdivide $[0, m_n]$ into $m_n 2^k$ intervals $[(j-1)2^{-k}, j2^{-k}]$, $j = 1, \dots, m_n 2^k$ with $E_{j,k} \equiv f^{-1}([(j-1)2^{-k}, j2^{-k}])$ and define the simple function as

$$f_{n,k} = \sum_{j=1}^{m_n 2^k} \frac{j-1}{2^k} \mathbb{1}_{\{x \in E_{j,k}\}} + m_n \mathbb{1}_{\{\mathbb{R}^d / f^{-1}[0, m_n]\}}.$$

Clearly, $f_{n,k} \leq \min\{f, n\} \leq f$ for all x , all n, k and $f_{n,k} \uparrow \min\{f, n\}$ as $k \rightarrow \infty$. Also

$$\begin{aligned} \int_{\mathbb{R}^d} \min(f(x), n) dx &= \lim_{k \rightarrow \infty} \text{simp} \int_{\mathbb{R}} f_{n,k}(x) dx \\ &= \left(\lim_{k \rightarrow \infty} \sum_{j=1}^{m_n 2^k} \frac{j-1}{2^k} m(E_{j,k}) \right) + m_n \left(\mathbb{R}^d / f^{-1}([0, m_n]) \right) \end{aligned}$$

If f is bounded above almost everywhere, for any $\epsilon > 0$, there exists $M \equiv \lceil \sup_x f(x) \rceil < \infty$, such that for all $n \geq M$, $\exists K$, for all $k \geq K$,

$$\begin{aligned} \int_{\mathbb{R}^d} f(x) dx - \int_{\mathbb{R}^d} \min(f(x), n) dx &\leq \int_{\mathbb{R}^d} f(x) dx - \sum_{j=1}^{M 2^k} \frac{j-1}{2^k} m(E_{j,k}) - m_n \left(\mathbb{R}^d / f^{-1}([0, m_n]) \right) \\ &\leq \epsilon \end{aligned}$$

If $f \geq M$ for any $M \geq 0$, $x \in E$, then $\int_{\mathbb{R}^d} f(x) dx = \infty$ and for $n \geq M$, $\int_{\mathbb{R}^d} \min(f(x), n) dx \geq M m(E) \rightarrow \infty$. So it completes the proof.

6. Denote $B_n = \{x : |x| \leq n\}$. We subdivide $[0, k]$ into $k 2^k$ intervals $[(j-1)2^{-k}, j2^{-k}]$, $j = 1, \dots, k 2^k$ with $E_{j,k} \equiv f^{-1}([(j-1)2^{-k}, j2^{-k}])$ and define the simple function as

$$f_k = \sum_{j=1}^{k 2^k} \frac{j-1}{2^k} \mathbb{1}_{\{E_{j,k}\}} + k \mathbb{1}_{\{\mathbb{R}^d / f^{-1}([0, k])\}}.$$

Thus $f_k \leq f$ for all k and $f_k \uparrow f$ for $k \rightarrow \infty$. Therefore

$$\begin{aligned} \int_{\mathbb{R}^d} f(x) dx &= \lim_{k \rightarrow \infty} \text{simp} \int_{\mathbb{R}} f_k(x) dx \\ \int_{\mathbb{R}^d} f(x) \mathbb{1}_{\{B_n\}} dx &= \lim_{k \rightarrow \infty} \text{simp} \int_{\mathbb{R}} f_k \mathbb{1}_{\{B_n\}} dx \\ &= \lim_{k \rightarrow \infty} \left\{ \sum_{j=1}^{k 2^k} \frac{j-1}{2^k} m(E_{j,k} \cap B_n) + k m(B_n / f^{-1}([0, k])) \right\} \end{aligned}$$

Since $n \rightarrow \infty$,

$$\begin{aligned} \lim_{n \rightarrow \infty} m(E_{j,k} \cap B_n) &= m(E_{j,k}), \quad \forall j, k \\ \lim_{n \rightarrow \infty} m(B_n / f^{-1}([0, k])) &= m(\mathbb{R}^d / f^{-1}([0, k])), \end{aligned}$$

then

$$\begin{aligned}\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} f(x) \mathbb{1}_{\{B_n\}} dx &= \lim_{k \rightarrow \infty} \left\{ \sum_{j=1}^{k2^k} \frac{j-1}{2^k} \left(\lim_{n \rightarrow \infty} m(E_{j,k} \cap B_n) \right) + k \lim_{n \rightarrow \infty} m(B_n / f^{-1}([0, k])) \right\} \\ &= \lim_{k \rightarrow \infty} \left\{ \sum_{j=1}^{k2^k} \frac{j-1}{2^k} m(E_{j,k}) + k m(\mathbb{R}^d / f^{-1}([0, k])) \right\} \\ &= \int_{\mathbb{R}^d} f(x) dx.\end{aligned}$$

To show $m(E \cap B_n) \rightarrow m(E)$ for any measurable set, we see that $\{E \cap B_n\}$ is monotone nondecreasing, and $E \cap B_n \uparrow E \cap \mathbb{R}^d = E$ for any $E \subseteq \mathbb{R}^d$. Then by upward convergence of the Lebesgue measure, the result holds. ■

- **Definition (Unsigned Lebesgue integral).**

If $f : \mathbb{R}^d \rightarrow [0, +\infty]$ is *measurable*, we define the *unsigned Lebesgue integral* $\int_{\mathbb{R}^d} f(x) dx$ of f to *equal* the *lower unsigned Lebesgue integral* $\int_{\mathbb{R}^d} f(x) dx$. (For non-measurable functions, we leave the unsigned Lebesgue integral undefined.)

- One nice feature of measurable functions is that *the lower and upper Lebesgue integrals can match*, if one also assumes some *boundedness*:

Exercise 1.13 Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be measurable, *bounded*, and *vanishing* outside of a set of *finite measure*. The *lower* and *upper* Lebesgue integrals of f *agree*.

- **Corollary 1.14 (Finite additivity of the Lebesgue integral).**

Let $f, g : \mathbb{R}^d \rightarrow [0, +\infty]$ be measurable. Then

$$\int_{\mathbb{R}^d} (f(x) + g(x)) dx = \int_{\mathbb{R}^d} f(x) dx + \int_{\mathbb{R}^d} g(x) dx.$$

Proof: From *the horizontal truncation property* and a *limiting argument*, we may assume that f, g are *bounded*. From the *vertical truncation property* and *another limiting argument*, we may assume that f, g are *supported inside a bounded set*. Under condition that f, g are both *measurability*, *bounded*, *vanishing outside a set of finite measure*, we now see that the lower and upper Lebesgue integrals of f, g , and $f + g$ agree. The claim now follows by *combining the superadditivity of the lower Lebesgue integral with the subadditivity of the upper Lebesgue integral*. ■

- **Exercise 1.15 (Upper Lebesgue integral and outer Lebesgue measure).**

Show that for any set $E \subseteq \mathbb{R}^d$, $\int_{\mathbb{R}^d} \mathbb{1}_E(x) dx = m^*(E)$. Conclude that the *upper* and *lower Lebesgue integrals* are *not necessarily additive* if *no measurability hypotheses* are assumed.

- **Exercise 1.16 (Area interpretation of integral).**

If $f : \mathbb{R}^d \rightarrow [0, +\infty]$ is *measurable*, show that $\int_{\mathbb{R}^d} f(x) dx$ is equal to the $(d+1)$ -dimensional *Lebesgue measure* of the *region*

$$\{(x, t) \in \mathbb{R}^d \times \mathbb{R} : 0 \leq t \leq f(x)\}$$

(This can be used as an *alternate*, and more *geometrically intuitive, definition* of the *unsigned Lebesgue integral*; it is a more convenient formulation for establishing the basic

convergence theorems, but not quite as convenient for establishing basic properties such as additivity.)

• **Proposition 1.17** (*Uniqueness of the Lebesgue integral*).

The **Lebesgue integration** $f \mapsto \int_{\mathbb{R}^d} f(x)dx$ is the only map from measurable unsigned functions $f : \mathbb{R}^d \rightarrow [0, +\infty]$ to $[0, \infty]$ that obeys the following properties for measurable $f, g : \mathbb{R}^d \rightarrow [0, +\infty]$:

1. (**Compatible with simple integration**): If f is simple, $\int_{\mathbb{R}^d} f dx = \text{simp} \int_{\mathbb{R}^d} f dx$
2. (**Finite Additivity**): $\int_{\mathbb{R}^d} (f(x) + g(x))dx = \int_{\mathbb{R}^d} f(x)dx + \int_{\mathbb{R}^d} g(x)dx$.
3. (**Horizontal truncation**) As $n \rightarrow \infty$,

$$\int_{\mathbb{R}^d} \min(f(x), n)dx \rightarrow \int_{\mathbb{R}^d} f(x)dx.$$

4. (**Vertical truncation**) As $n \rightarrow \infty$,

$$\int_{\mathbb{R}^d} f(x) \mathbf{1}_{\{|x| \leq n\}} dx \rightarrow \int_{\mathbb{R}^d} f(x)dx.$$

• **Exercise 1.18** (*Translation invariance*).

Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be measurable. Show that $\int_{\mathbb{R}^d} f(x+y)dx = \int_{\mathbb{R}^d} f(x)dx$ for any $y \in \mathbb{R}^d$.

• **Exercise 1.19** (*Linear change of variables*).

Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be measurable, and let $T : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be an invertible linear transformation. Show that

$$\int_{\mathbb{R}^d} f(T^{-1}(x))dx = |\det T| \int_{\mathbb{R}^d} f(x)dx,$$

or equivalently $\int_{\mathbb{R}^d} f(Tx)dx = |\det T|^{-1} \int_{\mathbb{R}^d} f(x)dx$.

• **Exercise 1.20** (*Compatibility with the Riemann integral*).

Let $f : [a, b] \rightarrow [0, +\infty]$ be Riemann integrable. If we extend f to \mathbb{R} by declaring f to equal zero outside of $[a, b]$, show that

$$\int_{\mathbb{R}^d} f(x)dx = \int_a^b f(x)dx.$$

• **Lemma 1.21** (*Markov's inequality*).

Let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be **measurable**. Then for any $0 < \lambda < \infty$, one has

$$m\left(\left\{x \in \mathbb{R}^d : f(x) \geq \lambda\right\}\right) \leq \frac{1}{\lambda} \int_{\mathbb{R}^d} f(x)dx$$

Proof: We have the trivial pointwise inequality

$$\lambda \mathbf{1}_{\left\{x \in \mathbb{R}^d : f(x) \geq \lambda\right\}} \leq f(x).$$

From the *definition* of the lower Lebesgue integral, we conclude that

$$\lambda m\left(\left\{x \in \mathbb{R}^d : f(x) \geq \lambda\right\}\right) \leq \int_{\mathbb{R}^d} f(x)dx$$

and the claim follows. ■

- By sending λ to infinity or to zero, we obtain the following important corollary:

Exercise 1.22 Let function $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be measurable. Then

1. Show that if $\int_{\mathbb{R}^d} f(x)dx < \infty$, f is finite almost everywhere. Give a counterexample to show that the converse statement is false.
2. Show that $\int_{\mathbb{R}^d} f(x)dx = 0$ if and only if $f = 0$ almost everywhere.

Proof: 1. Suppose that there exists subset $E \subset \mathbb{R}^d$, $m(E) > 0$ such that $f(\mathbf{x}) > M \geq 0$, $\mathbf{x} \in E$ for all $M \geq 0$. Let $g(\mathbf{x}) = M\mathbb{1}\{\mathbf{x} \in E\}$ be a unsigned simple function, then $f \geq g$ and $\int_{\mathbb{R}^d} f(x)dx \geq \int_{\mathbb{R}^d} g(x)dx = Mm(E) \rightarrow \infty$ for $M \rightarrow \infty$. Then $\int_{\mathbb{R}^d} f(x)dx = \infty$, which contradicts with the assumption.

For converse part, see that $f(\mathbf{x}) = \sum_{n=1}^{\infty} \frac{2^{-n}}{\|\mathbf{x} - \mathbf{q}_n\|^{1/2}}$, where $\mathbf{q}_n \in \mathbb{Q}^d \cap [0, 1]^d$ is finite everywhere $\mathbf{x} \in [0, 1]^d$, but $\int_{[0, 1]^d} f(x)dx \geq \int_{[0, 1]^d} \frac{1}{\|\mathbf{x} - \mathbf{q}_n\|} = \infty$. For f defined on all \mathbb{R} , simply let $f(\mathbf{x} + \mathbf{d}) = f(\mathbf{x})$ for all $\mathbf{d} \in \mathbb{N}^d$.

2. Suppose that there exists subset $E \subset \mathbb{R}^d$, $m(E) > 0$ such that for some $\epsilon > 0$, $f(\mathbf{x}) > \epsilon$, $\mathbf{x} \in E$. Then we can find a simple function $g(\mathbf{x}) = \epsilon\mathbb{1}\{\mathbf{x} \in E\}$ such that $f \geq g$ and $\int_{\mathbb{R}^d} g(x)dx = \epsilon m(E) > 0$. Thus $\int_{\mathbb{R}^d} f(x)dx > 0$, resulting in contradiction.

Conversely, if $f = 0$ outside a set of measure zero, then for any simple unsigned function $0 \leq g \leq f$, g is nonzero only for a set of measure zero. Therefore $\int_{\mathbb{R}^d} g(x)dx = 0$, so $\int_{\mathbb{R}^d} f(x)dx = 0$. ■

- **Remark** The use of the integral $\int_{\mathbb{R}^d} f(x)dx$ to control the *distribution* of f is known as **the first moment method**. One can also control this distribution using **higher moments** such as $\int_{\mathbb{R}^d} |f(x)|^p dx$ for various values of p , or **exponential moments** such as $\int_{\mathbb{R}^d} e^{t f(x)} dx$ or **the Fourier moments** $\int_{\mathbb{R}^d} e^{it f(x)} dx$ for various values of t ; such *moment methods* are *fundamental* to probability theory.

1.5 Absolute Integrable

- **Definition** (*Absolute integrability*).

An almost everywhere defined measurable function $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is said to be **absolutely integrable** if **the unsigned integral**

$$\|f\|_{L^1(\mathbb{R}^d)} := \int_{\mathbb{R}^d} |f(\mathbf{x})| d\mathbf{x} < \infty.$$

We refer to this quantity $\|f\|_{L^1(\mathbb{R}^d)}$ as **the $L^1(\mathbb{R}^d)$ norm of f** , and use $L^1(\mathbb{R}^d)$ or $L^1(\mathbb{R}^d \rightarrow \mathbb{C})$ to denote **the space of absolutely integrable functions**. If f is **real-valued** and absolutely integrable, we define **the Lebesgue integral** $\int_{\mathbb{R}^d} f(x)dx$ by the formula

$$\int_{\mathbb{R}^d} f(x)dx = \int_{\mathbb{R}^d} f_+(x)dx - \int_{\mathbb{R}^d} f_-(x)dx$$

where $f_+ = \max\{f, 0\}$ and $f_- = \max\{-f, 0\}$ are the magnitudes of the positive and negative components of f . (note that the two unsigned integrals on the right-hand side are finite, as f_+, f_- are pointwise dominated by $|f|$). If f is **complex-valued** and absolutely integrable,

we define the Lebesgue integral $\int_{\mathbb{R}^d} f(x)dx$ by the formula

$$\int_{\mathbb{R}^d} f(x)dx = \int_{\mathbb{R}^d} \Re f(x)dx + i \int_{\mathbb{R}^d} \Im f(x)dx,$$

where the two integrals *on the right* are interpreted as **real-valued absolutely integrable Lebesgue integrals**. It is easy to see that the unsigned, real-valued, and complex-valued Lebesgue integrals defined in this manner are *compatible* on their common domains of definition.

- **Remark** ($L^1(\mathbb{R}^d)$ is a Normed Vector Space)

From the pointwise triangle inequality $|f(x) + g(x)| \leq |f(x)| + |g(x)|$ we conclude the L^1 **triangle inequality**

$$\|f + g\|_{L^1(\mathbb{R}^d)} \leq \|f\|_{L^1(\mathbb{R}^d)} + \|g\|_{L^1(\mathbb{R}^d)},$$

for any *almost everywhere defined measurable* $f, g : \mathbb{R}^d \rightarrow \mathbb{C}$. It is also easy to see that

$$\|cf\|_{L^1(\mathbb{R}^d)} \leq |c| \|f\|_{L^1(\mathbb{R}^d)},$$

for any complex number c . As such, we see that $L^1(\mathbb{R}^d \rightarrow \mathbb{C})$ is a **complex vector space**.

- **Remark** (L^1 distance on $L^1(\mathbb{R}^d)$)

Given two functions $f, g \in L^1(\mathbb{R}^d \rightarrow \mathbb{C})$, we can define the L^1 distance $d_{L^1}(f, g)$ between them by the formula

$$d_{L^1}(f, g) := \|f - g\|_{L^1(\mathbb{R}^d)}.$$

Thanks to the triangle inequality, this distance obeys almost all the axioms of a metric on $L^1(\mathbb{R}^d)$, with **one exception**: it is possible for *two different functions* $f, g \in L^1(\mathbb{R}^d \rightarrow \mathbb{C})$ to have a *zero L^1 distance*, if they *agree almost everywhere*. As such, d_{L^1} is only **a semi-metric** (also known as a **pseudo-metric**) rather than a metric.

However, if one adopts the convention that any two functions that agree almost everywhere are considered **equivalent** (or more formally, one works in *the quotient space* of $L^1(\mathbb{R}^d)$ by **the equivalence relation of almost everywhere agreement**, which by abuse of notation is also denoted $L^1(\mathbb{R}^d)$, then *one recovers a genuine metric*.

- **Exercise 1.23** If $f : \mathbb{R}^d \rightarrow \mathbb{C}$ is absolutely integrable, then f is bounded almost everywhere by Markov inequality.
- **Exercise 1.24** $f : E \rightarrow \mathbb{C}$ is a function, we say that f is measurable (resp. absolutely integrable) if its extension $\hat{f} : \mathbb{R}^d \rightarrow \mathbb{C}$ is a (absolute) measurable function, s.t. $\hat{f} = f$ for all $x \in E$ and is zero outside. Then

$$\int_E f(x)dx \equiv \int_{\mathbb{R}^d} \hat{f}(x)dx.$$

- **Exercise 1.25** If E, F are disjoint sets, $f : E \cup F \rightarrow \mathbb{C}$, then

$$\int_E f(x)dx = \int_{E \cup F} f(x) \mathbb{1}_{\{x \in E\}} dx$$

and

$$\int_{E \cup F} f(x)dx = \int_E f(x)dx + \int_F f(x)dx.$$

• **Lemma 1.26** (*The Triangle Inequality*):

Let $f \in L^1(\mathbb{R}^d)$, then

$$\left| \int_{\mathbb{R}^d} f(x) dx \right| \leq \int_{\mathbb{R}^d} |f(x)| dx.$$

Proof: If f is real-valued, then $|f| = f_+ + f_-$ and the claim is obvious from triangle inequality.

When f is *complex-valued*, one *cannot argue quite so simply*; a naive mimicking of the real-valued argument would *lose a factor of 2*, giving the inferior bound

$$\left| \int_{\mathbb{R}^d} f(x) dx \right| \leq 2 \int_{\mathbb{R}^d} |f(x)| dx.$$

To do better, we exploit *the phase rotation invariance properties* of the *absolute value operation* and of the *integral*, as follows. Note that for any complex number z , one can write $|z|$ as $ze^{i\theta}$ for some real θ . In particular, we have

$$\left| \int_{\mathbb{R}^d} f(x) dx \right| = e^{i\theta} \int_{\mathbb{R}^d} f(x) dx = \int_{\mathbb{R}^d} e^{i\theta} f(x) dx$$

for some real θ . Taking *real parts of both sides*, we obtain

$$\left| \int_{\mathbb{R}^d} f(x) dx \right| = \int_{\mathbb{R}^d} \Re \left(e^{i\theta} f(x) \right) dx.$$

Since $\Re(e^{i\theta} f(x)) \leq |e^{i\theta} f(x)| = |f(x)|$, we obtain the claim. ■

2 Development of Lebesgue Integration

Table 1: Development on Lebesgue Integration

	<i>Unsigned Simple Function</i>	<i>Unsigned Measurable Function</i>	<i>Absolute Integrable Function</i>
Definition	$f = \sum_{i=k}^m c_k \mathbb{1}_{E_k}$	$\{f_n\} \rightarrow f$ pointwise $\{f_n\}$ unsigned simple	$\ f\ _{L^1(\mathbb{R}^d)} < \infty$ $ f $ unsigned
Integration	$\text{simp} \int_{\mathbb{R}^d} f(x) dx$ $= \sum_{i=k}^m c_k \mu(E_k)$	$\int_{\mathbb{R}^d} f(x) dx = \int_{\mathbb{R}^d} f(x) dx =$ $\sup_{0 \leq g \leq f, g \text{ simple}} \text{simp} \int_{\mathbb{R}^d} g(x) dx$	$\int_{\mathbb{R}^d} f(x) dx =$ $\int_{\mathbb{R}^d} f_+(x) dx - \int_{\mathbb{R}^d} f_-(x) dx$
<i>Compatibility</i>		✓	✓
<i>Compatibility to Riemann Integral</i>	✓	✓	✓
<i>Linearity</i>	Unsigned ✓	✓	✓
Equivalence	✓	✓	✓
<i>Vanishing</i>	✓	✓	✓
Monotonicity	✓	✓	✓
Superadditivity		lower Lebesgue integral ✓	
<i>Reflection</i>		lower Lebesgue integral ✓	
<i>Divisibility</i>		lower Lebesgue integral ✓	
Finite additivity	✓	✓	✓
Horizontal truncation		✓	✓
Vertical truncation		✓	✓
<i>Translation Invariance</i>		✓	✓

3 Examples

3.1 Unsigned Measureable Functions

- **Example** If $f : \mathbb{R}^d \rightarrow [0, +\infty]$ is measurable, then $f^{-1}(E)$ is *Lebesgue measurable* for many classes of sets E . However, we caution that it is **not necessarily** the case that $f^{-1}(E)$ is *Lebesgue measurable* if E is *Lebesgue measurable*.

Proof: We let C be *the Cantor set*

$$C := \left\{ \sum_{j=1}^{\infty} a_j 3^{-j} : a_j \in \{0, 2\}, \forall j \right\}$$

and let $f : \mathbb{R}^d \rightarrow [0, +\infty]$ be the function defined by setting

$$f(x) = \sum_{j=1}^{\infty} 2b_j 3^{-j}$$

whenever $x \in [0, 1]$ is *not a terminating binary decimal*, and so has a *unique binary expansion* $x = \sum_{j=1}^{\infty} b_j 2^{-j}$ for some $b_j \in \{0, 1\}$, and $f(x) \equiv 0$ otherwise. We thus see that f takes values in C , and is *bijective* on the set A of *non-terminating decimals* in $[0, 1]$.

Using Lemma 1.7, it is not difficult to show that f is measurable. On the other hand, by modifying the construction from the previous notes, we can find a subset F of A which is *non-measurable*. If we set $E := f(F)$, then E is a subset of *the null set* C and is thus itself *a null set*; but $f^{-1}(E) = F$ is *non-measurable*, and so the *inverse image of a Lebesgue measurable set by a measurable function need not remain Lebesgue measurable*. However, we will later see that it is still true that $f^{-1}(E)$ is *Lebesgue measurable* if E has a *slightly stronger measurability property* than *Lebesgue measurability*, namely *Borel measurability*.

References

Terence Tao. *An introduction to measure theory*, volume 126. American Mathematical Soc., 2011.