

Lecture 0: Summary (part 1)

Tianpei Xie

Oct. 24th., 2022

Contents

1	Topology	2
1.1	Topological Space	2
1.2	Subspaces and Quotients	3
1.3	Connectedness and Compactness	3
2	Smooth Manifolds and Smooth Maps	5
2.1	From Topological Manifolds to Smooth Manifolds	5
2.2	Manifolds with boundary	7
2.3	Smooth Maps on Manifolds	8
2.4	Einstein summation convention	9
3	Submersions, Immersions and Embeddings	9
3.1	Definitions	11
3.2	Local Diffeomorphisms	11
3.3	The Rank Theorem	11
3.4	Embeddings	12
3.5	Submersions	13
4	Submanifolds	14
4.1	Embedded Submanifolds	15
4.1.1	Definitions	15
4.1.2	Slice Charts for Embedded Submanifolds	16
4.1.3	Level Sets	16
4.2	Immersed Submanifolds	18
4.2.1	Definitions and Examples	18
4.3	Restricting Maps to Submanifolds	19
4.3.1	Restricting Domains and Codomains	19
4.3.2	Uniqueness of Smooth Structures on Submanifolds	20
4.3.3	Extending Functions from Submanifolds	20
4.4	The Tangent Space to a Submanifold	20
4.5	Vector Fields and Submanifolds	22
4.6	Restricting Covector Fields to Submanifolds	22

1 Topology

1.1 Topological Space

- **Definition** Let X be a set. A **topology** on X is a collection \mathcal{T} of subsets of X , called **open subsets**, satisfying
 1. X and \emptyset are *open*.
 2. The **union** of *any family* of open subsets is open.
 3. The **intersection** of *any finite family* of open subsets is open.

A pair (X, \mathcal{T}) consisting of a set X together with a topology \mathcal{T} on X is called a **topological space**.

- **Definition** A map $F : X \rightarrow Y$ is said to be **continuous** if for every open subset $U \subseteq Y$, the **preimage** $F^{-1}(U)$ is **open** in X .
- **Definition** A **continuous bijective** map $F : X \rightarrow Y$ with **continuous inverse** is called a **homeomorphism**. If there exists a **homeomorphism** from X to Y , we say that X and Y are **homeomorphic**.
- **Definition** A map $F : X \rightarrow Y$ (continuous or not) is said to be an **open map** if for every *open* subset $U \subseteq X$, the image set $F(U)$ is *open* in Y , and a **closed map** if for every *closed* subset $K \subseteq X$, the image $F(K)$ is *closed* in Y .
- **Definition** A topological space X is said to be a **Hausdorff space** if for every pair of **distinct** points $p, q \in X$, there exist **disjoint open subsets** $U, V \subseteq X$ such that $p \in U$ and $q \in V$.
- **Definition** Suppose X is a topological space. A collection \mathcal{B} of open subsets of X is said to be a **basis** for the topology of X (plural: **bases**) if every open subset of X is the *union of some collection of elements* of \mathcal{B} .

More generally, suppose X is merely a set, and \mathcal{B} is a collection of *subsets* of X satisfying the following conditions:

1. $X = \bigcup_{B \in \mathcal{B}} B$.
2. If $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, then there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.

Then the collection of **all unions** of elements of \mathcal{B} is a topology on X , called **the topology generated by \mathcal{B}** , and \mathcal{B} is a **basis** for this topology.

- **Definition** See the following definitions
 1. A set is said to be **countably infinite** if it admits a *bijection* with the set of *positive integers*, and
 2. **countable** if it is *finite* or *countably infinite*.
 3. A topological space X is said to be **first-countable** if there is a **countable neighborhood basis** at each point, and
 4. **second-countable** if there is a **countable basis** for its topology.

1.2 Subspaces and Quotients

- **Definition** If X is a topological space and $S \subseteq X$ is an arbitrary subset, we define *the subspace topology* on S (sometimes called *the relative topology*) by declaring a subset $U \subseteq S$ to be *open* in S if and only if there exists an open subset $V \subseteq X$ such that $U = V \cap S$.

Any subset of X endowed with the subspace topology is said to be *a subspace of X* .

- **Definition** If X and Y are topological spaces, a continuous injective map $F : X \rightarrow Y$ is called a *topological embedding* if it is a *homeomorphism* onto its image $F(X) \subseteq Y$ in the subspace topology.
- **Definition** If X is a topological space, Y is a set, and $\pi : X \rightarrow Y$ is a **surjective** map, *the quotient topology* on Y determined by π is defined by declaring a subset $U \subseteq Y$ to be *open* if and only if $\pi^{-1}(U)$ is *open* in X .

If X and Y are topological spaces, a map $\pi : X \rightarrow Y$ is called *a quotient map* if it is *surjective* and *continuous* and Y has the quotient topology determined by π .

- **Definition** The following construction is the most common way of producing quotient maps. A *relation* on a set X is called *an equivalence relation* if it is
 1. *reflexive*: $x \sim x$ for all $x \in X$,
 2. *symmetric*: $x \sim y$ implies $y \sim x$,
 3. *transitive*: $x \sim y$ and $y \sim z$ imply $x \sim z$.

If $R \subseteq X \times X$ is any *relation* on X , then *the intersection of all equivalence relations on X containing R* is an *equivalence relation*, called *the equivalence relation generated by R* .

Remark If \sim is an equivalence relation on X , then for each $x \in X$, *the equivalence class of x* , denoted by $[x]$, is the *set of all $y \in X$ such that $y \sim x$* . The set of *all equivalence classes* is a *partition* of X : a collection of disjoint nonempty subsets whose union is X .

- **Definition** Suppose X is a topological space and \sim is an equivalence relation on X . Let X/\sim denote *the set of equivalence classes* in X , and let $\pi : X \rightarrow X/\sim$ be the *natural projection* sending each *point* to its *equivalence class*. Endowed with *the quotient topology* determined by π , the space X/\sim is called *the quotient space* (or *identification space*) of X determined by π .
- **Definition** If $\pi : X \rightarrow Y$ is a map, a subset $U \subseteq X$ is said to be *saturated* with respect to π if U is the *entire preimage* of its *image*: $U = \pi^{-1}(\pi(U))$.
Given $y \in Y$, the *fiber* of π over y is the set $\pi^{-1}(y)$.

1.3 Connectedness and Compactness

- **Definition** A topological space X is said to be *disconnected* if it has two *disjoint nonempty open subsets* whose union is X , and it is *connected* otherwise. Equivalently, X is connected if and only if the only subsets of X that are *both open and closed* are \emptyset and X itself.

- **Definition** Recall that a topological space X is
 - **connected** if there do not exist two *disjoint, nonempty, open* subsets of X whose union is X ;
 - **path-connected** if every pair of points in X can be *joined by a path* in X , and
 - ***locally path-connected*** if X has a **basis** of *path-connected open subsets*.
- **Definition** A ***maximal connected subset*** of X (i.e., a connected subset that is not properly contained in any larger connected subset) is called a **component** (or **connected component**) of X .
- **Definition** A topological space X is said to be **compact** if every open cover of X has a *finite subcover*. A **compact subset** of a topological space is one that is a compact space in the subspace topology.
- **Definition** If X and Y are topological spaces, a map $F : X \rightarrow Y$ (continuous or not) is said to be **proper** if for every **compact** set $K \subseteq Y$, the **preimage** $F^{-1}(K)$ is **compact**.
- **Definition** A topological space X is said to be **locally compact** if every point has a *neighborhood* contained in a **compact subset** of X .

A subset of X is said to be **precompact** in X if its **closure** in X is **compact**.

-
- For a **Hausdorff space** X , the following are equivalent:
 1. X is **locally compact**.
 2. Each point of X has a **precompact** neighborhood.
 3. X has a basis of **precompact** open subsets.

2 Smooth Manifolds and Smooth Maps

2.1 From Topological Manifolds to Smooth Manifolds

- **Definition** Suppose M is a *topological space*. We say that M is a *topological manifold* of dimension n or a *topological n -manifold* if it has the following properties:

1. M is a *Hausdorff space*: for every pair of distinct points $p, q \in M$, there are disjoint open subsets $U, V \subseteq M$ such that $p \in U$ and $q \in V$.
2. M is *second-countable*: there exists a *countable basis* for the topology of M .
3. M is *locally Euclidean of dimension n* : each point of M has a neighborhood that is *homeomorphic* to an open subset of \mathbb{R}^n .

- The third property means, more specifically, that for each $p \in M$ we can find

- an open subset $U \subseteq M$ containing p ,
- an open subset $\widehat{U} \subseteq \mathbb{R}^n$, and
- a *homeomorphism* $\varphi : U \rightarrow \widehat{U}$.

- **Proposition 2.1** (*Manifolds Are Locally Compact*).

Every topological manifold is *locally compact*.

- **Definition** Let M be a topological n -manifold. A *coordinate chart* (or just a *chart*) on M is a *pair* (U, φ) , where U is an open subset of M and $\varphi : U \rightarrow \widehat{U}$ is a *homeomorphism* from U to an open subset $\widehat{U} = \varphi(U) \subseteq \mathbb{R}^n$.

- **Definition** Given a chart (U, φ) , we call the set U a *coordinate domain*, or a *coordinate neighborhood* of each of its points. The map φ is called a *(local) coordinate map*, and the *component functions* (x^1, \dots, x^n) of φ , defined by $\varphi(p) = (x^1(p), \dots, x^n(p))$, are called *local coordinates* on U .

- **Remark** We sometimes write things such as “ (U, φ) is a chart containing p ” as shorthand for “ (U, φ) is a chart whose domain U contains p .” If we wish to emphasize the coordinate function (x^1, \dots, x^n) instead of coordinate map φ , we sometimes denote the chart by $(U, (x^1, \dots, x^n))$ or $(U, (x^i))$.

- **Definition** If U and V are open subsets of Euclidean spaces \mathbb{R}^n and \mathbb{R}^m , respectively, a function $F : U \rightarrow V$ is said to be *smooth* (or C^∞ , or *infinitely differentiable*) if each of its component functions has continuous partial derivatives of *all orders*.

If in addition F is *bijective* and has a *smooth inverse map*, it is called a *diffeomorphism*. A *diffeomorphism* is, in particular, a *homeomorphism*.

- **Definition** Let M be a topological n -manifold. If (U, φ) , (V, ψ) are two charts such that $U \cap V \neq \emptyset$, the composite map $\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$ is called the *transition map* from φ to ψ . It is a *homeomorphism*.

Two charts (U, φ) and (V, ψ) are said to be *smoothly compatible* if either $U \cap V = \emptyset$ or the transition map $\psi \circ \varphi^{-1}$ is a *diffeomorphism*.

- **Definition** We define an *atlas* \mathcal{A} for M to be a collection of charts whose domains *cover*

M , i.e. $\mathcal{A} := \{(U_\alpha, \varphi_\alpha)\}_{\alpha \in A}$ such that $M = \bigcup_{\alpha \in A} U_\alpha$. An atlas \mathcal{A} is called a **smooth atlas** if any two charts in \mathcal{A} are **smoothly compatible** with each other.

- **Definition** A smooth atlas \mathcal{A} on M is **maximal** if it is **not** properly contained in **any larger** smooth atlas. This just means that any chart that is *smoothly compatible* with *every* chart in \mathcal{A} is already in \mathcal{A} . (Such a smooth atlas is also said to be **complete**.)
- **Definition** If M is a topological manifold, a **smooth structure** on M is a **maximal smooth atlas**.

A **smooth manifold** is a pair (M, \mathcal{A}) , where M is a **topological manifold** and \mathcal{A} is a **smooth structure** on M .

- **Remark** When the smooth structure is understood, we usually omit mention of it and just say “ M is a smooth manifold.” Smooth structures are also called **differentiable structures** or C^∞ **structures** by some authors. We also use the term **smooth manifold structure** to mean a manifold topology together with a smooth structure.
- **Remark** When defining smooth manifold, we ask for any two coordinate charts $(U, \varphi), (V, \psi)$, **the transition map** $\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$ is a diffeomorphism. This is **stronger** than *the coordinate map* φ itself being a diffeomorphism.
- **Remark** In practice, instead of specifying the maximal smooth atlas, it is sufficient to specify *some* smooth atlas to verify if M is smooth manifold

Proposition 2.2 *Let M be a topological manifold.*

1. *Every smooth atlas \mathcal{A} for M is contained in a **unique** maximal smooth atlas, called **the smooth structure determined by \mathcal{A}** .*
2. *Two smooth atlases for M determine the same smooth structure if and only if **their union is a smooth atlas**.*

- **Definition** If M is a smooth manifold, any chart (U, φ) contained in the given maximal smooth atlas is called a **smooth chart**, and the corresponding coordinate map φ is called a **smooth coordinate map**.

It is useful also to introduce the terms **smooth coordinate domain** or **smooth coordinate neighborhood** for the domain of a smooth coordinate chart.

- The following result is a generalization of *second-countable* definition for the general topological manifold.

Proposition 2.3 *Every smooth manifold has a **countable basis of regular coordinate balls**.*

- **Remark** Each coordinate map φ maps a smooth neighborhood $U \subseteq M$ to a neighborhood in Euclidean space $\tilde{U} \subseteq \mathbb{R}^n$. Under this map, we can **(locally) represent** a point $p \in U$ by its **coordinates** $(x^1, \dots, x^n) = \varphi(p)$, and think of this n -tuple as *being* the point p . We typically express this by saying “ (x^1, \dots, x^n) is **the (local) coordinate representation** for p ” or “ $p = (x^1, \dots, x^n)$ **in local coordinates**.”
- **Remark** As we see, a smooth manifold M does not come with any predetermined **choice of coordinates**. Thus any objects we wish to define globally on a manifold **shall not depend on a particular choice of coordinates**.

There are generally two ways of doing this:

- either by writing down a **coordinate-dependent definition** and then proving that the **definition gives the same results in any coordinate chart**,
- or by writing down a **definition that is manifestly coordinate-independent** (often called an **invariant definition**).

2.2 Manifolds with boundary

- **Definition** The **closed n -dimensional upper half-space** $\mathbb{H}^n \subseteq \mathbb{R}^n$ is defined as

$$\mathbb{H}^n = \{(x^1, \dots, x^n) \in \mathbb{R}^n : x^n \geq 0\}.$$

We will use the notations $\text{Int } \mathbb{H}^n$ and $\partial \mathbb{H}^n$ to denote the **interior** and **boundary** of \mathbb{H}^n , respectively, as a subset of \mathbb{R}^n . When $n > 0$, this means

$$\begin{aligned}\text{Int } \mathbb{H}^n &= \{(x^1, \dots, x^n) \in \mathbb{R}^n : x^n > 0\}, \\ \partial \mathbb{H}^n &= \{(x^1, \dots, x^n) \in \mathbb{R}^n : x^n = 0\}.\end{aligned}$$

- **Definition** An **n -dimensional topological manifold with boundary** is a *second-countable Hausdorff space* M in which every point has a neighborhood **homeomorphic** either to an *open subset* of \mathbb{R}^n or to a (*relatively*) *open subset* of \mathbb{H}^n .
- **Proposition 2.4** *Let M be a topological n -manifold with boundary.*
 1. *$\text{Int } M$ is an **open subset** of M and a topological n -manifold **without boundary**.*
 2. *∂M is a **closed subset** of M and a **topological $(n-1)$ -manifold without boundary**.*
 3. *M is a topological manifold if and only if $\partial M = \emptyset$.*
 4. *If $n = 0$, then $\partial M = \emptyset$ and M is a 0 -manifold.*
- **Definition** Now let M be a topological manifold **with boundary**. As in the manifold case, a **smooth structure** for M is defined to be a *maximal smooth atlas* – a collection of charts whose domains cover M and whose transition maps (and their inverses) are smooth in the sense just described. With such a structure, M is called a **smooth manifold with boundary**.
- **Remark** Note that, despite their name, **manifolds with boundary are not in general manifolds**, because boundary points do not have locally Euclidean neighborhoods. Moreover, a manifold with boundary might have **empty boundary** – there is nothing in the definition that requires the boundary to be a nonempty set.

On the other hand, **a manifold is also a manifold with boundary**, whose boundary is empty. Thus, every manifold is a manifold with boundary, but a manifold with boundary is a manifold if and only if its boundary is empty.

- **Remark** We will often use redundant phrases such as **manifold without boundary** if we wish to emphasize that we are talking about a manifold in the original sense, and **manifold with or without boundary** to refer to a manifold with boundary if we wish emphasize that the boundary might be empty.

2.3 Smooth Maps on Manifolds

- **Definition** Suppose M is a smooth n -manifold, k is a nonnegative integer, and $f : M \rightarrow \mathbb{R}^k$ is any function. We say that f is a **smooth function** if for every $p \in M$, there exists a **smooth chart** (U, φ) for M whose domain contains p and such that the **composite function** $f \circ \varphi^{-1}$ is smooth on the open subset $\widehat{U} = \varphi(U) \subseteq \mathbb{R}^n$.

If M is a smooth manifold **with boundary**, the definition is exactly the same, except that $\varphi(U)$ is now an open subset of either \mathbb{R}^n or \mathbb{H}^n , and in the latter case we interpret smoothness of $f \circ \varphi^{-1}$ to mean that each point of $\varphi(U)$ has a neighborhood (in \mathbb{R}^n) on which $f \circ \varphi^{-1}$ **extends to a smooth function** in the ordinary sense.

- **Definition** Given a function $f : M \rightarrow \mathbb{R}^k$ and a chart (U, φ) for M , the function $\widehat{f} : \varphi(U) \rightarrow \mathbb{R}^k$ defined by $\widehat{f}(x) = f \circ \varphi^{-1}(x)$ is called the **(local) coordinate representation** of f .
- **Remark** With the help of coordinate chart (U, φ) , we can generalize a lot of concepts from Euclidean space to Manifolds. The process of applying φ^{-1} is called **(local) parameterization** and $f \circ \varphi^{-1}$ is the local coordinate representation of the function f , which is **parametric**.

Even though many objects in this course is **independent of the choice of coordinates**, in practice, one need to use *the coordinate representation* of these objects to **compute** associated quantities.

- The definition of smooth functions generalizes easily to maps *between* manifolds.

Definition Let M, N be *smooth manifolds*, and let $F : M \rightarrow N$ be any map. We say that F is a **smooth map** if for every $p \in M$, there exist *smooth charts* (U, φ) containing p and (V, ψ) containing $F(p)$ such that $F(U) \subseteq V$ and **the composite map** $\psi \circ F \circ \varphi^{-1}$ is **smooth** from $\varphi(U)$ to $\psi(V)$.

If M and N are smooth manifolds **with boundary**, smoothness of F is defined in exactly the same way, with the usual understanding that a map whose domain is a subset of \mathbb{H}^n is smooth if it admits an extension to a smooth map in a neighborhood of each point, and a map whose codomain is a subset of \mathbb{H}^n is smooth if it is smooth as a map into \mathbb{R}^n .

- **Definition** If $F : M \rightarrow N$ is a *smooth map*, and (U, φ) and (V, ψ) are any smooth charts for M and N , respectively, we call $\widehat{F} = \psi \circ F \circ \varphi^{-1} : \varphi(U \cap F^{-1}(V)) \rightarrow \psi(V)$ the **coordinate representation** of F with respect to the given coordinates.
- **Remark** Use $F^{-1}(V) \cap U$ is a safer way to make sure the neighborhood V in coordinate chart in N is covered, as compared to using $F(U) \cap V$.
- **Remark** In practice, $\widehat{F} = \psi \circ F \circ \varphi^{-1}$ is the function we used in computation involving F .
- **Proposition 2.5 (Equivalent Characterizations of Smoothness)** [Lee, 2003.]

Suppose M and N are smooth manifolds with or without boundary, and $F : M \rightarrow N$ is a map. Then F is **smooth** if and only if either of the following conditions is satisfied:

1. For every $p \in M$, there exist **smooth charts** (U, φ) containing p and (V, ψ) containing $F(p)$ such that $U \cap F^{-1}(V)$ is **open** in M and the composite map $\psi \circ F \circ \varphi^{-1}$ is **smooth** from $\varphi(U \cap F^{-1}(V))$ to $\psi(V)$.
2. F is **continuous** and there exist **smooth atlases** $\{(U_\alpha, \varphi_\alpha)\}$ and $\{(V_\beta, \psi_\beta)\}$ for M and N , respectively, such that for **each** α and β , $\psi_\beta \circ F \circ \varphi_\alpha^{-1}$ is a smooth map from

$\varphi_\alpha(U_\alpha \cap F^{-1}(V_\beta))$ to $\psi_\beta(V_\beta)$.

- **Definition** If M and N are smooth manifolds with or without boundary, a **diffeomorphism** from M to N is a **smooth bijective map** $F : M \rightarrow N$ that has a **smooth inverse**. We say that M and N are **diffeomorphic** if there exists a *diffeomorphism* between them. Sometimes this is symbolized by $M \approx N$.

2.4 Einstein summation convention

- **Remark** We interpret any such expression according to the *following rule*, called the **Einstein summation convention**:
 - if the **same index name** (such as i in the expression above) appears exactly **twice** in any **monomial term**, once as an **upper index** and once as a **lower index**, that term is understood to be summed over *all possible values of that index*, generally from 1 to the dimension of the space in question.
 - **Basis (vector, functions)** (such as E_i) uses **lower indices**. The coordinate vector of tangent space ($\frac{\partial}{\partial x^i}$) are considered as lower indices.
 - **Coefficients** of a linear combination, **coordinate or component (functions)** of a vector with respect to a basis (such as x^i) use **upper indices**
 - **Basis of covectors (cotangent vectors)** (such as ϵ^i, dx^i) use **upper indices**, while the component of covector with respect to a basis (such as ω_i) use **lower indices**. That is, vector and covector notation rules are switched.
 - Any index that is implicitly summed over is a “**dummy index**,” meaning that the value of such an expression is *unchanged* if a *different name is substituted for each dummy index*. For example, $x^i E_i$ and $x^j E_j$ mean exactly the same thing.

3 Submersions, Immersions and Embeddings

- **Remark** The following two chapters concerns about the local properties of smooth function F on manifolds by analyzing its differentials dF_p . “**The differential of dF_p of a function F is its best linear approximation in the neighborhood of p .**” This statement can be generalized from \mathbb{R}^n to n -dimensional smooth manifold. These two chapters discuss results that centered around this idea.
 1. The first theorem is **the Inverse Function Theorem for Manifolds**. This is a direct generalization of the existing results from Euclidean space, since this theorem only concerns the property of the function **locally** and the manifold is diffeomorphic to Euclidean space locally. The result of this theorem confirms that for a smooth map with invertible differential at a point p , in the neighborhood of this point, this map is **an open map** and **has smooth inverse**.
 2. The second theorem concerns about the smooth map **with constant rank**. Note that the rank of a smooth map is **a local property** since it is about the rank of dF_p at point p . But when we enforce the rank to be constant all over the space, we generalize the

local properties to *global*.

3. **The Rank Theorem** for smooth map *with constant rank* reflects a **stronger** result than *the Inverse Function Theorem*. It states that we *only need to know the rank* of the differential dF_p to determine **a local representation of the smooth map F , regardless of the form of function itself**. In fact, all smooth maps *with constant rank* can be *locally* represented as **coordinate projection with zero padding**. The *rank* of map determines **the number of coordinate maintained** and the others are all zero-padded.
 4. *The Rank Theorem* confirms that the smooth function F with constant rank is **locally linear**, thus is best represented by the differential dF_p .
- **Remark** Another important topic that is primarily discussed in the following two chapters is two special type of smooth map with constant ranks: the **smooth submersion** and **smooth immersion**. Both of these properties characterize **the local differential properties** of a smooth function F **with full rank** (the *shape* of the differential matrix dF_p is determined by the dimension of the manifolds in domain and codomain).

1. The **smooth submersion** corresponds to F with **surjective differential** $dF_p \in \mathbb{R}^{m \times n}$ with $m \geq n$, i.e. the “fat” differential matrix. dF_p has *full column rank*.

A *smooth submersion* can be *locally* represented as **a natural coordinate projection**. Therefore it is critical when studying **projections** and its **section** between manifolds. It is also closely associated with **quotient map** and **the quotient manifolds**. The *pre-image* of a smooth submersion is a submanifold itself, making it important to define new subspace structure.

2. Similarly, **a smooth immersion** corresponds to F with **injective differential** $dF_p \in \mathbb{R}^{m \times n}$ with $m \leq n$, i.e. the differential matrix is a “tall” matrix with *full row rank*.

A smooth immersion can be locally represented as **a natural coordinate inclusion**, or a zero-padding function. It is critical when we try to *put a manifold inside another manifold* and makes it a submanifold of the latter.

- A sub-class of smooth immersion is **smooth embedding**. In addition being smooth immersion, a smooth embedding is also **locally homemorphic to its image**. In other word, it allows one to build an **equivalence relationship** between two *local regions* in *topological sense*.
- Like smooth immersion, a smooth embedding locally is an **inclusion** but the embedding map globally is an **injective** map with **smooth inverse locally**.
- *Smooth embedding map* is commonly used to generate k -dimensional **embedded submanifolds**, which can be seen as a subset equipped with the subspace topology and **locally homemorphic to a k -dimensional subspace in \mathbb{R}^n** .

3.1 Definitions

- **Definition** Suppose M and N are smooth manifolds with or without boundary. Given a smooth map $F : M \rightarrow N$ and a point $p \in M$, we define the rank of F at p to be **the rank of the linear map** $dF_p : T_p M \rightarrow T_{F(p)} N$; it is **the rank of the Jacobian matrix** of F in any smooth chart, or **the dimension of** $\text{Im } dF_p \subseteq T_{F(p)} N$. If F has the same rank r at every point, we say that it has constant rank, and write $\text{rank } F = r$.

- **Definition** Note that $\text{rank } dF_p \leq \min \{\dim M, \dim N\}$. If the rank of dF_p is equal to this upper bound, we say that F **has full rank at p** , and if F has full rank everywhere, we say F has full rank.

- **Definition** The most important *constant-rank maps* are those of *full rank*. A smooth map $F : M \rightarrow N$ is called a smooth submersion if its differential is surjective at each point (or equivalently, if $\text{rank } F = \dim N$).

It is called a smooth immersion if its differential is injective at each point (equivalently, $\text{rank } F = \dim M$).

- **Proposition 3.1** Suppose $F : M \rightarrow N$ is a smooth map and $p \in M$. If dF_p is **surjective**, then p has a neighborhood U such that $F|_U$ is a **submersion**. If dF_p is **injective**, then p has a neighborhood U such that $F|_U$ is an **immersion**.
- **Remark** F is a **surjective/injective** is different from F is a smooth **submersion/immersion**. The latter is the property of the differential map dF_p at each p not the property of the map itself. But F is **a smooth embedding** $\Rightarrow F$ is **an injective smooth immersion**. The converse is not true since F also need to *have continuous inverse* from its image to its domain.

3.2 Local Diffeomorphisms

- **Definition** If M and N are smooth manifolds with or without boundary, a map $F : M \rightarrow N$ is called a local diffeomorphism if every point $p \in M$ has a neighborhood U such that $F(U)$ is **open** in N and the restriction $F|_U : U \rightarrow F(U)$ is a **diffeomorphism**.
- The next theorem is the key to the most important properties of local diffeomorphisms.

Theorem 3.2 (Inverse Function Theorem for Manifolds). [Lee, 2003.]

Suppose M and N are smooth manifolds, and $F : M \rightarrow N$ is a **smooth map**. If $p \in M$ is a point such that dF_p is **invertible**, then there are **connected** neighborhoods U_0 of p and V_0 of $F(p)$ such that $F|_{U_0} : U_0 \rightarrow V_0$ is a **diffeomorphism**.

3.3 The Rank Theorem

- **Theorem 3.3 (Rank Theorem).** [Lee, 2003.]
Suppose M and N are smooth manifolds of dimensions m and n , respectively, and $F : M \rightarrow N$ is a smooth map **with constant rank r** . For each $p \in M$ there exist smooth charts (U, φ) for M centered at p and (V, ψ) for N centered at $F(p)$ such that $F(U) \subseteq V$, in which F has

a coordinate representation of the form

$$\widehat{F}(x^1, \dots, x^r, x^{r+1}, \dots, x^m) = (x^1, \dots, x^r, 0, \dots, 0). \quad (1)$$

In particular, if F is a smooth submersion, this becomes

$$\widehat{F}(x^1, \dots, x^n, x^{n+1}, \dots, x^m) = (x^1, \dots, x^n). \quad (2)$$

and if F is a smooth immersion, it is

$$\widehat{F}(x^1, \dots, x^m) = (x^1, \dots, x^m, 0, \dots, 0). \quad (3)$$

- **Corollary 3.4** Let M and N be smooth manifolds, let $F : M \rightarrow N$ be a smooth map, and suppose M is connected. Then the following are **equivalent**:

1. For each $p \in M$ there exist smooth charts containing p and $F(p)$ in which **the coordinate representation of F is linear**.
2. F has **constant rank**.

- The rank theorem is a purely **local statement**. However, it has the following powerful **global consequence**.

Theorem 3.5 (Global Rank Theorem). [Lee, 2003.]

Let M and N be smooth manifolds, and suppose $F : M \rightarrow N$ is a smooth map of **constant rank**.

1. If F is **surjective**, then it is a **smooth submersion**.
2. If F is **injective**, then it is a **smooth immersion**.
3. If F is **bijective**, then it is a **diffeomorphism**.

3.4 Embeddings

- One special kind of **immersion** is particularly important.

Definition If M and N are smooth manifolds with or without boundary, a smooth embedding of M into N is a smooth immersion $F : M \rightarrow N$ that is **also a topological embedding**, i.e., a **homeomorphism** onto its image $F(M) \subseteq N$ in the subspace topology.

- **Remark A smooth embedding** is a map that is **both a topological embedding** and a **smooth immersion**, not just a topological embedding that happens to be smooth. Also the map need to be **injective** and its inverse from $F(U)$ to U needs to be continuous.
- **Proposition 3.6** Suppose M and N are smooth manifolds with or without boundary, and $F : M \rightarrow N$ is an **injective smooth immersion**. If **any** of the following holds, then F is a **smooth embedding**.

1. F is an **open** or **closed** map. (i.e. it maps an open/closed set to an open/closed set)
2. F is a **proper map**. (i.e. the preimage of every compact set is compact)
3. M is **compact**.

4. M has empty boundary and $\dim M = \dim N$

• **Theorem 3.7 (Local Embedding Theorem).**

Suppose M and N are smooth manifolds with or without boundary, and $F : M \rightarrow N$ is a smooth map. Then F is a **smooth immersion** if and only if every point in M has a neighborhood $U \subseteq M$ such that $F|_U : U \rightarrow N$ is a **smooth embedding**.

3.5 Submersions

- **Definition** If $\pi : M \rightarrow N$ is any continuous map, a **section** of π is a **continuous right inverse** for π , i.e., a continuous map $\sigma : N \rightarrow M$ such that $\pi \circ \sigma = \text{Id}_N$:

$$\begin{array}{ccc} M & \xrightarrow{\pi} & N \\ & \searrow \sigma & \\ & & \end{array}$$

- **Definition** A **local section** of π is a continuous map $\sigma : U \rightarrow M$ defined on some open subset $U \subseteq N$ and satisfying the analogous relation $\pi \circ \sigma = \text{Id}_U$
- **Theorem 3.8 (Local Section Theorem).** [Lee, 2003.]
Suppose M and N are smooth manifolds and $\pi : M \rightarrow N$ is a smooth map. Then π is a **smooth submersion** if and only if every point of M is in the **image** of a **smooth local section** of π .
- **Proposition 3.9 (Properties of Smooth Submersions).**
Let M and N be smooth manifolds, and suppose $\pi : M \rightarrow N$ is a smooth submersion. Then π is **an open map**, and if it is **surjective** it is a **quotient map**.
- The next three theorems provide important tools that we will use frequently when studying submersions.

Theorem 3.10 (Characteristic Property of Surjective Smooth Submersions).

Suppose M and N are smooth manifolds, and $\pi : M \rightarrow N$ is a **surjective smooth submersion**. For any smooth manifold P with or without boundary, a map $F : N \rightarrow P$ is **smooth** if and only if $F \circ \pi$ is **smooth**:

$$\begin{array}{ccc} M & & \\ \pi \downarrow & \searrow F \circ \pi & \\ N & \xrightarrow{F} & P. \end{array}$$

Theorem 3.11 (Passing Smoothly to the Quotient).

Suppose M and N are smooth manifolds and $\pi : M \rightarrow N$ is a **surjective smooth submersion**. If P is a smooth manifold with or without boundary and $F : M \rightarrow P$ is a smooth map that is **constant on the fibers of π** , then there exists a **unique smooth map $\tilde{F} : N \rightarrow P$** such that $\tilde{F} \circ \pi = F$:

$$\begin{array}{ccc} M & & \\ \pi \downarrow & \searrow F & \\ N & \xrightarrow{\tilde{F}} & P. \end{array}$$

Theorem 3.12 (Uniqueness of Smooth Quotients).

Suppose that M, N_1 , and N_2 are smooth manifolds, and $\pi_1 : M \rightarrow N_1$ and $\pi_2 : M \rightarrow N_2$ are

surjective smooth submersions that are constant on each other's fibers. Then there exists a unique diffeomorphism $F : N_1 \rightarrow N_2$ such that $F \circ \pi_1 = \pi_2$:

$$\begin{array}{ccc} & M & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ N_1 & \xrightarrow{\quad F \quad} & N_2. \end{array}$$

4 Submanifolds

- **Remark** A great number of manifolds of interest can be considered as a submanifold of some other (simpler) manifolds. This chapter mainly concerns about *the embedded submanifolds* and then extends to *immersed submanifolds*.
- **Remark** The definition of *embedded submanifold* S of M consists of three parts:
 1. $S \subseteq M$ has *the subspace topology* (inherited from the topology of the ambient manifold M);
 2. S is endowed with *a smooth structure*;
 3. Under this smooth structure, *the inclusion map* $S \hookrightarrow M$ is a *smooth embedding*, i.e. it has injective differential everywhere and is *locally homemorphic* to its image.
- **Remark** Besides the definition, there are two other ways to identify embedded submanifold:
 1. **Local Slice Criterion:** If a subset S under local representation of M is *homemorphic* to an open subset of $\mathbb{R}^k \subseteq \mathbb{R}^n$. *The Local Slice Criterion* relies on *the topology and the smooth structure of the ambient manifold* M to determine both the topology and the smooth structure of submanifold S .
 2. **Level Set Criterion:** If every point of subset S has a neighborhood in M so that $S \cap U$ is a level set of some *smooth submersion*, then this set is a k -dimensional embedded submanifold. In particular, *every embedded submanifold admits a local defining function in a neighborhood of each of its points*. This result provides *a constructive way* to build embedded submanifold by defining map.
- **Remark** An important result of *smooth map with constant rank* is that each of its *level set* is a *properly embedded submanifold* with *codimension* equal to the *rank* of smooth map. In particular,
 1. If the map is a *submersion*, then the level set is *a regular level set*, which is a properly embedded submanifold. Moreover, the *dimension* of the submanifold is determined by the difference of dimension between domain and codomain.
 2. The *image of immersion* is also a submanifold, *the immersed submanifold*. An immersed submanifold may not be an embedded submanifold. But *locally* it is *an embedded submanifold*.
- **Remark** The final important remark on embedded submanifold is that it has a *unique topology and smooth structure*, which is *the subspace topology* and the coordinate map that makes the $S \cap U$ is a *k-slice* of U .

4.1 Embedded Submanifolds

4.1.1 Definitions

- **Definition** Suppose M is a smooth manifold with or without boundary. An **embedded submanifold** of M is a subset $S \subseteq M$ that is a *manifold* (without boundary) in the **subspace topology**, endowed with a **smooth structure** with respect to which the **inclusion map** $S \hookrightarrow M$ is a **smooth embedding**. Embedded submanifolds are also called **regular submanifolds**.
- **Definition** If S is an *embedded submanifold* of M , the **difference** $\dim M - \dim S$ is called **the codimension** of S in M , and the **containing manifold** M is called **the ambient manifold** for S .

An embedded **hypersurface** is an embedded submanifold of codimension 1. The *empty set* is an embedded submanifold of *any dimension*.

- **Proposition 4.1 (Open Submanifolds).** [Lee, 2003.]
Suppose M is a smooth manifold. The embedded submanifolds of **codimension 0** in M are exactly the **open submanifolds**.
- There are several other ways to create submanifolds:

Proposition 4.2 (Images of Embeddings as Submanifolds). [Lee, 2003.]

Suppose M is a smooth manifold with or without boundary, N is a smooth manifold, and $F : N \rightarrow M$ is a **smooth embedding**. Let $S = F(N)$. With the subspace topology, S is a topological manifold, and it has a **unique smooth structure** making it into an **embedded submanifold** of M with the property that F is a **diffeomorphism** onto its image.

- **Proposition 4.3 (Slices of Product Manifolds).** [Lee, 2003.]
Suppose M and N are smooth manifolds. For each $p \in N$, the subset $M \times \{p\}$ (called a **slice of the product manifold**) is an **embedded submanifold** of $M \times N$ diffeomorphic to M .
- **Proposition 4.4 (Graphs as Submanifolds).** [Lee, 2003.]
Suppose M is a smooth m -manifold (without boundary), N is a smooth n -manifold with or without boundary, $U \subseteq M$ is open, and $f : U \rightarrow N$ is a smooth map. Let $\Gamma(f) \subseteq M \times N$ denote **the graph of f** :

$$\Gamma(f) = \{(x, y) \in M \times N : x \in U, y = f(x)\}.$$

Then $\Gamma(f)$ is an **embedded m -dimensional** submanifold of $M \times N$

- **Definition** An embedded submanifold $S \subseteq M$ is said to be **properly embedded** if the inclusion $S \hookrightarrow M$ is a **proper map**.
- **Proposition 4.5** Suppose M is a smooth manifold with or without boundary and $S \subseteq M$ is an embedded submanifold. Then S is **properly embedded** if and only if it is a **closed** subset of M .
- **Corollary 4.6** Every **compact** embedded submanifold is **properly embedded**.
- **Proposition 4.7 (Global Graphs Are Properly Embedded).** [Lee, 2003.]
Suppose M is a smooth manifold, N is a smooth manifold with or without boundary, and $f : M \rightarrow N$ is a **smooth map**. With the smooth manifold structure as above, the graph of f $\Gamma(f)$ is **properly embedded** in $M \times N$.

4.1.2 Slice Charts for Embedded Submanifolds

- **Definition** if U is an open subset of \mathbb{R}^n and $k \in \{0, \dots, n\}$, a ***k-dimensional slice*** of U (or simply a *k-slice*) is any subset of the form

$$S = \left\{ (x^1, \dots, x^k, x^{k+1}, \dots, x^n) \in U : x^{k+1} = c^{k+1}, \dots, x^n = c^n \right\}$$

for some constants c^{k+1}, \dots, c^n . (When $k = n$, this just means $S = U$.) Clearly, ***every k-slice is homeomorphic to an open subset of \mathbb{R}^k*** .

- **Definition** Let M be a smooth n -manifold, and let (U, φ) be a smooth chart on M . If S is a subset of U such that $\varphi(S)$ is a k -slice of $\varphi(U)$, then we say that S ***is a k-slice of U*** .
- **Definition** Given a subset $S \subseteq M$ and a nonnegative integer k , we say that S ***satisfies the local k-slice condition*** if *each point* of S is contained in the domain of a smooth chart (U, φ) for M such that $S \cap U$ is a ***single k-slice in U*** . Any such chart is called ***a slice chart for S in M*** , and the corresponding coordinates (x^1, \dots, x^n) are called ***slice coordinates***.
- **Remark** The key to understand the ***the local k-slice condition*** for $S \subseteq M$:
 1. It is a condition on the ***subset S*** only; it does ***not presuppose*** any particular ***topology*** or ***smooth structure*** on S . All it needs is the topology and smooth structure from the ambient manifold M .
 2. The *local neighborhood* $U \subseteq M$ is a ***neighborhood of p in the ambient manifold M*** not a neighborhood in S (since we do not define such topology);
 3. The k -slice representation is for the ***intersection $S \cap U$*** under ***the smooth chart (U, φ)*** of ***the ambient manifold M*** .
- **Theorem 4.8 (Local Slice Criterion for Embedded Submanifolds)** [Lee, 2003.].
Let M be a smooth n -manifold. If $S \subseteq M$ is an embedded k -dimensional submanifold, then S satisfies the local k -slice condition. **Conversely**, if $S \subseteq M$ is a subset that ***satisfies the local k-slice condition***, then with the subspace topology, S is a topological manifold of dimension k , and it ***has a smooth structure*** making it into a k -dimensional embedded submanifold of M .
- **Theorem 4.9** If M is a smooth n -manifold with boundary, then with the subspace topology, ∂M is a topological $(n - 1)$ -dimensional manifold (without boundary), and has a smooth structure such that it is a properly ***embedded submanifold*** of M .

4.1.3 Level Sets

- **Remark** In practice, embedded submanifolds are most often presented as ***solution sets*** of equations or systems of equations.
- **Definition** If $\Phi : M \rightarrow N$ is any map and c is any point of N , we call the set $\Phi^{-1}(c)$ ***a level set of Φ*** (Fig. ??). (In the special case $N = \mathbb{R}^k$ and $c = 0$, the level set $\Phi^{-1}(0)$ is usually called ***the zero set of Φ*** .)
- **Remark** It is easy to find *level sets of smooth functions* that are *not smooth submanifolds*.

$$\Theta(x, y) = x^2 - y, \quad \Phi(x, y) = x^2 - y^2, \quad \Psi(x, y) = x^2 - y^3.$$

(Note that the zero set $\Theta^{-1}(0)$ is an embedded submanifold in \mathbb{R}^2 but not for others.) In fact, **every closed subset of M** can be expressed as **the zero set** of some smooth real-valued function.

- **Theorem 4.10 (Constant-Rank Level Set Theorem).** [Lee, 2003.]

Let M and N be smooth manifolds, and let $\Phi : M \rightarrow N$ be a smooth map **with constant rank r** . **Each level set of Φ is a properly embedded submanifold of codimension r in M .**

- **Corollary 4.11 (Submersion Level Set Theorem).** [Lee, 2003.]

If M and N are smooth manifolds and $\Phi : M \rightarrow N$ is a **smooth submersion**, then each level set of Φ is a **properly embedded** submanifold whose **codimension** is equal to the **dimension of N** .

- **Remark** This result should be compared to the corresponding result in linear algebra: if $L : \mathbb{R}^m \rightarrow \mathbb{R}^r$ is a surjective linear map, then the kernel of L is a linear subspace of codimension r by **the rank-nullity law**. The vector equation $Lx = 0$ is equivalent to r linearly independent scalar equations, each of which can be thought of as cutting down one of the degrees of freedom in \mathbb{R}^m , leaving a subspace of codimension r .

In the context of smooth manifolds, the analogue of a *surjective linear map* is a **smooth submersion**, each of whose **(local) component functions cuts down the dimension by one**.

- **Definition** If $\Phi : M \rightarrow N$ is a smooth map, a point $p \in M$ is said to be **a regular point** of Φ if $d\Phi_p : T_p M \rightarrow T_{\Phi(p)} N$ is **surjective**; it is **a critical point** of Φ otherwise.

This means, in particular, that **every point** of M is **critical** if $\dim M < \dim N$, and every point is **regular** if and only if Φ is a **submersion**.

- **Definition** A point $c \in N$ is said to be **a regular value** of Φ if **every point of the level set $\Phi^{-1}(c)$ is a regular point**, and **a critical value** otherwise. In particular, if $\Phi^{-1}(c) = \emptyset$, then c is a **regular value**. Finally, a level set $\Phi^{-1}(c)$ is called **a regular level set** if c is a regular value of Φ ; in other words, a regular level set is a level set consisting **entirely of regular points** of Φ (points p such that $d\Phi_p$ is surjective).

- **Remark** Every properly embedded submanifold $M = \Phi^{-1}(c)$ is a regular level set. The following theorem shows that the converse is true as well.

- **Theorem 4.12 (Regular Level Set Theorem).** [Lee, 2003.]

Every regular level set of a smooth map between smooth manifolds is a properly embedded submanifold whose codimension is equal to the dimension of the codomain.

- **Proposition 4.13 (Local Level Set Criterion for Smooth Embedded Submanifolds)**
Let S be a subset of a smooth m -manifold M . Then S is an **embedded k -submanifold** of M if and only if every point of S has a neighborhood U in M such that $U \cap S$ is a **level set** of a **smooth submersion** $\Phi : U \rightarrow \mathbb{R}^{m-k}$.

- **Proposition 4.14 (Local Level Set Criterion for Smooth Embedded Submanifolds)**
Let S be a subset of a smooth m -manifold M . Then S is an **embedded k -submanifold** of M if and only if every point of S has a neighborhood U in M such that $U \cap S$ is a **level set** of a **smooth submersion** $\Phi : U \rightarrow \mathbb{R}^{m-k}$.

- **Definition** If $S \subseteq M$ is an embedded submanifold, a smooth map $\Phi : M \rightarrow N$ such that S is

a **regular level set** of Φ is called **a defining map for S** . In the special case $N = \mathbb{R}^{m-k}$ (so that Φ is a real-valued or vector-valued function), it is usually called **a defining function**.

More generally, if U is an open subset of M and $\Phi : U \rightarrow N$ is a smooth map such that $S \cap U$ is a regular level set of Φ , then Φ is called **a local defining map (or local defining function) for S** .

4.2 Immersed Submanifolds

4.2.1 Definitions and Examples

- **Definition** Let M be a smooth manifold with or without boundary. An **immersed submanifold** of M is a subset $S \subseteq M$ endowed with a *topology* (not necessarily *the subspace topology*) with respect to which it is **a topological manifold** (without boundary), and a *smooth structure* with respect to which *the inclusion map* $S \hookrightarrow M$ is **a smooth immersion**.

As for embedded submanifolds, we define the **codimension** of S in M to be $\dim M - \dim S$.

- **Remark** Immersed submanifolds *do not require the submanifold topology to be the subspace topology* which is more general than embedded submanifold.
- The immersed submanifolds arise in natural way:

Proposition 4.15 (Images of Immersions as Submanifolds). [Lee, 2003.]

Suppose M is a smooth manifold with or without boundary, N is a smooth manifold, and $F : N \rightarrow M$ is an **injective smooth immersion**. Let $S = F(N)$. Then S has a unique topology and smooth structure such that it is a **smooth submanifold** of M and such that $F : N \rightarrow S$ is a **diffeomorphism** onto its image.

- **Example (Immersed Submanifold but Not an Embedded Submanifold)**
Both examples of *The Figure-Eight* and *the Dense Curve on the Torus* are *images of injective smooth immersions*, they are **immersed submanifolds** when given appropriate topologies and smooth structures. As smooth manifolds, they are **diffeomorphic** to \mathbb{R} . *They are not embedded submanifolds*, because *neither* one has *the subspace topology*. In fact, their image sets cannot be made into embedded submanifolds even if we are allowed to change their topologies and smooth structures. ■

- **Remark** Suppose M is a smooth manifold and $S \subseteq M$ is **an immersed submanifold**. It can be shown that every subset of S that is **open** in the *subspace topology* is also **open** in its given *submanifold topology*; and the **converse** is true if and only if S is **embedded**.
- **Proposition 4.16 (Immersed Submanifolds Are Locally Embedded).** [Lee, 2003.]
If M is a smooth manifold with or without boundary, and $S \subseteq M$ is an **immersed submanifold**, then for each $p \in S$ there exists a neighborhood U of p **in S** that is an **embedded submanifold** of M .

Note that a smooth immersion is locally a smooth embedding.

- **Remark** It is important to be clear about what this proposition does and does not say: given an immersed submanifold $S \subseteq M$ and a point $p \in S$, it is possible to find *a neighborhood U*

of p in S such that U is *embedded*; but it may not be possible to find a *neighborhood* V of p in M such that $V \cap S$ is embedded.

- **Definition** Suppose $S \subseteq M$ is an immersed k -dimensional submanifold. A **local parametrization** of S is a continuous map $X : U \rightarrow M$ whose domain is an **open subset** $U \subseteq \mathbb{R}^k$, whose image is an **open subset** of S , and which, considered as a map into S , is a **homeomorphism onto its image**. It is called a **smooth local parametrization** if it is a **diffeomorphism** onto its image (with respect to S 's smooth manifold structure). If the image of X is all of S , it is called a **global parametrization**.
- **Remark** For a smooth chart (U, φ) of M , $\varphi : U \rightarrow \widehat{U} \subseteq \mathbb{R}^n$ is a diffeomorphism, its inverse $\varphi^{-1} : \widehat{U} \rightarrow U \subseteq M$ is a **smooth local parameterization** (in fact $X = \text{Id}_M \circ \varphi^{-1}$).
- **Proposition 4.17** Suppose M is a smooth manifold with or without boundary, $S \subseteq M$ is an immersed k -submanifold, $\iota : S \hookrightarrow M$ is the inclusion map, and U is an open subset of \mathbb{R}^k . A map $X : U \rightarrow M$ is a **smooth local parametrization** of S **if and only if** there is a smooth coordinate chart (V, φ) for S such that $X = \iota \circ \varphi^{-1}$. Therefore, every point of S is in the image of some local parametrization.

4.3 Restricting Maps to Submanifolds

4.3.1 Restricting Domains and Codomains

- **Remark** Given a smooth map $F : M \rightarrow N$, it is important to know whether F is still smooth when its **domain** or **codomain** is restricted to a submanifold. See Fig. ??.

1. **Restricting Domains:** The answer is **yes**.

Theorem 4.18 (Restricting the Domain of a Smooth Map). [Lee, 2003.]

If M and N are smooth manifolds with or without boundary, $F : M \rightarrow N$ is a smooth map, and $S \subseteq M$ is an **immersed or embedded submanifold**, then $F|_S : S \rightarrow N$ is smooth.

2. **Restricting Codomains:** We provides **sufficient conditions**: the **image** of the smooth map should be **contained in the submanifold**.

- **Immersed Submanifolds:**

Theorem 4.19 (Restricting the Codomain of a Smooth Map). [Lee, 2003.]

Suppose M is a smooth manifold (without boundary), $S \subseteq M$ is an immersed submanifold, and $F : N \rightarrow M$ is a smooth map whose **image is contained in** S . If F is **continuous** as a map from N to S , then $F : N \rightarrow S$ is smooth.

- **Embedded Submanifolds:**

Corollary 4.20 (Embedded Case).

Let M be a smooth manifold and $S \subseteq M$ be an embedded submanifold. Then every smooth map $F : N \rightarrow M$ whose **image is contained in** S is also **smooth** as a map from N to S .

3. We can generalize the corollary above as the definition of weakly embedded submanifold.

Definition If M is a smooth manifold and $S \subseteq M$ is an immersed submanifold, then S is said to be **weakly embedded** in M if every smooth map $F : N \rightarrow M$ **whose image lies in S** is **smooth** as a map from N to S . (*Weakly embedded submanifolds* are called *initial submanifolds* by some authors.)

4.3.2 Uniqueness of Smooth Structures on Submanifolds

- **Theorem 4.21** *Suppose M is a smooth manifold and $S \subseteq M$ is an **embedded submanifold**. The subspace topology on S and the smooth structure from the local k -slice condition are **the only topology and smooth structure** with respect to which S is an embedded or immersed submanifold.*
- **Remark** Thanks to this uniqueness result, we now know that a subset $S \subseteq M$ is an *embedded submanifold* **if and only if** it satisfies the local slice condition, and if so, its topology and smooth structure are **uniquely determined**.

Because the local slice condition is **a local condition**, if every point $p \in S$ has a neighborhood $U \subseteq M$ such that $U \cap S$ is an embedded k -submanifold of U , then S is an embedded k -submanifold of M .

- **Theorem 4.22** *Suppose M is a smooth manifold and $S \subseteq M$ is an **immersed submanifold**. For the **given topology** on S , there is **only one smooth structure** making S into an immersed submanifold.*
- **Theorem 4.23** *If M is a smooth manifold and $S \subseteq M$ is a **weakly embedded submanifold**, then S has **only one topology and smooth structure** with respect to which it is an immersed submanifold.*

4.3.3 Extending Functions from Submanifolds

- **Remark** Complementary to the restriction problem is the problem of extending smooth functions from a submanifold to the ambient manifold. Here we say $f \in C^\infty(S)$ for submanifold $S \subseteq M$, when f is considered as a function on the manifold S .
- **Lemma 4.24** (*Extension Lemma for Functions on Submanifolds*).
Suppose M is a smooth manifold, $S \subseteq M$ is a smooth submanifold, and $f \in C^\infty(S)$.
 1. If S is **embedded**, then there exist a **neighborhood** U of S in M and a smooth function $\tilde{f} \in C^\infty(U)$ such that $\tilde{f}|_S = f$.
 2. If S is **properly embedded**, then the neighborhood U above can be taken to be **all** of M .

4.4 The Tangent Space to a Submanifold

- **Remark** The *tangent space to a smooth submanifold* of an abstract smooth manifold can be viewed as a subspace of *the tangent space to the ambient manifold*, once we make appropriate identifications. The following proof is based on the differential of *the inclusion map* as a smooth immersion.

Proof: Let M be a smooth manifold with or without boundary, and let $S \subseteq M$ be an immersed or embedded submanifold. Since the inclusion map $\iota : S \hookrightarrow M$ is a **smooth immersion**, at each point $p \in S$ we have an *injective linear map* $d\iota_p : T_p S \rightarrow T_p M$. In terms of **derivations**, this injection works in the following way: for any vector $v \in T_p S$, the image vector $\tilde{v} = d\iota_p(v) \in T_p M$ acts on smooth functions on M by

$$\tilde{v}f = d\iota_p(v)f = v(f \circ \iota) = v(f|_S).$$

We adopt the convention of **identifying** $T_p S$ with **its image** under this map, thereby thinking of $T_p S$ as a certain linear subspace of $T_p M$. This identification makes sense regardless of whether S is *embedded* or *immersed*. ■

- There are several *alternative* ways to *characterize* the tangent space of a submanifold

1. **Smooth curve on submanifold.**

Proposition 4.25 Suppose M is a smooth manifold with or without boundary, $S \subseteq M$ is an immersed or embedded submanifold, and $p \in S$. A vector $v \in T_p M$ is in $T_p S$ if and only if there is a smooth curve $\gamma : J \rightarrow M$ whose **image is contained in** S , and which is also **smooth** as a map into S , such that $0 \in J$, $\gamma(0) = p$, and $\gamma'(0) = v$.

2. **Derivations on functions whose restriction on submanifold are constant zero.**

Proposition 4.26 Suppose M is a smooth manifold, $S \subseteq M$ is an embedded submanifold, and $p \in S$. As a subspace of $T_p M$, the tangent space $T_p S$ is characterized by

$$T_p S = \{v \in T_p M : vf = 0 \text{ whenever } f \in C^\infty(M) \text{ and } f|_S = 0\}.$$

3. **Kernel subspace of differential of local defining map.**

Proposition 4.27 Suppose M is a smooth manifold and $S \subseteq M$ is an embedded submanifold. If $\Phi : U \rightarrow N$ is any **local defining map** for S , then $T_p S = \mathbf{Ker}(d\Phi_p) : T_p M \rightarrow T_{\Phi(p)} N$ for each $p \in S \cap U$.

Note that $S \cap U = (\Phi \circ \iota)^{-1}(c)$ is the level set of $\Phi \circ \iota$ thus it is constant for $\Phi \circ \iota$. So $d\Phi_p \circ d\iota_p = 0$.

Corollary 4.28 Suppose $S \subseteq M$ is a **level set** of a **smooth submersion** $\Phi = (\Phi^1, \dots, \Phi^k) : M \rightarrow \mathbb{R}^k$. A vector $v \in T_p M$ is tangent to S if and only if $v\Phi^1 = \dots = v\Phi^k = 0$.

- **Remark** If M is a smooth manifold **with boundary** and $p \in \partial M$, the vectors in $T_p M$ can be separated into **three classes**:

1. those **tangent to the boundary**;
2. those pointing **inward**;
3. those pointing **outward**.

Definition If $p \in \partial M$, a vector $v \in T_p M \setminus T_p \partial M$ is said to be **inward-pointing** if for some $\epsilon > 0$ there exists a smooth curve $\gamma : [0, \epsilon) \rightarrow M$ such that $\gamma(0) = p$ and $\gamma'(0) = v$, and it is **outward-pointing** if there exists such a curve whose domain is $(-\epsilon, 0]$.

Proposition 4.29 (*Characterization of Tangent Vectors on Boundary using Component Functions*)

Suppose M is a smooth n -dimensional manifold with boundary, $p \in \partial M$, and (x^i) are any smooth boundary coordinates defined on a neighborhood of p . The **inward-pointing vectors** in $T_p M$ are precisely those with **positive x^n -component**, the **outward-pointing** ones are those with **negative x^n -component**, and the ones **tangent to ∂M** are those with **zero x^n -component**. Thus, $T_p M$ is the **disjoint union** of $T_p \partial M$, the set of inward-pointing vectors, and the set of outward-pointing vectors, and $v \in T_p M$ is inward-pointing if and only if $-v$ is outward-pointing.

4.5 Vector Fields and Submanifolds

- **Remark** If $S \subseteq M$ is an immersed or embedded submanifold (with or without boundary), a vector field X on M does **not necessarily** restrict to a vector field on S , because X_p may not lie in the subspace $T_p S \subseteq T_p M$ at a point $p \in S$.
- **Definition** Given a point $p \in S$, a vector field X on M is said to **be tangent to S** at p if $X_p \in T_p S \subseteq T_p M$. It is **tangent to S** if it is tangent to S at every point of S .
- **Proposition 4.30** Let M be a smooth manifold, $S \subseteq M$ be an **embedded submanifold** with or without boundary, and X be a smooth vector field on M . Then X is **tangent to S** if and only if $(Xf)|_S = 0$ for **every $f \in C^\infty(M)$ such that $f|_S \equiv 0$** .
- **Remark** Suppose $S \subseteq M$ is an **immersed submanifold** with or without boundary, and Y is a smooth vector field on M . If there is a vector field $X \in \mathfrak{X}(S)$ that is **ι -related to Y** , where $\iota : S \hookrightarrow M$ is the inclusion map, then clearly Y is **tangent to S** , because $Y_p = d\iota_p(X_p)$ is in the image of $d\iota_p$ for each $p \in S$.

The converse is true as well.

Proposition 4.31 (Restricting Vector Fields to Submanifolds). [Lee, 2003.]

Let M be a smooth manifold, let $S \subseteq M$ be an **immersed submanifold** with or without boundary, and let $\iota : S \hookrightarrow M$ denote the inclusion map. If $Y \in \mathfrak{X}(M)$ is **tangent to S** , then there is a **unique smooth vector field** on S , denoted by $Y|_S$, that is **ι -related to Y** .

4.6 Restricting Covector Fields to Submanifolds

- **Remark** Compare to restricting vector fields to submanifolds, the restriction of covector fields to submanifolds is much simpler.
- **Remark (The Pullback of Covector Field by the Inclusion Map is a Covector Field on Submanifold)**
Suppose M is a smooth manifold with or without boundary, $S \subseteq M$ is an **immersed submanifold** with or without boundary, and $\iota : S \hookrightarrow M$ is the inclusion map. If ω is any smooth covector field on M , **the pullback by ι yields a smooth covector field $\iota^*\omega$ on S** .

To see what this means, let $v \in T_p S$ be arbitrary, and compute

$$(\iota^*\omega)_p(v) = \omega_p(d\iota_p(v)) = \omega_p(v).$$

since $d\iota_p : T_p S \rightarrow T_p M$ is just the inclusion map, under our usual identification of $T_p S$ with a subspace of $T_p M$. Thus, $\iota^*\omega$ is just the restriction of ω to vectors tangent to S . For this reason, $\iota^*\omega$ is often called **the restriction of ω to S** .

Be warned, however, that $\iota^*\omega$ might equal **zero** at a given point of S , even though **considered as a covector field on M , ω might not vanish there**.

- **Example** ($\omega \neq 0$ but $\iota^*\omega = 0$)

Let $\omega = dy$ on \mathbb{R}^2 , and let S be the x -axis, considered as an embedded submanifold of \mathbb{R}^2 . As a covector field on \mathbb{R}^2 , ω is **nonzero** everywhere, because one of its component functions is **always** 1. However, the restriction $\iota^*\omega$ is **identically zero**, because y vanishes identically on S :

$$\iota^*\omega = \iota^*dy = d(y \circ \iota) = 0.$$

- **Remark** One usually says that “ ω **vanishes along S** ” or “ ω **vanishes at points of S** ” if $\omega_p = 0$ for every point $p \in S$.

The **weaker condition** that $\iota^*\omega = 0$ is expressed by saying that “the restriction of ω to S vanishes”, or “the pullback of ω to S vanishes”.

References

John Marshall Lee. *Introduction to smooth manifolds*. Graduate texts in mathematics. Springer, New York, Berlin, Heidelberg, 2003. ISBN 0-387-95448-1.