

Summary Part 2: Concentration of Measure and Functional Methods

Tianpei Xie

Jan. 26th., 2023

Contents

1	Logarithmic Sobolev Inequality	2
1.1	Functional Form of Logarithmic Sobolev Inequality	2
1.2	Bernoulli Logarithmic Sobolev Inequality	3
1.3	Gaussian Logarithmic Sobolev Inequality	4
1.4	Modified Logarithmic Sobolev Inequalities	4
2	Isoperimetric Inequalities and Concentration of Measure	5
2.1	Brunn-Minkowski Inequality	5
2.2	Classical Isoperimetric Problem on Euclidean Space \mathbb{R}^n	6
2.3	Isoperimetric Problem on Unit Sphere	7
2.4	Concentration via Isoperimetric Inequalities	8
2.5	Convex Distance Inequality	10
2.6	Concentration of Convex Lipschitz Functions	10
3	Concentration of Gaussian Measure	11
3.1	Gaussian Isoperimetric Theorem and Gaussian Concentration Theorem	11
3.2	Lipschitz Functions of Gaussian Variables	13
3.3	Gaussian Logarithmic Sobolev Inequality	14
3.4	Gaussian Transportation Inequality	14
3.5	Gaussian Hypercontractivity	15
3.6	Suprema of Gaussian Process	15
4	Concentration of Bernoulli Measure on the Binary Hypercube	15
4.1	Edge Isoperimetric Inequality on the Binary Hypercube	15
4.2	Bobkov's Inequality	17
4.3	Vertex Isoperimetric Inequality on the Binary Hypercube	17
4.4	Hypercontractivity: The Bonami-Beckner Inequality	19
4.5	Influence Function	19
4.6	Monotone Sets	19
4.7	Threshold Phenomena	19

1 Logarithmic Sobolev Inequality

1.1 Functional Form of Logarithmic Sobolev Inequality

- From functional analysis, we have *the Sobolev inequality*,

Remark (*The Sobolev Inequality*) [Evans, 2010]

The Sobolev inequality states for smooth function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ in *Sobolev space* where $n \geq 3$ and $p = \frac{2n}{n-2} > 2$

$$\|f\|_p^2 \leq C_n \int_{\mathbb{R}^n} |\nabla f|^2 dx.$$

The inequality is sharp when the constant

$$C_n := \frac{1}{\pi n(n-2)} \left(\frac{\Gamma(n)}{\Gamma(n/2)} \right)^{2/n}$$

- **Proposition 1.1 (*Euclidean Logarithmic Sobolev Inequality*).**

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function and m be Lebesgue measure on \mathbb{R}^n , then

$$\begin{aligned} \text{Ent}_m(f^2) &\leq \frac{n}{2} \log \left(\frac{2}{n\pi e} \mathbb{E}_m \left[\|\nabla f\|_2^2 \right] \right) \\ \Leftrightarrow \int f^2 \log \left(\frac{f^2}{\int f^2 dx} \right) dx &\leq \frac{n}{2} \log \left(\frac{2}{n\pi e} \int |\nabla f|^2 dx \right) \end{aligned} \quad (1)$$

- **Definition (*Logarithmic Sobolev Inequality for General Probability Measure*).**

A probability measure μ on \mathbb{R}^n is said to satisfy the logarithmic Sobolev inequality for some constant $C > 0$ if for any smooth function f

$$\text{Ent}_\mu(f^2) \leq C \mathbb{E}_\mu \left[\|\nabla f\|_2^2 \right] \quad (2)$$

holds for any **continuous differentiable** function $f : \mathbb{R}^n \rightarrow \mathbb{R}$. The left-hand side is called **the entropy functional**, which is defined as

$$\begin{aligned} \text{Ent}(f^2) &:= \mathbb{E}_\mu [f^2 \log f^2] - \mathbb{E}_\mu [f^2] \log \mathbb{E}_\mu [f^2] \\ &= \int f^2 \log \left(\frac{f^2}{\int f^2 d\mu} \right) d\mu. \end{aligned}$$

The right-hand side is defined as

$$\mathbb{E}_\mu \left[\|\nabla f\|_2^2 \right] = \int \|\nabla f\|_2^2 d\mu.$$

Thus we can rewrite *the logarithmic Sobolev inequality* in *functional form*

$$\int f^2 \log \left(\frac{f^2}{\int f^2 d\mu} \right) d\mu \leq C \int \|\nabla f\|_2^2 d\mu \quad (3)$$

- **Remark (*Logarithmic Sobolev Inequality*)**

For non-negative function f , we can replace $f \rightarrow \sqrt{f}$, so that the *logarithmic Sobolev inequality* becomes

$$\text{Ent}_\mu(f) \leq C \int \frac{\|\nabla f\|_2^2}{f} d\mu \quad (4)$$

- **Remark (*Modified Logarithmic Sobolev Inequality via Convex Cost and Duality*)**

For some **convex non-negative cost** $c : \mathbb{R}^n \rightarrow \mathbb{R}_+$, the **convex conjugate** of c (Legendre transform of c) is defined as

$$c^*(x) := \sup_y \{ \langle x, y \rangle - c(y) \}$$

Then we can obtain the *modified logarithmic Sobolev inequality*

$$\text{Ent}_\mu(f) \leq \int f^2 c^* \left(\frac{\nabla f}{f} \right) d\mu \quad (5)$$

1.2 Bernoulli Logarithmic Sobolev Inequality

- **Remark (*Setting*)**

Consider a **uniformly distributed binary vector** $Z = (Z_1, \dots, Z_n)$ on the hypercube $\{-1, +1\}^n$. In other words, the components of X are *independent, identically distributed random sign (Rademacher) variables* with $\mathbb{P}\{Z_i = -1\} = \mathbb{P}\{Z_i = +1\} = 1/2$ (i.e. *symmetric Bernoulli random variables*).

Let $f : \{-1, +1\}^n \rightarrow \mathbb{R}$ be a real-valued function on **binary hypercube**. $X := f(Z)$ is an induced real-valued random variable. Define $\tilde{Z}^{(i)} = (Z_1, \dots, Z_{i-1}, Z'_i, Z_{i+1}, \dots, Z_n)$ be the sample Z with i -th component replaced by an *independent copy* Z'_i . Since $Z, \tilde{Z}^{(i)} \in \{-1, +1\}^n$, $\tilde{Z}^{(i)} = (Z_1, \dots, Z_{i-1}, -Z_i, Z_{i+1}, \dots, Z_n)$, i.e. the i -th sign is **flipped**. Also denote the i -th Jackknife sample as $Z_{(i)} = (Z_1, \dots, Z_{i-1}, Z_{i+1}, \dots, Z_n)$ by *leaving out* the i -th component. $\mathbb{E}_{(-i)}[X] := \mathbb{E}[X|Z_{(i)}]$.

Denote the i -th component of **discrete gradient** of f as

$$\nabla_i f(z) := \frac{1}{2} \left(f(z) - f(\tilde{z}^{(i)}) \right)$$

and $\nabla f(z) = (\nabla_1 f(z), \dots, \nabla_n f(z))$

- **Proposition 1.2 (*Logarithmic Sobolev Inequality for Rademacher Random Variables*)**. [Boucheron et al., 2013]

If $f : \{-1, +1\}^n \rightarrow \mathbb{R}$ be an arbitrary real-valued function defined on the n -dimensional **binary hypercube** and assume that Z is **uniformly distributed** over $\{-1, +1\}^n$. Then

$$\text{Ent}(f^2) \leq \mathcal{E}(f) \quad (6)$$

$$\Leftrightarrow \text{Ent}(f^2(Z)) \leq 2\mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right] \quad (7)$$

- **Remark (*Logarithmic Sobolev Inequality \Rightarrow Efron-Stein Inequality*)**. [Boucheron et al., 2013]

Note that for f non-negative,

$$\text{Var}(f(Z)) \leq \text{Ent}(f^2(Z)).$$

Thus *logarithmic Sobolev inequality* (6) implies

$$\text{Var}(f(Z)) \leq \mathcal{E}(f)$$

which is the *Efron-Stein inequality*.

- **Corollary 1.3** (*Logarithmic Sobolev Inequality for Asymmetric Bernoulli Random Variables*). [Boucheron et al., 2013]

If $f : \{-1, +1\}^n \rightarrow \mathbb{R}$ be an arbitrary real-valued function and $Z = (Z_1, \dots, Z_n) \in \{-1, +1\}^n$ with $p = \mathbb{P}\{Z_i = +1\}$. Then

$$\text{Ent}(f^2) \leq c(p) \mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right] \quad (8)$$

where

$$c(p) = \frac{1}{1-2p} \log \frac{1-p}{p}$$

Note that $\lim_{p \rightarrow 1/2} c(p) = 2$.

1.3 Gaussian Logarithmic Sobolev Inequality

- **Theorem 1.4** (*Gaussian Logarithmic Sobolev Inequality*). [Boucheron et al., 2013]
Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuous differentiable** function and let $Z = (Z_1, \dots, Z_n)$ be a vector of n **independent standard Gaussian** random variables. Then

$$\text{Ent}(f^2(Z)) \leq 2 \mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right]. \quad (9)$$

1.4 Modified Logarithmic Sobolev Inequalities

- **Proposition 1.5** (*A Modified Logarithmic Sobolev Inequalities for Moment Generating Function*) [Boucheron et al., 2013]

Consider independent random variables Z_1, \dots, Z_n taking values in \mathcal{X} , a real-valued function $f : \mathcal{X}^n \rightarrow \mathbb{R}$ and the random variable $X = f(Z_1, \dots, Z_n)$. Also denote $Z_{(-i)} = (Z_1, \dots, Z_{i-1}, Z_{i+1}, \dots, Z_n)$ and $X_{(-i)} = f_i(Z_{(-i)})$ where $f_i : \mathcal{X}^{n-1} \rightarrow \mathbb{R}$ is an arbitrary function. Let $\phi(x) = e^x - x - 1$. Then for all $\lambda \in \mathbb{R}$,

$$\text{Ent}(e^{\lambda X}) := \mathbb{E} \left[\lambda X e^{\lambda X} \right] - \mathbb{E} \left[e^{\lambda X} \right] \log \mathbb{E} \left[e^{\lambda X} \right] \leq \sum_{i=1}^n \mathbb{E} \left[e^{\lambda X} \phi(-\lambda(X - X_{(-i)})) \right] \quad (10)$$

- **Proposition 1.6** (*Symmetrized Modified Logarithmic Sobolev Inequalities*) [Boucheron et al., 2013]

Consider independent random variables Z_1, \dots, Z_n taking values in \mathcal{X} , a real-valued function $f : \mathcal{X}^n \rightarrow \mathbb{R}$ and the random variable $X = f(Z_1, \dots, Z_n)$. Also denote $\tilde{X}^{(i)} = f(Z_1, \dots, Z_{i-1}, Z'_i, Z_{i+1}, \dots, Z_n)$. Let $\phi(x) = e^x - x - 1$. Then for all $\lambda \in \mathbb{R}$,

$$\lambda \mathbb{E} \left[X e^{\lambda X} \right] - \mathbb{E} \left[e^{\lambda X} \right] \log \mathbb{E} \left[e^{\lambda X} \right] \leq \sum_{i=1}^n \mathbb{E} \left[e^{\lambda X} \phi(-\lambda(X - \tilde{X}^{(i)})) \right] \quad (11)$$

Moreover, denoting $\tau(x) = x(e^x - 1)$, for all $\lambda \in \mathbb{R}$,

$$\begin{aligned} \lambda \mathbb{E} \left[X e^{\lambda X} \right] - \mathbb{E} \left[e^{\lambda X} \right] \log \mathbb{E} \left[e^{\lambda X} \right] &\leq \sum_{i=1}^n \mathbb{E} \left[e^{\lambda X} \tau(-\lambda(X - \tilde{X}^{(i)})_+) \right], \\ \lambda \mathbb{E} \left[X e^{\lambda X} \right] - \mathbb{E} \left[e^{\lambda X} \right] \log \mathbb{E} \left[e^{\lambda X} \right] &\leq \sum_{i=1}^n \mathbb{E} \left[e^{\lambda X} \tau(\lambda(\tilde{X}^{(i)} - X)_-) \right]. \end{aligned}$$

2 Isoperimetric Inequalities and Concentration of Measure

2.1 Brunn-Minkowski Inequality

- **Definition (Minkowski Sum of Sets)**

Consider sets $A, B \subseteq \mathbb{R}^n$ and define the Minkowski sum of A and B as the set of all vectors in \mathbb{R}^n formed by sums of elements of A and B :

$$A + B := \{x + y : x \in A, y \in B\}$$

Similarly, for $c \in \mathbb{R}$, let $cA = \{cx : x \in A\}$. Denote by $\text{Vol}(A)$ the **Lebesgue measure** of a (measurable) set $A \subset \mathbb{R}^n$.

- **Theorem 2.1 (The Prékopa-Leindler Inequality).** [Boucheron et al., 2013, Wainwright, 2019]

Let $\lambda \in (0, 1)$, and let $f, g, h : \mathbb{R}^n \rightarrow [0, \infty)$ be **non-negative measurable functions** such that for all $x, y \in \mathbb{R}^n$,

$$h(\lambda x + (1 - \lambda)y) \geq f(x)^\lambda g(y)^{1-\lambda}.$$

Then

$$\int_{\mathbb{R}^n} h(x) dx \geq \left(\int_{\mathbb{R}^n} f(x) dx \right)^\lambda \left(\int_{\mathbb{R}^n} g(x) dx \right)^{1-\lambda}. \quad (12)$$

- **Corollary 2.2 (Weaker Brunn-Minkowski Inequality)** [Boucheron et al., 2013, Wainwright, 2019]

Let $A, B \subset \mathbb{R}^n$ be **non-empty compact sets**. Then for all $\lambda \in [0, 1]$,

$$\text{Vol}(\lambda A + (1 - \lambda)B) \geq \text{Vol}(A)^\lambda \text{Vol}(B)^{1-\lambda}. \quad (13)$$

- **Theorem 2.3 (Brunn-Minkowski Inequality)** [Boucheron et al., 2013, Vershynin, 2018, Wainwright, 2019]

Let $A, B \subset \mathbb{R}^n$ be **non-empty compact sets**. Then for all $\lambda \in [0, 1]$,

$$\text{Vol}(\lambda A + (1 - \lambda)B)^{\frac{1}{n}} \geq \lambda \text{Vol}(A)^{\frac{1}{n}} + (1 - \lambda) \text{Vol}(B)^{\frac{1}{n}}. \quad (14)$$

2.2 Classical Isoperimetric Problem on Euclidean Space \mathbb{R}^n

- **Definition (*Blowup of Sets*)**

For any $t > 0$, and any (measurable) sets $A \subset \mathbb{R}^n$, the t -blowup (or, t -enlargement) of A is defined by

$$A_t := \{x \in \mathbb{R}^n : d(x, A) < t\} = A + tB$$

where $B = \{x \in \mathbb{R}^n : d(0, x) < 1\}$ is an *open unit ball* and $d(x, A) = \inf_{y \in A} d(x, y)$.

- **Definition (*Surface Area of Sets*)**

let $A \subset \mathbb{R}^n$ be a measurable set and denote by $\text{Vol}(A)$ its *Lebesgue measure*. The surface area of A is defined by

$$\text{Vol}(\partial A) = \lim_{t \rightarrow 0} \frac{\text{Vol}(A_t) - \text{Vol}(A)}{t}.$$

provided that the limit exists. Here A_t denotes *the t -blowup* of A .

- **Remark (*Isoperimetry Theorem*)**

The classical isoperimetric theorem in \mathbb{R}^n states that, among all sets with **a given volume**, the Euclidean unit ball minimizes the surface area. This theorem can be formally stated as below:

- **Theorem 2.4 (*Isoperimetry Theorem*)** [Boucheron et al., 2013, Vershynin, 2018, Wainwright, 2019]

Let $A \subset \mathbb{R}^n$ be such that $\text{Vol}(A) = \text{Vol}(B)$ where $B := \{x \in \mathbb{R}^n : d(0, x) < 1\}$ is an unit ball. Then for any $t > 0$,

$$\text{Vol}(A_t) \geq \text{Vol}(B_t) \tag{15}$$

Moreover, if $\text{Vol}(\partial A)$ exists, then

$$\text{Vol}(\partial A) \geq \text{Vol}(\partial B). \tag{16}$$

- **Example (*Concentration of Lebesgue Measure in \mathbb{R}^n and Isoperimetric Inequality*)**

Note that the volume of a t -ball in \mathbb{R}^n is

$$\text{Vol}(tB) = \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2} + 1)} t^n \equiv c_n t^n$$

Thus the radius of ball B with the same volume of A is

$$r := \left(\frac{\text{Vol}(A)}{c_n} \right)^{\frac{1}{n}}.$$

The classical isoperimetric inequality states that

$$\begin{aligned} \text{Vol}(A_t) &\geq \left((r + t) \text{Vol}(B)^{1/n} \right)^n \\ \Leftrightarrow \text{Vol}(A_t) &\geq c_n \left(\left(\frac{\text{Vol}(A)}{c_n} \right)^{\frac{1}{n}} + t \right)^n \\ \Leftrightarrow \left(\frac{\text{Vol}(A_t)}{c_n} \right)^{\frac{1}{n}} &\geq \left(\frac{\text{Vol}(A)}{c_n} \right)^{\frac{1}{n}} + t \end{aligned} \tag{17}$$

- **Definition (*Isoperimetric Function of Probability Measure*)**

Define *the isoperimetric function* of the Lebesgue measure space (\mathbb{R}^n, μ) as

$$\lambda(u) := \left(\frac{u}{c_n} \right)^{\frac{1}{n}}$$

so the classical isoperimetric inequality is equivalent to the concentration of Lebesgue measure

$$\lambda(\mu(A_t)) \geq \lambda(\mu(A)) + t.$$

2.3 Isoperimetric Problem on Unit Sphere

- **Definition (*Spherical Cap and its t -Blowup*)**

Let $\mathbb{S}^{n-1} := \{x \in \mathbb{R}^n : \|x\| = 1\}$ be the $(n-1)$ -dimensional **unit sphere**. The **intersection** of a **half-space** and \mathbb{S}^{n-1} is called a **spherical cap**. In particular, for some $y \in \mathbb{R}^n$, denote the associated spherical cap as

$$H_y := \{x \in \mathbb{S}^{n-1} : \langle x, y \rangle \leq 0\}$$

With some simple geometry, it can be shown that its t -blowup corresponds to the set

$$H_y^t := \{x \in \mathbb{S}^{n-1} : \langle x, y \rangle < \sin(t)\}$$

- **Theorem 2.5 (*Isoperimetry Theorem on Unit Sphere*)** [Boucheron et al., 2013, Vershynin, 2018, Wainwright, 2019]

Let A be a subset of the sphere \mathbb{S}^{n-1} , and let σ denote the **normalized area** on that sphere. Let $t > 0$. Then, among all sets $A \subset \mathbb{S}^{n-1}$ with given area $\sigma(A)$, the **spherical caps minimize the area of the neighborhood** $\sigma(A_t)$, where

$$A_t := \{x \in \mathbb{S}^{n-1} : \exists y \in A \text{ such that } \|x - y\| < t\}$$

- **Remark** Define a *metric* ρ on sphere \mathbb{S}^{n-1} as

$$\rho(x, y) := \arccos(\langle x, y \rangle)$$

Thus (\mathbb{S}^{n-1}, ρ) is a **metric space**. Let \mathbb{P} be uniform distribution on \mathbb{S}^{n-1} so that $((\mathbb{S}^{n-1}, \rho), \mathbb{P})$ is a probability space.

- **Proposition 2.6 (*Isoperimetric Inequalities for Uniform Distribution over Sphere*)** [Boucheron et al., 2013, Vershynin, 2018, Wainwright, 2019]

Let $\mathbb{S}^{n-1} := \{x \in \mathbb{R}^n : \|x\| = 1\}$ be the $(n-1)$ -dimensional **unit sphere**. For any $t \in [0, 1]$,

$$\alpha_{\mathbb{S}^{n-1}}(t) \leq c \exp\left(-\frac{nt^2}{2}\right) \tag{18}$$

for some constant c .

- By Levy's inequality, we have the following proposition

Proposition 2.7 (*Lipschitz Function on \mathbb{S}^{n-1}*) [Wainwright, 2019]

For any 1-Lipschitz function f defined on the sphere \mathbb{S}^{n-1} , we have the two-sided bound

$$\mathbb{P}\{|f(Z) - \text{Med}(f(Z))| \geq t\} \leq \sqrt{2\pi} \exp\left(-\frac{nt^2}{2}\right) \quad (19)$$

Moreover, replacing median by the mean, we have

$$\mathbb{P}\{|f(Z) - \mathbb{E}[f(Z)]| \geq t\} \leq 2\sqrt{2\pi} \exp\left(-\frac{nt^2}{8}\right) \quad (20)$$

• **Exercise 2.8** (*The Blow-Up Phenomenon*)

Let A be a subset of the sphere $\sqrt{n}\mathbb{S}^{n-1}$ such that

$$\mathbb{P}(A) > 2 \exp(-cs^2) \text{ for some } s > 0;$$

1. Prove that $\mathbb{P}(A_s) > 1/2$.
2. Deduce from this that for any $t \geq s$,

$$\mathbb{P}(A_{2t}) > 1 - 2 \exp(-ct^2).$$

Here $c > 0$ is the absolute constant in upper bound of concentration function.

2.4 Concentration via Isoperimetric Inequalities

• **Definition** (*Isoperimetry Problem*) [Boucheron et al., 2013]

Given a **metric space** \mathcal{X} with corresponding *distance* d , consider **the measure space** formed by \mathcal{X} , the σ -algebra of all **Borel sets** of \mathcal{X} , and a probability measure \mathbb{P} . Let X be a *random variable* taking values in \mathcal{X} , distributed according to \mathbb{P} .

The isoperimetric problem in this case is the following: given $p \in (0, 1)$ and $t > 0$, **determine the sets** A with $\mathbb{P}[X \in A] \geq p$ for which **the measure**

$$\mathbb{P}[d(X, A) \geq t]$$

is **maximal**.

• **Remark** (*Isoperimetric Inequalities*)

Even though the exact solution is only known in a few special cases, useful *bounds* for $\mathbb{P}[d(X, A) \geq t]$ can be derived under remarkably general circumstances. *Such bounds are usually referred to as **isoperimetric inequalities**.*

• **Definition** (*Concentration Function*) [Boucheron et al., 2013, Wainwright, 2019]

The concentration function $\alpha : [0, \infty) \rightarrow \mathbb{R}_+$ associated with **metric measure space** $((\mathcal{X}, d), \mathbb{P})$ is given by

$$\alpha_{\mathbb{P}, (\mathcal{X}, d)}(t) := \sup_{A \subset \mathcal{X}: \mathbb{P}(A) \geq \frac{1}{2}} \mathbb{P}[d(X, A) \geq t] = \sup_{A \subset \mathcal{X}: \mathbb{P}(A) \geq \frac{1}{2}} \mathbb{P}(A_t^c)$$

where $A_t := A + tB = \{x \in \mathcal{X} : d(x, A) < t\}$ is the *t-blowup* of $A \subset \mathcal{X}$. We simply denote it as $\alpha(t)$.

Thus the optimal A^* for isoperimetry problem is the one that attains the $\alpha(t) = \mathbb{P}(A_t^c)$.

- **Theorem 2.9 (Levy's Inequalities)** [Boucheron et al., 2013, Wainwright, 2019]

For any Lipschitz function $f : \mathcal{X} \rightarrow \mathbb{R}$,

$$\begin{aligned}\mathbb{P}\{f(X) \geq \text{Med}(f(X)) + t\} &\leq \alpha_{\mathbb{P}}(t) \\ \mathbb{P}\{f(X) \leq \text{Med}(f(X)) - t\} &\leq \alpha_{\mathbb{P}}(t).\end{aligned}\tag{21}$$

where $\text{Med}(f(X))$ is the median of $f(X)$, i.e.

$$\mathbb{P}\{f(X) \leq \text{Med}(f(X))\} \geq \frac{1}{2}, \quad \text{and} \quad \mathbb{P}\{f(X) \geq \text{Med}(f(X))\} \geq \frac{1}{2}.$$

Conversely, if $\beta : \mathbb{R}_+ \rightarrow [0, 1]$ is a function such that for **every Lipschitz function** $f : \mathcal{X} \rightarrow \mathbb{R}$

$$\mathbb{P}\{f(X) - \text{Med}(f(X)) \geq t\} \leq \beta(t).\tag{22}$$

then $\beta(t) \geq \alpha_{\mathbb{P}}(t)$.

- **Corollary 2.10 (Concentration of Measure on Hamming Metric Space)** [Boucheron et al., 2013]

Consider independent random variables Z_1, \dots, Z_n taking their values in a (measurable) set \mathcal{X} and denote the vector of these variables by $Z = (Z_1, \dots, Z_n)$ taking its value in \mathcal{X}^n . For an arbitrary (measurable) set $A \subset \mathcal{X}^n$, we write $\mathbb{P}(A) = \mathbb{P}(Z \in A)$. The **Hamming distance** $d_H(x, y)$ between the vectors $x, y \in \mathcal{X}^n$ is defined as **the number of coordinates in which x and y differ**. Then for any $t > 0$,

$$\mathbb{P}\left\{d_H(x, A) \geq \sqrt{\frac{n}{2} \log \frac{1}{\mathbb{P}(A)}} + t\right\} \leq \exp\left(-\frac{2t^2}{n}\right)\tag{23}$$

- **Remark (Equivalent Form)**

From above isoperimetric inequality,

$$\mathbb{P}\left\{d_H(x, A) \geq \sqrt{\frac{n}{2} \log \frac{1}{\mathbb{P}(A)}} + t\right\} \leq \exp\left(-\frac{2t^2}{n}\right)$$

Denote $u := \sqrt{\frac{n}{2} \log \frac{1}{\mathbb{P}(A)}}$. By change of variable, for any $t \geq u$,

$$\mathbb{P}\{d_H(x, A) \geq t\} \leq \exp\left(-\frac{2(t-u)^2}{n}\right).$$

On the one hand, if $t \leq 2u = \sqrt{-2n \log \mathbb{P}(A)}$, then $\mathbb{P}(A) \leq \exp(-t^2/(2n))$. On the other hand, since $(t-u)^2 \geq t^2/4$ for $t \geq 2u = \sqrt{-2n \log \mathbb{P}(A)}$, the inequality above implies $\mathbb{P}\{d_H(x, A) \geq t\} \leq \exp(-t^2/(2n))$. Thus, for all $t > 0$, we have **the concentration of measure in Hamming metric space**:

$$\mathbb{P}(A)\mathbb{P}\{d_H(x, A) \geq t\} \leq \min\{\mathbb{P}(A), \mathbb{P}\{d_H(x, A) \geq t\}\} \leq \exp\left(-\frac{t^2}{2n}\right)\tag{24}$$

- **Proposition 2.11 (Levy's Inequalities for Mean)** [Boucheron et al., 2013, Wainwright, 2019]

If $\beta : \mathbb{R}_+ \rightarrow [0, 1]$ is a function such that for **every Lipschitz function** $f : \mathcal{X} \rightarrow \mathbb{R}$

$$\mathbb{P}\{f(X) - \mathbb{E}[f(X)] \geq t\} \leq \beta(t).\tag{25}$$

then $\beta(t) \geq \alpha_{\mathbb{P}}(t/2)$.

2.5 Convex Distance Inequality

- **Definition (Weighted Hamming Distance)**

Given $\alpha = (\alpha_1, \dots, \alpha_n)$, where $\alpha_i \geq 0$, *the weighed Hamming distance* between $x, y \in \mathcal{X}^n$ is defined as

$$d_\alpha(x, y) = \sum_{i=1}^n \alpha_i \mathbb{1}\{x_i \neq y_i\}.$$

- **Definition (Convex Distance)**

For any $x = (x_1, \dots, x_n) \in \mathcal{X}^n$, *the convex distance* of x from the set A by

$$d_T(x, A) := \sup_{\alpha \in \mathbb{R}_+^n : \|\alpha\|_2 = 1} d_\alpha(x, A)$$

- **Theorem 2.12 (Convex Distance Inequality)** [Boucheron et al., 2013]

For any subset $A \subset \mathcal{X}^n$ and $t > 0$,

$$\begin{aligned} \mathbb{P}(A) \mathbb{P}\{d_T(X, A) \geq t\} &\leq \exp\left(-\frac{t^2}{4}\right). \\ \Leftrightarrow \mathbb{P}(A) \mathbb{P}\left\{\sup_{\alpha \in \mathbb{R}_+^n : \|\alpha\|_2 = 1} \inf_{y \in A} \sum_{i=1}^n \alpha_i \mathbb{1}\{x_i \neq y_i\} \geq t\right\} &\leq \exp\left(-\frac{t^2}{4}\right). \end{aligned} \quad (26)$$

2.6 Concentration of Convex Lipschitz Functions

- **Theorem 2.13 (Concentration of Separately Convex Lipschitz Functions)** [Boucheron et al., 2013]

Let $Z := (Z_1, \dots, Z_n)$ be independent random variables taking values in the interval $[0, 1]$ and let $f : [0, 1]^n \rightarrow \mathbb{R}$ be a **separately convex function** (i.e. f is convex in each coordinate while the others are fixed) such that

$$|f(x) - f(y)| \leq \|x - y\| \quad \text{for all } x, y \in [0, 1]^n.$$

Then $X = f(Z_1, \dots, Z_n)$ satisfies, for all $t > 0$,

$$\mathbb{P}\{f(Z) - \mathbb{E}[f(Z)] \geq t\} \leq \exp\left(-\frac{t^2}{2}\right). \quad (27)$$

- **Remark** For $Z_i \in [a_i, b_i]$, and f being L -Lipschitz function, we have

$$\mathbb{P}\{f(Z) - \mathbb{E}[f(Z)] \geq t\} \leq \exp\left(-\frac{t^2}{2L^2 \sum_{i=1}^n (b_i - a_i)^2}\right)$$

- With convex distance inequality, we can improve *the concentration bound for convex Lipschitz functions*. First, we relate convex distance with the minimal distance to convex set

Lemma 2.14 (Convex Distance vs. Distance to Convex Set) [Boucheron et al., 2013]

Let $A \subset [0, 1]^n$ be a **convex set** and let $x = (x_1, \dots, x_n) \in [0, 1]^n$. Then

$$d(x, A) := \inf_{y \in A} \|x - y\|_2 \leq d_T(x, A). \quad (28)$$

- **Theorem 2.15 (Concentration of Convex Lipschitz Functions)** [Boucheron et al., 2013]

Let $Z := (Z_1, \dots, Z_n)$ be independent random variables taking values in the interval $[0, 1]$ and let $f : [0, 1]^n \rightarrow \mathbb{R}$ be a **quasi-convex function**; that is

$$\{z : f(z) \leq s\} \text{ is convex set for all } s \in \mathbb{R}.$$

Moreover, f is Lipschitz function satisfying

$$|f(x) - f(y)| \leq \|x - y\| \quad \text{for all } x, y \in [0, 1]^n.$$

Then $X = f(Z_1, \dots, Z_n)$ satisfies, for all $t > 0$,

$$\begin{aligned} \mathbb{P}\{f(Z) \geq \text{Med}(f(Z)) + t\} &\leq 2 \exp\left(-\frac{t^2}{4}\right), \\ \mathbb{P}\{f(Z) \leq \text{Med}(f(Z)) - t\} &\leq 2 \exp\left(-\frac{t^2}{4}\right). \end{aligned} \tag{29}$$

- **Remark** This result can be generalized to **convex Lipschitz function of sub-Gaussian random variables**. Note that in above theorems, $Z_i \in [0, 1]$ is **sub-Gaussian** as well.

3 Concentration of Gaussian Measure

3.1 Gaussian Isoperimetric Theorem and Gaussian Concentration Theorem

- **Remark (Gaussian Isoperimetric Problem)**

The Gaussian isoperimetric problem is to determine which (Borel) sets A have **minimal Gaussian boundary measure** among all sets in \mathbb{R}^n with a given probability p .

The Gaussian isoperimetric theorem states the beautiful fact that the extremal sets are linear half-spaces in all dimensions and for all p .

- **Definition (Gaussian Isoperimetric Function)**

Denote the cumulative distribution function of standard Normal distribution:

$$\Phi(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-\frac{t^2}{2}} dt := \int_{-\infty}^x \varphi(t) dt$$

where $\varphi(x) := \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} = (\Phi(x))'$ is the probability density function of standard normal distribution. $\Phi^{-1}(x)$ is the quantile function of normal distribution.

Define the Gaussian isoperimetric function as

$$\gamma(x) := \varphi(\Phi^{-1}(x)), \quad x \in (0, 1).$$

Also we define $\gamma(0) = \gamma(1) = 0$.

- **Remark** Note that

$$\begin{aligned} x &= \Phi(\Phi^{-1}(x)) \\ \Rightarrow 1 &= \varphi(\Phi^{-1}(x))(\Phi^{-1}(x))' = \gamma(x)(\Phi^{-1}(x))' \\ \Leftrightarrow 1/\gamma(x) &= (\Phi^{-1}(x))'. \end{aligned}$$

The quantity $1/\gamma(x) = (\Phi^{-1}(x))'$ is known as **quantile-density function of normal distribution**.

- **Proposition 3.1 (Basic Property of the Gaussian Isoperimetric Function)** [Boucheron et al., 2013]

The Gaussian isoperimetric function γ satisfies:

1.

$$\gamma'(x) = -\Phi^{-1}(x), \quad \text{for all } x \in (0, 1),$$

2.

$$\gamma(x)\gamma''(x) = -1, \quad \text{for all } x \in (0, 1),$$

3. $(\gamma')^2$ is convex over $(0, 1)$.

- **Lemma 3.2 (Asymptotic Behavior of Gaussian Isoperimetric Function)** [Boucheron et al., 2013]

For all $x \in [0, 1/2]$,

$$x\sqrt{\frac{1}{2}\log\frac{1}{x}} \leq \gamma(x) \leq x\sqrt{2\log\frac{1}{x}}.$$

Moreover,

$$\lim_{x \rightarrow 0} \frac{\gamma(x)}{x\sqrt{2\log\frac{1}{x}}} = 1$$

- **Proposition 3.3 (Bobkov's Gaussian Inequality)** [Boucheron et al., 2013]

Let $Z := (Z_1, \dots, Z_n)$ be a vector of **independent standard Gaussian** random variables. Let $f : \mathbb{R}^n \rightarrow [0, 1]$ be a differentiable function with gradient ∇f . Then

$$\gamma(\mathbb{E}[f(X)]) \leq \mathbb{E} \left[\sqrt{\gamma(f(X))^2 + \|\nabla f(X)\|_2^2} \right] \quad (30)$$

where $\gamma = \varphi \circ \Phi^{-1}$ is the **Gaussian isoperimetric function**.

- **Theorem 3.4 (Gaussian Isoperimetric Theorem)** [Boucheron et al., 2013] [Boucheron et al., 2013, Vershynin, 2018, Wainwright, 2019]

Let \mathbb{P} be the **standard Gaussian measure** on \mathbb{R}^n and let $A \subset \mathbb{R}^n$ be a Borel set. Then

$$\liminf_{t \rightarrow 0} \frac{\mathbb{P}(A_t) - \mathbb{P}(A)}{t} \geq \gamma(\mathbb{P}(A)), \quad (31)$$

where $A_t := \{x : d(x, A) < t\}$ be the t -blowup of A . Moreover, if A is a **half-space** defined by $A := \{x \in \mathbb{R}^n : x_1 \leq z\}$, then

$$\liminf_{t \rightarrow 0} \frac{\mathbb{P}(A_t) - \mathbb{P}(A)}{t} = \gamma(\mathbb{P}(A)) = \varphi(z), \quad (32)$$

where $\gamma := \varphi \circ \Phi^{-1}$ is the **Gaussian isoperimetric function**.

- **Proposition 3.5 (Differentiability of Measure of t -Blowup)** [Boucheron et al., 2013]
If A is a **finite union of open balls** in \mathbb{R}^n , then $\mathbb{P}(A_t)$ is a **differentiable** function of $t > 0$.
- Next we describe **an equivalent version of the Gaussian isoperimetric theorem** in the manner of *measure concentration*:

Theorem 3.6 (Gaussian Concentration Theorem) [Boucheron et al., 2013] [Boucheron et al., 2013, Vershynin, 2018, Wainwright, 2019]

Let \mathbb{P} be the **standard Gaussian measure** on \mathbb{R}^n and let $A \subset \mathbb{R}^n$ be a Borel set. Then for all $t \geq 0$,

$$\begin{aligned} \mathbb{P}(A_t) &\geq \Phi(\Phi^{-1}(\mathbb{P}(A)) + t) . \\ \Leftrightarrow \Phi^{-1}(\mathbb{P}(A_t)) &\geq \Phi^{-1}(\mathbb{P}(A)) + t \end{aligned} \quad (33)$$

Equality holds if A is a **half-space**.

- **Remark (Gaussian Concentration Theorem \equiv Gaussian Isoperimetric Theorem)**
The Gaussian concentration theorem is equivalent to the Gaussian isoperimetric theorem since

$$\begin{aligned} \liminf_{t \rightarrow 0} \frac{\mathbb{P}(A_t) - \mathbb{P}(A)}{t} &\geq \liminf_{t \rightarrow 0} \frac{\Phi(\Phi^{-1}(\mathbb{P}(A)) + t) - \Phi(\Phi^{-1}(\mathbb{P}(A)))}{t} \\ &= \Phi'(\Phi^{-1}(\mathbb{P}(A))) \\ &= \varphi(\Phi^{-1}(\mathbb{P}(A))) \\ &= \gamma(\mathbb{P}(A)). \end{aligned}$$

- **Exercise 3.7 (From Isoperimetry to Concentration)** [Boucheron et al., 2013]
Assume that a probability distribution \mathbb{P} on \mathbb{R}^n satisfies, for all Borel sets $A \subset \mathbb{R}^n$,

$$\liminf_{t \rightarrow 0} \frac{\mathbb{P}(A_t) - \mathbb{P}(A)}{t} \geq c f(F^{-1}(\mathbb{P}(A))),$$

where $c \in (0, 1]$ is a constant, F is a continuously differentiable distribution function and $f = F'$ is its derivative. Prove that for all Borel set A and all $t \geq 0$,

$$\mathbb{P}(A_t) \geq F(F^{-1}(\mathbb{P}(A)) + ct) .$$

3.2 Lipschitz Functions of Gaussian Variables

- **Theorem 3.8 (Rademacher Theorem).**
If $f : U \rightarrow \mathbb{R}$ is a L -Lipschitz function where $U \subseteq \mathbb{R}^n$, then f is **differentiable almost everywhere** in U and the **essential supremum of the norm of its derivative is bounded** by its **Lipschitz constant**.
- **Theorem 3.9 (Lipschitz Functions of Gaussian Variables)** [Boucheron et al., 2013]
Let $Z = (Z_1, \dots, Z_n)$ be a vector of n **independent standard normal** random variables. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ denote an **L -Lipschitz function**, that is, there exists a constant $L > 0$ such that for all $x, y \in \mathbb{R}^n$,

$$|f(x) - f(y)| \leq L \|x - y\| .$$

Then, for all $\lambda \in \mathbb{R}$,

$$\psi_{f(Z) - \mathbb{E}[f(Z)]}(\lambda) := \log \mathbb{E} \left[e^{\lambda(f(Z) - \mathbb{E}[f(Z)])} \right] \leq \frac{L^2 \lambda^2}{2} \quad (34)$$

- **Theorem 3.10 (Gaussian Concentration Inequality) (The Tsirelson-Ibragimov-Sudakov Inequality)** [Boucheron et al., 2013, Wainwright, 2019]

Let $Z = (Z_1, \dots, Z_n)$ be a vector of n **independent standard normal** random variables. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ denote an **L -Lipschitz function**. Then, for all $t > 0$,

$$\mathbb{P} \{f(Z) - \mathbb{E}[f(Z)] \geq t\} \leq \exp \left(-\frac{t^2}{2L^2} \right). \quad (35)$$

- As a direct consequence of the *Gaussian isoperimetric inequality*, we have the improved result for *Gaussian concentration inequality*:

Theorem 3.11 (Gaussian Concentration Inequality, Sharp Bound) [Boucheron et al., 2013, Wainwright, 2019]

Let $Z = (Z_1, \dots, Z_n)$ be a vector of n **independent standard normal** random variables. Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ denote an **L -Lipschitz function**. Then, for all $t > 0$,

$$\mathbb{P} \{f(Z) - \text{Med}(f(Z)) \geq t\} \leq 1 - \Phi \left(\frac{t}{L} \right). \quad (36)$$

where $\Phi(t)$ is the cumulative distribution function of standard normal random variable.

- **Remark** Note that by **Gordon's inequality**

$$1 - \Phi(t) \leq \left(\frac{1}{\sqrt{2\pi}} \right) \frac{1}{t} e^{-\frac{t^2}{2}} = \frac{1}{t} \varphi(t)$$

The Gaussian concentration inequality fails to capture the corrective factor t^{-1} . The inequality above cannot be improved in general as for $f(x) = n^{-1/2} \sum_{i=1}^n x_i$, equality is achieved for all $t > 0$.

3.3 Gaussian Logarithmic Sobolev Inequality

- **Theorem 3.12 (Gaussian Logarithmic Sobolev Inequality).** [Boucheron et al., 2013]
Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuous differentiable function** and let $Z = (Z_1, \dots, Z_n)$ be a vector of n **independent standard Gaussian** random variables. Then

$$\text{Ent}(f^2(Z)) \leq 2\mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right]. \quad (37)$$

3.4 Gaussian Transportation Inequality

- **Theorem 3.13 (Talagrand's Gaussian Transportation Inequality)** [Boucheron et al., 2013]

Let \mathbb{P} be the standard Gaussian probability measure on \mathbb{R}^n , and let \mathbb{Q} be a probability measure absolutely continuous with respect to \mathbb{P} . Define two random vectors $X = (X_1, \dots, X_n), Y = (Y_1, \dots, Y_n)$ in \mathcal{X}^n with distribution \mathbb{P} and \mathbb{Q} respectively. Then

$$\begin{aligned} \mathcal{W}_{2,d}(\mathbb{Q}, \mathbb{P}) &:= \sqrt{\min_{\gamma \in \Pi(\mathbb{Q}, \mathbb{P})} \sum_{i=1}^n \mathbb{E}_{\gamma} [(X_i - Y_i)^2]} \leq \sqrt{2\text{KL}(\mathbb{Q} \parallel \mathbb{P})} \\ &\Leftrightarrow \min_{\gamma \in \Pi(\mathbb{Q}, \mathbb{P})} \sum_{i=1}^n \mathbb{E}_{\gamma} [(X_i - Y_i)^2] \leq 2\text{KL}(\mathbb{Q} \parallel \mathbb{P}) \end{aligned} \quad (38)$$

3.5 Gaussian Hypercontractivity

3.6 Suprema of Gaussian Process

- **Definition (Gaussian Process)**

Let T be a metric space. A stochastic process $(X_t)_{t \in T}$ is a **Gaussian process indexed by T** if for any finite collection $\{t_1, \dots, t_n\} \subset T$, the vector $(X_{t_1}, \dots, X_{t_n})$ has a jointly Gaussian distribution.

In addition, we assume that T is **totally bounded** (i.e. for every $t > 0$ it can be covered by finitely many balls of radius t) and that the Gaussian process is **almost surely continuous**, that is, with probability 1, X_t is a continuous function of t .

- **Theorem 3.14 (Concentration of Suprema of Gaussian Process)** [Boucheron et al., 2013, Vershynin, 2018, Wainwright, 2019, Giné and Nickl, 2021]
Let $(X_t)_{t \in T}$ be an **almost surely continuous centered Gaussian process indexed by a totally bounded set T** . If

$$\sigma^2 := \sup_{t \in T} \mathbb{E} [X_t^2],$$

then $Z = \sup_{t \in T} X_t$ satisfies $\text{Var}(Z) \leq \sigma^2$, and for all $u > 0$,

$$\mathbb{P} \{Z - \mathbb{E} [Z] \geq u\} \leq \exp \left(-\frac{u^2}{2\sigma^2} \right) \quad (39)$$

and

$$\mathbb{P} \{\mathbb{E} [Z] - Z \geq u\} \leq \exp \left(-\frac{u^2}{2\sigma^2} \right) \quad (40)$$

4 Concentration of Bernoulli Measure on the Binary Hypercube

4.1 Edge Isoperimetric Inequality on the Binary Hypercube

- **Remark (Binary Hypercube as Nearest Neighbor Graph with respect to Hamming Distance)**

Consider binary hypercube $\{-1, 1\}^n$ with *Hamming distance metric*

$$d_H(x, y) = \sum_{i=1}^n \mathbb{1}\{x_i \neq y_i\}$$

The elements x of the binary n -cube may be considered as **vertices** of a graph $\mathcal{G} := (\mathcal{V}, \mathcal{E})$ in which two elements x and x' of $\{-1, 1\}^n$ are **adjacent** if and only if **their Hamming distance is 1**; i.e.

$$\mathcal{E} = \{(x, y) \in \{-1, 1\}^n \times \{-1, 1\}^n : d_H(x, y) = 1\}.$$

The graph structure has $|\mathcal{V}| := N = 2^n$ vertices and $|\mathcal{E}| = n2^{n-1}$ undirected edges. Its **density** (the ratio between the number of edges and the number of vertices) is thus $n/2 = (\log_2 N)/2$.

- **Remark (Maximum Density of Subgraph)**

A remarkable property of the binary n -cube is that for any subset $A \subseteq \{-1, 1\}^n$, the **density** of the subgraph induced by A is at most $(\log_2 |A|)/2$. Note that **equality** is achieved if the graph induced by A is a **lower-dimensional hypercube**, since if A is a hypercube of dimension $d \leq n$, then the subgraph induced by A has 2^d vertices and $\mathcal{E}(A) = d2^{d-1}$ edges.

Theorem 4.1 (Maximum Density of Subgraph) [Boucheron et al., 2013]

Let A be a subset of $\{-1, 1\}^n$. Let $\mathcal{E}(A)$ denote the set of edges of the subgraph induced by A , that is, the collection of (unordered) pairs (x, x') with $x, x' \in A$ such that $d_H(x, x') = 1$. Then

$$|\mathcal{E}(A)| \leq \frac{|A|}{2} \log_2(|A|). \quad (41)$$

- **Definition (Influence of Binary Variable with respect to Set)**

Let the binary random vector $X = (X_1, \dots, X_n)$ be uniformly distributed over $\{-1, 1\}^n$ and denote by $\tilde{X}^{(i)} = (X_1, \dots, X_{i-1}, -X_i, X_{i+1}, \dots, X_n)$ the vector obtained by **flipping the i -th bit** of X . For any $A \subseteq \{-1, 1\}^n$, the influence of the i -th variable is defined by

$$\begin{aligned} I_i(A) &= \mathbb{P} \left\{ \mathbb{1}\{X \in A\} \neq \mathbb{1}\{\tilde{X}^{(i)} \in A\} \right\} \\ &= \mathbb{P} \left\{ (X \in A \wedge \tilde{X}^{(i)} \notin A) \vee (X \notin A \wedge \tilde{X}^{(i)} \in A) \right\} \end{aligned}$$

If $\mathbb{1}\{X \in A\} \neq \mathbb{1}\{\tilde{X}^{(i)} \in A\}$, then the i -th variable is said to be **pivotal** for A . Thus, the influence $I_i(A)$ is just **the probability that the i -th variable is pivotal for A** .

The total influence is defined by the *sum of individual influences*

$$I(A) := \sum_{i=1}^n I_i(A)$$

- **Definition (Edge Boundary of Subset)**

Let A be a subset of $\{-1, 1\}^n$. Let $\mathcal{E}(A)$ denote the set of edges of the subgraph induced by A . The edge boundary of A , $\partial\mathcal{E}(A)$, is define as

$$\partial\mathcal{E}(A) := \{(x, y) : x \in A, y \in A^c, d_H(x, y) = 1\}.$$

Thus the total number of edges connects to all of vertices in A can be computed as

$$n|A| = 2|\mathcal{E}(A)| + |\partial\mathcal{E}(A)| \quad (42)$$

where each vertex connects to exactly n edges, and every edge with both endpoints in A is counted twice.

- **Theorem 4.2 (Edge Isoperimetric Theorem of Binary Hypercube)** [Boucheron et al., 2013]

For any $A \subset \{-1, 1\}^n$, let $\mathbb{P}(A)$ denote $\mathbb{P}\{X \in A\} = |A|/2^n$. Then

$$I(A) \geq 2\mathbb{P}(A) \log_2 \left(\frac{1}{\mathbb{P}(A)} \right) \quad (43)$$

4.2 Bobkov's Inequality

- **Proposition 4.3 (Bobkov's Inequality)** [Boucheron et al., 2013]

Suppose Z is uniformly distributed over $\{-1, 1\}^n$. Then for all $n \geq 1$ and for all functions $f : \{-1, 1\}^n \rightarrow [0, 1]$,

$$\gamma(\mathbb{E}[f(Z)]) \leq \mathbb{E} \left[\sqrt{\gamma(f(Z))^2 + \|\nabla f(Z)\|_2^2} \right] \quad (44)$$

4.3 Vertex Isoperimetric Inequality on the Binary Hypercube

- **Definition (Vertex Boundary of Graph)** [Boucheron et al., 2013]

Consider a graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ and let $\mathcal{A} \subset \mathcal{V}$ be a set of its vertices. **The vertex boundary** of \mathcal{A} is defined as the set of those vertices, **not in** \mathcal{A} , which are **connected** to some vertex in \mathcal{V} by an edge. We denote the vertex boundary of \mathcal{A} by $\partial V(\mathcal{A})$.

- **Remark (Binary Hypercube as Nearest Neighbor Graph with Hamming Distance)**

Consider the graph as binary hypercube $\{-1, +1\}^n$ in which two vertices are connected by an edge if and only if their **Hamming distance** equals 1. Define the **norm** as the **Hamming distance** to $-1^n = (-1, \dots, -1)$

$$\|x\|_H := \sum_{i=1}^n \mathbb{1}\{x_i = 1\} = d_H(x, -1^n)$$

- **Definition (Simplicial Order)**

We define the so-called **simplicial order** of the elements of the binary hypercube. We say that $x = (x_1, \dots, x_n) \in \{-1, 1\}^n$ **precedes** $y = (y_1, \dots, y_n) \in \{-1, 1\}^n$ in the simplicial order if either $\|x\|_H < \|y\|_H$ (where $\|x\|_H := \sum_{i=1}^n \mathbb{1}\{x_i = 1\} = d_H(x, -1^n)$) or $\|x\|_H = \|y\|_H$ and $x_i = 1$ and $y_i = -1$ for the smallest i for which $x_i \neq y_i$. That is

$$\begin{aligned} & x \prec y \\ \Leftrightarrow & \{(x, y) : \|x\|_H < \|y\|_H \vee (\|x\|_H = \|y\|_H \wedge (x_i = 1 \wedge y_i = -1, \text{ where } i = \min\{k : x_k \neq y_k\}))\} \end{aligned}$$

In other words, the vector with **less** 1's **precedes** the vector with more 1's. If the number of 1's are the same, then the first 1's on the leftmost position is preferred.

Theorem 4.4 (*Harp's Vertex Isoperimetric Theorem*) [Boucheron et al., 2013]

For $N = 1, \dots, 2^n$, let S_N denote the set of **first** N **elements** of $\{-1, +1\}^n$ in the **simplicial order**. For any subset $A \subset \{-1, +1\}^n$, where $|A| = N$,

$$|\partial V(A)| \geq |\partial V(S_N)|$$

- **Remark** Note that if $N = \sum_{i=0}^k \binom{n}{i}$, for $k = 0, \dots, n$, then

$$S_N = \{x \in \{-1, +1\}^n : d_H(x, -1^n) \leq k\} = B_H(-1^n, k)$$

In other words, S_N is a **Hamming ball** centered at the vector $-1^n = (-1, \dots, -1)$.

- **Definition** (*t-Blowup of Set A in Binary Hypercube*)

For any $A \subset \{-1, +1\}^n$ and $x \in \{-1, +1\}^n$, let $d_H(x, A) = \min_{y \in A} d_H(x, y)$ be the *Hamming distance* of x to the set A . Also, denote by

$$A_t := \{x \in \{-1, +1\}^n : d_H(x, A) < t\}$$

the *t-blowup* of the set A , that is, the set of points whose Hamming distance from A is at most t .

- **Corollary 4.5** (*Isoperimetric Inequality in Binary Hypercube*) [Boucheron et al., 2013]

Let $A \subset \{-1, +1\}^n$ such that $|A| \geq \sum_{i=0}^k \binom{n}{i}$. Then for any $t = 1, 2, \dots, n - k + 1$,

$$|A_t| \geq \sum_{i=0}^{k+1-t} \binom{n}{i}. \quad (45)$$

In particular, if $|A|/2^n \geq 1/2$ then we may take $k = \lfloor n/2 \rfloor$ in the corollary above and

$$\frac{|A_t|}{2^n} \geq \mathbb{P}\{X < \mathbb{E}[X] + t\} \geq 1 - \exp\left(-\frac{2t^2}{n}\right) \quad (46)$$

where $X \sim \text{Ber}(1/2)$ is a symmetric Bernoulli random variable taking values in $\{-1, +1\}$ with $\mathbb{P}\{X = 1\} = \mathbb{P}\{X = -1\} = 1/2$.

- **Proof:** (*Proof of Bounded Difference Inequality on Binary Hypercube*)

Note that any function with **bounded difference property** is **Lipschitz function** with respect to **Hamming distance**.

$$\begin{aligned} & \sup_{x \in \mathcal{X}^n, y_i \in \mathcal{X}} |f(x_1, \dots, x_n) - f(x_1, \dots, x_{i-1}, y_i, x_{i+1}, \dots, x_n)| \\ & \leq c_i = c_i d_H((x_1, \dots, x_n), (x_1, \dots, x_{i-1}, y_i, x_{i+1}, \dots, x_n)), \quad 1 \leq i \leq n \\ \Rightarrow |f(x) - f(y)| &= \left| \sum_{i=1}^n (f(x_{(i-1)}) - f(x_{(i)})) \right| \\ & \leq \sum_{i=1}^n |f(x_{(i-1)}) - f(x_{(i)})| \\ & \leq \sum_{i=1}^n L_i \mathbb{1}\{x_{(i-1)}[i] \neq x_{(i)}[i]\} \\ & = d_{H,L}(x, y) \end{aligned}$$

where $x_{(i)}$ is replicate of $x_{(i-1)}$ except for i -th component, which is replaced by y_i . Note that $x_{(0)} = x$ and $x_{(n)} = y$.

The Harp's isoperimetric theorem suggests that the concentration function

$$\alpha_{\mathbb{P},(\{-1,+1\}^n, d_{H,L})}(t) := \sup_{A: \mathbb{P}\{A\} \geq 1/2} \mathbb{P}\{A_t\} \leq \exp\left(-\frac{2t^2}{\sum_i L_i^2}\right)$$

where \mathbb{P} is uniform distribution on $\{-1, +1\}^n$. Thus by *Levy's inequality*, we prove that for $Z \in \{-1, 1\}^n$ and Lipschitz function $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ with respect to weighted Hamming distance $d_{H,L}$,

$$\mathbb{P}\{|f(Z) - \text{Med}(f(Z))| \geq t\} \leq 2 \exp\left(-\frac{2t^2}{\sum_i L_i^2}\right). \quad \blacksquare$$

- **Remark (*Concentration of Uniform Measure on Binary Hypercube*)**

Consider any set A containing at least half of the points of $\{-1, +1\}^n$. According to the corollary above, *the fraction of those points which **cannot be obtained** by changing **at most** $c\sqrt{n}$ bits of some point in A is **at most** e^{2c^2}* . In other words, ***an immense majority** of the points in $\{-1, +1\}^n$ is **within Hamming distance of the order of \sqrt{n}** of A*

- **Definition (*Monotone Set*)**

A set $A \subset \{-1, +1\}^n$ is **monotone** if $\mathbb{1}\{x \in A\} \geq \mathbb{1}\{y \in A\}$ for all $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ in $\{-1, +1\}^n$ such that $x_i \geq y_i$ **for all** i .

- **Theorem 4.6 (*Isoperimetric Theorem for Bernoulli Distribution*)** [Boucheron et al., 2013]

Let $k \in \{0, \dots, n\}$ and let $S = \{x \in \{-1, +1\}^n : \|x\| < t\}$ be a **Hamming ball of radius** k . \mathbb{P} is Bernoulli distribution on $\{-1, 1\}^n$ with parameter p such that

$$\mathbb{P}(x) := p^{\|x\|} (1-p)^{1-\|x\|}.$$

If $A \subset \{-1, +1\}^n$ is a **monotone set** such that $\mathbb{P}(A) \geq \mathbb{P}(S)$ then

$$\mathbb{P}(\partial V(A)) \geq \mathbb{P}(\partial V(S)).$$

4.4 Hypercontractivity: The Bonami-Beckner Inequality

4.5 Influence Function

4.6 Monotone Sets

4.7 Threshold Phenomena

References

- Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.
- Lawrence C Evans. *Partial differential equations*, volume 19. American Mathematical Soc., 2010.
- Evarist Giné and Richard Nickl. *Mathematical foundations of infinite-dimensional statistical models*. Cambridge university press, 2021.
- Roman Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.
- Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge University Press, 2019.