

Lecture 3: Topology Review

Tianpei Xie

Jul. 8th., 2015

Contents

1	Review of Topology	2
1.1	Set Theory Basis	2
1.2	Topological Space	6
1.3	Limit Point and Closure	7
1.4	Subspace, Product and Quotient Topologies	7
1.4.1	Subspace Topology	7
1.4.2	Product Topology	8
1.4.3	Quotient Topology	9
1.5	Constructing Continuous Functions	10
1.6	Metric Topology	11
1.7	Connectedness and Compactness	13
1.7.1	Connectedness and Local Connectedness	13
1.7.2	Compactness and Local Compactness	14
1.8	Countability and Separability	16
1.8.1	Countability Axioms	16
1.8.2	Separability Axioms	17
1.9	Important Results and Theorems on Normal Space	18
1.10	Metrization	20
1.11	Nets and Convergence in Topological Space	20
2	Special Space	21
3	Summary of Preservation of Topological Properties	23

1 Review of Topology

1.1 Set Theory Basis

- **Definition** Given a set X , the collection of all subsets of X , denoted as 2^X , is defined as

$$2^X := \{E : E \subseteq X\}$$

- **Remark** The followings are basic operation on 2^X : For $A, B \in 2^X$,

1. **Inclusion**: $A \subseteq B$ if and only if $\forall x \in A, x \in B$.
2. **Union**: $A \cup B = \{x : x \in A \vee x \in B\}$.
3. **Intersection**: $A \cap B = \{x : x \in A \wedge x \in B\}$.
4. **Difference**: $A \setminus B = \{x : x \in A \wedge x \notin B\}$.
5. **Complement**: $A^c = X \setminus A = \{x : x \in X \wedge x \notin A\}$.
6. **Symmetric Difference**: $A \Delta B = (A \setminus B) \cup (B \setminus A) = \{x \in X : x \notin A \vee x \notin B\}$.

We have *deMorgan's laws*:

$$\left(\bigcup_{a \in A} U_a \right)^c = \bigcap_{a \in A} U_a^c, \quad \left(\bigcap_{a \in A} U_a \right)^c = \bigcup_{a \in A} U_a^c$$

- **Remark** Note that the following equality is useful:

$$A \Delta B = (A \cup B) \setminus (A \cap B)$$

- **Definition** A **rule of assignment** is a subset r of the cartesian product $C \times D$ of two sets, having the property that each element of C appears as the first coordinate **at most one ordered pair belonging to r** . Thus, a subset r of $C \times D$ is a *rule of assignment* if

$$[(c, d) \in r \text{ and } (c, d') \in r] \Rightarrow [d = d'].$$

Given a rule of assignment r , **the domain** of r is defined to be the *subset* of C consisting of *all first coordinates of elements* of r , and **the image** set of r is defined as the *subset* of D consisting of *all second coordinates of elements* of r .

A **function** f is a rule of assignment r , together with a set B that *contains the image set* of r .

The **pre-image** of f is defined as

$$f^{-1}(E) = \{x \in X : f(x) \in E\}.$$

- **Remark** The pre-image operation *commutes* with *all basic set operations*:

$$\begin{aligned}
A \subseteq B &\Rightarrow f^{-1}(A) \subseteq f^{-1}(B) \\
f^{-1}\left(\bigcup_{\alpha \in A} E_{\alpha}\right) &= \bigcup_{\alpha \in A} f^{-1}(E_{\alpha}) \\
f^{-1}\left(\bigcap_{\alpha \in A} E_{\alpha}\right) &= \bigcap_{\alpha \in A} f^{-1}(E_{\alpha}) \\
f^{-1}(A \setminus B) &= f^{-1}(A) \setminus f^{-1}(B) \\
f^{-1}(E^c) &= (f^{-1}(E))^c
\end{aligned}$$

- **Remark** The image operation *commutes* with only *inclusion and union operations*:

$$\begin{aligned}
A \subseteq B &\Rightarrow f(A) \subseteq f(B) \\
f\left(\bigcup_{\alpha \in A} E_{\alpha}\right) &= \bigcup_{\alpha \in A} f(E_{\alpha})
\end{aligned}$$

For the other operations:

$$\begin{aligned}
f\left(\bigcap_{\alpha \in A} E_{\alpha}\right) &\subseteq \bigcap_{\alpha \in A} f(E_{\alpha}) \\
f(A \setminus B) &\supseteq f(A) \setminus f(B)
\end{aligned}$$

- **Definition** A map $f : X \rightarrow Y$ is **surjective, or, onto**, if for every $y \in Y$, there exists a $x \in X$ such that $y = f(x)$. In set theory notation:

$$f : X \rightarrow Y \text{ is surjective} \Leftrightarrow f^{-1}(Y) \subseteq X.$$

A map $f : X \rightarrow Y$ is **injective**, if for every $x_1 \neq x_2 \in X$, their map $f(x_1) \neq f(x_2)$, or equivalently, $f(x_1) = f(x_2)$ only if $x_1 = x_2$.

If a map $f : X \rightarrow Y$ is both *surjective* and *injective*, we say f is a **bijective**, or there exists an *one-to-one correspondence* between X and Y . Thus $Y = f(X)$.

- **Remark**

$$\begin{aligned}
f^{-1}(f(B)) &\supseteq B, \quad \forall B \subseteq X \\
f(f^{-1}(E)) &\subseteq E, \quad \forall E \subseteq Y \\
f : X \rightarrow Y \text{ is surjective} &\Leftrightarrow f^{-1}(Y) \subseteq X. \\
&\Rightarrow f(f^{-1}(E)) = E. \\
f : X \rightarrow Y \text{ is injective} &\Rightarrow f^{-1}(f(B)) = B \\
&\Rightarrow f\left(\bigcap_{\alpha \in A} E_{\alpha}\right) = \bigcap_{\alpha \in A} f(E_{\alpha}) \\
&\Rightarrow f(A \setminus B) = f(A) \setminus f(B)
\end{aligned}$$

- **Proposition 1.1** *The following statements for composite functions are true:*

1. If f, g are both *injective*, then $g \circ f$ is *injective*.
2. If f, g are both *surjective*, then $g \circ f$ is *surjective*.
3. Every **injective** map $f : X \rightarrow Y$ can be written as $f = \iota \circ f_R$ where $f_R : X \rightarrow f(X)$ is a **bijective** map and ι is the **inclusion map**.
4. Every **surjective** map $f : X \rightarrow Y$ can be written as $f = f_p \circ \pi$ where $\pi : X \rightarrow (X/\sim)$ is a **quotient map** (projection $x \mapsto [x]$) for the equivalent relation $x \sim y \Leftrightarrow f(x) = f(y)$ and $f_p : (X/\sim) \rightarrow Y$ is defined as $f_p([x]) = f(x)$ **constant** in each coset $[x]$.
5. If $g \circ f$ is *injective*, then f is *injective*.
6. If $g \circ f$ is *surjective*, then g is *surjective*.

- **Definition** A **relation** on a set A is a subset R of the cartesian product $A \times A$.

If R is a relation on A , we use the notation xRy to mean the same thing as $(x, y) \in R$. We read it “ x is in the relation R to y .”

- **Remark** A **rule of assignment** r for a function $f : A \rightarrow A$ is also a subset of $A \times A$. But it is a subset of a *very special kind*: namely, one such that **each element** of A appears as the **first coordinate** of an element of r **exactly once**. **Any subset** of $A \times A$ is a relation on A .

- **Definition** **An equivalence relation** on X is a relation R on X such that

1. (**Reflexivity**): xRx for all $x \in X$;
2. (**Symmetry**): xRy if and only if yRx for all $x, y \in X$;
3. (**Transitivity**): xRy and yRz then xRz for all $x, y, z \in X$.

We usually denote the equivalence relation R as \sim .

- **Definition** (**Equivalence Class**)

The equivalence class of an element x is denoted as $[x] := \{y \in X : xRy\}$.

- **Definition** A relation C on a set A is called **an order relation** (or **a simple order**, or **a linear order**) if it has the following properties:

1. (**Comparability**) For every x and y in A for which $x \neq y$, either xCy or yCx .
2. (**Nonreflexivity**) For no x in A does the relation xCx hold.
3. (**Transitivity**) If xCy and yCz , then xCz .

We denote order relation as $>$ or $<$. We shall use the notation $x \leq y$ to stand for the statement “either $x < y$ or $x = y$ ”; and we shall use the notation $y > x$ to stand for the statement “ $x < y$.” We write $x < y < z$ to mean “ $x < y$ and $y < z$ ”

- **Definition** Suppose that A is a set ordered by the relation $<$. Let A_0 be a subset of A . We say that the element b is the largest element of A_0 if $b \in A_0$ and $x \leq b$ for every $x \in A_0$.

Similarly, we say that a is the smallest element of A_0 if $a \in A_0$ and if $a \leq x$ for every $x \in A_0$.

- **Remark** It is easy to see that a set has *at most one* largest element and *at most one* smallest element.

- **Definition (The Upper Bound and The Supremum of Subset)**

We say that *the subset A_0 of A is **bounded above*** if there is *an element b of A such that $x \leq b$ for every $x \in A_0$* ; the element $b \in A$ is called **an upper bound for A_0** .

If *the set of all upper bounds for A_0 has a **smallest element***, that element is called **the least upper bound**, or **the supremum**, of A_0 . It is denoted by $\sup A_0$, it may or may not belong to A_0 . If it *does*, it is **the largest element** of A_0 .

- **Definition (The Lower Bound and The Infimum of Subset)**

Similarly, we say that *the subset A_0 of A is **bounded below*** if there is *an element a of A such that $a \leq x$ for every $x \in A_0$* ; the element $a \in A$ is called **a lower bound for A_0** .

If *the set of all lower bounds for A_0 has a **largest element***, that element is called **the greatest lower bound**, or **the infimum**, of A_0 . It is denoted by $\inf A_0$, it may or may not belong to A_0 . If it *does*, it is **the smallest element** of A_0 .

- **Definition (The Least Upper Bound Property and The Greatest Lower Bound Property)**

An ordered set A is said to have **the least upper bound property** if *every nonempty subset A_0 of A that is bounded above has a least upper bound*.

Analogously, the set A is said to have **the greatest lower bound property** if *every nonempty subset A_0 of A that is bounded below has a greatest lower bound*.

- **Definition (Well-Ordered Set)**

A set A with an order relation $<$ is said to be ***well-ordered*** if *every nonempty subset of A has a **smallest element***.

- **Definition (Strict Partial Order)**

Given a set A , a relation \prec on A is called a **strict partial order** on A if it has the following two properties;

1. (**Nonreflexivity**) The relation $a \prec a$ never holds.
2. (**Transitivity**) If $a \prec b$ and $b \prec c$, then $a \prec c$.

Moreover, suppose that we define $a \preceq b$ either $a \prec b$ or $a = b$. Then the relation \preceq is called **a partial order** on A .

- **Definition (Upper Bound and Maximal Element for Strict Partial Order)**

Let A be a set and let \prec be a *strict partial order* on A . If B is a subset of A , **an upper bound** on B is an element c of A such that for every b in B , either $b = c$ or $b \prec c$.

A maximal element of A is an element m of A such that for **no element a of A does the relation $m \prec a$ hold**.

- **Theorem 1.2 (Zorn's Lemma).** [Munkres, 2000]

*Let A be a set that is **strictly partially ordered**. If every **simply ordered subset** of A has an **upper bound** in A , then A has a **maximal element**.*

- **Principle 1.3 (The Axiom of Choice).**

If $\{X_\alpha\}_{\alpha \in A}$ is a nonempty collection of nonempty sets, then $\prod_{\alpha \in A} X_\alpha$ is non-empty.

- **Corollary 1.4** If $\{X_\alpha\}_{\alpha \in A}$ is a **disjoint** collection of nonempty sets, there is a set $Y \subset \bigcup_{\alpha \in A} X_\alpha$ such that $Y \cap X_\alpha$ contains **precisely one element** for each $\alpha \in A$.

1.2 Topological Space

- **Definition** Let X be a set. A **topology** on X is a collection \mathcal{T} of subsets of X , called **open subsets**, satisfying

1. X and \emptyset are *open*.
2. The **union** of **any family** of open subsets is open.
3. The **intersection** of **any finite family** of open subsets is open.

A pair (X, \mathcal{T}) consisting of a set X together with a topology \mathcal{T} on X is called a **topological space**.

- **Definition** A map $F : X \rightarrow Y$ is said to be **continuous** if for every open subset $U \subseteq Y$, the **preimage** $F^{-1}(U)$ is **open** in X .
- **Definition** A **continuous bijective** map $F : X \rightarrow Y$ with **continuous inverse** is called a **homeomorphism**. If there exists a *homeomorphism* from X to Y , we say that X and Y are **homeomorphic**.
- **Definition** Suppose X is a topological space. A collection \mathcal{B} of open subsets of X is said to be a **basis** for the topology of X (plural: **bases**) if every open subset of X is the *union of some collection of elements of \mathcal{B}* .

More generally, suppose X is merely a set, and \mathcal{B} is a collection of *subsets* of X satisfying the following conditions:

1. $X = \bigcup_{B \in \mathcal{B}} B$.
2. If $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, then there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq B_1 \cap B_2$.

Then the collection of **all unions** of elements of \mathcal{B} is a topology on X , called **the topology generated by \mathcal{B}** , and \mathcal{B} is a **basis** for this topology.

- **Lemma 1.5 (Obtaining Basis from Given Topology).** [Munkres, 2000]
Let X be a topological space. Suppose that \mathcal{C} is a collection of open sets of X such that for each open set U of X and each x in U , there is an element C of \mathcal{C} such that $x \in C \subset U$. Then \mathcal{C} is a basis for the topology of X .
- **Lemma 1.6 (Topology Comparison via Bases).** [Munkres, 2000]
Let \mathcal{B} and \mathcal{B}' be bases for the topologies \mathcal{T} and \mathcal{T}' , respectively, on X . Then the following are equivalent:
 1. \mathcal{T}' is **finer** than \mathcal{T} .
 2. For each $x \in X$ and each basis element $B \in \mathcal{B}$ containing x , there is a basis element $B' \in \mathcal{B}'$ such that $x \in B' \subset B$.

- **Definition (Subbasis)**
A **subbasis** \mathcal{S} for a topology on X is a collection of subsets of X whose union equals X .

The topology generated by the *subbasis* \mathcal{S} is defined to be the collection \mathcal{T} of **all unions of finite intersections of elements of \mathcal{S}** .

- **Remark (*Basis from Subbasis*)**

For a *subbasis* \mathcal{S} , the collection \mathcal{B} of **all finite intersections** of elements of \mathcal{S} is a **basis**,

1.3 Limit Point and Closure

- **Definition** A subset A of a topological space X is said to be **closed** if the set $X \setminus A$ is *open*.
- **Definition** Given a subset A of a topological space X , **the interior of A** is defined as *the union of all open sets contained in A* , and **the closure of A** is defined as *the intersection of all closed sets containing A* .

The interior of A is denoted by $\text{Int } A$ or by $\overset{\circ}{A}$ and **the closure of A** is denoted by $\text{Cl } A$ or by \bar{A} . Obviously $\overset{\circ}{A}$ is an *open set* and \bar{A} is a *closed set*; furthermore,

$$\overset{\circ}{A} \subseteq A \subseteq \bar{A}.$$

If A is **open**, $A = \overset{\circ}{A}$; while if A is **closed**, $A = \bar{A}$.

- **Proposition 1.7 (*Characterization of Closure in terms of Basis*)** [Munkres, 2000]
Let A be a subset of the topological space X .

1. Then $x \in \bar{A}$ if and only if every **open set U containing x intersects A** .
2. Supposing the topology of X is given by a **basis**, then $x \in \bar{A}$ if and only if every **basis element B containing x intersects A** .

- **Remark** We can say “ U is a **neighborhood of x** ” if “ U is an open set containing x ”.

- **Definition (*Limit Point*)**

If A is a subset of the topological space X and if x is a point of X , we say that x is a **limit point** (or “**cluster point**,” or “**point of accumulation**”) of A if **every neighborhood of x intersects A in some point other than x itself**.

Said differently, x is a **limit point** of A if it belongs to **the closure of $A \setminus \{x\}$** . The point x may lie in A or not; for this definition it does not matter.

- **Theorem 1.8 (*Decomposition of Closure*)**

Let A be a subset of the topological space X ; let A' be the set of **all limit points of A** . Then

$$\bar{A} = A \cup A'.$$

- **Corollary 1.9** A subset of a topological space is **closed** if and only if it contains all its **limit points**.

1.4 Subspace, Product and Quotient Topologies

1.4.1 Subspace Topology

- **Definition** If X is a topological space and $S \subseteq X$ is an arbitrary subset, we define **the subspace topology** on S (sometimes called **the relative topology**) by declaring a subset

$U \subseteq S$ to be *open in S* if and only if there exists an open subset $V \subseteq X$ such that $U = V \cap S$.

Any subset of X endowed with the subspace topology is said to be **a subspace of X** .

- **Lemma 1.10 (*Basis of Subspace Topology*)**

If \mathcal{B} is a basis for the topology of X then the collection

$$\mathcal{B}_S = \{B \cap S : B \in \mathcal{B}\}$$

is a **basis** for the subspace topology on $S \subset X$.

- **Proposition 1.11** *Let Y be a subspace of X . If A is closed in Y and Y is closed in X , then A is closed in X .*

- **Proposition 1.12 (*Closure in Subspace Topology*)**

Let Y be a subspace of X ; let A be a subset of Y ; let \bar{A} denote the closure of A in X . Then the closure of A in Y equals $\bar{A} \cap Y$.

1.4.2 Product Topology

- **Definition (*J-tuples*)**

Let J be an index set. Given a set X , we define a ***J-tuple*** of elements of X to be a function $x : J \rightarrow X$. If α is an element of J , we often denote **the value of X at α** by X_α rather than $x(\alpha)$; we call it **the α -th coordinate** of x . And we often denote the function x itself by the symbol

$$(x_\alpha)_{\alpha \in J}$$

which is as close as we can come to a “*tuple notation*” for an arbitrary index set J . We denote **the set of all J -tuples** of elements of X by X^J .

- **Definition (*Arbitrary Cartesian Products*)**

Let $\{A_\alpha\}_{\alpha \in J}$ be an *indexed* family of sets; let $X = \bigcup_{\alpha \in J} A_\alpha$. The **cartesian product** of this indexed family, denoted by

$$\prod_{\alpha \in J} A_\alpha$$

is defined to be the set of all J -tuples $(x_\alpha)_{\alpha \in J}$ of elements of X such that $x_\alpha \in A_\alpha$ for each $\alpha \in J$. That is, it is the set of all functions

$$x : J \rightarrow \bigcup_{\alpha \in J} A_\alpha$$

such that $x(\alpha) \in A_\alpha$ for each $\alpha \in J$.

- **Definition (*Projection Mapping or Coordinate Projection*)**

Let

$$\pi_\beta : \prod_{\alpha \in J} X_\alpha \rightarrow X_\beta$$

be the function assigning to each element of the product space its β -th coordinate,

$$\pi_\beta((x_\alpha)_{\alpha \in J}) = x_\beta;$$

it is called **the projection mapping** associated with the index β .

- **Definition (*Product Topology*)**

Let \mathcal{S}_β denote the collection

$$\mathcal{S}_\beta = \left\{ \pi_\beta^{-1}(U_\beta) : U_\beta \text{ open in } X_\beta \right\},$$

and let \mathcal{S} denote *the union of these collections*,

$$\mathcal{S} = \bigcup_{\beta \in J} \mathcal{S}_\beta.$$

The topology generated by *the subbasis* \mathcal{S} is called *the product topology*. In this topology $\prod_{\alpha \in J} X_\alpha$ is called *a product space*.

- **Remark (*Product Topology = Weak Topology by Coordinate Projections*)**

The product topology on $\prod_{\alpha \in J} X_\alpha$ is *the weak topology* generated by a family of projection mappings $(\pi_\beta)_{\beta \in J}$. It is *the coarsest (weakest) topology such that $(\pi_\beta)_{\beta \in J}$ are continuous*.

A typical element of the basis from the product topology is *the finite intersection of subbasis* where the *index is different*:

$$\pi_{\beta_1}^{-1}(V_{\beta_1}) \cap \dots \cap \pi_{\beta_n}^{-1}(V_{\beta_n})$$

Thus a *neighborhood* of x in *the product topology* is

$$N(x) = \{(x_\alpha)_{\alpha \in J} : x_{\beta_1} \in V_{\beta_1}, \dots, x_{\beta_n} \in V_{\beta_n}\}$$

where there is *no restriction* for $\alpha \in \{\beta_1, \dots, \beta_n\}$.

Note that for *the box topology*, a neighborhood of x is

$$N_b(x) = \{(x_\alpha)_{\alpha \in J} : x_\alpha \in U_\alpha, \forall \alpha \in J\} \subset N(x)$$

Thus *the box topology* is *finer* than *the product topology*. Moreover, for *finite product* $\prod_{\alpha=1}^n X_\alpha$, the box topology and the product topology is the *same*.

- **Definition** If X and Y are topological spaces, a continuous injective map $F : X \rightarrow Y$ is called a *topological embedding* if it is a *homeomorphism* onto its image $F(X) \subseteq Y$ in the subspace topology.

1.4.3 Quotient Topology

- **Definition (*Quotient Map*)**

Let X and Y be topological spaces; let $\pi : X \rightarrow Y$ be a *surjective map*. The map π is said to be *a quotient map* provided a subset U of Y is *open* in Y *if and only if* $\pi^{-1}(U)$ is *open* in X .

- **Remark (*Quotient Map = Strong Continuity*)**

The condition of quotient map is *stronger* than continuity (it is called *strong continuity* in some literature).

$$\text{continuity : } U \text{ is open in } Y \Rightarrow \pi^{-1}(U) \text{ is open in } X$$

$$\text{open map : } \pi(V) \text{ is open in } Y \Leftarrow V \text{ is open in } X$$

$$\text{quotient map : } U \text{ is open in } Y \Leftrightarrow \pi^{-1}(U) \text{ is open in } X$$

An equivalent condition is to require that a subset A of K be **closed** in Y if and only if $\pi^{-1}(A)$ is **closed** in X . Equivalence of the two conditions follows from equation

$$\pi^{-1}(Y \setminus B) = X \setminus \pi^{-1}(B).$$

- **Definition (Saturated Set and Fiber)**

If $\pi : X \rightarrow Y$ is a **surjective map**, a subset $U \subseteq X$ is said to be **saturated** with respect to π if U contains every set $\pi^{-1}(\{y\})$ that it **intersects**. Thus U is **saturated** if it equals to the **entire preimage** of its **image**: $U = \pi^{-1}(\pi(U))$.

Given $y \in Y$, the **fiber** of π over y is the set $\pi^{-1}(\{y\})$.

- **Definition (Quotient Map via Saturated Set)**

A surjective map $\pi : X \rightarrow Y$ is a **quotient map** if π is **continuous** and π maps **saturated open sets** of X to **open sets** of Y (or **saturated closed sets** of X to **closed sets** of Y).

- **Definition (Open Map and Closed Map)**

A map $f : X \rightarrow Y$ (continuous or not) is said to be an **open map** if for every **open** subset $U \subseteq X$, the image set $f(U)$ is **open** in Y , and a **closed map** if for every **closed** subset $K \subseteq X$, the image $f(K)$ is **closed** in Y .

- **Definition (Quotient Topology)**

If X is a space and A is a set and if $\pi : X \rightarrow A$ is a **surjective** map, then there exists **exactly one topology** \mathcal{T} on A relative to which π is a quotient map; it is called **the quotient topology** induced by π .

- **Definition (Quotient Space)**

Suppose X is a topological space and \sim is an **equivalence relation** on X . Let X/\sim denote **the set of equivalence classes** in X , and let $\pi : X \rightarrow X/\sim$ be the **natural projection** sending each **point** to its **equivalence class**. Endowed with **the quotient topology** determined by π , the space X/\sim is called **the quotient space** (or **identification space**) of X determined by π .

1.5 Constructing Continuous Functions

- **Proposition 1.13 (Rules for Constructing Continuous Functions).** [Munkres, 2000]

Let X , Y , and Z be topological spaces.

1. **(Constant Function)** If $f : X \rightarrow Y$ maps all of X into the **single point** y_0 of Y , then f is **continuous**.
2. **(Inclusion)** If A is a subspace of X , the **inclusion function** $\iota : A \xrightarrow{X}$ is **continuous**.
3. **(Composites)** If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous, then the map $g \circ f : X \rightarrow Z$ is **continuous**.
4. **(Restricting the Domain)** If $f : X \rightarrow Y$ is **continuous**, and if A is a subspace of X , then **the restricted function** $f|_A : A \rightarrow Y$ is **continuous**.
5. **(Restricting or Expanding the Range)** Let $f : X \rightarrow Y$ be **continuous**. If Z is a **subspace** of Y containing the **image** set $f(X)$, then the function $g : X \rightarrow Z$ obtained

by restricting the range of f is **continuous**. If Z is a space having Y as a **subspace**, then the function $h : X \rightarrow Z$ obtained by **expanding the range** of f is **continuous**.

6. (**Local Formulation of Continuity**) The map $f : X \rightarrow Y$ is **continuous** if X can be written as the **union of open sets** U_α such that $f|_{U_\alpha}$ is **continuous** for each α .

- **Theorem 1.14 (The Pasting Lemma / Gluing Lemma)**. [Munkres, 2000]
Let $X = A \cup B$, where A and B are **closed** in X . Let $f : A \rightarrow Y$ and $g : B \rightarrow Y$ be **continuous**. If $f(x) = g(x)$ for **every** $x \in A \cap B$, then f and g combine to give a **continuous function** $h : X \rightarrow Y$, defined by setting $h|_A = f$, and $h|_B = g$.

- **Remark** The set A and B can be open sets, and the gluing lemma comes “**Local Formulation of Continuity**”.

- **Remark** Notice the condition for the *gluing lemma*:

1. The domain X is a union of two **closed sets (or open sets)** A and B
2. The two functions f and g are **continuous** each of closed domain sets, respectively
3. f and g **agree on the intersection** of two sets $A \cap B$.

- **Theorem 1.15 (Maps into Products)**. [Munkres, 2000]
Let $f : A \rightarrow X \times Y$ be given by the equation

$$f(a) = (f_1(a), f_2(a)).$$

Then f is **continuous** if and only if the functions

$$f_1 : A \rightarrow X \quad \text{and} \quad f_2 : A \rightarrow Y$$

are **continuous**. The maps f_1 and f_2 are called the coordinate functions of f .

1.6 Metric Topology

- **Definition (Metric Space)**

A **metric space** is a set M and a real-valued function $d(\cdot, \cdot) : M \times M \rightarrow \mathbb{R}$ which satisfies:

1. (**Non-Negativity**) $d(x, y) \geq 0$
2. (**Definiteness**) $d(x, y) = 0$ if and only if $x = y$
3. (**Symmetric**) $d(x, y) = d(y, x)$
4. (**Triangle Inequality**) $d(x, z) \leq d(x, y) + d(y, z)$

The function d is called a **metric** on M . The metric space M equipped with metric d is denoted as (M, d) .

- **Definition (ϵ -Ball)**

Given a metric d on X , the number $d(x, y)$ is often called the **distance** between x and y in the metric d . Given $\epsilon > 0$, consider the set

$$B_d(x, \epsilon) = \{y : d(x, y) < \epsilon\}$$

of all points y whose distance from x is less than ϵ . It is called ***the ϵ -ball centered at x*** . Sometimes we omit the metric d from the notation and write this ball simply as $B(x, \epsilon)$, when no confusion will arise.

- **Definition (*Metric Topology*)**

If d is a *metric* on the set X , then *the collection of all ϵ -balls $B_d(x, \epsilon)$, for $x \in X$ and $\epsilon > 0$, is a **basis** for a topology on X , called ***the metric topology induced by d*** .*

- **Definition (*Metrizability*)**

If X is a topological space, X is said to be ***metrizable*** if *there exists a metric d on the set X that induces the topology of X* . A ***metric space*** is a metrizable space X together with a specific metric d that gives the topology of X .

- **Remark (*Metrizability as Inverse Problem*)**

Given a *metric* d on X , we can generate a *metric topology* using ϵ -balls as basis. ***Conversely, given a topology \mathcal{T} on X , is \mathcal{T} a metric topology for some unknown metric d ?***

This is the question that ***the metrization theory*** is trying to answer.

- **Theorem 1.16 (*ϵ - δ Definition of Continuous Function in Metric Space*)**. [Munkres, 2000]

Let $f : X \rightarrow Y$; let X and Y be ***metrizable*** with metrics d_x and d_y , respectively. Then ***continuity*** of f is ***equivalent*** to the requirement that given $x \in X$ and given $\epsilon > 0$, there exists $\delta > 0$ such that

$$d_x(x, y) < \delta \Rightarrow d_y(f(x), f(y)) < \epsilon.$$

- **Remark** To use ϵ - δ definition, both ***domain*** and ***codomain*** need to be ***metrizable***.

- **Lemma 1.17 (*The Sequence Lemma*)**. [Munkres, 2000]

Let X be a topological space; let $A \subseteq X$. If there is a sequence of points of A ***converging*** to x , then $x \in \bar{A}$; the ***converse*** holds if X is ***metrizable***.

- **Proposition 1.18** Let $f : X \rightarrow Y$. If the function f is ***continuous***, then for every ***convergent*** sequence $x_n \rightarrow x$ in X , the sequence $f(x_n)$ ***converges*** to $f(x)$. The ***converse*** holds if X is ***metrizable***.

- **Remark** To show the converse part, i.e. “if $x_n \rightarrow x \Rightarrow f(x_n) \rightarrow f(x)$ then f is continuous”, we just need the space X to be ***first countable***. That is, at each point x , there is a ***countable collection*** $(U_n)_{n \in \mathbb{Z}_+}$ of ***neighborhoods*** of x such that any neighborhood U of x contains at least one of the sets U_n .

- **Proposition 1.19 (*Arithmetic Operations of Continuous Functions*)**.

If X is a topological space, and if $f, g : X \rightarrow Y$ are continuous functions, then $f + g$, $f - g$, and $f \cdot g$ are continuous. If $g(x) \neq 0$ for all x , then f/g is continuous.

- **Definition (*Uniform Convergence*)**

Let $f_n : X \rightarrow Y$ be a sequence of functions from the ***set*** X to ***the metric space*** Y . Let d be the metric for Y . We say that the sequence (f_n) ***converges uniformly*** to the function $f : X \rightarrow Y$ if given $\epsilon > 0$, there exists an integer N such that

$$d(f_n(x), f(x)) < \epsilon$$

for all $n > N$ and ***all x in X*** .

- **Theorem 1.20 (Uniform Limit Theorem).** [Munkres, 2000]

Let $f_n : X \rightarrow Y$ be a sequence of **continuous** functions from the **topological** space X to the **metric** space Y . If (f_n) converges **uniformly** to f , then f is **continuous**.

1.7 Connectedness and Compactness


Remark *Connectedness* and *compactness* are basic **topological properties**. Both of them are defined based on a collection of open subsets.

1. **Connectedness** is a **global topological property**: a topological space is *connected* if it cannot be partitioned by two *disjoint nonempty open* subsets. *Connectedness* reveals the information of **entire space not just within a neighborhood**. Connectedness is **compatible** with the **continuity** of functions as it implies **the intermediate value theorem**, which in turn, can be used to construct *inverse function*. Moreover, *connectedness* defines **an equivalence relationship** which allows a **partition** of the space into **components**.
2. **Connectedness** is a **local-to-global topological property**: a topological space is *compact* if every open cover have a finite sub-cover. Using **finite sub-cover**, **local properties** defined *within each neighborhood* can be **generalized globally** to entire space. Concept of functions that are closely related to compactness is **the uniform continuity** and **the maximum value theorem**. The compactness allows us to drop dependency on each individual point x .

Compared to *connectedness*, **compactness** is usually a **strong condition** on the topological space.

1.7.1 Connectedness and Local Connectedness

- **Concepts Related to Connectedness**



- **Definition (Separation and Connectedness)**

Let X be a topological space. A **separation** of X is a pair U, V of **disjoint nonempty open subsets** of X whose union is X .

The space X is said to be **connected** if there *does not exist* a separation of X .

- **Definition** Equivalently, X is **connected** if and only if the only subsets of X that are **both open and closed** are \emptyset and X itself.

- **Remark (*Proof of Connectedness*)**

As the definition suggests, the proof of connectedness is done *by contradiction*. One first assume that the set X has a *separation*; it can be separated into two *disjoint nonempty open* sets such that $X = A \cup B$. Then we proof by contradiction using *existing connectedness conditions* and the *property of open subsets (basis, continuity etc.)*.


- **Definition** Recall that a topological space X is

- **connected** if there do not exist two *disjoint, nonempty, open* subsets of X whose union is X ;
- **path-connected** if every pair of points in X can be *joined by a path* in X , and
- *locally path-connected* if X has a *basis* of *path-connected open subsets*.

- **Definition** A *maximal connected subset* of X (i.e., a connected subset that is not properly contained in any larger connected subset) is called a **component** (or **connected component**) of X .

1.7.2 Compactness and Local Compactness

- **Concepts Related to Compactness**



- **Definition (*Covering of Set and Open Covering of Topological Set*)**

A collection \mathcal{A} of subsets of a space X is said to cover X , or to be a covering of X , if the union of the elements of \mathcal{A} is equal to X .

It is called an open covering of X if its elements are *open subsets* of X .

- **Definition (*Compactness*)**

A topological space X is said to be compact if *every open covering* \mathcal{A} of X contains a *finite subcollection* that also *covers* X .

- To prove *compactness*, the following property is useful:

Definition (*Finite Intersection Property*)

A collection \mathcal{C} of subsets of X is said to have the finite intersection property if for *every*

finite subcollection

$$\{C_1, \dots, C_n\}$$

of \mathcal{C} , the **intersection** $C_1 \cap \dots \cap C_n$ is **nonempty**.

- **Proposition 1.21 (Equivalent Definition of Compactness)** [Munkres, 2000]
Let X be a topological space. Then X is **compact if and only if** for every collection \mathcal{C} of **closed** sets in X having the **finite intersection property**, the intersection $\bigcap_{C \in \mathcal{C}} C$ of all the elements of \mathcal{C} is **nonempty**.
- **Definition** If X and Y are topological spaces, a map $F : X \rightarrow Y$ (continuous or not) is said to be **proper** if for every **compact** set $K \subseteq Y$, the **preimage** $F^{-1}(K)$ is **compact**.
- **Definition** A topological space X is said to be **locally compact** if every point has a **neighborhood** contained in a **compact subset** of X .

A subset of X is said to be **precompact** in X if its **closure** in X is **compact**.

- If X is not a compact Hausdorff space, then under what conditions is X homeomorphic with a **subspace** of a compact Hausdorff space ?

Theorem 1.22 (Unique One-Point Compactification) [Munkres, 2000]

Let X be a space. Then X is **locally compact Hausdorff** if and only if there exists a space Y satisfying the following conditions:

1. X is a subspace of Y .
2. The set $Y \setminus X$ consists of a **single point** (which is the limit point of X).
3. Y is a **compact Hausdorff** space.

If Y and Y' are two spaces satisfying these conditions, then there is a **homeomorphism** of Y with Y' that equals **the identity map** on X .

- **Definition (One-Point Compactification)**
If Y is a **compact Hausdorff** space and X is a proper subspace of Y whose **closure** equals Y , then Y is said to be a **compactification** of X .

If $Y \setminus X$ equals a **single point**, then Y is called **the one-point compactification** of X .

- **Proposition 1.23 (Locally Compact Hausdorff = Precompact Basis)** [Munkres, 2000]
Let X be a **Hausdorff** space. Then X is **locally compact if and only if** given x in X , and given a neighborhood U of x , there is a neighborhood V of x such that \bar{V} is **compact** and $\bar{V} \subseteq U$.
- **Corollary 1.24 (Closed or Open Subspace)** [Munkres, 2000]
Let X be locally compact Hausdorff; let A be a subspace of X . If A is **closed** in X or **open** in X , then A is locally compact.
- **Corollary 1.25** [Munkres, 2000]
A space X is **homeomorphic** to an **open** subspace of a **compact Hausdorff** space **if and only if** X is **locally compact Hausdorff**.
- For a **Hausdorff** space X , the following are equivalent:


1. X is **locally compact**.

2. Each point of X has a **precompact** neighborhood.
 3. X has a basis of **precompact** open subsets.
- **Theorem 1.26 (Tychonoff Theorem).** [Munkres, 2000]
An arbitrary product of compact spaces is **compact** in the product topology.

1.8 Countability and Separability

1.8.1 Countability Axioms

- **Concepts Related to Countability Axioms**



- **Definition (Countability)**
A topological space X is said to be
 1. **first-countable** if there is a **countable neighborhood basis** at each point,
 2. **second-countable** if there is a **countable basis** for its topology.
- **Proposition 1.27 (Limit Point Detected by Convergent Sequence)** [Munkres, 2000]
Let X be a topological space.
 1. Let A be a subset of X . If there is a sequence of points of A converging to x , then $x \in \bar{A}$; the **converse** holds if X is **first-countable**.
 2. Let $f : X \rightarrow Y$. If f is continuous, then for every convergent sequence $x_n \rightarrow x$ in X , the sequence $f(x_n)$ converges to $f(x)$. The **converse** holds if X is **first-countable**.
- **Definition (Dense Subset)**
A subset A of a space X is said to be **dense** in X if $\bar{A} = X$. (That is, every point in X is a limit point of A .)
- **Definition (Separability)**
A topological space X is called **separable** if and only if it has a **countable dense set**.
- **Definition (Lindelöf Space)**
A space for which every open covering contains a **countable subcovering** is called a **Lindelöf space**.
- **Proposition 1.28 (Properties of Second-Countability)** [Munkres, 2000]
Suppose that X has a **countable basis**. Then:
 1. Every **open covering** of X contains a **countable subcollection** covering X . (X is **Lindelöf space**)


2. There exists a **countable** subset of X that is **dense** in X . (X is **separable**)

• **Proposition 1.29** (*Metric Space Countability and Separability*)

1. Every **metric space** is **first countable**.
2. A **metric space** is **second countable** if and only if it is **separable**.
3. Any **second countable** topological space is **separable**.

1.8.2 Separability Axioms

• *Concepts Related to Separation Axioms*



• **Definition** (*Separation Axioms*)

1. A topological space is called a **T_1 space** if and only if for all x and y , $x \neq y$, there is an **open set** U with $y \in U$, $x \notin U$.

Equivalently, a space is T_1 if and only if $\{x\}$ is **closed** for each x .

2. A topological space is called **Hausdorff** (or T_2) if and only if for all x and y , $x \neq y$, there are **open sets** U , V such that $x \in U$, $y \in V$, and $U \cap V = \emptyset$.
3. A topological space is called **regular** (or T_3) if and only if it is T_1 and for all x and C , **closed**, with $x \notin C$, there are **open sets** U , V such that $x \in U$, $C \subset V$, and $U \cap V = \emptyset$.

Equivalently, a space is T_3 if the **closed neighborhoods** of any point are a **neighborhood base**.

4. A topological space is called **normal** (or T_4) if and only if it is T_1 and for all C_1 , C_2 , **closed**, with $C_1 \cap C_2 = \emptyset$, there are **open sets** U , V with $C_1 \subset U$, $C_2 \subset V$, and $U \cap V = \emptyset$.

• **Proposition 1.30**

$$T_4 \Rightarrow T_3 \Rightarrow T_2 \Rightarrow T_1$$

- **Proposition 1.31** (*Limit Point in T_1 Axiom*). [Munkres, 2000]
Let X be a space satisfying the T_1 axiom; let A be a subset of X . Then the point x is a **limit point** of A if and only if every **neighborhood** of x contains **infinitely many points** of A .
- **Proposition 1.32** (*Limit Point is Unique in Hausdorff Space*). [Munkres, 2000]
If X is a **Hausdorff space**, then a sequence of points of X **converges to at most one point** of X .
- **Lemma 1.33** Let X be a topological space. Let one-point sets in X be closed.
 1. X is **regular** if and only if given a point x of X and a neighborhood U of x , there is a **neighborhood** V of x such that $\bar{V} \subseteq U$.
 2. X is **normal** if and only if given a **closed** set A and an open set U containing A , there is an **open set** V containing A such that $\bar{V} \subseteq U$.
- **Proposition 1.34** [Munkres, 2000]
Every **locally compact Hausdorff** space is **regular**.

1.9 Important Results and Theorems on Normal Space

- **Theorem 1.35** (*Regular + Second-Countable \Rightarrow Normal*)[Munkres, 2000]
Every **regular space with a countable basis** is **normal**.
- **Proposition 1.36** (*Regular + Lindelöf \Rightarrow Normal*)[Munkres, 2000]
Every **regular Lindelöf space** is **normal**.
- **Theorem 1.37** [Munkres, 2000]
Every **metrizable space** is **normal**.
- **Theorem 1.38** [Munkres, 2000, Reed and Simon, 1980]
Every **compact Hausdorff space** X is **normal**.
- **Theorem 1.39** [Munkres, 2000]
Every **well-ordered** set X is **normal** in the **order topology**.
- **Theorem 1.40** (*Urysohn Lemma*). [Munkres, 2000]
Let X be a **normal space**; let A and B be **disjoint closed subsets** of X . Let $[a, b]$ be a **closed interval** in the real line. Then there exists a **continuous map**

$$f : X \rightarrow [a, b]$$

such that $f(x) = a$ for **every** x in A , and $f(x) = b$ for **every** x in B .

- **Definition** (*Separation by Continuous Function*)
If A and B are two subsets of the topological space X , and if there is a **continuous function** $f : X \rightarrow [0, 1]$ such that $f(A) = \{0\}$ and $f(B) = \{1\}$, we say that A and B can be **separated by a continuous function**.
- **Definition** (*Completely Regular*)
A space X is **completely regular** if **one-point sets** are **closed** in X and if for each point x_0 and each **closed** set A not containing x_0 , there is a **continuous function** $f : X \rightarrow [0, 1]$ such that $f(x_0) = 1$ and $f(A) = \{0\}$.

- **Remark**

normal \Rightarrow completely regular \Rightarrow regular

- **Theorem 1.41 (Urysohn Lemma, Locally Compact Version).** [Folland, 2013]
Let X be a **locally compact Hausdorff** space and $K \subseteq U \subseteq X$ where K is **compact** and U is **open**. Then there exists a **continuous** map

$$f : X \rightarrow [0, 1]$$

such that $f(x) = 1$ for **every** $x \in K$, and $f(x) = 0$ for x outside a **compact subset** of U .

- **Corollary 1.42** [Folland, 2013]
Every **locally compact Hausdorff** space is **completely regular**.

- **Proposition 1.43** [Reed and Simon, 1980]
Let $\mathcal{C}(X)$ be the set of all complex-valued **continuous functions** on X and $\mathcal{C}_{\mathbb{R}}(X) \subseteq \mathcal{C}(X)$ be the set of all **real-valued continuous functions** on X . Also define $\mathcal{C}^b(X)$ as the set of all complex-valued **bounded continuous functions** on X . When X is a **compact space**, $\mathcal{C}^b(X) = \mathcal{C}(X)$. Define the norm as

$$\|f\|_{\infty} = \sup_{x \in X} |f(x)|.$$

Then for **compact Hausdorff space** X , $\mathcal{C}(X)$ is a (complex) **Banach space** and $\mathcal{C}(X)$ is a (real) **Banach space**.

- **Theorem 1.44 (Embedding Theorem).** [Munkres, 2000]
Let X be a space in which one-point sets are closed. Suppose that $\{f_{\alpha}\}_{\alpha \in J}$ is an indexed family of **continuous functions** $f_{\alpha} : X \rightarrow \mathbb{R}$ satisfying the requirement that for each point x_0 of X and each neighborhood U of x_0 , there is an index α such that f_{α} is **positive** at x_0 and **vanishes outside** U . Then the function $F : X \rightarrow \mathbb{R}^J$ defined by

$$F(x) = (f_{\alpha}(x))_{\alpha \in J}$$

is a **topological embedding** of X in \mathbb{R}^J . If f_{α} maps X into $[0, 1]$ for each α then F **embeds** X in $[0, 1]^J$.

- **Definition (Separation of Points From Closed Set by Continuous Functions)**
A family of **continuous functions** that satisfies the hypotheses of the embedding theorem above is said to **separate points from closed sets in** X .

The existence of such a family is readily seen to be *equivalent*, for a space X in which one-point sets are *closed*, to the requirement that X be *completely regular*.

- **Corollary 1.45 (Embedding Equivalent Definition of Completely Regular)** [Munkres, 2000]
A space X is **completely regular** if and only if it is **homeomorphic** to a subspace of $[0, 1]^J$ for some J .

- **Theorem 1.46 (Tietze Extension Theorem)** [Munkres, 2000, Reed and Simon, 1980]
Let X be a **normal space**; let A be a **closed subspace** of X .

1. Any **continuous** map of A into the **closed interval** $[a, b]$ of \mathbb{R} may be **extended** to a **continuous** map of **all of** X into $[a, b]$.
2. Any **continuous** map of A into \mathbb{R} may be **extended** to a **continuous** map of **all of** X into \mathbb{R} .

- **Theorem 1.47 (Tietze Extension Theorem, Locally Compact Version)** [Folland, 2013]
Let X be a **locally compact Hausdorff space**; let K be a **compact subspace** of X . If $f \in C(K)$ is a **continuous** map of K into \mathbb{R} , there exists a **continuous** extension $F \in C(X)$ of **all of** X into \mathbb{R} such that $F|_K = f$. Moreover, F may be taken to **vanish outside a compact set**.

1.10 Metrization

- **Theorem 1.48 (The Urysohn Metrization Theorem).** [Munkres, 1975, Folland, 2013]
Every **second countable normal space** is **metrizable**.

1.11 Nets and Convergence in Topological Space

- **Definition (Directed System of Index Set)**

A **directed system** is an **index set** I together with an **ordering** \prec which satisfies:

1. If $\alpha, \beta \in I$ then there exists $\gamma \in I$ so that $\gamma \succ \alpha$ and $\gamma \succ \beta$.
2. \prec is a **partial ordering**.

- **Definition (Net)**

A **net** in a topological space X is a mapping from a **directed system** I to X ; we denote it by $\{x_\alpha\}_{\alpha \in I}$

- **Remark (Net vs. Sequence)**

Net is a generalization and abstraction of **sequence**. The directed system I is **not necessarily countable**. So $\{x_\alpha\}_{\alpha \in I}$ may not be a countable sequence. A **sequence** is a **net with countable index set** $I \subseteq \mathbb{N}$. The directed system can be any set e.g. a graph.

- **Definition** If $P(\alpha)$ is a **proposition** depending on an **index** α in a **directed set** I we say $P(\alpha)$ is **eventually true** if there is a β in I with $P(\alpha)$ **true** if for all $\alpha \succ \beta$.

We say $P(\alpha)$ is **frequently true** if it is **not eventually false**, that is, if for any β there exists an $\alpha \succ \beta$ with $P(\alpha)$ **true**.

- **Definition (Convergence)**

A **net** $\{x_\alpha\}_{\alpha \in I}$ in a topological space X is said to **converge** to a point $x \in X$ (written $x_\alpha \rightarrow x$) if for **any neighborhood** N of x , **there exists** a $\beta \in I$ so that $x_\alpha \in N$ if $\alpha \succ \beta$. The point x that being converged to is called **the limit point** of x_α .

Note that if $x_\alpha \rightarrow x$, then x_α is **eventually in all neighborhoods of** x . If x_α is **frequently in any neighborhood of** x , we say that x is a **cluster point** of x_α .

- **Remark** This definition *generalizes* the ϵ - δ language for convergence in metric space. Notice

that the notions of *limit* and *cluster point* generalize the same notions for sequences in a metric space..

- **Proposition 1.49** [Reed and Simon, 1980]

Let A be a set in a topological space X . Then, a point x is in the **closure** of A if and only if there is a net $\{x_\alpha\}_{\alpha \in I}$ with $x_\alpha \in A$, So that $x_\alpha \rightarrow x$.

- **Proposition 1.50** [Reed and Simon, 1980]

1. (**Continuous Function**): A function f from a topological space X to a topological space Y is **continuous** if and only if for **every convergent net** $\{x_\alpha\}_{\alpha \in I}$ **in** X , with $x_\alpha \rightarrow x$, the net $\{f(x_\alpha)\}_{\alpha \in I}$ **converges in** Y to $f(x)$.

2. (**Uniqueness of Limit Point for Hausdorff Space**): Let X be a **Hausdorff space**. Then a net $\{x_\alpha\}_{\alpha \in I}$ in X can have **at most one limit**; that is, if $x_\alpha \rightarrow x$ and $x_\alpha \rightarrow y$, then $x = y$.

- **Definition** A net $\{x_\alpha\}_{\alpha \in I}$ is a **subnet** of a net $\{y_\beta\}_{\beta \in J}$ if and only if there is a function $F : I \rightarrow J$ such that

1. $x_\alpha = y_{F(\alpha)}$ for each $\alpha \in I$.
2. For all $\beta' \in J$, there is an $\alpha' \in I$ such that $\alpha \succ \alpha'$ implies $F(\alpha) \succ \beta'$ (that is, $F(\alpha)$ is **eventually larger** than any fixed $\beta \in J$).

- **Proposition 1.51** A point x in a topological space X is a **cluster point** of a **net** $\{x_\alpha\}_{\alpha \in I}$ if and only if **some subnet** of $\{x_\alpha\}_{\alpha \in I}$ **converges** to x .

- **Theorem 1.52 (The Bolzano-Weierstrass Theorem)** [Reed and Simon, 1980]
A space X is **compact** if and only if **every net in** X **has a convergent subnet**.

2 Special Space

- **Remark (Metric Space and Compact Hausdorff Space)**

Two of the most well-behaved classes of spaces to deal with in mathematics are **the metrizable spaces** and **the compact Hausdorff spaces**.

1. **Metrizable space** (X, d) :

- **subspace** of metrizable space is **metrizable**;
- **compact subspace** of **metric space** is **bounded** in that metric and is **closed**;
- every metrizable space is **normal** (T_4);
- **compactness** = **sequential compactness** = **limit point compactness**;
- **sequence lemma**: for $A \subset X$, $x \in \bar{A}$ if and only if there exists a squence of points in A that converges to x . (\Rightarrow need X being metric space);
- f is **continuous** at x if and only if $x_n \rightarrow x$ leads to $f(x_n) \rightarrow f(x)$ (\Leftarrow part holds for metric space)
- **uniform limit theorem**: If the range of f_n is a **metric space** and f_n are **continuous**,

then $f_n \rightarrow f$ *uniformly* means that f is a *continuous* function.

- **uniform continuity theorem**: if f is a *countinuous* map between two *metric spaces*, and the domain is **compact**, then f is **uniformly continuous**.
- every metric space is **first-countable**.

2. Compact Hausdorff Space:

- **subspace** of *compact Hausdorff space* is *compact Hausdorff* if and only if it is **closed**.
- **closed subspace** of *compact space* is **compact**;
- **compact subspace** of *Hausdorff space* is **closed**;
- *compact Hausdorff space* X is **normal** (T_4), thus it is **completely regular**;
- **arbitrary product** of *compact (Hausdorff) space* is *compact (Hausdorff)*;
- **compactness** \Rightarrow **sequential compactness**;
- **compactness** = **net compactness**, i.e. every *net* has a convergence *subnet*;
- **image** of *compact space* under continuous map f is *compact*;
- **continuous bijection** between two *compact Hausdorff spaces* is a **homemorphism** (and is a **closed map**);
- **closed graph theorem**: f is **continuous** if and only if its **graph** is **closed**;
- **uncountability**: for *compact Hausdorff space*, if the space has *no isolated points*, then it is *uncountable*;
- if *compact Hausdorff space* is **second-countable**, then it is **metrizable**.

3 Summary of Preservation of Topological Properties

Table 1: Summary of Preservation of Topological Properties Under Transformations

	<i>subspace</i>	<i>product space</i>	<i>image of continuous function</i>
<i>connected</i>	✓	✓ under <i>product topology</i>	✓
<i>locally connected</i>	if <i>open and connected</i> subspace, ✓	if <i>all but finitely many</i> of spaces are <i>connected</i> , ✓	in general ×
<i>compact</i>	if <i>closed</i> subspace, ✓;	for <i>arbitrary</i> product, ✓	✓
<i>locally compact</i>	if <i>closed or open</i> subspace and Hausdorff, ✓	if <i>finite</i> product, ✓; if <i>infinite</i> product ×	if <i>f</i> is a <i>perfect map</i> , then ✓; in general ×
<i>first-countable</i>	✓	if <i>countable</i> product, ✓	if <i>f</i> is a <i>open map</i> , then ✓; in general ×
<i>second-countable</i>	✓	if <i>countable</i> product, ✓	if <i>f</i> is a <i>open map or perfect map</i> , then ✓; in general ×
<i>separable</i>	if metrizable, then ✓; in general ×	if <i>countable</i> product, ✓	✓
<i>Lindelöf</i>	if metrizable, then ✓; in general ×	×	✓
<i>T₁ axiom</i>	✓	for <i>arbitrary</i> product, ✓	in general ×
<i>Hausdorff T₂</i>	✓	for <i>arbitrary</i> product, ✓	if <i>f</i> is a <i>perfect map</i> , then ✓; in general ×
<i>regular T₃</i>	✓	for <i>arbitrary</i> product, ✓	if <i>f</i> is a <i>perfect map</i> , then ✓; in general ×
<i>completely regular</i>	✓	for <i>arbitrary</i> product, ✓	in general ×
<i>normal T₄</i>	×	×	×
<i>paracompact</i>	if <i>closed</i> subspace, ✓;	×	×
<i>topologically complete</i>	for <i>closed or open</i> subspace, ✓	if <i>countable</i> product, ✓	×

References

- Gerald B Folland. *Real analysis: modern techniques and their applications*. John Wiley & Sons, 2013.
- James R Munkres. Topology: a first course. *Englewood Cliffs, New Jersey*, 1975.
- James R Munkres. *Topology, 2nd*. Prentice Hall, 2000.
- Michael Reed and Barry Simon. *Methods of modern mathematical physics: Functional analysis*, volume 1. Gulf Professional Publishing, 1980.