

Lecture 0: Summary (part 3)

Tianpei Xie

Oct. 27th., 2022

Contents

1	Tensors	2
1.1	Abstract Tensor Product	3
1.2	Covariant and Contravariant Tensors on a Vector Space	5
1.3	Symmetric and Alternating Tensors	7
1.3.1	Symmetric Tensors	7
1.3.2	Alternating Tensors	8
1.4	Tensor Fields and Tensor Bundle	9
1.5	Pullbacks of Tensor Fields	11
1.6	Contraction	11
2	Differential Forms	13
2.1	Elementary k -covectors	13
2.2	Wedge Product	14
2.3	Interior Product	16
2.4	Differential Forms on Manifolds	16
2.5	Exterior Derivatives of Differential Forms	19
2.6	An Invariant Formula for the Exterior Derivative	20
3	Directional Derivatives of Vector Fields	21
3.1	Connections	21
3.2	Connections in the Tangent Bundle	22
3.3	Total Covariant Derivatives	24
3.4	Vector and Tensor Fields Along Curves	26
3.5	Geodesics	27
3.6	Parallel Transport	28
4	Lie Derivatives	30
5	Lie Group and Lie Algebra	31

1 Tensors

- **Definition** Suppose V_1, \dots, V_k , and W are *vector spaces*. A map $F : V_1 \times \dots \times V_k \rightarrow W$ is said to be **multilinear** if it is **linear** as a function of *each variable separately* when the others are held **fixed**: for each i ,

$$F(v_1, \dots, av_i + a'v'_i, \dots, v_k) = aF(v_1, \dots, v_i, \dots, v_k) + a'F(v_1, \dots, v'_i, \dots, v_k).$$

A *multilinear function of one variable* is just a **linear function**, and a multilinear function of *two variables* is generally called **bilinear**.

- **Remark** Let us write $L(V_1, \dots, V_k; W)$ for *the set of all multilinear maps from $V_1 \times \dots \times V_k$ to W* . It is a **vector space** under the usual operations of *pointwise addition* and *scalar multiplication*:

$$\begin{aligned}(F' + F)(v_1, \dots, v_i, \dots, v_k) &= F(v_1, \dots, v_i, \dots, v_k) + F'(v_1, \dots, v_i, \dots, v_k), \\ (aF)(v_1, \dots, v_i, \dots, v_k) &= aF(v_1, \dots, v_i, \dots, v_k).\end{aligned}$$

- **Example** (*Some Familiar Multilinear Functions*).

1. The **dot product**, $\langle \cdot, \cdot \rangle$ in \mathbb{R}^n is a **scalar-valued bilinear function** of two vectors, used to compute **lengths** of vectors and **angles** between them.
2. The **cross product**, $(\cdot \times \cdot)$ in \mathbb{R}^3 is a **vector-valued bilinear function** of two vectors, used to compute **areas** of parallelograms and to find a third vector **orthogonal** to two given ones.
3. The **determinant**, $\det(\cdot)$ is a **real-valued multilinear function** of n vectors in \mathbb{R}^n , used to detect **linear independence** and to compute the **volume** of the parallelepiped spanned by the vectors.
4. The **bracket in a Lie algebra** \mathfrak{g} , $[\cdot, \cdot]$ is a **\mathfrak{g} -valued bilinear function** of two elements of \mathfrak{g} .

- **Definition** let $V_1, \dots, V_k, W_1, \dots, W_l$ be real vector spaces, and suppose $F \in L(V_1, \dots, V_k; \mathbb{R})$ and $G \in L(W_1, \dots, W_l; \mathbb{R})$. Define a function $F \otimes G : V_1 \times \dots \times V_k \times W_1 \times \dots \times W_l \rightarrow \mathbb{R}$ by

$$(F \otimes G)(v_1, \dots, v_k, w_1, \dots, w_l) = F(v_1, \dots, v_k)G(w_1, \dots, w_l) \quad (1)$$

It follows from the multilinearity of F and G that $(F \otimes G)(v_1, \dots, v_k, w_1, \dots, w_l)$ depends *linearly* on each argument v_i or w_j *separately*, so $F \otimes G$ is an element of $L(V_1, \dots, V_k, W_1, \dots, W_l; \mathbb{R})$ called **the tensor product of F and G** .

- **Remark** If $\omega^j \in V_j^*$ for $j = 1, \dots, k$, then $\omega^1 \otimes \dots \otimes \omega^k \in L(V_1, \dots, V_k; \mathbb{R})$ is the **multilinear function** given by

$$(\omega^1 \otimes \dots \otimes \omega^k)(v_1, \dots, v_k) = \omega^1(v_1) \dots \omega^k(v_k). \quad (2)$$

We can see that $\omega^1 \otimes \dots \otimes \omega^k$ is a multilinear extension of the linear functional ω .

• **Proposition 1.1** (*A Basis for the Space of Multilinear Functions*).

Let V_1, \dots, V_k be real vector spaces of dimensions n_1, \dots, n_k , respectively. For each $j \in \{1, \dots, k\}$, let $(E_1^{(j)}, \dots, E_{n_j}^{(j)})$ be a **basis** for V_j , and let $(\epsilon_{(j)}^1, \dots, \epsilon_{(j)}^{n_j})$ be the corresponding **dual basis** for V_j^* . Then the set

$$\mathfrak{B} = \left\{ \epsilon_{(1)}^{i_1} \otimes \dots \otimes \epsilon_{(k)}^{i_k} : 1 \leq i_j \leq n_j, j = 1, \dots, k \right\}$$

is a **basis** for $L(V_1, \dots, V_k; \mathbb{R})$, which therefore has **dimension equal to** $n_1 \dots n_k$.

1.1 Abstract Tensor Product

- **Remark** Intuitively, we want to define the tensor product $v_1 \otimes \dots \otimes v_k$ by concatenating all vectors into k -tuple (v_1, \dots, v_k) . But this naive construction is not enough. We have the following challenges:

1. The product space $V_1 \times \dots \times V_k$ is *not necessarily a vector space* since we have not define the addition and scalar product for k -tuple (v_1, \dots, v_k)
2. We want the **multilinearity holds** for the operator on k -tuple (v_1, \dots, v_k) , i.e. we want

$$(v_1, \dots, a v_i' + b v_i'', v_k) = a (v_1, \dots, v_i', v_k) + b (v_1, \dots, v_i'', v_k) \quad (3)$$

for any $i \in \{1, \dots, k\}$ and any $a, b \in \mathbb{R}$.

The above constructions aim to solve these challenges:

1. Instead of defining the algebraic structure on product space $V_1 \times \dots \times V_k$, we extend it to **the free vector space** $\mathcal{F}(V_1 \times \dots \times V_k)$, the set of *all linear combinations* of k -tuples (v_1, \dots, v_k) . By construction $\mathcal{F}(V_1 \times \dots \times V_k) \supseteq V_1 \times \dots \times V_k$ and *it is a vector space without defining the algebraic structure* since it use an indicator function to map to \mathbb{R} .
2. Instead of enforcing the *multilinearity* to hold, we **partition the space** $\mathcal{F}(V_1 \times \dots \times V_k)$ **according to the multilinearity rule**. That is, the set of tuples $(v_1, \dots, a v_i' + b v_i'', v_k)$ and $(v_1, \dots, v_i', v_k), (v_1, \dots, v_i'', v_k)$ that satisfies the equation (3) *will be grouped together* via the equivalence relationship. The rule is actually a set of linear combinations of (difference of) tuples, denoted as \mathcal{R} .

Now we instead focusing on the equivalent class itself. By construction, **the equivalence class will satisfies the multilinearity rule** (3) (The representer of the equivalence class follow the rule). Thsu $V_1 \otimes \dots \otimes V_k = \mathcal{F}(V_1 \times \dots \times V_k) / \mathcal{R}$ is the tensor product space that we wants.

- **Definition** Now let V_1, \dots, V_k be real vector spaces. We begin by forming **the free vector space** $\mathcal{F}(V_1 \times \dots \times V_k)$, which is the set of all finite formal linear combinations of k -tuples (v_1, \dots, v_k) with $v_i \in V_i$ for $i = 1, \dots, k$. Let \mathcal{R} be the **subspace** of $\mathcal{F}(V_1 \times \dots \times V_k)$ *spanned* by all elements of the following forms:

$$\begin{aligned} & (v_1, \dots, a v_i, \dots, v_k) - a (v_1, \dots, v_i, \dots, v_k) \\ & (v_1, \dots, v_i + v_i', \dots, v_k) - (v_1, \dots, v_i, \dots, v_k) - (v_1, \dots, v_i', \dots, v_k) \end{aligned} \quad (4)$$

with $v_j, v_j' \in V_j$, $i \in \{1, \dots, k\}$, and $a \in \mathbb{R}$.

Define the tensor product of the spaces V_1, \dots, V_k , denoted by $V_1 \otimes \dots \otimes V_k$, to be the following quotient vector space:

$$V_1 \otimes \dots \otimes V_k = \mathcal{F}(V_1 \times \dots \times V_k) / \mathcal{R}$$

and let $\Pi : \mathcal{F}(V_1 \times \dots \times V_k) \rightarrow V_1 \otimes \dots \otimes V_k$ be the natural projection. The equivalence class of an element (v_1, \dots, v_k) in $V_1 \otimes \dots \otimes V_k$ is denoted by

$$v_1 \otimes \dots \otimes v_k = \Pi(v_1, \dots, v_k) \quad (5)$$

and is called the (abstract) tensor product of (v_1, \dots, v_k) .

It follows from the definition that abstract tensor products satisfy

$$\begin{aligned} v_1 \otimes \dots \otimes (a v_i) \otimes \dots \otimes v_k &= a(v_1 \otimes \dots \otimes v_i \otimes \dots \otimes v_k), \\ v_1 \otimes \dots \otimes (v_i + v'_i) \otimes \dots \otimes v_k &= (v_1 \otimes \dots \otimes v_i \otimes \dots \otimes v_k) + (v_1 \otimes \dots \otimes v'_i \otimes \dots \otimes v_k) \end{aligned}$$

• **Proposition 1.2 (Characteristic Property of the Tensor Product Space).**

Let V_1, \dots, V_k be finite-dimensional real vector spaces. If $A : V_1 \times \dots \times V_k \rightarrow X$ is **any multilinear map** into a vector space X , then there is a **unique linear map** $\tilde{A} : V_1 \otimes \dots \otimes V_k \rightarrow X$ such that the following diagram commutes:

$$\begin{array}{ccc} V_1 \times \dots \times V_k & \xrightarrow{A} & X \\ \pi \downarrow & \nearrow \tilde{A} & \\ V_1 \otimes \dots \otimes V_k & & \end{array}, \quad (6)$$

where π is the map $\pi(v_1, \dots, v_k) = v_1 \otimes \dots \otimes v_k$.

- **Remark** The *characteristic property of the tensor product space* states that *any multilinear function* $\tau : V_1 \times \dots \times V_k \rightarrow \mathbb{R}$ descends into a *linear map* $\tilde{\tau} : V_1 \otimes \dots \otimes V_k \rightarrow \mathbb{R}$ so that any linear combinations of tensor products $v_{i_1} \otimes \dots \otimes v_{i_k}$ is expressed as

$$\tilde{\tau}(a^{i_1 \dots i_k} v_{i_1} \otimes \dots \otimes v_{i_k}) = a^{i_1 \dots i_k} \tau(v_{i_1}, \dots, v_{i_k})$$

• **Proposition 1.3 (A Basis for the Tensor Product Space).**

Suppose V_1, \dots, V_k are real vector spaces of dimensions $n_1 \dots n_k$, respectively. For each $j = 1, \dots, k$, suppose $(E_1^{(j)}, \dots, E_{n_j}^{(j)})$ is a **basis** for V_j . Then the set

$$\mathfrak{E} = \left\{ E_{i_1}^{(1)} \otimes \dots \otimes E_{i_k}^{(k)} : 1 \leq i_j \leq n_j, j = 1, \dots, k \right\}$$

is a **basis** for $V_1 \otimes \dots \otimes V_k$, which therefore has **dimension equal to** $n_1 \dots n_k$.

• **Proposition 1.4 (Abstract vs. Concrete Tensor Products).** [Lee, 2003.]

If V_1, \dots, V_k are finite-dimensional vector spaces, there is a canonical isomorphism

$$V_1^* \otimes \dots \otimes V_k^* \simeq L(V_1, \dots, V_k; \mathbb{R}) \quad (7)$$

under which the **abstract tensor product** defined by (5) corresponds to the **tensor product of covectors** defined by (2).

- **Remark** Since we are assuming the vector spaces are all finite-dimensional, we can also identify each V_j with its second dual space V_j^{**} , and thereby obtain *another canonical identification*

$$V_1 \otimes \dots \otimes V_k \simeq L(V_1^*, \dots, V_k^*; \mathbb{R}) \quad (8)$$

- **Remark (Kronecker Product vs. Tensor Product)**

The two notions represent operations on different objects: Kronecker product on matrices; tensor product on linear maps between vector spaces. But there is a connection: Given two matrices, we can think of them as representing linear maps between vector spaces equipped with a chosen basis. *The Kronecker product of the two matrices then represents the tensor product of the two linear maps.* (This claim makes sense because the tensor product of two vector spaces with distinguished bases comes with a distinguished basis.)

- **Remark** As we see, the space of tensor product defines a set of *parallel linear systems*. All *sub-systems* are *independent*. Each sub-system has its own *basis*, its own *linear operations* and its own *representation*. The tensor product operation *group* these independent linear systems together and *consider them as a whole*.

For *the whole system perspective*, its representations are collected locally and then concatenated together. The linear map on *the concatenated representation* is essentially the same as *applying linear map* in each sub-system and *multiplying* them together. This is the same as computing the joint distribution by product of marginal distributions. *The multiplication principle* is applied when counting the size of the whole system.

The space of tensor product $V_1 \otimes \dots \otimes V_k$ reflect a *divide-and-conquer strategy* and a *local-global strategy* to study the complex functions such as *multilinear functionals* $\alpha(v_1, \dots, v_k)$. In the former, we study it by *perturbing the input of each sub-system*. In the latter, we think of it as a *linear map* on the k -tensors $v_1 \otimes \dots \otimes v_k$.

1.2 Covariant and Contravariant Tensors on a Vector Space

- **Definition** Let V be a finite-dimensional real vector space. If k is a positive integer, *a covariant k -tensor* on V is an element of the *k -fold tensor product* $V^* \otimes \dots \otimes V^*$, which we typically think of as *a real-valued multilinear function of k elements of V* :

$$\alpha : \underbrace{V \times \dots \times V}_k \rightarrow \mathbb{R}$$

The number k is called *the rank of α* . A 0-tensor is, by convention, just a real number (a real-valued function depending multilinearly on *no vectors*!).

We denote *the vector space of all covariant k -tensors on V* by the shorthand notation

$$T^k V^* = \underbrace{V^* \otimes \dots \otimes V^*}_k$$

- **Example (Covariant Tensors).**

Let V be a finite-dimensional vector space.

1. Every linear functional $\omega : V \rightarrow \mathbb{R}$ is multilinear, so **a covariant 1-tensor is just a covector**. Thus, $T^1(V^*)$ is equal to V^* .
 2. A covariant 2-tensor on V is a real-valued **bilinear function** of two vectors, also called **a bilinear form**. One example is the dot product on \mathbb{R}^n . More generally, **every inner product is a covariant 2-tensor**.
 3. The determinant, thought of as a function of n vectors, is **a covariant n -tensor on \mathbb{R}^n** .
- **Definition** For any finite-dimensional real vector space V , we define the space of **contravariant tensors on V of rank k** to be the vector space

$$T^k V = \underbrace{V \otimes \dots \otimes V}_k$$

In particular, $T^1(V) = V$, and by convention $T^0(V) = \mathbb{R}$. Because we are assuming that V is finite-dimensional, it is possible to identify this space with *the set of multilinear functionals of k covectors*:

$$T^k V = \left\{ \text{multilinear functionals } \alpha : \underbrace{V^* \times \dots \times V^*}_k \rightarrow \mathbb{R} \right\}$$

- **Definition** Even more generally, for any nonnegative integers k, l , we define the space of **mixed tensors on V of type (k, l)** as

$$T^{(k,l)} V = \underbrace{V \otimes \dots \otimes V}_k \otimes \underbrace{V^* \otimes \dots \otimes V^*}_l$$

- **Corollary 1.5** Let V be an n -dimensional real vector space. Suppose (E_i) is any basis for V and (ϵ^j) is the dual basis for V^* . Then the following sets constitute bases for the tensor spaces over V :

$$\begin{aligned} & \{ \epsilon^{i_1} \otimes \dots \otimes \epsilon^{i_k} : 1 \leq i_s \leq n, s = 1, \dots, k \} \text{ is basis for } T^k V^*; \\ & \{ E_{i_1} \otimes \dots \otimes E_{i_k} : 1 \leq i_s \leq n, s = 1, \dots, k \} \text{ is basis for } T^k V; \\ & \{ E_{i_1} \otimes \dots \otimes E_{i_k} \otimes \epsilon^{j_1} \otimes \dots \otimes \epsilon^{j_l} : 1 \leq i_1, \dots, i_k, j_1, \dots, j_l \leq n \} \text{ is basis for } T^{(k,l)} V; \end{aligned} \quad (9)$$

Therefore, $\dim T^k V^* = \dim T^k V = n^k$ and $\dim T^{(k,l)} V = n^{k+l}$

- **Remark (Coordinate Representation of Covariant k -Tensor)**

In particular, once a basis is chosen for V , every **covariant k -tensor** $\alpha \in T^k(V^*)$ can be written uniquely in the form

$$\alpha = \alpha_{i_1, i_2, \dots, i_k} \epsilon^{i_1} \otimes \dots \otimes \epsilon^{i_k} \quad (10)$$

where the n^k coefficients $\alpha_{i_1, i_2, \dots, i_k}$ are determined by

$$\alpha_{i_1, i_2, \dots, i_k} = \alpha(E_{i_1}, \dots, E_{i_k}) \quad (11)$$

For instance, covariant 2-tensor is bilinear form. Every *bilinear form* can be written as $\beta = \beta_{i,j} \epsilon^i \otimes \epsilon^j$, for some uniquely determined $n \times n$ matrix $(\beta_{i,j})$.

1.3 Symmetric and Alternating Tensors

1.3.1 Symmetric Tensors

- **Definition** Let V be a finite-dimensional vector space. A *covariant k -tensor* α on V is said to be **symmetric** if its value is *unchanged* by **interchanging** any pair of arguments:

$$\alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) = \alpha(v_1, \dots, v_j, \dots, v_i, \dots, v_k)$$

whenever $i \leq i < j \leq k$.

- **Definition** The set of **symmetric covariant k -tensors** is a linear subspace of the space $T^k(V^*)$ of all covariant k -tensors on V ; we denote this subspace by $\underline{\Sigma^k(V^*)}$

There is a **natural projection** from $T^k(V^*)$ to $\Sigma^k(V^*)$ defined as follows. First, let S_k denote **the symmetric group on k elements**, that is, the group of **permutations** of the set $\{1, \dots, k\}$. Given a k -tensor α and a permutation $\sigma \in S_k$, we define a new k -tensor ${}^\sigma\alpha$ by

$${}^\sigma\alpha(v_1, \dots, v_k) = \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)})$$

Note that ${}^\tau({}^\sigma\alpha) = {}^{\tau\sigma}\alpha$ where $\tau\sigma$ represents the composition of τ and σ , that is, $\tau\sigma(i) = \tau(\sigma(i))$. (This is the reason for putting σ before α in the notation ${}^\sigma\alpha$ instead of after it.)

We define a **projection** $\text{Sym} : T^k(V^*) \rightarrow \Sigma^k(V^*)$ called **symmetrization** by

$$\text{Sym } \alpha = \frac{1}{k!} \sum_{\sigma \in S_k} {}^\sigma\alpha$$

More explicitly, this means that

$$\text{Sym } \alpha(v_1, \dots, v_k) = \frac{1}{k!} \sum_{\sigma \in S_k} \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)})$$

- **Proposition 1.6 (Properties of Symmetrization).**

Let α be a covariant tensor on a finite-dimensional vector space.

1. $\text{Sym } \alpha$ is symmetric.
2. $\text{Sym } \alpha = \alpha$ if and only if α is symmetric.

- **Definition** If $\alpha \in \Sigma^k(V^*)$ and $\beta \in \Sigma^l(V^*)$, we define their **symmetric product** to be the $(k+l)$ -tensor $\alpha\beta$ (denoted by juxtaposition with no intervening product symbol) given by

$$\alpha\beta = \text{Sym } (\alpha \otimes \beta)$$

More explicitly, the action of $\alpha\beta$ on vectors v_1, \dots, v_{k+l} is given by

$$\alpha\beta(v_1, \dots, v_{k+l}) = \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \beta(v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)})$$

- **Proposition 1.7 (Properties of the Symmetric Product).**

1. The symmetric product is **symmetric** and **bilinear**: for all symmetric tensors α, β, γ and all $a, b \in \mathbb{R}$,

$$\begin{aligned}\alpha \beta &= \beta \alpha \\ (a \alpha + b \beta) \gamma &= a \alpha \gamma + b \beta \gamma = \gamma (a \alpha + b \beta)\end{aligned}$$

2. If α and β are **covectors**, then

$$\alpha \beta = \frac{1}{2} (\alpha \otimes \beta + \beta \otimes \alpha).$$

1.3.2 Alternating Tensors

- **Definition** Assume that V is a finite-dimensional real vector space. A **covariant k -tensor** α on V is said to be **alternating** (or **antisymmetric** or **skew-symmetric**) if it **changes sign** whenever two of its arguments are *interchanged*. This means that for all vectors $v_1, \dots, v_k \in V$ and every pair of distinct indices i, j it satisfies

$$\alpha(v_1, \dots, v_i, \dots, v_j, \dots, v_k) = -\alpha(v_1, \dots, v_j, \dots, v_i, \dots, v_k)$$

Alternating covariant k -tensors are also variously called **exterior forms, multicovectors,** or **k -covectors.**

The subspace of **all alternating covariant k -tensors** on V is denoted by $\Lambda^k(V^*) \subseteq T^k(V^*)$.

- **Definition** Recall that for any permutation $\sigma \in S_k$, **the sign of σ** , denoted by $\text{sgn } \sigma$, is equal to $+1$ if σ is **even** (i.e., can be written as a composition of an **even** number of transpositions), and -1 if σ is **odd**.
- **Lemma 1.8** The following statements are equivalent for a covariant k -tensor α :

1. α is **alternating**;
2. For any vectors $v_1, \dots, v_k \in V$, and **any permutation** $\sigma \in S_k$

$$\alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}) = (\text{sgn } \sigma) \alpha(v_1, \dots, v_k)$$

3. With respect to any basis, the components α_{i_1, \dots, i_k} of α change sign whenever two indices are interchanged.

- **Lemma 1.9** Let α be a covariant k -tensor on a finite-dimensional vector space V . The following are equivalent:

1. α is **alternating**.
2. $\alpha(v_1, \dots, v_k) = 0$ whenever the k -tuple (v_1, \dots, v_k) is **linearly dependent**.
3. α gives the value zero whenever **two of its arguments** are equal:

$$\alpha(v_1, \dots, w, \dots, w, v_k) = 0.$$

- **Definition** We define a projection $\text{Alt} : T^k(V^*) \rightarrow \Lambda^k(V^*)$, called **alternation**, as follows:

$$\text{Alt } \alpha = \frac{1}{k!} \sum_{\sigma \in S_k} (\text{sign } \{\sigma\}) (\sigma \alpha)$$

where S_k is the symmetric group on k elements. More explicitly, this means

$$\text{Alt } \alpha(v_1, \dots, v_k) = \frac{1}{k!} \sum_{\sigma \in S_k} (\text{sign } \{\sigma\}) \alpha(v_{\sigma(1)}, \dots, v_{\sigma(k)}).$$

- **Proposition 1.10** (*Properties of Alternation*).

Let α be a covariant tensor on a finite-dimensional vector space.

1. Alt α is alternating.
2. Alt $\alpha = \alpha$ if and only if α is alternating.

1.4 Tensor Fields and Tensor Bundle

- **Definition** Now let M be a smooth manifold with or without boundary. We define the **bundle of covariant k -tensors** on M by

$$T^k T^* M = \bigsqcup_{p \in M} T^k (T_p^* M)$$

Analogously, we define **the bundle of contravariant k -tensors** by

$$T^k T M = \bigsqcup_{p \in M} T^k (T_p M)$$

and **the bundle of mixed tensors of type (k, l)** by

$$T^{(k, l)} T M = \bigsqcup_{p \in M} T^{(k, l)} (T_p M)$$

- **Remark** There are natural identifications

$$\begin{aligned} T^{(0, 0)} T M &= T^0 T^* M = T^0 T M = M \times \mathbb{R}; \\ T^{(0, 1)} T M &= T^1 T^* M = T^* M; \\ T^{(1, 0)} T M &= T^1 T M = T M; \\ T^{(0, k)} T M &= T^k T^* M; \\ T^{(k, 0)} T M &= T^k T M. \end{aligned}$$

Any one of these bundles is called **a tensor bundle over M** . (Thus, the tangent and cotangent bundles are special cases of tensor bundles.)

- **Definition** A **section** of a tensor bundle is called a (**covariant, contravariant, or mixed**) **tensor field** on M . A **smooth tensor field** is a section that is smooth in the usual sense of smooth sections of vector bundles.
- **Remark** The **spaces of smooth sections** of these tensor bundles, $\Gamma(T^k T^* M)$, $\Gamma(T^k T M)$, and $\Gamma(T^{(k, l)} T M)$, are **infinite-dimensional vector spaces over \mathbb{R}** , and **modules** over $\mathcal{C}^\infty(M)$. We also denote the **space of smooth covariant tensor fields** as

$$\mathcal{T}^k(M) = \Gamma(T^k T^* M).$$

- **Remark (Coordinate Representation of Tensor Fields)**

In any smooth local coordinates (x^i) , sections of these bundles can be written (using the summation convention) as

$$A = \begin{cases} A_{i_1, \dots, i_k} dx^{i_1} \otimes \dots \otimes dx^{i_k}, & A \in \Gamma(T^k T^* M); \\ A^{i_1, \dots, i_k} \frac{\partial}{\partial x^{i_1}} \otimes \dots \otimes \frac{\partial}{\partial x^{i_k}}, & A \in \Gamma(T^k TM); \\ A_{j_1, \dots, j_l}^{i_1, \dots, i_k} \frac{\partial}{\partial x^{i_1}} \otimes \dots \otimes \frac{\partial}{\partial x^{i_k}} \otimes dx^{j_1} \otimes \dots \otimes dx^{j_l} & A \in \Gamma(T^{(k,l)} TM); \end{cases} \quad (12)$$

The functions A_{i_1, \dots, i_k} , A^{i_1, \dots, i_k} , or $A_{j_1, \dots, j_l}^{i_1, \dots, i_k}$ are called the **component functions** of A in the chosen coordinates.

- **Proposition 1.11 (Smoothness Criteria for Tensor Fields).**

Let M be a smooth manifold with or without boundary, and let $A : M \rightarrow T^k T^* M$ be a rough section. The following are equivalent.

1. A is smooth.
2. In **every** smooth coordinate chart, the **component functions** of A are smooth.
3. Each point of M is contained in **some** coordinate chart in which A has **smooth component functions**.
4. If $X_1, \dots, X_k \in \mathfrak{X}(M)$, then the function $A(X_1, \dots, X_k) : M \rightarrow \mathbb{R}$, defined by

$$A(X_1, \dots, X_k)(p) = A_p(X_1|_p, \dots, X_k|_p) \quad (13)$$

is smooth

5. Whenever X_1, \dots, X_k are smooth vector fields defined on **some open subset** $U \subseteq M$, the function $A(X_1, \dots, X_k)$ is smooth on U .

- **Lemma 1.12 (Tensor Characterization Lemma).**[Lee, 2003.]

A map

$$A : \underbrace{\mathfrak{X}(M) \times \dots \times \mathfrak{X}(M)}_k \rightarrow C^\infty(M). \quad (14)$$

is **induced** by a **smooth covariant k -tensor field** A as in (13) if and only if it is **multilinear** over $C^\infty(M)$.

- **Definition** For symmetric and alternating tensor field, we have the following definition:

1. A **symmetric tensor field** on a manifold (with or without boundary) is simply a **covariant tensor field** whose value at each point is a **symmetric tensor**.

The **symmetric product** of two or more tensor fields is defined pointwise, just like the tensor product. Thus, for example, if A and B are **smooth covector fields**, their symmetric product is **the smooth 2-tensor field** AB , which is given by

$$AB = \frac{1}{2} (A \otimes B) + \frac{1}{2} (B \otimes A).$$

2. **Alternating tensor fields** are called **differential forms**;

1.5 Pullbacks of Tensor Fields

- **Definition** Suppose $F : M \rightarrow N$ is a smooth map. For any point $p \in M$ and any k -tensor $\alpha \in T^k(T_{F(p)}^*N)$, we define a tensor $dF_p^*(\alpha) \in T^k(T_p^*M)$, called the pointwise pullback of α by F at p , by

$$dF_p^*(\alpha)(v_1, \dots, v_k) = \alpha(dF_p(v_1), \dots, dF_p(v_k))$$

for any $v_1, \dots, v_k \in T_pM$.

- **Definition** If A is a covariant k -tensor field on N , we define a rough k -tensor field F^*A on M ; called the pullback of A by F , by

$$(F^*A)_p = dF_p^*(A_{F(p)}).$$

This tensor field acts on vectors $v_1, \dots, v_k \in T_pM$ by

$$(F^*A)_p(v_1, \dots, v_k) = A_{F(p)}(dF_p(v_1), \dots, dF_p(v_k)).$$

- **Proposition 1.13 (Properties of Tensor Pullbacks).**

Suppose $F : M \rightarrow N$ and $G : N \rightarrow P$ are smooth maps, A and B are covariant tensor fields on N , and f is a real-valued function on N .

1. $F^*(fB) = (f \circ F) F^*(B)$
2. $F^*(A \otimes B) = F^*A \otimes F^*(B)$
3. $F^*(A + B) = F^*A + F^*(B)$
4. $F^*(B)$ is a (continuous) tensor field, and is smooth if B is smooth.
5. $(G \circ F)^*B = F^*(G^*B)$.
6. $(Id_N)^*B = B$.

- **Corollary 1.14 (Coordinate Representation of Pullback Tensor Fields)**

Let $F : M \rightarrow N$ be smooth, and let B be a covariant k -tensor field on N . If $p \in M$ and (y^i) are smooth coordinates for N on a neighborhood of $F(p)$, then F^*B has the following expression in a neighborhood of p :

$$F^*(B_{i_1, \dots, i_k} dy^{i_1} \otimes \dots \otimes dy^{i_k}) = (B_{i_1, \dots, i_k} \circ F) d(y^{i_1} \circ F) \otimes \dots \otimes (dy^{i_k} \circ F).$$

- **Remark** F^*B is computed as follows: wherever you see y^i in the expression for B , just substitute the i th component function of F and expand.

1.6 Contraction

- **Proposition 1.15** Let V be a finite-dimensional vector space. There is a natural (basis-independent) **isomorphism** between $T^{(k+1, l)}V$ and the space of **multilinear** maps

$$\underbrace{V^* \times \dots \times V^*}_k \times \underbrace{V \times \dots \times V}_l \rightarrow V$$

- **Definition** We can use the result of Proposition 1.15 to define a natural operation called ***trace*** or ***contraction***, which *lowers the rank of a tensor by 2*.

For $F = v \otimes \omega \in T^{(1,1)}V$. Define the operator $\text{tr} : T^{(1,1)}V \rightarrow \mathbb{R}$ is just ***the trace of F*** for i.e. the sum of the diagonal entries of any matrix representation of F . More generally, we define $\text{tr} : T^{(k+1,l+1)}V \rightarrow T^{(k,l)}V$ by letting $\text{tr} F(\omega^1, \dots, \omega^k, v_1, \dots, v_l)$ be the ***trace*** of the ***(1,1)-tensor***

$$F(\omega^1, \dots, \omega^k, \cdot, v_1, \dots, v_l, \cdot) \in T^{(1,1)}V$$

In terms of a basis, the ***components*** of $\text{tr} F$ are

$$(\text{tr} F)_{j_1, \dots, j_l}^{i_1, \dots, i_k} = F_{j_1, \dots, j_l, m}^{i_1, \dots, i_k, m}.$$

In other words, just ***set the last upper and lower indices equal and sum***.

- **Remark** We consider a $(1,1)$ -tensor $F = v \otimes \omega$. Under standard basis, $v = v^i E_i$ and $\omega = \omega_j \epsilon^j$, F has representation

$$\begin{aligned} F &= v \otimes \omega \\ &= (v^i E_i) \otimes (\omega_j \epsilon^j) \\ &= (\omega_j v^i) E_i \otimes \epsilon^j := F_j^i E_i \otimes \epsilon^j \end{aligned}$$

There is an isomorphism $T^{(1,1)}V \rightarrow L(V; V)$ as $F \mapsto [F_j^i]_{j,i}$. Then the ***trace*** of F is

$$\begin{aligned} \text{tr}(v \otimes \omega) &= \omega(v) \\ &= \omega_i v^i \\ &= \text{tr} \left(\begin{bmatrix} \omega_1 v^1 & \dots & \omega_1 v^n \\ \vdots & \ddots & \vdots \\ \omega_n v^1 & \dots & \omega_n v^n \end{bmatrix} \right) = \text{tr} [F_j^i]_{j,i} \end{aligned}$$

- **Remark** We have the formula for a (k,l) -tensor field F

$$F(\omega^1, \dots, \omega^k, V_1, \dots, V_l) = \underbrace{\text{tr} \circ \dots \circ \text{tr}}_{k+l} \left(F \otimes \omega^1 \otimes \dots \otimes \omega^k \otimes V_1 \otimes \dots \otimes V_l \right), \quad (15)$$

where each trace operator acts on an upper index of F and the lower index of the corresponding 1-form, or a lower index of F and the upper index of the corresponding vector field.

For instance, for covariant 2-tensor field $g = \omega^1 \otimes \omega^2$:

$$\begin{aligned} g(X, Y) &= \text{tr}(\text{tr}(\omega^1 \otimes \omega^2 \otimes X \otimes Y)) \\ &= \text{tr}(\text{tr}(\omega^2 \otimes Y) \omega^1 \otimes X) \\ &= \text{tr}((\omega^2(Y)) \omega^1 \otimes X) \\ &= (\omega^2(Y)) \text{tr}(\omega^1 \otimes X) \\ &= (\omega^2(Y)) (\omega^1(X)) \end{aligned}$$

2 Differential Forms

2.1 Elementary k -covectors

- **Definition** Given a positive integer k , an **ordered k -tuple** $I = (i_1, \dots, i_k)$ of positive integers is called a **multi-index** of length k . If I is such a multi-index and $\sigma \in S_k$ is a permutation of $\{1, \dots, k\}$, we write I for the following multi-index:

$$I_\sigma = (i_{\sigma(1)}, \dots, i_{\sigma(k)}).$$

Note that $I_{\sigma\tau} = (I_\sigma)_\tau$ for $\sigma, \tau \in S_k$.

- **Definition** Let V be an n -dimensional vector space, and suppose $(\epsilon^1, \dots, \epsilon^n)$ is any basis for V^* . We now define a collection of k -covectors on V that generalize the determinant function on \mathbb{R}^n .

For each multi-index $I = (i_1, \dots, i_k)$ of length k such that $1 \leq i_1 \leq \dots \leq i_k \leq n$, define a **covariant k -tensor** $\epsilon^I = \epsilon^{i_1, \dots, i_k}$ by

$$\epsilon^I(v_1, \dots, v_k) = \det \begin{bmatrix} \epsilon^{i_1}(v_1) & \dots & \epsilon^{i_1}(v_k) \\ \vdots & \ddots & \vdots \\ \epsilon^{i_k}(v_1) & \dots & \epsilon^{i_k}(v_k) \end{bmatrix} = \det \begin{bmatrix} v_1^{i_1} & \dots & v_k^{i_1} \\ \vdots & \ddots & \vdots \\ v_1^{i_k} & \dots & v_k^{i_k} \end{bmatrix}. \quad (16)$$

In other words, if \mathbf{V} denotes the $n \times k$ matrix whose columns are the components of the vectors v_1, \dots, v_k with respect to the basis (E_i) dual to (ϵ^i) , then $\epsilon^I(v_1, \dots, v_k)$ is the **determinant of the $k \times k$ submatrix** consisting of rows i_1, \dots, i_k of \mathbf{V} . Because the determinant changes sign whenever two columns are interchanged, it is clear that ϵ^I is an **alternating k -tensor**. We call ϵ^I an **elementary alternating tensor** or **elementary k -covector**.

- **Definition** If I and J are multiindices of length k , we define the Kronecker delta function:

$$\delta_J^I = \det \begin{bmatrix} v_{j_1}^{i_1} & \dots & v_{j_k}^{i_1} \\ \vdots & \ddots & \vdots \\ v_{j_1}^{i_k} & \dots & v_{j_k}^{i_k} \end{bmatrix}$$

(I represent the row number, J represent the column number.)

- **Remark** The following is the property of Kronecker delta

$$\delta_J^I = \begin{cases} \text{sign}\{\sigma\} & \text{if neither } I \text{ nor } J \text{ has a repeated index, } J = I_\sigma, \sigma \in S_k \\ 0 & \text{if } I \text{ or } J \text{ has a repeated index or } J \text{ is not a permutation of } I \end{cases}$$

- **Lemma 2.1 (Properties of Elementary k -Covectors).**

Let (E_i) be a basis for V , let (ϵ^i) be the dual basis for V^* , and let ϵ^I be as defined above.

1. If I has a repeated index, then $\epsilon^I = 0$.
2. If $J = I_\sigma$ for some $\sigma \in S_k$, then $\epsilon^I = \text{sign}\{\sigma\} \epsilon^J$.
3. The result of evaluating ϵ^I on a sequence of basis vectors is

$$\epsilon^I(E_{j_1}, \dots, E_{j_k}) = \delta_J^I.$$

- **Definition** A multi-index $I = (i_1, \dots, i_k)$ is said to be **increasing** if $i_1 < \dots < i_k$. It is useful to use a primed summation sign to denote a sum over *only increasing multi-indices*

$$\sum_I' a_I \epsilon^I = \sum_{\{I: i_1 < \dots < i_k\}} a_I \epsilon^I.$$

- **Proposition 2.2 (A Basis for $\Lambda^k(V^*)$)**

Let V be an n -dimensional vector space. If (ϵ^i) is any basis for V^* , then for each positive integer $k \leq n$, the collection of k -covectors

$$\mathcal{E} = \{\epsilon^I : I \text{ is an increasing multi-index of length } k\}$$

is a basis for $\Lambda^k(V^*)$. Therefore,

$$\dim \Lambda^k(V^*) = \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

If $k > n$, then $\dim \Lambda^k(V^*) = 0$.

- **Remark** In particular, for an n -dimensional vector space V , this proposition implies that $\Lambda^n(V^*)$ is 1-**dimensional** and is spanned by $\epsilon^{1, \dots, n}$.
- **Proposition 2.3** Suppose V is an n -dimensional vector space and $\omega \in \Lambda^n(V^*)$. If $T : V \rightarrow V$ is any **linear map** and v_1, \dots, v_n are arbitrary vectors in V , then

$$\omega(Tv_1, \dots, Tv_n) = (\det T) \omega(v_1, \dots, v_n). \quad (17)$$

2.2 Wedge Product

- **Definition** Let V be a finite-dimensional real vector space. Given $\omega \in \Lambda^k(V^*)$ and $\eta \in \Lambda^l(V^*)$, we define their **wedge product** or **exterior product** to be the following $(k+l)$ -covector:

$$\omega \wedge \eta = \frac{(k+l)!}{k!l!} \text{Alt}(\omega \otimes \eta) = \frac{1}{k!l!} \sum_{\sigma \in S_{k+l}} (\text{sign}\{\sigma\}) (\sigma(\omega \otimes \eta)) \quad (18)$$

- The coefficients come from the following lemma:

Lemma 2.4 Let V be an n -dimensional vector space and let $(\epsilon^1, \dots, \epsilon^n)$ be a basis for V^* . For any multi-indices $I = (i_1, \dots, i_k)$ and $J = (j_1, \dots, j_l)$,

$$\epsilon^I \wedge \epsilon^J = \epsilon^{IJ} \quad (19)$$

where $IJ = (i_1, \dots, i_k, j_1, \dots, j_l)$ is obtained by **concatenating** I and J .

- **Proposition 2.5 (Properties of the Wedge Product).**

Suppose $\omega, \omega', \eta, \eta'$ and ξ are **multicovectors** on a finite-dimensional vector space V .

1. (**Bilinearity**): For $a, a' \in \mathbb{R}$,

$$\begin{aligned} (a\omega + a'\omega') \wedge \eta &= a(\omega \wedge \eta) + a'(\omega' \wedge \eta), \\ \eta \wedge (a\omega + a'\omega') &= a(\eta \wedge \omega) + a'(\eta \wedge \omega'). \end{aligned}$$

2. (**Associativity**):

$$\omega \wedge (\eta \wedge \xi) = (\omega \wedge \eta) \wedge \xi$$

3. (**Anticommutativity**): For $\omega \in \Lambda^k(V^*)$ and $\eta \in \Lambda^l(V^*)$,

$$\omega \wedge \eta = (-1)^{kl} \eta \wedge \omega \quad (20)$$

4. If (ϵ^i) is any basis for V^* and $I = (i_1, \dots, i_k)$ is any multi-index, then

$$\epsilon^{i_1} \wedge \dots \wedge \epsilon^{i_k} = \epsilon^I \quad (21)$$

5. For any covectors $\omega^1, \dots, \omega^k$ and vectors v_1, \dots, v_k ,

$$(\omega^1 \wedge \dots \wedge \omega^k)(v_1, \dots, v_k) = \det(\omega^j(v_i)) \quad (22)$$

- **Remark** Because of part (4) of this lemma, henceforth we generally use the notations ϵ^I and $\epsilon^{i_1} \wedge \dots \wedge \epsilon^{i_k}$ **interchangeably**
- **Definition** A k -covector η is said to be **decomposable** if it can be expressed in the form $\eta = \omega^1 \wedge \dots \wedge \omega^k$, where $\omega^1, \dots, \omega^k$ are *covectors*.
- **Remark** It is important to be aware that not every k -covector is decomposable when $k > 1$; however, it follows from Proposition 2.2 and above Proposition 2.5 (4) that **every k -covector can be written as a linear combination of decomposable ones**.
- **Definition** For any n -dimensional vector space V , define a **vector space** $\Lambda(V^*)$ by

$$\Lambda(V^*) = \bigoplus_{k=0}^n \Lambda^k(V^*).$$

It follows from Proposition 2.2 that $\dim \Lambda(V^*) = 2^n$. The wedge product turns $\Lambda(V^*)$ into an **associative algebra**, called the exterior algebra (or **Grassmann algebra**) of V .

- **Remark** For any covectors $\omega^1, \dots, \omega^k$ and vectors v_1, \dots, v_k , **the exterior product** is considered as the **determinant function** of a $k \times k$ submatrix

$$(\omega^1 \wedge \dots \wedge \omega^k)(v_1, \dots, v_k) = \det \begin{bmatrix} \omega^1(v_1) & \dots & \omega^1(v_k) \\ \vdots & \ddots & \vdots \\ \omega^k(v_1) & \dots & \omega^k(v_k) \end{bmatrix}$$

where **vectors** v_1, \dots, v_k forms **column vector**, and **covectors** $\omega^1, \dots, \omega^k$ form the **row vector**.

In other words, we can think of **exterior product of covectors** as an abstraction of determinant operation.

2.3 Interior Product

- **Definition** Let V be a finite-dimensional vector space. For each $v \in V$, we define a *linear map* $\iota_v : \Lambda^k(V^*) \rightarrow \Lambda^{k-1}(V^*)$, called **interior multiplication (interior product)** by v , as follows:

$$(\iota_v \omega)(w_1, \dots, w_{k-1}) = \omega(v, w_1, \dots, w_{k-1}).$$

In other words, $(\iota_v \omega)$ is obtained from ω by **inserting v into the first slot**. By convention, we interpret $(\iota_v \omega)$ to be **zero** when ω is a **0-covector** (i.e., a **number**). Another common notation is

$$v \lrcorner \omega = (\iota_v \omega).$$

This is often read “ **v into ω** .”

- **Proposition 2.6** Let V be a finite-dimensional vector space and $v \in V$.

$$1. \quad \iota_v \circ \iota_v = 0.$$

$$2. \quad \text{If } \omega \in \Lambda^k(V^*) \text{ and } \eta \in \Lambda^l(V^*),$$

$$\iota_v(\omega \wedge \eta) = \iota_v(\omega) \wedge \eta + (-1)^k \omega \wedge \iota_v(\eta) \quad (23)$$

- **Remark** It is easy to verify the following form

$$\iota_v(\omega^1 \wedge \dots \wedge \omega^k) = v \lrcorner (\omega^1 \wedge \dots \wedge \omega^k) = \sum_{i=1}^k (-1)^{i-1} \omega^i(v) (\omega^1 \wedge \dots \wedge \widehat{\omega}^i \wedge \dots \wedge \omega^k) \quad (24)$$

$$\Leftrightarrow (\omega^1 \wedge \dots \wedge \omega^k)(v, v_2, \dots, v_k) = \sum_{i=1}^k (-1)^{i-1} \omega^i(v) (\omega^1 \wedge \dots \wedge \widehat{\omega}^i \wedge \dots \wedge \omega^k)(v_2, \dots, v_k)$$

where the hat indicates that ω^i is **omitted**. In *determinant form*, it can be written as

$$\det \mathbf{V} = \sum_{i=1}^k (-1)^{i-1} \omega^i(v) \det \mathbf{V}_1^i \quad (25)$$

where \mathbf{V}_j^i denote the $(k-1) \times (k-1)$ submatrix of \mathbf{V} obtained by **deleting** the i -th row and j -th column. This is just **the expansion of $\det \mathbf{V}$ by minors** along the first column, and therefore is equal to $\det \mathbf{v}$.

- **Remark** The *exterior product* **increase** the rank of tensor, while the *interior product* **decrease** the rank of tensor by 1.

2.4 Differential Forms on Manifolds

- **Definition** Let $T^k T^* M$ be the *bundle* of all covariant k -tensors on M . The subset of $T^k T^* M$ consisting of **alternating tensors** is denoted by $\Lambda^k(T^* M)$:

$$\Lambda^k(T^* M) = \bigsqcup_{p \in M} \Lambda^k(T_p^* M).$$

$\Lambda^k(T^* M)$ is a *smooth subbundle* of $T^k T^* M$, so it is a *smooth vector bundle* of rank $\binom{n}{k}$.

- **Remark** $\Lambda^k(T^*M)$ is *the bundle of all alternating covariant k -tensors (exterior forms, k -covectors)* on M .
- **Definition** A *section* of $\Lambda^k(T^*M)$ is called a differential k -form, or just a k -form; this is a (continuous) tensor field whose value at each point is an alternating tensor. The integer k is called the *degree of the form*. We denote the vector space of **smooth k -forms** by

$$\Omega^k(M) = \Gamma\left(\Lambda^k(T^*M)\right).$$

- **Remark** A k -form is just an alternating covariant k -tensor fields.
- **Remark** The *wedge product* of two differential forms is defined *pointwise*: $(\omega \wedge \eta)_p = \omega_p \wedge \eta_p$. Thus, the wedge product of a k -form with an l -form is a $(k+l)$ -form. If f is a 0-form (i.e. a smooth function) and ω is a k -form, we interpret the wedge product $f \wedge \omega$ to mean the ordinary product $f\omega$.
- **Remark** The direct sum of all vector spaces of smooth k -forms for $k \leq n$ is

$$\Omega^*(M) = \bigoplus_{k=0}^n \Omega^k(M). \quad (26)$$

Then the wedge product turns $\Omega^*(M)$ into an **associative, anticommutative graded algebra**.

- **Remark (Duality of Basis)**
The basis of differential k -forms $(dx^{i_1} \wedge \dots \wedge dx^{i_k})$ in $\Gamma(\Lambda^k(T^*M))$ acts on the local coordinate frames $(\partial/\partial x^i)$ in TM

$$(dx^{i_1} \wedge \dots \wedge dx^{i_k}) \left(\frac{\partial}{\partial x^{j_1}}, \dots, \frac{\partial}{\partial x^{j_k}} \right) = \delta_J^I$$

- **Remark (Coordinate Representation of k -Forms)**
In any smooth chart, a k -form ω can be written locally as

$$\omega = \sum_I \omega_I dx^I := \sum_I \omega_I dx^{i_1} \wedge \dots \wedge dx^{i_k}$$

where the coefficients ω^I are **continuous functions** defined on the coordinate domain, and we use dx^I as an abbreviation for $dx^{i_1} \wedge \dots \wedge dx^{i_k}$ (not to be mistaken for the differential of a real-valued function x^I). Also $\sum_I \epsilon^I$ means that sum with increasing multi-indices. **The component function** ω_I is computed as

$$\omega_I = \omega \left(\frac{\partial}{\partial x^{j_1}}, \dots, \frac{\partial}{\partial x^{j_k}} \right).$$

Note that ω_I is the determinant of a $k \times k$ principal sub-matrix (i.e. principal minors) whose rows and columns are indexed by increasing multi-index I .

- **Example** The followings are some basic differential k -forms:

1. Any smooth function $f \in \mathcal{C}^\infty(M)$ is a **0-form**;

2. A **differential 1-form** is the covariant vector field df

$$df = \sum_i \frac{\partial f}{\partial x^i} dx^i$$

3. A **differential 2-form** is written as

$$\omega = \sum_{i < j} \omega_{i,j} dx^i \wedge dx^j$$

- **Definition** If $F : M \rightarrow N$ is a smooth map and ω is a **differential form** on N , the **pullback** F^* is a **differential form on** M ; defined as for any covariant tensor field:

$$(F^*\omega)_p(v_1, \dots, v_k) = \omega_p(dF_p(v_1), \dots, dF_p(v_k)).$$

- **Lemma 2.7** Suppose $F : M \rightarrow N$ is smooth.

1. $F^* : \Omega^k(N) \rightarrow \Omega^k(M)$ is **linear** over \mathbb{R} .
2. $F^*(\omega \wedge \eta) = (F^*\omega) \wedge (F^*\eta)$.
3. In any smooth chart,

$$F^* \left(\sum_I \omega_I dy^{i_1} \wedge \dots \wedge dy^{i_k} \right) = \sum_I (\omega_I \circ F) d(y^{i_1} \circ F) \wedge \dots \wedge d(y^{i_k} \circ F) \quad (27)$$

- **Proposition 2.8 (Pullback Formula for Top-Degree Forms).**

Let $F : M \rightarrow N$ be a smooth map between n -manifolds with or without boundary. If (x^i) and (y^j) are smooth coordinates on open subsets $U \subseteq M$ and $V \subseteq N$, respectively, and u is a continuous real-valued function on V , then the following holds on $U \cap F^{-1}(V)$:

$$F^*(u dy^1 \wedge \dots \wedge dy^n) = (u \circ F) (\det(DF)) dx^1 \wedge \dots \wedge dx^n \quad (28)$$

where DF represents the **Jacobian matrix of** F in these coordinates.

Note that $d(y^i \circ F) = dF^i = \det(DF)_j^i dx^j$

- **Corollary 2.9 (Change of Coordinates for Differential Forms)**

If $(U, (x^i))$ and $(\tilde{U}, (\tilde{x}^j))$ are overlapping smooth coordinate charts on M , then the following identity holds on $U \cap \tilde{U}$:

$$d\tilde{x}^1 \wedge \dots \wedge d\tilde{x}^n = \det \left(\frac{\partial \tilde{x}^j}{\partial x^i} \right) dx^1 \wedge \dots \wedge dx^n. \quad (29)$$

- **Remark** The equation (28) provides a computational formula for pullback of differential forms under coordinate systems for domain and codomain. And the equation (29) provides the fomula for change of variables of differential forms.
- **Definition Interior multiplication** also extends naturally to **vector fields** and **differential forms**, simply by letting it act *pointwise*: if $X \in \mathfrak{X}(M)$ and $\omega \in \Omega^k(M)$, define a $(k-1)$ -form $X \lrcorner \omega = \iota_X \omega$ by

$$(X \lrcorner \omega)_p = X_p \lrcorner \omega_p.$$

- **Proposition 2.10** *Let X be a smooth vector field on M .*

1. *If ω is a smooth differential form, then $\iota_X \omega$ is smooth.*
2. *$\iota_X : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$ is **linear** over $\mathcal{C}^\infty(M)$ and therefore corresponds to a **smooth bundle homomorphism** $\iota_X : \Lambda^k(T^*M) \rightarrow \Lambda^{k-1}(T^*M)$.*

2.5 Exterior Derivatives of Differential Forms

- **Remark** For each smooth manifold M with or without boundary, we will show that there is **a differential operator** $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ satisfying $d(d\omega) = 0$ for all ω . Thus, it will follow that *a necessary condition* for a smooth k -form ω to be equal to $d\eta$ for some $(k-1)$ -form η is that $d\omega = 0$.
- **Definition** If $\omega = \sum_J' \omega_J dx^J$ is a smooth k -form on an open subset $U \subseteq \mathbb{R}^n$ or \mathbb{H}^n , we define its **exterior derivative** $d\omega$ to be the following $(k+1)$ -form:

$$d\omega := d \left(\sum_J' \omega_J dx^J \right) = \sum_J' d\omega_J \wedge dx^J, \quad (30)$$

where $d\omega_J$ is the differential of the function ω_J . In somewhat more detail, this is

$$d\omega := d \left(\sum_J' \omega_J dx^J \right) = \sum_J' \sum_i \frac{\partial \omega_J}{\partial x^i} dx^i \wedge dx^{j_1} \wedge \dots \wedge dx^{j_k}. \quad (31)$$

- **Remark** The *exterior derivatives* of a k -form is **a linear combination** of $(k+1)$ -forms. Its component function is **the principal minor of Jacobian matrix of component functions** $\left(\frac{\partial \omega_j}{\partial x^i} \right)$.
- **Remark** When ω is a 1-form, this becomes

$$\begin{aligned} d\omega &= d \left(\sum_j \omega_j dx^j \right) = \sum_j d\omega_j \wedge dx^j \\ &= \sum_j \sum_i \frac{\partial \omega_j}{\partial x^i} dx^i \wedge dx^j \\ &= \sum_{i < j} \frac{\partial \omega_j}{\partial x^i} dx^i \wedge dx^j + \sum_{i > j} \frac{\partial \omega_j}{\partial x^i} dx^i \wedge dx^j \\ &= \sum_{i < j} \left(\frac{\partial \omega_j}{\partial x^i} - \frac{\partial \omega_i}{\partial x^j} \right) dx^i \wedge dx^j. \end{aligned}$$

Note that the component is the determinant of a 2×2 sub-matrix of Jacobian $\left(\frac{\partial \omega_j}{\partial x^i} \right)$.

- **Remark** The **exterior differentiation** defines the differential of k -form. It is an **extension** of **differentiation** to **determinant function**.
- **Definition** If $A = \oplus_k A^k$ is a *graded algebra*, a *linear map* $T : A \rightarrow A$ is said to be a map **of degree m** if $T(A^k) \subseteq A^{k+m}$ for each k . It is said to be an **antiderivation** if it satisfies $T(xy) = (Tx)y + (-1)^k x(Ty)$ whenever $x \in A^k$ and $y \in A^l$.

- **Remark** (*The Exterior Differentiation vs. The Interior Multiplication*)
 1. The **exterior differentiation** $d : \Omega^k(M) \rightarrow \Omega^{k+1}(M)$ is an **antiderivation** of degree $+1$ whose **square is zero**.
 2. On the other hand, the **interior multiplication** $\iota_X : \Omega^k(M) \rightarrow \Omega^{k-1}(M)$ is an **antiderivation** of degree -1 whose **square is zero**, where $X \in \mathfrak{X}(M)$.
- Another important feature of the exterior derivative is that it **commutes with all pullbacks**.

Proposition 2.11 (*Naturality of the Exterior Derivative*).

If $F : M \rightarrow N$ is a smooth map, then for each k the pullback map $F^* : \Omega^k(N) \rightarrow \Omega^k(M)$ commutes with d : for all $\omega \in \Omega^k(N)$,

$$F^*(d\omega) = d(F^*\omega). \quad (32)$$

2.6 An Invariant Formula for the Exterior Derivative

- **Proposition 2.12** (*Exterior Derivative of a 1-Form*).
For any smooth 1-form ω and smooth vector fields X and Y ,

$$d\omega(X, Y) = X(\omega(Y)) - Y(\omega(X)) - \omega([X, Y]). \quad (33)$$

Proof: Since any smooth 1-form can be expressed locally as a sum of terms of the form $u dv$ for smooth functions u and v , it suffices to consider that case. Suppose $\omega = u dv$, and X, Y are smooth vector fields. The LHS of (33)

$$\begin{aligned} d(u dv)(X, Y) &= (du \wedge dv)(X, Y) = du(X)dv(Y) - du(Y)dv(X) \\ &= X(u)Y(v) - Y(u)X(v) \end{aligned}$$

The RHS is

$$\begin{aligned} &= X(u dv(Y)) - Y(u dv(X)) - u dv([X, Y]) \\ &= X(u Y(v)) - Y(u X(v)) - u [X, Y](v) \\ &= X(u)Y(v) + u XY(v) - Y(u)X(v) - u YX(v) - u ((XY - YX)v) \\ &= X(u)Y(v) - Y(u)X(v) + u (XY(v) - YX(v)) - u (XY(v) - YX(v)) \\ &= X(u)Y(v) - Y(u)X(v). \end{aligned}$$

Thus (33) holds. ■

- **Proposition 2.13** Let M be a smooth n -manifold with or without boundary, let (E_i) be a smooth local frame for M , and let (ϵ^i) be the dual coframe. For each i , let $b_{j,k}^i$ denote the **component functions of the exterior derivative of ϵ^i in this frame**, and for each j, k , let $c_{j,k}^i$ be the **component functions of the Lie bracket $[E_j, E_k]$** :

$$d\epsilon^i = \sum_{j < k} b_{j,k}^i \epsilon^j \wedge \epsilon^k; \quad [E_j, E_k] = c_{j,k}^i E_i$$

Then $b_{j,k}^i = -c_{j,k}^i$.

- **Proposition 2.14** (*Invariant Formula for the Exterior Derivative*).

Let M be a smooth manifold with or without boundary, and $\omega \in \Omega^k(M)$. For any smooth vector fields X_1, \dots, X_{k+1} on M ,

$$\begin{aligned} d\omega(X_1, \dots, X_{k+1}) &= \sum_{1 \leq i \leq k+1} (-1)^i X_i \left(\omega(X_1, \dots, \widehat{X}_i, \dots, X_{k+1}) \right) \\ &+ \sum_{1 \leq i < j \leq k+1} (-1)^{i+j} \omega([X_i, X_j], X_1, \dots, \widehat{X}_i, \dots, \widehat{X}_j, \dots, X_{k+1}), \end{aligned} \quad (34)$$

where the hats indicate **omitted** arguments.

- **Remark** The proof of formula (34) and (33) is only based on the definition of k -form and vector fields, and it does not involve any specific coordinate system. Thus it can be used to give an **invariant definition** of the **exterior differentiation** d .

3 Directional Derivatives of Vector Fields

3.1 Connections

- **Remark** There are *two alternatives* for the definition of **geodesics**:
 - Geodesics is the “**shortest**” path that connects two points on the surface; This definition is hard since the definition of manifold is abstract.
 - Geodesics is the curve on the surface that has **zero tangential acceleration**. This is the motivation to introduce the concept of **connections**.
- **Remark** Although the **velocity** of a curve γ in a manifold M is well defined, the **acceleration** of the curve on M is **not** since it requires comparison between tangent vectors in two different tangent spaces $T_{\gamma(t)}M$ and $T_{\gamma(t+\Delta)}M$.
- **Remark** To do so, we need a way to **compare values of the vector field at different points**, or intuitively, to “**connect**” **nearby tangent spaces**. This is where a connection comes in: it will be an additional piece of data on a manifold, a **rule** for **computing directional derivatives of vector fields**.
- **Remark** A **connection** is a **coordinate-independent** set of rules for taking **directional derivatives of vector fields**.
- **Definition** Let $\pi : E \rightarrow M$ be a smooth vector bundle over a smooth manifold M with or without boundary, and let $\Gamma(E)$ denote the space of smooth sections of E . A **connection** in E is a map

$$\nabla : \mathfrak{X}(M) \times \Gamma(E) \rightarrow \Gamma(E),$$

written $(X, Y) \mapsto \nabla_X Y$, satisfying the following properties:

1. $\nabla_X Y$ is **linear** over $\mathcal{C}^\infty(M)$ **in** X : for $f_1, f_2 \in \mathcal{C}^\infty(M)$ and $X_1, X_2 \in \mathfrak{X}(M)$,

$$\nabla_{(f_1 X_1 + f_2 X_2)} Y = f_1 \nabla_{X_1} Y + f_2 \nabla_{X_2} Y$$

2. $\nabla_X Y$ is **linear** over \mathbb{R} **in** Y : for $a_1, a_2 \in \mathbb{R}$ and $Y_1, Y_2 \in \Gamma(E)$,

$$\nabla_X(a_1 Y_1 + a_2 Y_2) = a_1 \nabla_X Y_1 + a_2 \nabla_X Y_2$$

3. ∇ satisfies the following **product rule**: for $f \in C^\infty(M)$,

$$\nabla_X(fY) = f \nabla_X Y + (Xf)Y$$

The symbol ∇ is read “**del**” or “**nabla**,” and $\nabla_X Y$ is called the covariant derivative of Y in the direction X .

- **Remark** There is a *variety of types of connections* that are useful in different circumstances. The type of connection we have defined here is sometimes called a **Koszul connection** to distinguish it from other types.

- **Lemma 3.1 (Locality).** [Lee, 2018]

Suppose ∇ is a connection in a smooth vector bundle $E \rightarrow M$. For every $X \in \mathfrak{X}(M)$, $Y \in \Gamma(E)$, and $p \in M$, the covariant derivative $\nabla_X Y|_p$ depends **only** on the values of X and Y in an arbitrarily **small neighborhood** of p . More precisely, if $X = \tilde{X}$ and $Y = \tilde{Y}$ on a neighborhood of p , then $\nabla_X Y|_p = \nabla_{\tilde{X}} \tilde{Y}|_p$.

- **Proposition 3.2 (Restriction of a Connection).** [Lee, 2018]

Suppose ∇ is a connection in a smooth vector bundle $E \rightarrow M$. For every open subset $U \subseteq M$, there is a **unique connection** ∇^U on the **restricted bundle** $E|_U$ that satisfies the following relation for every $X \in \mathfrak{X}(M)$ and $Y \in \Gamma(E)$:

$$\nabla_{(X|_U)}^U(Y|_U) = (\nabla_X Y)|_U. \quad (35)$$

- **Proposition 3.3** Under the hypotheses of Lemma 3.1, $\nabla_X Y|_p$ depends **only** on the **values** of Y in a **neighborhood** of p and the **value** of X **at** p .
- **Remark** In the situation of these two propositions, we typically just refer to the *restricted connection* as ∇ instead of ∇^U ; the proposition guarantees that there is no ambiguity in doing so. Thus if X is a vector field defined in a neighborhood of p ,

$$\nabla_v Y = \nabla_X Y|_p, \quad \text{for } v = X_p.$$

3.2 Connections in the Tangent Bundle

- We focus on the connection in tangent bundle.

Definition Suppose M is a smooth manifold with or without boundary. By the definition we just gave, a *connection in TM* is a map

$$\nabla : \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M),$$

satisfying properties (1)-(3) above. A *connection in the tangent bundle TM* is often called simply **a connection on M** . (The terms **affine connection** and **linear connection** are also sometimes used in this context.)

- **Definition** For computations, we need to examine how a connection appears in terms of a *local frame*. Let (E_i) be a *smooth local frame* for TM on an open subset $U \subseteq M$. For every choice of the indices i and j , we can expand the vector field $\nabla_{E_i} E_j$ in terms of this same frame:

$$\nabla_{E_i} E_j = \Gamma_{i,j}^k E_k. \quad (36)$$

As i, j , and k range from 1 to $n = \dim M$, this defines n^3 smooth functions $\Gamma_{i,j}^k : U \rightarrow \mathbb{R}$, called **the connection coefficients of ∇ with respect to the given frame**.

- The following proposition shows that the connection is completely determined in U by its connection coefficients.

Proposition 3.4 (Coordinate Representation of Connection) [Lee, 2018]

Let M be a smooth manifold with or without boundary, and let ∇ be a connection in TM . Suppose (E_i) is a smooth local frame over an open subset $U \subseteq M$, and let $\{\Gamma_{i,j}^k\}$ be the connection coefficients of ∇ with respect to this frame. For smooth vector fields $X, Y \in \mathfrak{X}(M)$, written in terms of the frame as $X = X^i E_i$, $Y = Y^j E_j$, one has

$$\nabla_X Y = \left(X(Y^k) + X^i Y^j \Gamma_{i,j}^k \right) E_k. \quad (37)$$

- **Remark** The n^3 functions $\{\Gamma_{i,j}^k\}$ are called **the Christoffel symbols** under the metric connections. [do Carmo Valero, 1976]
- **Remark** The smooth function $\Gamma_{i,j}^k \in C^\infty(M)$ has three indices: *two lower indices* (i, j) corresponds to the index of **component X^i for the directional vector field X** , and the index of **component Y^j for the differentiated vector field Y** in $\nabla_X Y$; *the one upper index k* corresponds to the index of the **basis** vector field $\partial/\partial x^k$ which spans the space of vector fields.
- **Remark** The *first term* of (37) accounts for the **change of position relative to the local frame** when moving Y from one tangent space to another along the direction of X . The second term accounts for the **additional “rotation” of frames**. For Euclidean space, the basis is fixed when moving along the tangent direction (i.e. no *rotation* just *translation*).
- **Proposition 3.5 (Transformation Law for Connection Coefficients)**. [Lee, 2018]
Let M be a smooth manifold with or without boundary, and let ∇ be a connection in TM . Suppose we are given two smooth local frames (E_i) and (\tilde{E}_j) for TM on an open subset $U \subseteq M$, related by $\tilde{E}_i = A_i^j E_j$ for some matrix of functions (A_i^j) . Let $\Gamma_{i,j}^k$ and $\tilde{\Gamma}_{i,j}^k$ denote the connection coefficients of ∇ with respect to these two frames. Then

$$\tilde{\Gamma}_{i,j}^k = (A^{-1})_t^k A_i^r A_j^s \Gamma_{r,s}^t + (A^{-1})_t^k A_i^s E_s(A_j^t) \quad (38)$$

- **Lemma 3.6** Suppose M is a smooth n -manifold with or without boundary, and M admits a **global frame** (E_i) . Formula (37) gives a **one-to-one correspondence** between connections in TM and choices of n^3 smooth real-valued functions $\{\Gamma_{i,j}^k\}$ on M .
- **Proposition 3.7** The tangent bundle of every smooth manifold with or without boundary admits a connection.

- **Proposition 3.8** (*The Difference Tensor*).

Let M be a smooth manifold with or without boundary. For any two connections ∇^0 and ∇^1 in TM , define a map $D : \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$ by

$$D(X, Y) = \nabla_X^0 Y - \nabla_X^1 Y.$$

Then D is **bilinear** over $\mathcal{C}^\infty(M)$, and thus defines a $(1, 2)$ -**tensor field** called **the difference tensor between ∇^0 and ∇^1** .

- **Theorem 3.9** Let M be a smooth manifold with or without boundary, and let ∇^0 be any connection in TM . Then **the set $\mathcal{A}(TM)$ of all connections in TM is equal to the following affine space:**

$$\mathcal{A}(TM) = \left\{ \nabla^0 + D : D \in \Gamma(T^{(1,2)}TM) \right\},$$

where $D \in \Gamma(T^{(1,2)}TM)$ is interpreted as a map from $\mathfrak{X}(M) \times \mathfrak{X}(M)$ to $\mathfrak{X}(M)$, and $\nabla^0 + D : \mathfrak{X}(M) \times \mathfrak{X}(M) \rightarrow \mathfrak{X}(M)$ is defined by

$$(\nabla^0 + D)(X, Y) = \nabla_X^0 Y + D(X, Y).$$

- **Remark** Finally we can define **the covariant derivative of every 1-form ω** based on connection on TM . In particular, **the connection on 1-form $\nabla : \mathfrak{X}(M) \times \mathfrak{X}^*(M) \rightarrow \mathfrak{X}^*(M)$** can be defined as

$$\begin{aligned} \langle \nabla_X \omega, Y \rangle &= \nabla_X \langle \omega, Y \rangle - \langle \omega, \nabla_X Y \rangle \\ \Rightarrow (\nabla_X \omega)(Y) &= X(\omega(Y)) - \omega(\nabla_X Y). \end{aligned} \quad (39)$$

where $\langle \omega, Y \rangle = \omega(Y)$ is a natural pairing. The coordinate representation of connection on 1-form is

$$\nabla_X \omega = (X(\omega_k) - X^j \omega_i \Gamma_{j,k}^i) \epsilon^k \quad (40)$$

where (ϵ^i) are coframes and $\omega = \omega_k \epsilon^k$, $X = X^i E_i$.

- **Remark** For a covariant 2-tensor field $g = g_{i,j} dx^i \otimes dx^j$, the covariant derivative of g in direction of Z is

$$(\nabla_{(Z)} g)(X, Y) = Z(g(X, Y)) - g(\nabla_Z X, Y) - g(X, \nabla_Z Y)$$

3.3 Total Covariant Derivatives

- **Proposition 3.10** (*The Total Covariant Derivative*). [Lee, 2018]

Let M be a smooth manifold with or without boundary and let ∇ be a connection in TM . For every $F \in \Gamma(T^{(k,l)}TM)$, the map

$$\nabla F : \underbrace{\Omega^1(M) \times \dots \times \Omega^1(M)}_k \times \underbrace{\mathfrak{X}(M) \times \dots \times \mathfrak{X}(M)}_{l+1} \rightarrow \mathcal{C}^\infty(M)$$

given by

$$\nabla F(\omega_1, \dots, \omega_k, Y_1, \dots, Y_l, X) = (\nabla_X F)(\omega_1, \dots, \omega_k, Y_1, \dots, Y_l) \quad (41)$$

defines a **smooth $(k, l+1)$ -tensor field on M called the total covariant derivative of F .**

- **Remark** The total covariant derivative of $Y \in \mathfrak{X}(M) := \Gamma(T^{(1,0)}TM)$ is a $(1,1)$ -**tensor field**

$$\nabla Y(\omega, X) = (\nabla_X Y)(\omega) = \omega(\nabla_X Y).$$

Similarly, the total covariant derivative of $\omega \in \mathfrak{X}^*(M) = \Omega^1(M) = \Gamma(T^{(0,1)}TM)$ is a $(0,2)$ -**tensor field**

$$\nabla \omega(Y, X) = (\nabla_X \omega)(Y) = X(\omega(Y)) - \omega(\nabla_X Y)$$

- **Remark** It can be verified that the following formula for total covariant derivative holds

$$\nabla_Y F = \text{tr}(\nabla F \otimes Y) \quad (42)$$

- **Definition** Given vector fields $X, Y \in \mathfrak{X}(M)$, let us introduce the notation $\nabla_{X,Y}^2 F$ for the (k,l) -tensor field obtained by inserting X, Y in the last two slots of $\nabla^2 F = \nabla(\nabla F)$:

$$\nabla_{X,Y}^2 F(\dots) = \nabla^2 F(\dots, Y, X)$$

- **Proposition 3.11** *Let M be a smooth manifold with or without boundary and let ∇ be a connection in TM . For every smooth vector field or tensor field F ,*

$$\nabla_{X,Y}^2 F = \nabla_X(\nabla_Y F) - \nabla_{(\nabla_X Y)} F. \quad (43)$$

- **Example (The Covariant Hessian).**

Let u be a smooth function on M .

- The total covariant derivative of a smooth function is equal to its 1-form $\nabla u = du \in \Omega^1(M) = \Gamma(T^{(0,1)}TM)$ since

$$\nabla u(X) = \nabla_X u = Xu = du(X)$$

- The 2-tensor $\nabla^2 u = \nabla(du)$ is called the covariant Hessian of u . Its action on smooth vector fields X, Y can be computed by the following formula:

$$\nabla^2 u(Y, X) = \nabla_{X,Y}^2 u = \nabla_X \nabla_Y u - \nabla_{(\nabla_X Y)} u = X(Yu) - (\nabla_X Y)(u) \quad (44)$$

In any local coordinates, it is

$$\nabla^2 u = u_{;i,j} dx^i \otimes dx^j$$

where

$$u_{;i,j} = \frac{\partial}{\partial x^j} \frac{\partial u}{\partial x^i} - \Gamma_{j,i}^k \frac{\partial u}{\partial x^k}$$

3.4 Vector and Tensor Fields Along Curves

- **Definition** Let M be a smooth manifold with or without boundary. Given a smooth curve $\gamma : I \rightarrow M$, a vector field along γ is a *continuous* map $V : I \rightarrow TM$ such that $V(t) \in T_{\gamma(t)}M$ for every $t \in I$; it is a smooth vector field along γ if it is *smooth* as a map from I to TM .

We let $\mathfrak{X}(\gamma)$ denote *the set of all smooth vector fields along γ* . It is a *real vector space* under pointwise vector addition and multiplication by constants, and it is a module over $C^\infty(I)$ with multiplication defined pointwise:

$$(fX)(t) = f(t)X(t).$$

- **Remark (Construction of A Smooth Vector Field Along the Curve)**

Suppose $\gamma : I \rightarrow M$ is a smooth curve and $\tilde{V} \in \mathfrak{X}(M)$ is a smooth vector field on an open subset of M containing the image of γ . The smooth vector field along the curve γ , $V = \tilde{V} \circ \gamma$:

$$V(t) = \tilde{V}_{\gamma(t)} \in T_{\gamma(t)}M.$$

A smooth vector field along γ is said to be *extendible* if there *exists* a smooth vector field \tilde{V} on a neighborhood of the image of γ that is related to V in this way.

Not every vector field along a curve need be extendible; for example, if $\gamma(t_1) = \gamma(t_2)$ but $\gamma'(t_1) \neq \gamma'(t_2)$, then γ' is not extendible.

- **Definition** More generally, a tensor field along γ is a *continuous* map σ from I to some tensor bundle $T^{(k,l)}TM$ such that $\sigma(t) \in T^{(k,l)}T_{\gamma(t)}M$ for each $t \in I$.

It is a *smooth tensor field along γ* if it is *smooth* as a map from I to $T^{(k,l)}TM$, and it is *extendible* if there is a smooth tensor field $\tilde{\sigma}$ on a neighborhood of $\gamma(I)$ such that $\tilde{\sigma} = \sigma \circ \gamma$.

- **Theorem 3.12 (Covariant Derivative Along a Curve).**

Let M be a smooth manifold with or without boundary and let ∇ be a connection in TM . For each smooth curve $\gamma : I \rightarrow M$, the *connection* determines *a unique operator*

$$D_t : \mathfrak{X}(\gamma) \rightarrow \mathfrak{X}(\gamma)$$

called the covariant derivative along γ , satisfying the following properties:

1. (**Linearity over \mathbb{R}**):

$$D_t(aV + bW) = aD_t(V) + bD_t(W), \quad \text{for } a, b \in \mathbb{R}.$$

2. (**Product Rule**):

$$D_t(fV) = f'V + fD_t(V), \quad \text{for } f \in C^\infty(I).$$

3. If $V \in \mathfrak{X}(\gamma)$ is *extendible*, then for every extension \tilde{V} of V ,

$$D_t(V(t)) = \nabla_{\gamma'(t)}\tilde{V}.$$

There is **an analogous operator** on the space of **smooth tensor fields** of any type along γ .

- **Remark (Coordinate Representation for Covariant Derivatives Along a Curve)**
Choose smooth coordinates (x^i) for M in a neighborhood of $\gamma(t_0)$, and write

$$V(t) = V^i(t) \frac{\partial}{\partial x^i} \Big|_{\gamma(t)}$$

for t near t_0 , where V^1, \dots, V^n are *smooth real-valued functions* defined on some neighborhood of t_0 in I . By the properties of D_t , since each $\frac{\partial}{\partial x^i}$ is extendible,

$$\begin{aligned} D_t(V_t) &= \dot{V}^i(t) \frac{\partial}{\partial x^i} \Big|_{\gamma(t)} + V^i(t) \nabla_{\gamma'(t)} \frac{\partial}{\partial x^i} \Big|_{\gamma(t)} \\ &= \left(\dot{V}^k(t) + \dot{\gamma}^i(t) V^j(t) \Gamma_{i,j}^k(\gamma(t)) \right) \frac{\partial}{\partial x^k} \Big|_{\gamma(t)} \end{aligned} \quad (45)$$

3.5 Geodesics

- **Definition** Let M be a smooth manifold with or without boundary and let ∇ be a connection in TM . For every smooth curve $\gamma : I \rightarrow M$, we define the **acceleration** of γ to be **the vector field $D_t(\gamma')$ along γ** .
- **Definition** A smooth curve γ is called a **geodesic** (**with respect to ∇**) if *its acceleration is zero*: $D_t(\gamma'(t)) = 0$.
- **Remark** **Geodesic** is the curve whose **tangential acceleration is zero**. From *the connection ∇ point of view*, it specifies both the directional vector field and the target vector field equal to $\gamma'(t)$. That is, the *tangential acceleration along a curve γ* is

$$\nabla_{\gamma'(t)} \gamma'(t).$$

- **Remark (The Ordinary Differential Equations for the Geodesic)**
In terms of smooth coordinates (x^i) , if we write the component functions of γ as $\gamma(t) = (x^1(t), \dots, x^n(t))$. From (45) and $D_t(\gamma'(t))$, we have a set of ordinary differential equations called **the geodesic equations**:

$$\ddot{x}^k(t) + \dot{x}^i(t) \dot{x}^j(t) \Gamma_{i,j}^k(x(t)) = 0, \quad k = 1, \dots, n. \quad (46)$$

where $x(t) := (x^1(t), \dots, x^n(t))$. A (parameterized) curve γ is a geodesic **if and only if** its component functions satisfy *the geodesic equations*. This is a set of **second-order nonlinear ODEs**.

- **Theorem 3.13 (Existence and Uniqueness of Geodesics).** [Lee, 2018]
Let M be a smooth manifold and ∇ a connection in TM . For every $p \in M$, $w \in T_p M$, and $t_0 \in \mathbb{R}$, there **exist** an open interval $I \subseteq \mathbb{R}$ containing t_0 and a **geodesic** $\gamma : I \rightarrow M$ satisfying $\gamma(t_0) = p$ and $\gamma'(t_0) = w$. Any two such geodesics **agree** on their common domain.
- **Remark** From the geodesic equation, we see that **the only parameters of the ODE that determines the geodesic is the coefficients of the connection $\{\Gamma_{i,j}^k\}$** . That is, *the geodesic is solely determined by the connection ∇* . Thus we also call it a **∇ -geodesic**.

- **Remark** The *geodesic equation under the initial boundary condition* can be written in the following form:

$$\dot{x}^k(t) = v^k(t) \quad (47)$$

$$\dot{v}^k(t) = -v^i(t)v^j(t)\Gamma_{i,j}^k(x(t)) \quad (48)$$

Treating $(x^1, \dots, x^n, v^1, \dots, v^n)$ as coordinates on $U \times \mathbb{R}^n$, we can recognize (48) as the equations for the **flow of the vector field** $G \in \mathfrak{X}(U \times \mathbb{R}^n)$ given by

$$G_{(x,v)} = v^k \frac{\partial}{\partial x^k} \Big|_{(x,v)} - v^i v^j \Gamma_{i,j}^k(x) \frac{\partial}{\partial v^k} \Big|_{(x,v)}. \quad (49)$$

The importance of G stems from the fact that it actually defines **a global vector field on the total space of TM** , called **the geodesic vector field**. It can be verified that the components of G under a change of coordinates *take the same form in every coordinate chart*.

Note that G acts on a function $f \in \mathcal{C}^\infty(U \times \mathbb{R}^n)$ as

$$Gf(p, v) = \frac{d}{dt} \Big|_{t=0} f(\gamma_v(t), \gamma'_v(t)). \quad (50)$$

- **Definition** A geodesic $\gamma : I \rightarrow M$ is said to be **maximal** if it *cannot be extended* to a geodesic on a *larger interval*, that is, if there does not exist a geodesic $\tilde{\gamma} : \tilde{I} \rightarrow M$ defined on an interval \tilde{I} properly containing I and satisfying $\tilde{\gamma}|_I = \gamma$.

A **geodesic segment** is a geodesic whose domain is a **compact interval**.

- **Corollary 3.14** Let M be a smooth manifold and let ∇ be a connection in TM . For each $p \in M$ and $v \in T_p M$, there is a **unique maximal geodesic** $\gamma : I \rightarrow M$ with $\gamma(0) = p$ and $\gamma'(0) = v$, defined on some open interval I containing 0.
- **Definition** The **unique maximal geodesic** γ with $\gamma(0) = p$ and $\gamma'(0) = v$ is often called simply **the geodesic with initial point p and initial velocity v** , and is denoted by γ_v . (Note that we can always find $p = \pi(v)$ where $\pi : TM \rightarrow M$ is the natural projection.)

3.6 Parallel Transport

- **Definition** Let M be a smooth manifold and let ∇ be a connection in TM . A *smooth vector or tensor field V along a smooth curve γ* is said to be **parallel along γ** (with respect to ∇) if $D_t(V) \equiv 0$.
- **Remark** A *geodesic* can be characterized as a curve whose **velocity vector field is parallel along the curve**.
- **Remark (Coordinate Representation of Vector Field Parallel Along a Curve)**
Given a smooth curve γ with a local coordinate representation $\gamma(t) = (\gamma^1(t), \dots, \gamma^n(t))$, formula (45) shows that a vector field V is parallel along γ if and only if

$$\dot{V}^k(t) + \dot{\gamma}^i(t)V^j(t)\Gamma_{i,j}^k(\gamma(t)) = 0, \quad k = 1, \dots, n \quad (51)$$

This is a set of **linear ordinary differential equations** with respect to $(V^1(t), \dots, V^n(t))$.

- **Theorem 3.15** (*Existence and Uniqueness of Parallel Transport*).

Suppose M is a smooth manifold with or without boundary, and ∇ is a connection in TM . Given a smooth curve $\gamma : I \rightarrow M$, $t_0 \in I$, and a vector $v \in T_{\gamma(t_0)}M$ or tensor $v \in T^{(k,l)}T_{\gamma(t_0)}M$, there exists a **unique parallel vector or tensor field** V along γ such that $V(t_0) = v$.

- **Remark** The vector or tensor field whose existence and uniqueness are proved in Theorem above is called **the parallel transport of v along γ** .
- **Definition** For each $t_0, t_1 \in I$, we define a map

$$P_{t_0, t_1}^\gamma : T_{\gamma(t_0)}M \rightarrow T_{\gamma(t_1)}M, \quad (52)$$

called **the parallel transport map**, by setting

$$P_{t_0, t_1}^\gamma(v) = V(t_1), \quad \forall v \in T_{\gamma(t_0)}M$$

where V is the parallel transport of v along γ .

This map is **linear**, because the equation of parallelism is linear. It is in fact an **isomorphism**, because P_{t_1, t_0}^γ is an **inverse** for it.

- **Remark** (*Parallel Frames Along a Curve*)

Given any basis (b_1, \dots, b_n) for $T_{\gamma(t_0)}M$, we can **parallel transport the vectors b_i along γ** , thus obtaining an n -tuple of **parallel vector fields** (E_1, \dots, E_n) along γ . Because each parallel transport map is an **isomorphism**, **the vectors $(E_i(t))$ form a basis for $T_{\gamma(t)}M$ at each point $\gamma(t)$** . Such an n -tuple of vector fields along γ is called **a parallel frame along γ** .

Every smooth (or piecewise smooth) vector field along γ can be expressed in terms of such a frame as

$$V(t) = V^i(t) E_i(t),$$

and then the properties of covariant derivatives along curves, together with the fact that the E_i 's are parallel, imply

$$D_t(V_t) = \dot{V}^i(t) E_i(t) \quad (53)$$

wherever V and γ are smooth. This means that **a vector field is parallel along γ if and only if its component functions with respect to the frame (E_i) are constants**.

- **Theorem 3.16** (*Parallel Transport Determines Covariant Differentiation*). [Lee, 2018]

Let M be a smooth manifold with or without boundary, and let ∇ be a connection in TM . Suppose $\gamma : I \rightarrow M$ is a smooth curve and V is a smooth vector field along γ . For each $t_0 \in I$,

$$D_t V(t_0) = \lim_{\Delta t \rightarrow 0} \frac{P_{(t_0+\Delta t), t_0}^\gamma(V(t_0 + \Delta t)) - V(t_0)}{\Delta t} \quad (54)$$

- **Corollary 3.17** (*Parallel Transport Determines the Connection*). [Lee, 2018]

Let M be a smooth manifold with or without boundary, and let ∇ be a connection in TM . Suppose X and Y are smooth vector fields on M . For every $p \in M$,

$$\nabla_X Y \Big|_p = \lim_{t \rightarrow 0} \frac{P_{t, 0}^\gamma(Y_{\gamma(t)}) - Y_p}{t}, \quad (55)$$

where $\gamma : I \rightarrow M$ is any smooth curve such that $\gamma(0) = p$ and $\gamma'(0) = X_p$.

- **Remark** See similarity between (55) and the definition of Lie derivatives:

$$(\mathcal{L}_X Y)_p = \lim_{t \rightarrow 0} \frac{d(\theta_{-t})_{\theta_t(p)} (Y_{\theta_t(p)}) - Y_p}{t},$$

where θ is the **flow of** X in the neighborhood of p such that $\theta_0(p) = p$, $(\theta^{(p)})'(0) = X_p$.

- **Remark** A smooth vector or tensor field on M is said to be **parallel** (with respect to ∇) if it is *parallel along every smooth curve* in M .
- **Proposition 3.18** Suppose M is a smooth manifold with or without boundary, ∇ is a connection in TM , and A is a **smooth vector or tensor field** on M . Then A is parallel on M if and only if $\nabla A \equiv 0$.
- **Remark** It is always possible to extend a vector at a point to a parallel vector field along any given curve. However, it may not be possible in general to extend it to a **parallel vector field** on an open subset of the manifold. The impossibility of finding such extensions is intimately connected with the phenomenon of **curvature**.
- **Remark** We see that both the concept of **connections** and the concept of **parallel transport along a curve** can be derived from each other.

$$\nabla \rightleftharpoons P_{t_0, t_1}^\gamma$$

They both define a way that “connects” the tangent space $T_p M$ at $p = \gamma(t_0)$ and the tangent space $T_q M$ at $q = \gamma(t_0 + \Delta t)$ in p ’s close neighborhood. The former begins with **a set of rules for a mapping** and the latter begins with **covariant derivatives along a curve**.

4 Lie Derivatives

- **Definition** Suppose M is a smooth manifold, V is a *smooth vector field* on M ; and θ is the **flow of** V . For any smooth vector field W on M , define **a rough vector field** on M , denoted by $\mathcal{L}_V W$ and called the **Lie derivative of W with respect to V** , by

$$\begin{aligned} (\mathcal{L}_V W)_p &= \lim_{t \rightarrow 0} \frac{d(\theta_{-t})_{\theta_t(p)} (W_{\theta_t(p)}) - W_p}{t} \\ &= \left. \frac{d}{dt} \right|_{t=0} d(\theta_{-t})_{\theta_t(p)} (W_{\theta_t(p)}), \end{aligned} \tag{56}$$

provided the derivative exists. For small $t \neq 0$, at least the difference quotient makes sense: θ_t is defined in a neighborhood of p , and θ_{-t} is the inverse of θ_t , so both $d(\theta_{-t})_{\theta_t(p)} (W_{\theta_t(p)})$ and W_p are elements of $T_p M$.

- **Remark** If M has nonempty boundary, this definition of $\mathcal{L}_V W$ makes sense as long as V is tangent to ∂M so that its flow exists.
- **Lemma 4.1** Suppose M is a smooth manifold with or without boundary, and $V, W \in \mathfrak{X}(M)$. If $\partial M \neq \emptyset$, assume in addition that V is tangent to ∂M . Then $(\mathcal{L}_V W)_p$ exists for every $p \in M$, and $\mathcal{L}_V W$ is a **smooth vector field**.
- The following theorem is critical to understand the **Lie derivatives** and **Lie bracket**.

Theorem 4.2 *If M is a smooth manifold and $V, W \in \mathfrak{X}(M)$, then $\mathcal{L}_V W = [V, W]$.*

- **Remark** This theorem allows us to extend the definition of the **Lie derivative** to arbitrary *smooth vector fields* on a smooth manifold M with boundary. Given $V, W \in \mathfrak{X}(M)$ we define $(\mathcal{L}_V W)_p$ for $p \in \partial M$ by embedding M in a smooth manifold \widetilde{M} without boundary (such as the double of M), extending V and W to smooth vector fields on \widetilde{M} , and computing the Lie derivative there. By virtue of the preceding theorem, $(\mathcal{L}_V W)_p = [V, W]_p$ is independent of the choice of extension.
- **Remark** This theorem also gives us a **geometric interpretation** of the Lie bracket of two vector fields: it is the **directional derivative** of the second vector field along the **flow** of the first.
- **Corollary 4.3** *Suppose M is a smooth manifold with or without boundary, and $V, W, X \in \mathfrak{X}(M)$.*
 1. (**Anti-symmetric**) $\mathcal{L}_V W = -\mathcal{L}_W V$.
 2. (**Product Rule**) $\mathcal{L}_V [W, X] = [\mathcal{L}_V W, X] + [W, \mathcal{L}_V X]$.
 3. $\mathcal{L}_{[V, W]} X = \mathcal{L}_V \mathcal{L}_W X - \mathcal{L}_W \mathcal{L}_V X$.
 4. If $g \in C^\infty(M)$, then $\mathcal{L}_V (gW) = (Vg)W + g\mathcal{L}_V W$.
 5. (**Pushforward**) If $F : M \rightarrow N$ is a **diffeomorphism**, then $F_*(\mathcal{L}_V X) = \mathcal{L}_{F_*V} F_*X$.
- **Remark** Note that the Lie derivative is **not linear over $C^\infty(M)$ in V** , i.e.

$$\mathcal{L}_{fV} W \neq f \mathcal{L}_V W$$

- **Remark** If V and W are vector fields on M and θ is the flow of V , the Lie derivative $(\mathcal{L}_V W)_p$, by definition, expresses the t -derivative of the **time-dependent vector** $d(\theta_{-t})_{\theta_t(p)}(W_{\theta_t(p)}) \in T_p M$ at $t = 0$. The next proposition shows how it can also be used to compute the derivative of this expression at other times.
- **Proposition 4.4** *Suppose M is a smooth manifold with or without boundary and $V, W \in \mathfrak{X}(M)$. If $\partial M \neq \emptyset$, assume also that V is tangent to ∂M . Let θ be the flow of V . For any (t_0, p) in the domain of θ ,*

$$\left. \frac{d}{dt} \right|_{t=t_0} d(\theta_{-t})_{\theta_t(p)}(W_{\theta_t(p)}) = d(\theta_{-t_0})_{\theta_{t_0}(p)}((\mathcal{L}_V W)_{\theta_{t_0}(p)}). \quad (57)$$

5 Lie Group and Lie Algebra

References

- Manfredo Perdigao do Carmo Valero. *Differential geometry of curves and surfaces*, volume 2. Prentice-hall Englewood Cliffs, 1976.
- John M Lee. *Introduction to Riemannian manifolds*, volume 176. Springer, 2018.
- John Marshall Lee. *Introduction to smooth manifolds*. Graduate texts in mathematics. Springer, New York, Berlin, Heidelberg, 2003. ISBN 0-387-95448-1.