

Lecture 4: The Entropy Methods

Tianpei Xie

Jan. 19th., 2023

Contents

1	Logarithmic Sobolev Inequality	2
1.1	Bernoulli Logarithmic Sobolev Inequality	2
1.2	Gaussian Logarithmic Sobolev Inequality	4
1.3	Information Theory Interpretation	6
1.4	Logarithmic Sobolev Inequality for General Probability Measures	7
1.5	Variants of Logarithmic Sobolev Inequalities	8
1.6	Poisson Logarithmic Sobolev Inequality	8
1.7	Applications	9
2	The Entropy Methods	9
2.1	Tensorization Property of Φ -Entropy	9
2.2	Herbst's Argument	9
2.3	Bounded Difference Inequality	9
2.4	Modified Logarithmic Sobolev Inequalities	9
2.5	Concentration of Convex Lipschitz Functions	9
2.6	Exponential Tail Bounds for Self-Bounding Functions	9

1 Logarithmic Sobolev Inequality

1.1 Bernoulli Logarithmic Sobolev Inequality

- **Remark (Setting)**

Consider a **uniformly distributed binary vector** $Z = (Z_1, \dots, Z_n)$ on the hypercube $\{-1, +1\}^n$. In other words, the components of X are *independent, identically distributed random sign (Rademacher) variables* with $\mathbb{P}\{Z_i = -1\} = \mathbb{P}\{Z_i = +1\} = 1/2$ (i.e. *symmetric Bernoulli random variables*).

Let $f : \{-1, +1\}^n \rightarrow \mathbb{R}$ be a real-valued function on **binary hypercube**. $X := f(Z)$ is an induced real-valued random variable. Define $\tilde{Z}^{(i)} = (Z_1, \dots, Z_{i-1}, Z'_i, Z_{i+1}, \dots, Z_n)$ be the sample Z with i -th component replaced by an *independent copy* Z'_i . Since $Z, \tilde{Z}^{(i)} \in \{-1, +1\}^n$, $\tilde{Z}^{(i)} = (Z_1, \dots, Z_{i-1}, -Z_i, Z_{i+1}, \dots, Z_n)$, i.e. *the i -th sign is flipped*. Also denote the i -th *Jackknife sample* as $Z_{(i)} = (Z_1, \dots, Z_{i-1}, Z_{i+1}, \dots, Z_n)$ by *leaving out* the i -th component. $\mathbb{E}_{(-i)}[X] := \mathbb{E}[X|Z_{(i)}]$.

Denote the i -th component of **discrete gradient** of f as

$$\nabla_i f(z) := \frac{1}{2} \left(f(z) - f(\tilde{z}^{(i)}) \right)$$

and $\nabla f(z) = (\nabla_1 f(z), \dots, \nabla_n f(z))$

- **Remark (Jackknife Estimate of Variance)**

Recall that *the Jackknife estimate of variance*

$$\begin{aligned} \mathcal{E}(f) &:= \mathbb{E} \left[\sum_{i=1}^n \left(f(Z) - \mathbb{E}_{(-i)}[f(\tilde{Z}^{(i)})] \right)^2 \right] \\ &= \frac{1}{2} \mathbb{E} \left[\sum_{i=1}^n \left(f(Z) - f(\tilde{Z}^{(i)}) \right)^2 \right]. \end{aligned}$$

Using the notation of discrete gradient of f , we see that

$$\mathcal{E}(f) := 2\mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right]$$

- **Remark (Entropy Functional)**

Recall that the entropy functional for f is defined as

$$H_\Phi(f(Z)) = \text{Ent}(f) := \mathbb{E}[f(Z) \log f(Z)] - \mathbb{E}[f(Z)] \log(\mathbb{E}[f(Z)]).$$

- **Proposition 1.1 (Logarithmic Sobolev Inequality for Rademacher Random Variables).** [Boucheron et al., 2013]

If $f : \{-1, +1\}^n \rightarrow \mathbb{R}$ be an arbitrary real-valued function defined on the n -dimensional **binary hypercube** and assume that Z is **uniformly distributed** over $\{-1, +1\}^n$. Then

$$\text{Ent}(f^2) \leq \mathcal{E}(f) \tag{1}$$

$$\Leftrightarrow \text{Ent}(f^2(Z)) \leq 2\mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right] \tag{2}$$

Proof: The key is to apply the tensorization property of Φ -entropy. Let $X = f(Z)$. By tensorization property,

$$\text{Ent}(X^2) \leq \sum_{i=1}^n \mathbb{E} [\text{Ent}_{(-i)}(X^2)]$$

where $\text{Ent}_{(-i)}(X^2) := \mathbb{E}_{(-i)} [X^2 \log X^2] - \mathbb{E}_{(-i)} [X^2] \log (\mathbb{E}_{(-i)} [X^2])$.

It thus suffice to show that for all $i = 1, \dots, n$,

$$\text{Ent}_{(-i)}(X^2) \leq \frac{1}{2} \mathbb{E}_{(-i)} \left[\left(f(Z) - f(\tilde{Z}^{(i)}) \right)^2 \right].$$

Given any fixed realization of $Z_{(-i)}$, $X = f(Z) = \tilde{f}(Z_i)$ can only takes two different values with equal probability. Call these two values a and b . See that

$$\begin{aligned} \text{Ent}_{(-i)}(X^2) &= \frac{1}{2} a^2 \log a^2 + \frac{1}{2} b^2 \log b^2 - \frac{1}{2} (a^2 + b^2) \log \left(\frac{a^2 + b^2}{2} \right) \\ \frac{1}{2} \mathbb{E}_{(-i)} \left[\left(f(Z) - f(\tilde{Z}^{(i)}) \right)^2 \right] &= \frac{1}{2} (a - b)^2. \end{aligned}$$

Thus we need to show

$$\frac{1}{2} a^2 \log a^2 + \frac{1}{2} b^2 \log b^2 - \frac{1}{2} (a^2 + b^2) \log \left(\frac{a^2 + b^2}{2} \right) \leq \frac{1}{2} (a - b)^2.$$

By symmetry, we may assume that $a \geq b$. Since $(|a| - |b|)^2 \leq (a - b)^2$, without loss of generality, we may further assume that $a, b \geq 0$.

Define

$$h(a) := \frac{1}{2} a^2 \log a^2 + \frac{1}{2} b^2 \log b^2 - \frac{1}{2} (a^2 + b^2) \log \left(\frac{a^2 + b^2}{2} \right) - \frac{1}{2} (a - b)^2$$

for $a \in [b, \infty)$. $h(b) = 0$. It suffice to check that $h'(b) = 0$ and that h is concave on $[b, \infty)$. Note that

$$\begin{aligned} h'(a) &= a \log a^2 + 1 - a \log \left(\frac{a^2 + b^2}{2} \right) - 1 - (a - b) \\ &= a \log \frac{2a^2}{(a^2 + b^2)} - (a - b). \end{aligned}$$

So $h'(b) = 0$. Moreover,

$$h''(a) = \log \frac{2a^2}{(a^2 + b^2)} + 1 - \frac{2a^2}{(a^2 + b^2)} \leq 0$$

due to inequality $\log(x) + 1 \leq x$. ■

- **Remark** (*Logarithmic Sobolev Inequality* \Rightarrow *Efron-Stein Inequality*). [Boucheron et al., 2013]

Note that for f non-negative,

$$\text{Var}(f(Z)) \leq \text{Ent}(f^2(Z)).$$

Thus *logarithmic Sobolev inequality* (1) implies

$$\text{Var}(f(Z)) \leq \mathcal{E}(f)$$

which is the *Efron-Stein inequality*.

- **Corollary 1.2** (*Logarithmic Sobolev Inequality for Asymmetric Bernoulli Random Variables*). [Boucheron et al., 2013]
If $f : \{-1, +1\}^n \rightarrow \mathbb{R}$ be an arbitrary real-valued function and $Z = (Z_1, \dots, Z_n) \in \{-1, +1\}^n$ with $p = \mathbb{P}\{Z_i = +1\}$. Then

$$\text{Ent}(f^2) \leq c(p) \mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right] \quad (3)$$

where

$$c(p) = \frac{1}{1-2p} \log \frac{1-p}{p}$$

Note that $\lim_{p \rightarrow 1/2} c(p) = 2$.

1.2 Gaussian Logarithmic Sobolev Inequality

- **Proposition 1.3** (*Gaussian Logarithmic Sobolev Inequality*). [Boucheron et al., 2013]
Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a **continuous differentiable** function and let $Z = (Z_1, \dots, Z_n)$ be a vector of n **independent standard Gaussian** random variables. Then

$$\text{Ent}(f^2(Z)) \leq 2 \mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right]. \quad (4)$$

Proof: We first prove for $n = 1$, where $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous differentiable and Z is standard Gaussian distribution. Without loss of generality, assume that $\mathbb{E}[f'(Z)] < \infty$ since it is trivial when $\mathbb{E}[f'(Z)] = \infty$. By density argument, it suffice to prove the proposition when f is twice differentiable with bounded support.

Now let $\epsilon_1, \dots, \epsilon_n$ be independent Rademacher random variables and introduce

$$S_n := \frac{1}{\sqrt{n}} \sum_{j=1}^n \epsilon_j.$$

Note that $\epsilon_i \in \{-1, +1\}$ with equal probability, thus

$$\begin{aligned} \mathbb{E}_{(-i)}[S_n] &= \frac{1}{2} \left[\left(\frac{1}{\sqrt{n}} \sum_{j \neq i} \epsilon_j + \frac{1}{\sqrt{n}} \right) + \left(\frac{1}{\sqrt{n}} \sum_{j \neq i} \epsilon_j - \frac{1}{\sqrt{n}} \right) \right] \\ &= \frac{1}{2} \left[\left(S_n + \frac{1 - \epsilon_i}{\sqrt{n}} \right) + \left(S_n - \frac{1 + \epsilon_i}{\sqrt{n}} \right) \right]. \end{aligned}$$

In the proof of Gaussian Poincaré inequality, we show that by *central limit theorem*,

$$\limsup_{n \rightarrow \infty} \mathbb{E} \left[\sum_{i=1}^n \left| f(S_n) - f\left(S_n - \frac{2\epsilon_i}{\sqrt{n}}\right) \right|^2 \right] = 4 \mathbb{E}[(f'(Z))^2].$$

On the other hands, for any *continuous uniformly bounded function* f , by *central limit theorem*,

$$\lim_{n \rightarrow \infty} \text{Ent} (f^2(S_n)) = \text{Ent}(f^2(Z))$$

The proof is then completed by invoking *the logarithmic Sobolev inequality* for *Rademacher random variables*

$$\begin{aligned} \text{Ent} (f^2(S_n)) &\leq \frac{1}{2} \mathbb{E} \left[\sum_{i=1}^n \left| f(S_n) - f \left(S_n - \frac{2\epsilon_i}{\sqrt{n}} \right) \right|^2 \right] \\ \Rightarrow \lim_{n \rightarrow \infty} \text{Ent} (f^2(S_n)) &\leq \frac{1}{2} \lim_{n \rightarrow \infty} \mathbb{E} \left[\sum_{i=1}^n \left| f(S_n) - f \left(S_n - \frac{2\epsilon_i}{\sqrt{n}} \right) \right|^2 \right] \\ &\Rightarrow \text{Ent}(f^2(Z)) \leq 2\mathbb{E} [(f'(Z))^2]. \end{aligned}$$

The extension of the result to dimension $n \geq 1$ follows easily from *the sub-additivity of entropy* which states that

$$\text{Ent}(f^2) \leq \sum_{i=1}^n \mathbb{E} [\mathbb{E}_{(-i)} [f^2(Z) \log f^2(Z)] - \mathbb{E}_{(-i)} [f^2(Z)] \log \mathbb{E}_{(-i)} [f^2(Z)]]$$

where $\mathbb{E}_{(-i)} [\cdot]$ denotes the integration with respect to i -th variable Z_i only. Thus by induction, for all i

$$\mathbb{E}_{(-i)} [f^2(Z) \log f^2(Z)] - \mathbb{E}_{(-i)} [f^2(Z)] \log \mathbb{E}_{(-i)} [f^2(Z)] \leq 2\mathbb{E}_{(-i)} [(\partial_i f(Z))^2].$$

Thus

$$\text{Ent}(f^2) \leq 2\mathbb{E} \left[\mathbb{E}_{(-i)} \left[\sum_{i=1}^n (\partial_i f(Z))^2 \right] \right] = 2\mathbb{E} [\|\nabla f(Z)\|_2^2]. \quad \blacksquare$$

- **Remark (*Dimension Free Property*).**

The *Gaussian logarithmic Sobolev inequality* has a constant $C = 2$ that is ***independent of dimension*** n :

$$\mathbb{E}_\mu [f^2] \leq 2\mathbb{E}_\mu [\|\nabla f\|_2^2].$$

This *dimension-free property* is related to the ***concentration of Gaussian measure*** μ . As a consequence, this inequality can be extended to functions of *Gaussian measure* on ***infinite dimensional space***, such as Gibbs measure, *Gaussian process* etc.

- **Remark (*Equivalent Form of Gaussian Logarithmic Sobolev Inequality*)**

Assume $f : \mathbb{R}^n \rightarrow (0, \infty)$ and $\int_{\mathbb{R}^n} f d\mu = 1$ under Gaussian measure μ . Substituting $f \rightarrow \sqrt{f}$, the *logarithmic Sobolev inequality* becomes

$$\text{Ent}_\mu(f) = \int f \log f d\mu \leq \frac{1}{2} \int \frac{\|\nabla f\|_2^2}{f} d\mu \quad (5)$$

- **Remark** (*Gaussian Logarithmic Sobolev Inequality \Rightarrow Gaussian Poincaré Inequality*). [Boucheron et al., 2013]

Recall that *the Gaussian Poincaré inequality*

$$\text{Var}(f(Z)) \leq \mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right]$$

Since

$$(1+t) \log(1+t) = t + \frac{t^2}{2} + o(t^2)$$

as $t \rightarrow 0$, we can get for Gaussian measures,

$$\text{Ent}_\mu(1 + \epsilon h) = \frac{\epsilon^2}{2} \text{Var}_\mu(h) + o(\epsilon^2).$$

Similarly,

$$\int \frac{\|\nabla(1 + \epsilon h)\|_2^2}{1 + \epsilon h} d\mu = \epsilon^2 \int \|\nabla h\|_2^2 d\mu + o(\epsilon^2).$$

Thus from *the Gaussian logarithmic Sobolev inequality*,

$$\begin{aligned} \text{Ent}_\mu(1 + \epsilon h) &\leq \frac{1}{2} \int \frac{\|\nabla(1 + \epsilon h)\|_2^2}{1 + \epsilon h} d\mu \\ \Leftrightarrow \frac{\epsilon^2}{2} \text{Var}_\mu(h) + o(\epsilon^2) &\leq \frac{\epsilon^2}{2} \int \|\nabla h\|_2^2 d\mu + o(\epsilon^2) \\ \Leftrightarrow \text{Var}(f(Z)) &\leq \mathbb{E} \left[\|\nabla f(Z)\|_2^2 \right] \quad \text{as } \epsilon \rightarrow 0. \end{aligned}$$

Thus *the Gaussian logarithmic Sobolev inequality* implies *the Gaussian Poincaré inequality*.

1.3 Information Theory Interpretation

- **Remark** (*Information Interpretation of Gaussian Logarithmic Sobolev Inequality*)

Let ν, μ be two probability measures on $(\mathcal{X}, \mathcal{F})$ and $\nu \ll \mu$. Define $f := \frac{d\nu}{d\mu}$ be the Radon-Nikodym derivative of ν with respect to μ (i.e f is the probability density function of ν with respect to μ). Then the entropy becomes *the relative entropy*

$$\text{Ent}_\mu(f) := \mathbb{E}_\mu[f \log f] = \text{KL}(\nu \parallel \mu)$$

since $\mathbb{E}_\mu[f] = \int_{\mathcal{X}^n} f d\mu = 1$.

On the other hand, *the (relative) Fisher information* is defined as

$$\begin{aligned} I(\nu \parallel \mu) &:= \mathbb{E}_\nu \left[\|\nabla \log f\|_2^2 \right] \\ &= \int \left\| \frac{\nabla f}{f} \right\|_2^2 d\nu = \int \frac{\|\nabla f\|_2^2}{f^2} d\nu \\ &= \int \frac{\|\nabla f\|_2^2}{f} d\mu \end{aligned}$$

Thus *the information interpretation* of the Gaussian logarithmic Sobolev inequality is

$$\text{KL}(\nu \parallel \mu) \leq \frac{1}{2} I(\nu \parallel \mu) \tag{6}$$

Note that *the Fisher information metric* is *the Riemannian metric* induced by the relative entropy.

1.4 Logarithmic Sobolev Inequality for General Probability Measures

- **Remark** (*The Sobolev Inequality*)

The Sobolev inequality states for smooth function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ where $n \geq 3$ and $p = \frac{2n}{n-2} > 2$

$$\|f\|_p^2 \leq C_n \int_{\mathbb{R}^n} |\nabla f|^2 dx$$

where the inequality is sharp when the constant

$$C_n := \frac{1}{\pi n(n-2)} \left(\frac{\Gamma(n)}{\Gamma(n/2)} \right)^{2/n}$$

- **Proposition 1.4** (*Euclidean Logarithmic Sobolev Inequality*).

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function and m be Lebesgue measure on \mathbb{R}^n , then

$$\begin{aligned} \text{Ent}_m(f^2) &\leq \frac{n}{2} \log \left(\frac{2}{n\pi e} \mathbb{E}_m \left[\|\nabla f\|_2^2 \right] \right) \\ \Leftrightarrow \int f^2 \log \left(\frac{f^2}{\int f^2 dx} \right) dx &\leq \frac{n}{2} \log \left(\frac{2}{n\pi e} \int |\nabla f|^2 dx \right) \end{aligned} \quad (7)$$

- **Definition** (*Logarithmic Sobolev Inequality for General Probability Measure*).

A probability measure μ on \mathbb{R}^n is said to satisfy the logarithmic Sobolev inequality for some constant $C > 0$ if for any smooth function f

$$\text{Ent}_\mu(f^2) \leq C \mathbb{E}_\mu \left[\|\nabla f\|_2^2 \right] \quad (8)$$

holds for any *continuous differentiable* function $f : \mathbb{R}^n \rightarrow \mathbb{R}$. The left-hand side is called *the entropy functional*, which is defined as

$$\begin{aligned} \text{Ent}(f^2) &:= \mathbb{E}_\mu [f^2 \log f^2] - \mathbb{E}_\mu [f^2] \log \mathbb{E}_\mu [f^2] \\ &= \int f^2 \log \left(\frac{f^2}{\int f^2 d\mu} \right) d\mu. \end{aligned}$$

The right-hand side is defined as

$$\mathbb{E}_\mu \left[\|\nabla f\|_2^2 \right] = \int \|\nabla f\|_2^2 d\mu.$$

Thus we can rewrite *the logarithmic Sobolev inequality* in functional form

$$\int f^2 \log \left(\frac{f^2}{\int f^2 d\mu} \right) d\mu \leq C \int \|\nabla f\|_2^2 d\mu \quad (9)$$

- **Remark** (*Logarithmic Sobolev Inequality*)

For non-negative function f , we can replace $f \rightarrow \sqrt{f}$, so that *the logarithmic Sobolev inequality* becomes

$$\text{Ent}_\mu(f) \leq \frac{1}{2} \int \frac{\|\nabla f\|_2^2}{f} d\mu \quad (10)$$

- **Remark (Modified Logarithmic Sobolev Inequality via Convex Cost and Duality)**
For some **convex non-negative cost** $c : \mathbb{R}^n \rightarrow \mathbb{R}_+$, the **convex conjugate** of c (Legendre transform of c) is defined as

$$c^*(x) := \sup_y \{ \langle x, y \rangle - c(y) \}$$

Then we can obtain **the modified logarithmic Sobolev inequality**

$$\text{Ent}_\mu(f) \leq \int f^2 c^* \left(\frac{\nabla f}{f} \right) d\mu \quad (11)$$

1.5 Variants of Logarithmic Sobolev Inequalities

- **Proposition 1.5 (A Modified Logarithmic Sobolev Inequalities for Moment Generating Function)** [Boucheron et al., 2013]

Consider independent random variables Z_1, \dots, Z_n taking values in \mathcal{X} , a real-valued function $f : \mathcal{X}^n \rightarrow \mathbb{R}$ and the random variable $X = f(Z_1, \dots, Z_n)$. Also denote $Z_{(-i)} = (Z_1, \dots, Z_{i-1}, Z_{i+1}, \dots, Z_n)$ and $X_{(-i)} = f_i(Z_{(-i)})$ where $f_i : \mathcal{X}^{n-1} \rightarrow \mathbb{R}$ is an arbitrary function. Let $\phi(x) = e^x - x - 1$. Then for all $\lambda \in \mathbb{R}$,

$$\lambda \mathbb{E} [X e^{\lambda X}] - \mathbb{E} [e^{\lambda X}] \log \mathbb{E} [e^{\lambda X}] \leq \sum_{i=1}^n \mathbb{E} [e^{\lambda X} \phi(-\lambda(X - X_{(-i)}))] \quad (12)$$

- **Proposition 1.6 (Symmetrized Modified Logarithmic Sobolev Inequalities)** [Boucheron et al., 2013]

Consider independent random variables Z_1, \dots, Z_n taking values in \mathcal{X} , a real-valued function $f : \mathcal{X}^n \rightarrow \mathbb{R}$ and the random variable $X = f(Z_1, \dots, Z_n)$. Also denote $\tilde{X}^{(i)} = f(Z_1, \dots, Z_{i-1}, Z'_i, Z_{i+1}, \dots, Z_n)$. Let $\phi(x) = e^x - x - 1$. Then for all $\lambda \in \mathbb{R}$,

$$\lambda \mathbb{E} [X e^{\lambda X}] - \mathbb{E} [e^{\lambda X}] \log \mathbb{E} [e^{\lambda X}] \leq \sum_{i=1}^n \mathbb{E} [e^{\lambda X} \phi(-\lambda(X - \tilde{X}^{(i)}))] \quad (13)$$

Moreover, denoting $\tau(x) = x(e^x - 1)$, for all $\lambda \in \mathbb{R}$,

$$\begin{aligned} \lambda \mathbb{E} [X e^{\lambda X}] - \mathbb{E} [e^{\lambda X}] \log \mathbb{E} [e^{\lambda X}] &\leq \sum_{i=1}^n \mathbb{E} [e^{\lambda X} \tau(-\lambda(X - \tilde{X}^{(i)})_+)], \\ \lambda \mathbb{E} [X e^{\lambda X}] - \mathbb{E} [e^{\lambda X}] \log \mathbb{E} [e^{\lambda X}] &\leq \sum_{i=1}^n \mathbb{E} [e^{\lambda X} \tau(\lambda(\tilde{X}^{(i)} - X)_-)]. \end{aligned}$$

1.6 Poisson Logarithmic Sobolev Inequality

- **Proposition 1.7 (Modified Logarithmic Sobolev Inequality for Bernoulli Random Variable).** [Boucheron et al., 2013]

Let $f : \{0, 1\} \rightarrow (0, \infty)$ be a **non-negative** real-valued function defined on the binary set $\{0, 1\}$. Define **the discrete derivative** of f at $x \in \{0, 1\}$ by

$$\nabla f := f(1 - x) - f(x).$$

Let X be a Bernoulli random variable with parameter $p \in (0, 1)$ (i.e. $\mathbb{P}\{X = 1\} = p$). Then

$$\text{Ent}(f(X)) \leq (p(1-p))\mathbb{E}[\nabla f(X)\nabla \log f(X)]. \quad (14)$$

and

$$\text{Ent}(f(X)) \leq (p(1-p))\mathbb{E}\left[\frac{|\nabla f(X)|^2}{f(X)}\right]. \quad (15)$$

- **Proposition 1.8 (Poisson Logarithmic Sobolev Inequality).** [Boucheron et al., 2013]
Let $f : \mathbb{N} \rightarrow (0, \infty)$ be a **non-negative** real-valued function defined on the set of non-negative integers \mathbb{N} . Define **the discrete derivative** of f at $x \in \mathbb{N}$ by

$$\nabla f := f(x+1) - f(x).$$

Let X be a Poisson random variable. Then

$$\text{Ent}(f(X)) \leq (\mathbb{E}[X])\mathbb{E}[\nabla f(X)\nabla \log f(X)]. \quad (16)$$

and

$$\text{Ent}(f(X)) \leq (\mathbb{E}[X])\mathbb{E}\left[\frac{|\nabla f(X)|^2}{f(X)}\right]. \quad (17)$$

1.7 Applications

2 The Entropy Methods

2.1 Tensorization Property of Φ -Entropy

- **Remark** Recall that the Φ -entropy for $\Phi(x) = x \log(x)$ as

$$H_\Phi(X) = \text{Ent}(X) := \mathbb{E}[X \log X] - \mathbb{E}[X] \log(\mathbb{E}[X]).$$

The variational formulation of $H_\Phi(X)$ is

$$\text{Ent}(X) = \sup_T \{X(\log(T) - \log(\mathbb{E}[T]))\}$$

•

2.2 Herbst's Argument

2.3 Bounded Difference Inequality

2.4 Modified Logarithmic Sobolev Inequalities

2.5 Concentration of Convex Lipschitz Functions

2.6 Exponential Tail Bounds for Self-Bounding Functions

References

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.