

Mathematical Induction (Lecture)

Worksheet 5: MI^s on Sequences, Pascal's Triangle, and Pie Fights¹

Date: 9/25/2021

MATH 74: Transition to Upper-Division Mathematics

with Roy Zhao, UC Berkeley

Read: *Session 6: Mathematical Induction* (vol. I)

- §4. Strong Induction, up to Exercise 14 (pp. 115-119)

Write: clearly. Supply your reasoning in words and/or symbols. Show calculations and relevant pictures.

- (Recurrences)** Prove the direct formulas by MI^s.
 - If $a_1 = 3$, $a_2 = 5$, and $a_{n+1} = 3a_n - 2a_{n-1}$ for all $n \geq 2$, then $a_n = 2^n + 1$ for all $n \in \mathbb{N}$.
 - (Fibonacci*)** If $f_0 = 0$, $f_1 = 1$, and $\forall n \geq 1$ $f_{n+1} = f_n + f_{n-1}$, then $\forall n \geq 0$:
$$f_n = \frac{1}{\sqrt{5}}(\phi^n - \bar{\phi}^n)$$
 for $\phi = \frac{1+\sqrt{5}}{2}$, $\bar{\phi} = \frac{1-\sqrt{5}}{2}$.
Is there a non-MI way to arrive at these formulas?
(Hint: Use some linear algebra: eigenvalues?)
- (Sequence Cycling)** Let $a_{n+1} = a_n - a_{n-1}$ for any $n \geq 2$, where $a_1 = 1$ and $a_2 = 2$.
 - Find a pattern for $\{a_n\}$ and prove it.
 - Prove that $a_n = -2 \cos \frac{(n+1)\pi}{3}$ for all $n \geq 1$.
 - (C-preview*)** How could we have arrived at the direct formula in (b) without guessing?
(Die-Hards Hint: If $x^2 = x - 1$ has roots $r_{1,2}$, set $a_n = Ar_1^n + Br_2^n$. With $n = 0, 1$, create a linear system to calculate that $A = -r_1$ and $B = -r_2$. Finally, use de Moivre's formula to rewrite r_1^{n+1} and r_2^{n+1} and simplify the direct formula for a_n .)
- Each statement below depends on two variables, or has a double index. Decide on what you will induct, and using MI^s, prove the statement.
 - (Pascal's Triangle)** The number in the n th row and k th diagonal of Pascal's Triangle equals the binomial coefficient $\binom{n}{k}$ for $n, k \geq 0$.
- (True/False Jeopardy)** Supply convincing reasoning for your answer.
 - T F For a recurrence sequence with $a_{n+1} = 3a_n - 4a_{n-1} + 2a_{n-2}$, to prove a conjectured equation for a_n using MI^s, we will have 3 base cases.
 - T F MI helps us come up with direct formulas for sequences.
 - T F A bank that has an unlimited supply of 4-peso and 10-peso bills can pay any even number of pesos greater than 6.
 - T F Sending two or more newly-wed couples on honeymoons to different planets proves that the Pie-Throwing Problem is false for an even number of people.
- (Hockey-Stick)** For $n \geq k \geq 0$:
$$\binom{k}{k} + \binom{k+1}{k} + \binom{k+2}{k} + \cdots + \binom{n}{k} = \binom{n+1}{k+1}.$$
- (Double Fibonacci)** For all $n, m \geq 0$:
$$f_{n+m+1} = f_n f_m + f_{n+1} f_{m+1}.$$

Re-prove this, using the direct formula for f_n .
- (Pie-Fight*)** There is an *odd* number of people (at least 3) and the distances between any two of them are all *different*. Every person throws a pie into the face of his/her closest neighbor.
 - Using MI, prove that there will be at least one "survivor" (a person not "pied" by another).
 - Why should the number of people be *odd*; e.g., what is wrong with having 100 people making the pie fight? What if some distances were the *same*? Does this change the problem?
- (Finite Induction)** Recall Problem 2, IMO '88: If the ratio $\frac{a^2+b^2}{1+ab} \in \mathbb{Z}$ for some $a, b \in \mathbb{N}$, then this ratio is a perfect square. In the solution, we created a chain of pairs (a_n, b_n) of non-negative numbers that were solutions for the same ratio k , and which steadily got "smaller" until a pair with a 0 was hit: $(0, b_n)$ or $(a_n, 0)$.
 - What quantity exactly "got smaller"?
(This is called a decreasing *monovariant*.)
 - What ensures that we reach a pair with a 0? Can't we go on forever? Was MI involved?

¹These worksheets are copyrighted and provided for the personal use of Fall 2021 MATH 74 students only. They may not be reproduced or posted anywhere without explicit written permission from the instructor.