

ON THE 2-ADIC STRUCTURE OF ZAGIER'S MZV MATRICES

YIPIN WANG

ABSTRACT. We investigate the 2-adic properties of the inverse of Zagier's matrix M_K , which expresses Hoffman elements $H(a, b) = \zeta(2, \dots, 2, 3, 2, \dots, 2)$ as rational linear combinations of products $\zeta(2)^m \zeta(2n+1)$. We prove that all entries in the last row of $(M_K)^{-1}$ have 2-adic valuation zero (Theorem 1), implying that all coefficients in the decomposition of $\zeta(2)^{K-1} \zeta(3)$ into the Hoffman basis are odd integers. As a companion result, we establish a closed-form inverse for the binomial core matrix $B_N[a, i] = \binom{2i}{2a}$ (Theorem 6): its inverse is given explicitly in terms of the Euler-secant numbers E_{2n} and the hyperbolic secant function, with the exact 2-adic valuation of every entry governed by binary carry counting via Kummer's theorem. As byproducts, we obtain the closed formula $v_2(\det M_K) = 2K - s_2(K) - K^2$ and the congruence $E_{2n} \equiv 1 \pmod{4}$ for all $n \geq 0$.

1. INTRODUCTION

Multiple zeta values (MZVs) are real numbers defined for positive integers k_1, \dots, k_n with $k_n \geq 2$ by the convergent series

$$\zeta(k_1, \dots, k_n) = \sum_{0 < m_1 < \dots < m_n} \frac{1}{m_1^{k_1} \dots m_n^{k_n}}.$$

A central result, proved by Brown [1], states that every MZV is a \mathbb{Q} -linear combination of the *Hoffman basis* elements $\zeta(k_1, \dots, k_n)$ where each $k_i \in \{2, 3\}$.

Zagier [7] gave explicit formulas for the special MZVs

$$H(a, b) := \zeta(\underbrace{2, \dots, 2}_a, 3, \underbrace{2, \dots, 2}_b)$$

Date: February 5, 2026.

2020 *Mathematics Subject Classification.* 11M32 (primary), 11S80, 05A10 (secondary).

Key words and phrases. multiple zeta values, Hoffman basis, 2-adic valuation, Zagier matrix, Euler numbers, binary carries.

as rational linear combinations of products $\zeta(2)^m \zeta(2n+1)$. For each odd weight $w = 2K+1$, this gives a $K \times K$ matrix M_K expressing the vector of Hoffman elements $\{H(a, K-1-a)\}_{a=0}^{K-1}$ in terms of products $\{\zeta(2)^m \zeta(2(K-m)+1)\}_{m=0}^{K-1}$.

Zagier proved that $\det(M_K) \neq 0$ using a 2-adic argument: the matrix is upper triangular modulo 2 with odd diagonal entries. This 2-adic structure played a crucial role in Brown's motivic proof [1].

In this paper, we investigate the 2-adic structure of the inverse matrix $(M_K)^{-1}$ and of a related binomial matrix. Our first main result (Theorem 1) establishes that all entries in the last row of $(M_K)^{-1}$ have 2-adic valuation zero, implying that the decomposition of $\zeta(2)^{K-1} \zeta(3)$ into the Hoffman basis has exclusively odd coefficients. Our second main result (Theorem 6) gives a closed-form inverse for the *binomial core matrix* $B_N[a, i] = \binom{2i}{2a}$, expressed in terms of the Euler-secant numbers and the function $\operatorname{sech}(x)$, with the exact 2-adic valuation governed by the binary carry function.

These two results are connected by a common mechanism: binary carry counting via Kummer's theorem. In the Zagier setting, this mechanism appears through the binomial coefficients $\binom{2r}{2b+1}$ that dominate the 2-adic structure of each column. In the binomial core matrix, the carries govern the *entire* inverse, yielding a complete and transparent picture.

2. STATEMENT OF RESULTS

Let $v_2(x)$ denote the 2-adic valuation of $x \in \mathbb{Q}^\times$, let $s_2(n)$ denote the number of 1-bits in the binary expansion of n , and define the *binary carry count*

$$\operatorname{carries}(a, b) := s_2(a) + s_2(b) - s_2(a+b),$$

which equals the number of carries when adding a and b in binary (Kummer's theorem gives $v_2\binom{a+b}{a} = \operatorname{carries}(a, b)$).

2.1. The Zagier matrix.

Theorem 1 (Uniform Cofactor Valuation). *For Zagier's matrix M_K of weight $2K+1$, all last-column cofactors have the same 2-adic valuation:*

$$v_2(C(j, K-1)) = v_2(\det M_K) \quad \text{for all } j \in \{0, \dots, K-1\},$$

where $C(j, K-1)$ is the $(j, K-1)$ cofactor of M_K .

Corollary 2 (Odd Last Row). *All entries in the last row of $(M_K)^{-1}$ have 2-adic valuation zero:*

$$v_2((M_K)^{-1}[K-1, j]) = 0 \quad \text{for all } j \in \{0, \dots, K-1\}.$$

Remark 3. The last row of $(M_K)^{-1}$ gives the coefficients expressing $\zeta(2)^{K-1}\zeta(3)$ in the Hoffman basis. Corollary 2 implies that these coefficients are all rationals with odd numerator and odd denominator (in lowest terms).

Remark 4. The proof yields the closed formula

$$v_2(\det M_K) = \sum_{r=2}^K (v_2(r) + 2 - 2r) = 2K - s_2(K) - K^2.$$

2.2. The binomial core matrix.

Definition 5. For $N \geq 1$, the *binomial core matrix* B_N is the $N \times N$ upper unitriangular matrix with entries

$$B_N[a, i] = \binom{2i}{2a} \quad \text{for } 0 \leq a, i \leq N-1.$$

Note that $B_N[a, i] = 0$ for $a > i$ and $B_N[a, a] = 1$, so B_N is indeed upper unitriangular.

Theorem 6 (Inverse via Euler–Secant Numbers). *The inverse of the binomial core matrix is given by*

$$B_N^{-1}[a, i] = (-1)^{i-a} \binom{2i}{2a} E_{2(i-a)},$$

where E_{2n} denotes the n -th (unsigned) Euler–secant number, defined by

$$\operatorname{sech}(x) = \frac{1}{\cosh(x)} = \sum_{n=0}^{\infty} (-1)^n E_{2n} \frac{x^{2n}}{(2n)!}.$$

The first several values are $E_0 = 1$, $E_2 = 1$, $E_4 = 5$, $E_6 = 61$, $E_8 = 1385$, $E_{10} = 50521$.

Corollary 7 (Carries Formula). *For all $0 \leq a \leq i \leq N-1$:*

$$v_2(B_N^{-1}[a, i]) = \operatorname{carries}(a, i-a) = s_2(a) + s_2(i-a) - s_2(i).$$

Corollary 8. *The Euler–secant numbers satisfy $E_{2n} \equiv 1 \pmod{4}$ for all $n \geq 0$. In particular, $v_2(E_{2n}) = 0$.*

Corollary 9 (Structural Properties of B_N^{-1}).

- (a) **Last row.** $B_N^{-1}[0, i] = (-1)^i E_{2i}$, which is always odd.
- (b) **Diagonal.** $B_N^{-1}[a, a] = 1$ for all a .
- (c) **Column stability.** The entries in column i of B_N^{-1} are independent of N for $N > i$.
- (d) **Maximum valuation.** $\max_{0 \leq a \leq i} v_2(B_N^{-1}[a, i]) = \lfloor \log_2 i \rfloor$.

3. NUMERICAL VERIFICATION

We have verified Theorem 1 and Corollary 2 for all $K \leq 10$.

K	$v_2(\det M_K)$	Cofactor v_2	Last row numerators (all odd)
2	-1	$[-1, -1]$	11, 9
3	-5	all -5	523, 597, 399
4	-9	all -9	23003, 30657, 28023, 16957
5	-17	all -17	15331307, 22114173, ...
6	-26	all -26	1706973557, 28435623213, ...
7	-38	all -38	3724076580251, ...
8	-49	all -49	66117499294929143, ...
9	-65	all -65	18191208729544328424257, ...
10	-82	all -82	152212394986525434594354065, ...

TABLE 1. Verification of uniform cofactor valuation for weights 5–21.

N	v_2 matrix of B_N^{-1}	Last row	Carries match
4	$\begin{pmatrix} 0 & 0 & 0 & 0 \\ \cdot & 0 & 1 & 0 \\ \cdot & \cdot & 0 & 0 \\ \cdot & \cdot & \cdot & 0 \end{pmatrix}$	1, -1, 5, -61	✓
5	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ \cdot & 0 & 1 & 0 & 2 \\ \cdot & \cdot & 0 & 0 & 1 \\ \cdot & \cdot & \cdot & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot & 0 \end{pmatrix}$	1, -1, 5, -61, 1385	✓

TABLE 2. The 2-adic valuation matrix of B_N^{-1} matches carries($a, i - a$) exactly, verified for all $N \leq 10$.

4. PROOF OF THEOREM 1

We recall Zagier's formula [7, Theorem 1]:

$$(1) \quad M_K[a, r] = 2 \binom{2r}{2a+2} - \frac{2(2^{2r} - 1)}{2^{2r}} \binom{2r}{2b+1},$$

where $b = K - 1 - a$ and $r \in \{1, \dots, K\}$. We write $M_K[a, r] = T_1(a, r) - T_2(a, r)$ with $T_1 = 2 \binom{2r}{2a+2}$ and $T_2 = \frac{2(2^{2r}-1)}{2^{2r}} \binom{2r}{2b+1}$.

Let M' denote the $K \times (K - 1)$ submatrix consisting of columns $r = 2, \dots, K$, and let column $j \in \{0, \dots, K - 2\}$ of M' correspond to $r = K - j$.

Lemma 10 (Sparse Last Column). *The last column of M_K (corresponding to $r = 1$) is $[-2, 0, 0, \dots, 0, 3]^T$.*

Proof. Set $r = 1$ in (1). Then $\binom{2}{2a+2} = 0$ for $a \geq 1$ and $\binom{2}{2b+1} = 0$ for $b \geq 1$. The only nonzero entries are $a = 0$ (giving -2) and $a = K - 1$ (giving 3). \square

Lemma 11 (Column Minimum). *For each column $j \in \{0, \dots, K - 2\}$ of M' , with $r = K - j$:*

$$\min_{0 \leq a \leq K-1} v_2(M'[a, j]) = v_2(r) + 2 - 2r.$$

Moreover, this minimum is achieved by both $a = j$ (diagonal) and $a = K - 1$ (last row).

Proof. Step 1: Last row achieves the minimum. When $a = K - 1$, we have $b = 0$, so T_2 involves $\binom{2r}{1} = 2r$. Since $v_2(2^{2r} - 1) = 0$, we get $v_2(T_2(K - 1, r)) = v_2(r) + 2 - 2r$. For $j > 0$, $\binom{2r}{2K} = 0$ (since $2K > 2r$), so $T_1 = 0$; for $j = 0$, $v_2(T_1) = 1$. In both cases $v_2(T_1) > v_2(T_2)$ for $r \geq 2$, so $v_2(M'[K - 1, j]) = v_2(r) + 2 - 2r$.

Step 2: Diagonal achieves the same value. When $a = j = K - r$, we have $b = r - 1$, so T_2 involves $\binom{2r}{2r-1} = 2r$. This gives the same valuation as Step 1.

Step 3: All other rows have $v_2 \geq v_2(r) + 2 - 2r$. For the T_2 term, applying Kummer's theorem gives

$$v_2\left(\binom{2r}{2b+1}\right) = 1 + v_2\left(\binom{r}{b}\right) + v_2(r - b).$$

By the identity $\binom{r}{b}(r - b) = r\binom{r-1}{b}$, we have $v_2\left(\binom{r}{b}\right) + v_2(r - b) \geq v_2(r)$. Therefore $v_2(T_2(a, r)) \geq v_2(r) + 2 - 2r$ for all a . Since $v_2(T_1) \geq 1 > v_2(r) + 2 - 2r$ for $r \geq 2$, the claim follows. \square

Proof of Theorem 1. For each $\ell \in \{0, \dots, K - 1\}$, let M'_ℓ denote the $(K - 1) \times (K - 1)$ minor obtained by removing row ℓ from M' .

Case $\ell = K - 1$. The minor M'_{K-1} uses rows $0, \dots, K - 2$. By Lemma 11, the diagonal permutation $\sigma(j) = j$ is the unique permutation achieving the minimum v_2 sum $\sum_j (v_2(K - j) + 2 - 2(K - j))$. Therefore $v_2(\det M'_{K-1})$ equals this sum.

Case $\ell < K - 1$. The minor M'_ℓ includes row $K - 1$. Since both the diagonal entry and row $K - 1$ achieve each column's minimum (Lemma 11), we construct a permutation achieving the minimum v_2 sum: assign $\sigma(j) = j$ for $j \neq \ell$ and $\sigma(\ell) = K - 1$. Computational verification for all $K \leq 10$ confirms that no 2-adic cancellation occurs among the minimum- v_2 terms, yielding $v_2(\det M'_\ell) = \sum_j (v_2(K - j) + 2 - 2(K - j))$.

Synthesis. By Lemma 10, expanding $\det M_K$ along the last column:

$$\det M_K = -2 \cdot C(0, K-1) + 3 \cdot C(K-1, K-1).$$

Both cofactors satisfy $v_2(C(\ell, K-1)) = S$ where $S = \sum_j (v_2(K-j) + 2 - 2(K-j))$. Since $v_2(-2 \cdot C(0, K-1)) = 1 + S > S = v_2(3 \cdot C(K-1, K-1))$, we get $v_2(\det M_K) = S$, and hence $v_2(C(\ell, K-1)) = v_2(\det M_K)$ for all ℓ . \square

5. THE BINOMIAL CORE MATRIX

In this section we prove Theorem 6 and its corollaries. The matrix $B_N[a, i] = \binom{2i}{2a}$ arises naturally in the study of Zagier's matrix: the 2-adic dominant term in each column of M_K involves binomial coefficients $\binom{2r}{2b+1}$, whose valuations are controlled by carries in the same way as $\binom{2i}{2a}$ (see Lemma 11).

The key idea is an exponential generating function (EGF) argument that reduces the matrix inversion to the identity $\cosh(x) \cdot \operatorname{sech}(x) = 1$.

Proof of Theorem 6. The matrix equation $B_N^{-1} \cdot B_N = I$ says that for each a ,

$$(2) \quad \sum_{k=a}^i B_N^{-1}[a, k] \binom{2i}{2k} = \delta_{a,i} \quad \text{for all } i \geq a.$$

Define the exponential generating function for row a :

$$W_a(x) := \sum_{k \geq a} B_N^{-1}[a, k] \frac{x^{2k}}{(2k)!}.$$

Then the left side of (2) is the coefficient of $\frac{x^{2i}}{(2i)!}$ in the Cauchy product $W_a(x) \cdot \cosh(x)$, since

$$[x^{2i}/(2i)!](W_a(x) \cdot \cosh(x)) = \sum_k B_N^{-1}[a, k] \binom{2i}{2k}.$$

The identity (2) therefore becomes $W_a(x) \cdot \cosh(x) = \frac{x^{2a}}{(2a)!}$, giving

$$(3) \quad W_a(x) = \frac{x^{2a}}{(2a)!} \operatorname{sech}(x).$$

Since $\operatorname{sech}(x) = \sum_{n \geq 0} (-1)^n E_{2n} \frac{x^{2n}}{(2n)!}$, extracting the coefficient of x^{2i} from (3):

$$B_N^{-1}[a, i] = (2i)! \cdot [x^{2i}] \frac{x^{2a}}{(2a)!} \operatorname{sech}(x) = \frac{(2i)!}{(2a)!} \cdot \frac{(-1)^{i-a} E_{2(i-a)}}{(2(i-a))!} = (-1)^{i-a} \binom{2i}{2a} E_{2(i-a)}.$$

\square

Proof of Corollary 8. The recurrence from $\cosh(x) \cdot \operatorname{sech}(x) = 1$ gives, for $n \geq 1$:

$$E_{2n} = \sum_{k=1}^n (-1)^{k+1} \binom{2n}{2k} E_{2(n-k)}.$$

We prove $E_{2n} \equiv 1 \pmod{4}$ by induction. The base case $E_0 = 1$ is clear. For $n \geq 1$, assuming $E_{2(n-k)} \equiv 1 \pmod{4}$ for all $k \geq 1$:

$$E_{2n} \equiv \sum_{k=1}^n (-1)^{k+1} \binom{2n}{2k} = 1 - \sum_{k=0}^n (-1)^k \binom{2n}{2k} = 1 - \operatorname{Re}((1+i)^{2n}) = 1 - 2^n \cos(n\pi/2) \pmod{4}.$$

For $n \geq 2$, $2^n \equiv 0 \pmod{4}$, giving $E_{2n} \equiv 1 \pmod{4}$. For $n = 1$: $E_2 = 1 \equiv 1 \pmod{4}$. \square

Proof of Corollary 7. By Theorem 6, $B_N^{-1}[a, i] = (-1)^{i-a} \binom{2i}{2a} E_{2(i-a)}$. Since $v_2(E_{2(i-a)}) = 0$ by Corollary 8, we have

$$v_2(B_N^{-1}[a, i]) = v_2\left(\binom{2i}{2a}\right).$$

By Kummer's theorem, $v_2\left(\binom{2i}{2a}\right) = \text{carries}(2a, 2i - 2a)$. Since multiplying both arguments by 2 shifts binary representations one bit left without introducing new carries, $\text{carries}(2a, 2i - 2a) = \text{carries}(a, i - a)$. \square

Proof of Corollary 9. Part (a): setting $a = 0$ gives $B_N^{-1}[0, i] = (-1)^i E_{2i}$, which is odd since $v_2(E_{2i}) = 0$.

Part (b): setting $a = i$ gives $B_N^{-1}[i, i] = \binom{2i}{2i} E_0 = 1$.

Part (c): by Theorem 6, $B_N^{-1}[a, i]$ depends only on a and i , not on N .

Part (d): by Corollary 7, $\max_{0 \leq a \leq i} \text{carries}(a, i - a) = \lfloor \log_2 i \rfloor$ (e.g., achieved when a and $i - a$ have maximal carry count in binary addition). \square

6. CONNECTION BETWEEN THE TWO RESULTS

The proof of Theorem 1 and the proof of Theorem 6 both depend on the same primitive: the interaction between 2-adic valuations and binary carry counting in binomial coefficients.

Specifically, in Lemma 11, the key estimate

$$v_2\left(\binom{2r}{2b+1}\right) = 1 + v_2\left(\binom{r}{b}\right) + v_2(r - b)$$

reduces to $v_2\left(\binom{r}{b}\right) = \text{carries}(b, r - b)$ by Kummer's theorem. The identity $\binom{r}{b}(r - b) = r\binom{r-1}{b}$ then shows that the minimum is achieved when $\text{carries}(b, r - b)$ is minimized, i.e., when $b = 0$ or $b = r - 1$.

In the binomial core matrix, the same mechanism operates transparently: the inverse $B_N^{-1}[a, i] = (-1)^{i-a} \binom{2i}{2a} E_{2(i-a)}$ factors into a binomial

coefficient (whose v_2 is a carry count) and an Euler number (which is a 2-adic unit). The carries formula $v_2(B_N^{-1}[a, i]) = \text{carries}(a, i - a)$ then gives the complete 2-adic structure.

This suggests that the 2-adic structure of $(M_K)^{-1}$ should be governed by a “carries-like” formula involving the more complex combinatorics of Zagier’s two-term expression (1).

7. DISCUSSION AND OPEN QUESTIONS

Question 12 (No-cancellation property). In Case 2 of the proof of Theorem 1, the absence of 2-adic cancellation is verified computationally for $K \leq 10$. Can this be proved in full generality? One approach would be to establish a factorization of M'_ℓ that makes the carry structure manifest, analogous to the EGF factorization used for B_N .

Question 13 (Full 2-adic structure of $(M_K)^{-1}$). What is the 2-adic valuation of *all* entries of $(M_K)^{-1}$, not just the last row? Computation suggests a pattern depending on both the row and column index, but we have not found a closed formula analogous to the carries formula for B_N^{-1} .

Question 14 (Odd primes). Does similar structure exist for odd primes p ? By Kummer’s theorem, $v_p\binom{m+n}{m} = \text{carries}_p(m, n)$ counts p -adic carries. The p -adic valuations of Zagier’s matrices for odd primes may reveal additional arithmetic structure.

Question 15 (Motivic interpretation). The 2-adic properties of Zagier’s matrices were essential in Brown’s proof [1] of the Hoffman conjecture. Can the carries formula for B_N^{-1} be given a motivic interpretation, perhaps in terms of the action of the motivic Galois group on the relevant component of the category of mixed Tate motives over \mathbb{Z} ?

Question 16 (Connection to q -zeta functions). The matrix B_N also appears in the study of Habiro’s q -series and completed q -zeta functions, where a triangular inversion with analogous 2-adic structure is required. In that setting, the generating function $\psi(\varepsilon) = \varepsilon/(e^\varepsilon - 1) = \sum B_k \varepsilon^k/k!$ involves Bernoulli numbers in place of the Euler numbers that appear here. Since $v_2(B_{2k}) = -1$ by the von Staudt–Clausen theorem while $v_2(E_{2k}) = 0$, the Habiro setting has a richer 2-adic structure. Can the methods of this paper be extended to give closed forms or carries formulas in the Habiro setting?

ACKNOWLEDGMENTS

The author thanks Don Zagier for providing the explicit formulas that made these computations possible, and Bruce Berndt for helpful support. Computations were performed using Python with exact rational arithmetic.

REFERENCES

- [1] F. Brown, *Mixed Tate motives over \mathbb{Z}* , Ann. of Math. **175** (2012), 949–976.
- [2] M. Hoffman, *The algebra of multiple harmonic series*, J. Algebra **194** (1997), 477–495.
- [3] K. Ihara, M. Kaneko, and D. Zagier, *Derivation and double shuffle relations for multiple zeta values*, Compos. Math. **142** (2006), 307–338.
- [4] E. E. Kummer, *Über die Ergänzungssätze zu den allgemeinen Reciprocitätsgesetzen*, J. Reine Angew. Math. **44** (1852), 93–146.
- [5] Z. Li, *Another proof of Zagier's evaluation formula of the multiple zeta values $\zeta(2, \dots, 2, 3, 2, \dots, 2)$* , Math. Res. Lett. **20** (2013), 947–950.
- [6] F. W. J. Olver et al., eds., *NIST Digital Library of Mathematical Functions*, Release 1.2.3, <https://dlmf.nist.gov/>, 2024.
- [7] D. Zagier, *Evaluation of the multiple zeta values $\zeta(2, \dots, 2, 3, 2, \dots, 2)$* , Ann. of Math. **175** (2012), 977–1000.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN
Email address: yipinw2@illinois.edu