


ECO326 Advanced Microeconomic Theory

A Course in Game Theory

Tianyu Du

December 21, 2018

This work is licensed under a Creative Commons
“Attribution-NonCommercial 4.0 International”
license.



Github Page https://github.com/TianyuDu/Spikey_UofT_Notes

Note Page TianyuDu.com/notes

Readme this note is based on the course content of *ECO326 Advanced Microeconomics - Game Theory*, this note contains all materials covered during lectures and mentioned in the course syllabus. However, notations, statements of theorems and proofs are following the book *A Course in Game Theory* by Osborne and Rubinstein, so they might be, to some extent, more mathematical than the required text for ECO326, *An Introduction to Game Theory*.

Contents

1	Lecture 1. Games and Dominant Strategies	2
2	Lecture 2. Iterated Elimination and Rationalizability	2
2.1	Iterated Elimination of Strictly Dominated Strategies(Actions)	2
2.2	Rationalizability	3
3	Lecture 3. Nash Equilibrium	3
4	Lecture 6. Extensive Form Games and Subgame Perfection	5
4.1	Extensive Form Game	5
4.2	Subgame Perfection	6

1 Lecture 1. Games and Dominant Strategies

Assumption 1.1 (pg.4). Assume that each decision-maker is *rational* in the sense that he is aware of his alternatives, forms expectation about any unknowns, has clear preferences, and chooses his action deliberately after some process of optimization.

Definition 1.1 (pg.4). A model of **rational choice** consists

- A set A of *actions*.
- A set C of *consequences*.
- A *consequence function* $g : A \rightarrow C$.
- A *preference relation* \succsim on C .

Definition 1.2 (pg.7). A **preference relation** is a complete reflexive and transitive binary relation.

Definition 1.3 (11.1). A **strategic game** consists of

- a finite set of **players** N .
- for each player $i \in N$, an **actions** $A_i \neq \emptyset$.
- for each player $i \in N$, a **preference relation** \succsim_i defined on $A \equiv \times_{i \in N} A_i$.

and can be written as a triple $\langle N, (A_i), (\succsim_i) \rangle$.

Definition 1.4 (pg.11). A strategic game $\langle N, (A_i), (\succsim_i) \rangle$ is **finite** if

$$|A_i| < \aleph_0 \quad \forall i \in N$$

2 Lecture 2. Iterated Elimination and Rationalizability

2.1 Iterated Elimination of Strictly Dominated Strategies(Actions)

Definition 2.1 (60.2). The set $X \subseteq A$ of outcomes of a finite strategic game $\langle N, (A_i), (u_i) \rangle$ **survives iterated elimination of strictly dominated actions** if $X = \times_{j \in N} X_j$ and there is a collection $((X_j^t)_{j \in N})_{t=0}^T$ of sets that satisfies the following conditions for each $j \in N$.

- $X_j^0 = A_j$ and $X_j^T = X_j$.
- $X_j^{t+1} \subseteq X_j^t$ for each $t = 0, \dots, T-1$.
- For each $t = 0, \dots, T-1$ every action of player j in $X_j^t \setminus X_j^{t+1}$ is strictly dominated in the game $\langle N, (X_i^t), (u_i^t) \rangle$, where u_i^t for each $i \in N$ is the function u_i restricted to $\times_{j \in N} X_j^t$.
- No action in X_t^T is strictly dominated in game $\langle N, (X_i^T), (u_i^T) \rangle$.

Proposition 2.1 (61.2). If $X = \times_{j \in N} X_j$ survives iterated elimination of strictly dominated actions in a finite strategic game $\langle N, (A_i), (u_i) \rangle$ then X_j is the set of player j 's rationalizable actions for each $j \in N$.

2.2 Rationalizability

Definition 2.2 (pg. 54). A **belief** of player i (about the actions of the other players) is a probability measure, μ_i , on $A_{-i} = \times_{j \in N \setminus \{i\}} A_j$.

Definition 2.3 (59.1). An action of player i in a strategic game is a **never best response** if it is not a best response to any belief of player i .

Definition 2.4 (59.2). The action $a_i \in A_i$ of player i in the strategic game $\langle N, (A_i), (u_i) \rangle$ is **strictly dominated** if there is a mixed strategy α_i of player i such that

$$U_i(a_{-i}, \alpha_i) > u_i(a_{-i}, a_i)$$

for all $a_{-i} \in A_{-i}$, where $U_i(a_{-i}, \alpha_i)$ is the payoff of player i if he uses the mixed strategy α_i and the other players' vector of actions is a_{-i} .

3 Lecture 3. Nash Equilibrium

Definition 3.1 (14.1). A **Nash equilibrium of a strategic game** $\langle N, (A_i), (\succsim_i) \rangle$ is a profile $a^* \in A$ of actions with property that for every player $i \in N$

$$(a_i^*, a_{-i}^*) \succsim_i (a_i, a_{-i}^*) \quad \forall a_i \in A_i$$

Definition 3.2 (pg.15). The **best-response function** for a player i is defined as

$$B_i(a_{-i}) = \{a_i \in A_i : (a_i, a_{-i}) \succsim_i (a'_i, a_{-i}) \quad \forall a'_i \in A_i\}$$

Remark 3.1. The best-response of a_{-i} can be written as

$$B_i(a_{-i}) = \bigcap_{a'_i \in A_i} \{a_i \in A_i : (a_i, a_{-i}) \succsim_i (a'_i, a_{-i})\}$$

where each of them is the upper contour set of a'_i .

Thus, if \succsim_i is quasi-concave, then $B_i(a_{-i})$ is an intersection of convex sets and therefore itself convex.

Remark 3.2 (pg.15). So a Nash equilibrium is a profile $a^* \in A$ such that

$$a_i^* \in B_i(a_{-i}^*) \quad \forall i \in N$$

Lemma 3.1 (pg.19). A strategic game $\langle N, (A_i), (\succsim_i) \rangle$ has a Nash equilibrium if equivalent to the following statement:

Define set-valued function $B : A \rightarrow A$ by

$$B(a) = \times_{i \in N} B_i(a_{-i})$$

and there exists $a^* \in A$ such that $a^* \in B(a^*)$.

Lemma 3.2 (20.1 Kakutani's fixed point theorem). Let X be a compact convex subset of \mathbb{R}^n and let $f : X \rightarrow X$ be a set-valued function for which

- for all $x \in X$ the set $f(x)$ is non-empty and convex.
- the graph of f is closed. (i.e. for all sequences $\{x_n\}$ and $\{y_n\}$ such that $y_n \in f(x_n)$ for all n , $x_n \rightarrow x$ and $y_n \rightarrow y$ then $y \in f(x)$)

Then there exists $x^* \in X$ such that $x^* \in f(x^*)$.

Definition 3.3 (pg.20). A preference relation \succsim_i over A is quasi-concave on A_i if for every $a^* \in A$ the upper contour set over a_i^* , given other players' strategies

$$\{a_i \in A_i : (a_{-i}^*, a_i) \succsim_i a^*\}$$

is convex.

Proposition 3.1 (20.3). The strategic game $\langle N, (A_i), (\succsim_i) \rangle$ has a Nash equilibrium if for all $i \in N$,

- the set A_i of actions of player i is a nonempty compact convex subset of a Euclidian space

and the preference relation \succsim_i is

- continuous
- quasi-concave on A_i .

Proof. Let $B : A \rightarrow A$ be a correspondence defined as

$$B(a) := \times_{i \in N} B_i(a_{-i})$$

Note that for each $a \in A$ and for each $i \in N$,

$B_i(a_{-i}) \neq \emptyset$ since preference \succsim_i is continuous and A_i is compact (EVT).

Also $B_i(a_{-i})$ is convex since it's basically an intersection of upper contour sets and each of those upper contour is convex since \succsim_i is quasi-concave.

So the Cartesian product of the finite collection of B_i is non-empty and convex.

Also the graph B is closed since \succsim_i is continuous.

So there exists $a^* \in A$ such that $a^* \in B(a^*)$.

So Nash equilibrium presents. ■

4 Lecture 6. Extensive Form Games and Subgame Perfection

4.1 Extensive Form Game

Definition 4.1 (89.1). An **extensive game with perfect information** has the following components.

- A set N of **players**.
- A set H of sequences (finite or infinite) of **histories** with properties:
 - $\emptyset \in H$.
 - For all $L < K$, $(a^k)_{k=1,2,\dots,K} \in H \implies (a^k)_{k=1,2,\dots,L} \in H$.
 - For infinite sequence $(a^k)_{k=1}^\infty$,
 $(a^k)_{k=1,2,\dots,L} \in H, \forall L \in \mathbb{Z}_{++} \implies (a^k)_{k=1}^\infty \in H$.

And each component of history $h \in H$ is an **action** taken by a player.

- A function $P : H \setminus Z \rightarrow N$, where for $h \in H$, $P(h) \in N$ is defined by the player who takes an action after the history h .
- For each player $i \in N$ a **preference relation** \succsim_i defined on Z .

Notation 4.1 (pg.90). An extensive game with perfect information can be represented by a 4-tuple, $\langle N, H, P, (\succsim_i) \rangle$. *Sometimes it is convenient to specify the structure of an extensive game without specifying the players' preference, as $\langle N, H, P \rangle$.*

Definition 4.2 (pg.90). A history $(a^k)_{k=1,2,\dots,K} \in H$ is **terminal** if

1. it is infinite,
2. or (i.e. it cannot be extended to another valid history sequence)

$$\forall a^{K+1}, (a^k)_{k=1,2,\dots,K+1} \notin H$$

The set of terminal histories is denoted by Z .

Notation 4.2 (pg.90, the action set). After any nonterminal history, $h \in H \setminus Z$, the player $P(h)$ chooses an action from set

$$A(h) = \{a : (h, a) \in H\}$$

Remark 4.1. Note that all player function, action set and player preference relation are defined on H . Thus, unlike a normal form game, which was *player oriented*, we'd better consider an extensive form game as *history oriented*.

Definition 4.3 (pg.90). We refer to the empty set, which is required to be an element of H , as the **initial history**.

Definition 4.4 (92.1). A **strategy of player** $i \in N$, s_i , in an extensive game with perfect information $\langle N, H, P, (\succsim_i) \rangle$ is a function that assigns an action in $A(h)$ to each nonterminal history $h \in H \setminus Z$ for which $P(h) = i$.

Remark 4.2 (pg.92). A strategy specifies the action chosen by a player for *every* history after which it is his turn to move, *even for histories that is, if the strategy is followed, are never reached*.

Definition 4.5 (pg.93). For each strategy profile $s = (s_i)_{i \in N}$ in the extensive game $\langle N, H, P, (\succsim_i) \rangle$, the **outcome** of s , $O(s)$, is defined as the terminal history that results when each player $i \in N$ follows the precepts of s_i . That is, $O(s)$ is the (possibly infinite) history

$$(a^1, \dots, a^K) \in Z$$

such that

$$\forall k \in \{0, 1, \dots, K-1\}, s_{P(a^1, \dots, a^k)}(a^1, \dots, a^k) = a^{k+1}$$

4.2 Subgame Perfection