

Notes on MAT137 Video Playlist 3

Tianyu Du

Saturday 14th April, 2018

Info.

Created: September, 2017

Last modified: Saturday 14th April, 2018

Revision: 00:40 Saturday 14th April, 2018

This work is licensed under a Creative Commons “Attribution-NonCommercial-ShareAlike 3.0 Unported” license.



Contents

1	Video Playlist 1	4
2	Video Playlist 2	4
3	Video Playlist 3	4
3.1	Define Derivate As Slope	4
3.2	Calculate $f'(x)$ by definition	5
3.3	Rate of Change	5
3.4	The Product Rule (Formal Version)	5
3.5	Differentiable \implies Continuous	5
3.6	Proof of product rule for derivative.	6
3.7	Partial proof of differentiation rule	6
3.8	Higher Order Derivatives: Notations	7
3.9	Continuous But Not differentiable	7
3.10	Chain Rule	8
3.11	Derivatives of Trig Functions	8
3.12	Implicit Differentiation	9
3.13	Derivative of Exponential Functions	10
3.14	Properties of logarithms	10
3.15	The derivatives of logarithm functions	11
3.16	Derivative of other exponentials	11
3.17	The power rule, complete proof	12
3.18	Logarithmic Differentiation	12

4	Video Playlist 4	12
4.1	Functions	12
4.2	Inverse Functions	13
4.3	Surjective Functions	13
4.4	Injective function	14
4.5	Some theorems	14
4.6	ArcSin	15
4.7	Derivative of ArcSin	15
4.8	Other inverse trig functions	16
4.8.1	$y = \cos(x)$	16
4.8.2	$y = \tan(x)$	16
5	Video Playlist 5	16
5.1	Usage of MVT	16
5.2	Local Extreme Theorem	17
5.3	Find Extremum	19
5.4	Rolle's Theorem	19
5.5	Application of Rolle's Theorem	19
5.6	(Lagrange)Mean Value Theorem	20
5.7	Proof. of MVT	21
5.8	Zero-derivative implies constant	21
5.9	Monotonicity of functions	21
6	Video Playlist 6	22
7	Video Playlist 7	22
7.1	Integral	22
7.2	Sigma Notation	22
7.3	Supremum and Infimum	22
7.4	Supremum and Infimum of a function	23
7.5	Definition of Integral (i)	23
7.6	Definition of Integral (ii): Properties of $U_P(f)$ and $L_P(f)$	24
7.7	Definition of Integral (iii): Upper Integral and Lower Integral . .	24
7.8	An example of integrable function	25
7.9	An example of non-integrable function	25
7.10	Integrals as limits	25
7.11	Riemann Sums	25
7.12	Properties of the integral	26
8	Video Playlist 8	27
8.1	Anti-derivatives	27
8.2	Functions Defined as Integrals	27
8.3	The Fundamental Theorem of Calculus: Part 1	27
8.4	A Proof of Part 1 of the FTC	28
8.5	The Fundamental Theorem of Calculus: Part 2	28
8.6	A Proof of Part 2 of the FTC	29

8.7	Summary: Definite and indefinite integrals, notation, definitions and theorems.	29
8.7.1	Definite Integral.	29
8.7.2	Indefinite Integral	30
8.7.3	Function Defined by an Integral.	30
9	Video Playlist 9	30
9.1	Integration By Substitution: derivation of the formula	30
9.2	Example 2	30
9.3	Example 3	30
9.4	Example 4	30
9.5	Integration by parts	30
9.6	Examples	31
9.7	Integration of products of trigonometric functions	31
10	Video Playlist 10	32
11	Video Playlist 11	32
11.1	What Is a Sequence	32
11.1.1	Conventions	32
11.1.2	Describe sequences	32
11.2	The Limit of a Sequence	33
11.3	Properties of Limits of Sequences	33
11.3.1	Sequence from a function	33
11.3.2	Composite of sequence and function	33
11.4	Monotonic and Bounded Sequences	34
11.4.1	Monotonic Sequences	34
11.4.2	Bounded Sequences	34
11.5	Proof: Every convergent sequence is bounded	35
11.6	The monotone convergence theorem of sequences	35
11.7	the Big theorem of sequences	36
12	Video Playlist 12	36
12.1	Improper Integral	36
12.1.1	Improper integral "type 1" (Unbounded domain)	36
12.2	The most important family if the improper integrals	37
12.3	Example	37
12.4	The most important family of improper integral	37
12.5	Example	37
12.6	Doubly improper integrals	37
12.7	Basic Comparison Test	38
12.8	Examples	38
12.9	Limit Comparison Test	38
12.10	Proof of LCT	38

13 Video Playlist 13	38
13.1 Infinite Sums	38
13.2 Definition of series	38
13.3 Example	38
13.4 Divergent Series Examples	39
13.5 Geometric Series	39
13.6 Linearity of series	39
13.7 the Tail of a series	40
13.8 A necessary condition for convergence of series	41
13.9 Positive series	41
13.10 The Integral Test	42
13.11 Examples	42
13.12 Comparison Tests for Series	42
14 Video Playlist 14	43
14.1 Power Series: Example	43
14.2 Main Theorem	43
14.3 Taylor Polynomial Definition 1	43
14.4 Taylor Polynomial Definition 2	44
14.5 Taylor Polynomial Definition 3	44

1 Video Playlist 1

2 Video Playlist 2

3 Video Playlist 3

3.1 Define Derivate As Slope

Definition Let $a \in \mathbb{R}$, and $f(x)$ is defined on $(a-\delta, a+\delta)$, then the **derivative** of $f(x)$ at a is,

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

Definition If function is **differentiable** at point $x = a$, if and only if, there exists,

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

Interpretation $f'(a)$ is the slope of tangent line at $x = a$.

3.2 Calculate $f'(x)$ by definition

Example $f(x) = 4x - x^2$, find $f'(1)$:

$$\begin{aligned} f'(1) &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = \lim_{h \rightarrow 0} \frac{4(h+1) - (h+1)^2 - 3}{h} \\ &= \lim_{h \rightarrow 0} \frac{4h + 4 - 3 - h^2 - 2h - 1}{h} = \lim_{h \rightarrow 0} \frac{-h^2 + 2h}{h} \\ &= \lim_{h \rightarrow 0} -h + 2 = 2 \end{aligned}$$

3.3 Rate of Change

Definition Define derivative as rate of change. Let $x = f(t)$, then $f'(x)$ can be represented as,

$$\lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} = f'(t) = \frac{dx}{dt}$$

3.4 The Product Rule (Formal Version)

Let $a \in \mathbb{R}$, f and g are functions defined at $(a - \delta, a + \delta)$, let $h(x) = f(x)g(x)$. Then, if $f(x), g(x)$ are differentiable at a , we have,

$$h'(a) = f'(a)g(a) + f(a)g'(a)$$

3.5 Differentiable \implies Continuous

Recall $f(x)$ is **differentiable** at a :

$$\exists \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \tag{1}$$

Recall $f(x)$ is **continuous** at a :

$$\lim_{x \rightarrow a} f(x) = f(a) \tag{2}$$

Proof.

Since $f(x)$ is differentiable at a

$$(1) \iff \exists \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

$$\text{And } \lim_{x \rightarrow a} (x - a) = 0$$

$$\implies \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \lim_{x \rightarrow a} (x - a) = 0$$

$$\implies \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} (x - a) = 0$$

$$\implies \lim_{x \rightarrow a} (f(x) - f(a)) = 0$$

$$\implies \lim_{x \rightarrow a} f(x) = f(a)$$

■

3.6 Proof of product rule for derivative.

$(fg)' = f'g + fg'$, see above for a formal definition.

Let $h = fg$

$$h'(a) = \lim_{x \rightarrow a} \frac{h(x) - h(a)}{x - a}$$

$$= \lim_{x \rightarrow a} \frac{f(x)g(x) - f(a)g(a)}{x - a}$$

$$= \lim_{x \rightarrow a} \frac{f(x)g(x) + f(a)g(x) - f(a)g(x) - f(a)g(a)}{x - a}$$

$$= \lim_{x \rightarrow a} \frac{g(x)(f(x) - f(a)) + f(a)(g(x) - g(a))}{x - a}$$

$$= \lim_{x \rightarrow a} g(x) \frac{f(x) - f(a)}{x - a} + \lim_{x \rightarrow a} f(a) \frac{g(x) - g(a)}{x - a}$$

$$= g(a) \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} + f(a) \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a}$$

$$= g(a)f'(a) + f(a)g'(a)$$

■

3.7 Partial proof of differentiation rule

WTS $\frac{d}{dx} x^c = cx^{c-1}$, $\forall c \in \mathbb{R}$

Here we only prove statements is true $\forall c \in \mathbb{Z}^+$

Proof.

Base: $c = 1$

$$\begin{aligned} f(x) &= x \\ f'(x) &= \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \\ &= \lim_{x \rightarrow a} 1 = 1 \end{aligned}$$

Induction step

$$\text{Assume } \frac{d}{dx}[x^k] = kx^{k-1}|_{x=a}$$

$$\text{For } f(x) = x^{k+1}$$

$$\begin{aligned} f'(x) &= \frac{d}{dx}[x * x^k] \\ &= x^k + xkx^{k-1} \\ &= (k+1)x^k \end{aligned}$$

■

3.8 Higher Order Derivatives: Notations

Original function: $f(x)$

- **Lagrange** notation: $f^{(n)}$
- **Leibnitz** notation: $\frac{d^n f}{dx^n}$

3.9 Continuous But Not differentiable

Definition Function $f(x)$ is **non-differentiable** at a .

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \text{ DNE}$$

Example 1 Corner/Kink $f(x) = |x|$ at 0.

$$\begin{aligned} \lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x} &= \lim_{x \rightarrow 0^-} \frac{|x|}{x} = -1 \\ \lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x} &= \lim_{x \rightarrow 0^+} \frac{|x|}{x} = 1 \\ \lim_{x \rightarrow 0^-} &\neq \lim_{x \rightarrow 0^+} \\ \implies \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x} &\text{ DNE} \end{aligned}$$

Example 2 Vertical Tangent Line $g(x) = x^{\frac{1}{3}}$ at 0,

$$g'(0) = \lim_{x \rightarrow 0} \frac{x^{\frac{1}{3}}}{x} = \lim_{x \rightarrow 0} \frac{1}{x^{\frac{2}{3}}} = \infty (\text{DNE})$$

Caution Difference between **vertical asymptote** and **vertical tangent line**

- Vertical asymptote: $f(a) = \infty$ ($f(a)$ is not defined)
- Vertical tangent line: $f(a)$ is defined, $f'(a)$ is undefined.

3.10 Chain Rule

Derivation

$$\begin{aligned} (g \circ f)'(a) &= \lim_{x \rightarrow a} \frac{g(f(x)) - g(f(a))}{x - a} \\ &= \lim_{x \rightarrow a} \frac{g(f(x)) - g(f(a))}{f(x) - f(a)} \frac{f(x) - f(a)}{x - a} \end{aligned}$$

Attention: we could only apply the operation above if $f(x) \neq f(a)$ during the process of $x \rightarrow a$.

This holds for majority of functions we operate in calculus.

$$\begin{aligned} &= \lim_{f(x) \rightarrow f(a)} \frac{g(f(x)) - g(f(a))}{x - a} f'(a) \\ &= g'(f(a)) \cdot f'(a) \end{aligned}$$

■

Formal Theorem of Chain Rule Let $a \in \mathbb{R}$, let f and g be functions. If f is differentiable at a and g is differentiable at $f(a)$, then, $(g \circ f)$ is differentiable at a ,

$$(g \circ f)'(a) = g'(f(a)) \cdot f'(a)$$

3.11 Derivatives of Trig Functions

Basic 6 results

1. $\frac{d}{dx} \sin(x) = \cos(x)$
2. $\frac{d}{dx} \cos(x) = -\sin(x)$
3. $\frac{d}{dx} \tan(x) = \sec^2(x)$
4. $\frac{d}{dx} \cot(x) = -\csc^2(x)$
5. $\frac{d}{dx} \sec(x) = \sec(x)\tan(x)$
6. $\frac{d}{dx} \csc(x) = -\csc(x)\cot(x)$

Proof. Prove (i) and (ii) and use (i), (ii) and quotient rule to derive (iii), (iv), (v) and (vi).

Proof. (i) WTS $f(x) = \sin(x)$, then $f'(x) = \cos(x)$

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin(x)\cos(h) + \cos(x)\sin(h) - \sin(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sin(x)(\cos(h) - 1) + \cos(x)\sin(h)}{h} \\
 &= \lim_{h \rightarrow 0} \cos(x) \frac{\sin(h)}{h} \\
 &= \cos(x)
 \end{aligned}
 \quad \blacksquare \quad (3)$$

Proof. (ii) WTS $f(x) = \cos(x)$, then $f'(x) = -\sin(x)$

$$\begin{aligned}
 f'(x) &= \lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cos(x)\cos(h) - \sin(h)\sin(x) - \cos(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(\cos(h) - 1)\cos(x) - \sin(h)\sin(x)}{h} \\
 &= \lim_{h \rightarrow 0} -\frac{\sin(h)}{h} \sin(x) \\
 &= -\sin(x)
 \end{aligned}
 \quad \blacksquare \quad (4)$$

Recall Compound angle formula:

1. $\sin(\alpha + \beta) = \sin(\alpha)\cos(\beta) + \sin(\beta)\cos(\alpha)$
2. $\sin(\alpha - \beta) = \sin(\alpha)\cos(\beta) - \sin(\beta)\cos(\alpha)$
3. $\cos(\alpha + \beta) = \cos(\alpha)\cos(\beta) - \sin(\alpha)\sin(\beta)$
4. $\cos(\alpha - \beta) = \cos(\alpha)\cos(\beta) + \sin(\alpha)\sin(\beta)$

3.12 Implicit Differentiation

Key Use chain rule.

3.13 Derivative of Exponential Functions

Let $f(x) = a^x$ ($a > 0$), find $f'(x)$, by definition,

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} \\ &= \lim_{h \rightarrow 0} \frac{a^x a^h - a^x}{h} \\ &= \lim_{h \rightarrow 0} \frac{(a^h - 1)a^x}{h} \end{aligned}$$

By property of limit, a^x is the only variable, so that a^x is a constant

$$= a^x \lim_{h \rightarrow 0} \frac{a^h - 1}{h}$$

(5)

Equivalently, $\frac{d}{dx}a^x = L_a a^x$

Definition e is the only positive number, such that,

$$\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1$$

So that, $\frac{d}{dx}e^x = e^x$

3.14 Properties of logarithms

Definition Let $a > 0, a \neq 1, x > 0, y \in \mathbb{R}$,

$$\log_a x = y \iff a^y = x$$

Properties

1. $\log_a 1 = 0$
2. $\log_a a = 1$
3. $\log_a x = \frac{\log_b x}{\log_b a}$
4. $\log_a xy = \log_a x + \log_a y$
5. $\log_a \frac{x}{y} = \log_a x - \log_a y$
6. $\log_a x^r = r \log_a x$

Proof. (i) let $a > 0, a \neq 1, \text{let } x, y > 0$, **WTS** $\log_a xy = \log_a x + \log_a y$

$$\text{Let } p = \log_a x \iff a^p = x$$

$$\text{Let } q = \log_a y \iff a^q = y$$

$$\text{We have } a^p a^q = xy$$

$$\iff a^{p+q} = xy$$

$$\iff \log_a xy = p + q = \log_a x + \log_a y$$

■

3.15 The derivatives of logarithm functions

For $\ln x \quad \frac{d}{dx} \ln x = \frac{1}{x}$

$$e^{\ln x} = x$$

$$\frac{d}{dx} e^{\ln x} = \frac{d}{dx} x$$

$$\frac{d}{d \ln x} e^{\ln x} \cdot \frac{d}{dx} \ln x = 1$$

$$x \frac{d \ln x}{dx} = 1$$

$$\frac{d}{dx} \ln x = \frac{1}{x}$$

■

3.16 Derivative of other exponentials

WTS $\frac{d}{dx} a^x = \ln a \cdot a^x$,

$$a^x = (e^{\ln a})^x = e^{x \ln a}$$

$$\frac{d}{dx} a^x = \frac{d}{dx} e^{x \ln a}$$

$$= \frac{d}{dx} e^{x \ln a} \cdot \frac{d}{dx} \ln a$$

$$= e^{x \ln a} \ln a$$

$$= \ln a \cdot a^x$$

■

3.17 The power rule, complete proof

WTS $x^c = cx^{c-1}$

$$\begin{aligned}
 x^c &= (e^{\ln x})^c = e^{c \ln x} \\
 \text{So that } \frac{d}{dx} x^c &= \frac{d}{dx} e^{c \ln x} \\
 &= \frac{de^{c \ln x}}{d \ln xc} \cdot \frac{\ln xc}{d \ln x} \cdot \frac{d \ln x}{dx} \\
 &= e^{c \ln x} \cdot c \cdot \frac{1}{x} \\
 &= c \cdot x^c \cdot \frac{1}{x} \\
 &= cx^{c-1}
 \end{aligned}$$

■

3.18 Logarithmic Differentiation

Example $f(x) = \cos(x)^{\sin(x)}(\star)$, find $f'(x)$

Step1. Take \ln on both sides of (\star)

$$\ln f(x) = \ln \cos(x)^{\sin(x)} = \sin(x) \ln \cos(x)$$

Step2. Take derivative.

$$\frac{f'(x)}{f(x)} = \cos(x) \ln \cos(x) - \sin^2(x) \frac{1}{\cos(x)}$$

Step3. Solve for $f'(x)$

$$f'(x) = \cos(x)^{\sin(x)} (\cos(x) \ln \cos(x) - \sin^2(x) \frac{1}{\cos(x)})$$

4 Video Playlist 4

4.1 Functions

In calculus We assume the domain is the largest subset of \mathbb{R} that makes sense. And assume the codomain is always \mathbb{R} .

Notations	Math	Computer Science
	Domain	Domain
	Codomain	Range
	Range	Image

4.2 Inverse Functions

Definition Let $f : A \rightarrow B$ be a function. Function $f^{-1} : B \rightarrow A$ is the **inverse function** if and only if

$$\forall x \in A, \forall y \in B, x = f^{-1}(y) \iff y = f(x)$$

Properties

- $\forall x \in A, f^{-1}(f(x)) = x$
- $\forall y \in B, f(f^{-1}(y)) = y$

Pre-condition Function f has inverse function f^{-1} if and only if f is **injective/one-to-one** function.

4.3 Surjective Functions

Why function don't have an inverse: Part 1.

Definition Function $f(x)$ is **surjective/onto** if $\text{codomain}(f(x)) = \text{range}(f(x))$.

Problem If $f(x)$ is not surjective, then some points in codomain has no corresponding point in domain, then f^{-1} is not a function.

Solution Shrink the codomain to range.

Example Let $f(x) = e^x$, $g(x) = \ln x$, then we have,

- $\text{Domain}(f(x)) = \mathbb{R}$
 $\text{Codomain}(f(x)) = \mathbb{R}$
 $\text{Range}(f(x)) = (0, \infty)$
- $\text{Domain}(g(x)) = (0, \infty)$
 $\text{Codomain}(g(x)) = \mathbb{R}$
 $\text{Range}(g(x)) = \mathbb{R}$

Definition Definition of inverse in calculus (*simplified, we don't consider codomain here.*)

Let $f(x)$ be a function, and $f^{-1}(x)$ be the **inverse** of it. Then,

- $\text{Domain}(f^{-1}(x)) = \text{Range}(f(x))$
- $\text{Range}(f^{-1}(x)) = \text{Domain}(f(x))$

also,

$$\forall x \in \text{Domain}(f(x)), \forall y \in \text{Range}(f(x)), x = f^{-1}(y) \iff y = f(x)$$

and,

$$\forall x \in \text{Domain}(f(x)), f^{-1}(f(x)) = x$$

$$\forall y \in \text{Range}(f(x)), f(f^{-1}(y)) = y$$

4.4 Injective function

Definition Let $f(x)$ be a function, with $\text{Domain}(f(x)) = A$, we say $f(x)$ is **injective/one-to-one** when,

$$\forall x_1, x_2 \in A, x_1 \neq x_2 \implies f(x_1) \neq f(x_2)$$

equivalently (contrapositive)

$$f(x_1) = f(x_2) \implies x_1 = x_2$$

Theorem Function f has an inverse if and only if f is **injective**.

Example $f(x) = x^2$ has no inverse, but we could take it's inverse by shrinking the domain.

- Take domain = $[0, \infty)$, $f^{-1}(x) = \sqrt{x}$
- Take domain = $(-\infty, 0]$, $f^{-1}(x) = -\sqrt{x}$

4.5 Some theorems

Let $f(x)$ be a function with domain I .

Theorem 1 Function f has an inverse function f^{-1} if and only if f is injective.

Theorem 2 For function f , if

1. f is **continuous** (*This means, f is continuous on its domain.*).
2. I is an **interval**.

then, $f^{-1}(x)$ is continuous.

Theorem 3 If

1. f is **differentiable**.
2. $\forall x \in I, f'(x) \neq 0$ (*This ensures the inverse function does not have a vertical tangent line, which causes non-differentiability*).

then, $f^{-1}(x)$ is differentiable.

Theorem 4 $\forall x \in I$ with $y = f(x)$, we have

$$(f^{-1})'(y) = \frac{1}{f'(x)}$$

Proof.

$$\begin{aligned} f(f^{-1}(y)) &= y \\ \frac{d}{dy} f(f^{-1}(y)) &= \frac{d}{dy} y \\ \frac{d}{dy} f(f^{-1}(y)) &= 1 \\ f'(f^{-1}(y)) \cdot (f^{-1})'(y) &= 1 \\ f'(x) \cdot (f^{-1})'(y) &= 1 \\ (f^{-1})'(y) &= \frac{1}{f'(x)} \end{aligned}$$

■

4.6 ArcSin

Note ArcSin is **NOT** the inverse of Sin . $y = \sin(x)$ has *domain* $= \mathbb{R}$ and *range* $= [-1, 1]$, so that, it is **not injective**.

Definition ArcSin is the inverse function to the **restriction** of \sin to $[-\frac{\pi}{2}, \frac{\pi}{2}]$. So that, $\text{Domain}(\text{ArcSin}) = \text{Range}(\text{Sin}) = [-1, 1]$, and, $\text{Range}(\text{ArcSin}) = \text{Domain}(\text{Sin}) = [-\frac{\pi}{2}, \frac{\pi}{2}]$.

Meaning $\text{ArcSin}(\frac{1}{2}) = t$ means:

$$\begin{cases} \sin(t) = \frac{1}{2} \\ -\frac{\pi}{2} \leq t \leq \frac{\pi}{2} \end{cases}$$

Composite

$$\begin{aligned} \forall x \in [-\frac{\pi}{2}, \frac{\pi}{2}], \text{ArcSin}(\text{Sin}(x)) &= x \\ \forall y \in [-1, 1], \text{Sin}(\text{ArcSin}(y)) &= y \end{aligned}$$

4.7 Derivative of ArcSin

Result

$$\frac{d\text{ArcSin}(x)}{dx} = \frac{1}{\sqrt{1-x^2}}$$

Derive.

$$\begin{aligned}
 & \forall x \in [-1, 1] \\
 & \sin(\arcsin(x)) = x \\
 & \frac{d}{dx} \sin(\arcsin(x)) = \frac{d}{dx} x \\
 & \cos(\arcsin(x)) \cdot \frac{d}{dx} \arcsin(x) = 1 \\
 & \frac{d}{dx} \arcsin(x) = \frac{1}{\cos(\arcsin(x))} \\
 & \text{Let } \theta = \arcsin(x) \\
 & \cos^2(\theta) = 1 - \sin^2(\theta) \\
 & \cos(\theta) = \pm \sqrt{1 - x^2} \\
 & \text{Since } \forall \theta \in [-\frac{\pi}{2}, \frac{\pi}{2}], \sin(\theta) \geq 0 \\
 & \implies \cos(\theta) = +\sqrt{1 - x^2} \\
 & \implies \frac{d}{dx} \arcsin(x) = \frac{1}{\sqrt{1 - x^2}}
 \end{aligned}$$

■

4.8 Other inverse trig functions

4.8.1 $y = \cos(x)$

Definition \arccos is the inverse function to the restriction of $\cos(x)$ to $[0, \pi]$, and,

$$\forall x \in [-1, 1], \forall y \in [0, \pi], x = \arccos(y) \iff \cos(y) = x$$

Result

$$\frac{d}{dx} \arccos(x) = -\frac{1}{\sqrt{1 - x^2}}$$

4.8.2 $y = \tan(x)$

Definition $\arctan(x)$ is the inverse function to the restriction of $\tan(x)$ to $[-\frac{\pi}{2}, \frac{\pi}{2}]$, and,

$$\forall y \in \mathbb{R}, \forall x \in [-\frac{\pi}{2}, \frac{\pi}{2}], x = \arctan(y) \iff \tan(x) = y$$

5 Video Playlist 5

5.1 Usage of MVT

Theorem Let I be an open interval. Let f be a function defined on I . If $\forall x \in I, f'(x) = 0$ then f is a constant function.

If we want to prove this theorem, we need mean value theorem

5.2 Local Extreme Theorem

Definition Let f be a function with domain I , let $c \in I$.

- f takes **maximum** at c if $\forall x \in I, f(x) \leq f(c)$.
- f takes **local maximum** at c if $\exists \delta > 0, \text{ s.t. } |x - c| < \delta \implies f(x) \leq f(c)$.

Definition Let f be a function with domain I , let $c \in I$.

- f takes **minimum** at c if $\forall x \in I, f(x) \geq f(c)$.
- f takes **local minimum** at c if $\exists \delta > 0, \text{ s.t. } |x - c| < \delta \implies f(x) \geq f(c)$.

End-point cannot be a local extremum since the definition of local extremum requires a open interval at both left and right sides around point c .

Theorem (Local EVT) Let f be a function with domain I as an interval. Let $c \in I$, then if,

1. $f(c)$ is an extremum.
2. c is an interior point.

then, $f'(c) = 0$ or DNE.

Definition Point $c \in I$ for function f is a **critical point** if $f'(c) = 0$ or it does not exist.

Proof. (Local EVT) Proof is in two parts: (1) f has maximum at c , (2) f has minimum at c .

Part1: $f(c)$ is a maximum

Take left and right side limits

$$\text{As } x \rightarrow c^+, x - c > 0$$

$$\text{As } x \rightarrow c^-, x - c < 0$$

$$\text{By definition of maximum } f(x) - f(c) \leq 0$$

Left limit

$$x - c < 0 \wedge f(x) - f(c) \leq 0$$

$$\implies \lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} \geq 0$$

Right limit

$$x - c > 0 \wedge f(x) - f(c) \leq 0$$

$$\implies \lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \leq 0$$

For limit to exist

$$\lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \leq 0 \wedge \lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} \geq 0$$

$$\implies \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = 0$$

$$\iff f'(c) = 0$$

Part2: $f(c)$ is a minimum

Take left and right side limits

$$\text{As } x \rightarrow c^+, x - c > 0$$

$$\text{As } x \rightarrow c^-, x - c < 0$$

$$\text{By definition of minimum } f(x) - f(c) \geq 0$$

Left limit

$$x - c < 0 \wedge f(x) - f(c) \geq 0$$

$$\implies \lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} \leq 0$$

Right limit

$$x - c > 0 \wedge f(x) - f(c) \geq 0$$

$$\implies \lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \geq 0$$

For limit to exist

$$\lim_{x \rightarrow c^+} \frac{f(x) - f(c)}{x - c} \geq 0 \wedge \lim_{x \rightarrow c^-} \frac{f(x) - f(c)}{x - c} \leq 0$$

$$\implies \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = 0$$

$$\iff f'(c) = 0$$

5.3 Find Extremum

Example find extremum of function $f(x) = x^3 - 3x^2 - 9x + 3$ for $I = [-4, 4]$
Steps

1. Ensure existence of extremum. f is polynomial and therefore continuous, and $[-4, 4]$ is a compact set. By EVT, extremum exist.
2. Find all *critical points* and *end-points*.
3. Compare values at candidate points.

5.4 Rolle's Theorem

Theorem let $a < b$, let f be a function defined on a closed interval $[a, b]$ (Compact set). Then, if,

1. $f(x)$ is continuous on $[a, b]$.
2. $(\wedge) f(x)$ is differentiable on (a, b) .
3. $(\wedge) f(a) = f(b)$.

then,

$$\exists c \in (a, b) \text{ s.t. } f'(c) = 0$$

Proof.

By EVT, $f(x)$ has extremum in $[a, b]$.

Case1 Interior Extremum Point. ($c \in (a, b)$)

By Local EVT, $f'(c) = 0 \vee f'(c) DNE$

By (ii) $f'(c) = 0$

Case2 End-point Extremum

Since (iii) $f(a) = f(b)$

$\forall x \in (a, b)$

$f(x) \leq \max(f(a), f(b))$

$f(x) \geq \min(f(a), f(b))$

$\implies f(x)$ is constant.

$\implies \forall c \in (a, b), f'(c) = 0$

■

5.5 Application of Rolle's Theorem

Application How many zeros does a function have.

Step 1 Use IVT to prove it has *at least* n zeros.

Step 2 Use Rolle's theorem to prove it has *at most* n zeros.

Example

$$g(x) = x^6 + x^2 + x - 2$$

IVT Applied

$$g(-2) = 64$$

$$g(0) = -2$$

$$g(1) = 1$$

So that, $g(x)$ has at least 2 zeros.

Rolle's theorem applied Assume $f(x_1) = f(x_2) = 0$, by Rolle's theorem, there must exist a $a \in (x_1, x_2)$ such that $f'(a) = 0$

Conclusion 1 Between any two zeros of f there must be *at least* one zero of f' .

Conclusion 2 $\#$ of zeros of $f' \geq \#$ of zeros of $f - 1$

Conclusion 2' $\#$ of zeros of $f \leq \#$ of zeros of $f' + 1$

$$g'(x) = 6x^5 + 2x + 1$$

$$g''(x) = 30x^4 + 2$$

$$g''(x) \text{ has no zeros}$$

5.6 (Lagrange)Mean Value Theorem

Theorem Let $a < b$, let f be a function defined on $[a, b]$, if,

1. f is continuous on $[a, b]$.
2. f is differentiable on (a, b) .

then,

$$\exists c \in (a, b) \text{ s.t. } f'(c) = \frac{f(b) - f(a)}{b - a}$$

5.7 Proof. of MVT

$$\text{Let } m = \frac{f(b) - f(a)}{b - a}$$

$$\text{Let } g(x) = f(x) - f(a) - m(x - a)$$

$$\text{Satisfies } g(a) = f(a) - f(a) - m(a - a) = 0$$

$$\wedge g(b) = f(b) - f(a) - m(b - a) = 0$$

By Rolle's Theorem

$$g(a) = g(b) = 0$$

$$\exists c \in (a, b) \text{ s.t. } g'(c) = 0$$

$$\implies \frac{d}{dx}[f(x) - f(a) - m(x - a)] = 0$$

$$\implies f'(c) = \frac{f(b) - f(a)}{b - a}$$

■

5.8 Zero-derivative implies constant

Theorem Let $a < b$. Let f be a function defined on $[a, b]$, then,

$$\forall x \in (a, b), f'(x) = 0 \wedge f \text{ is continuous on } [a, b] \implies f \text{ is constant on } [a, b].$$

proof.

$$\text{Let } x_1, x_2 \in [a, b] \wedge x_1 < x_2$$

$$\text{By MVT, } \exists c \in (x_1, x_2), \text{ s.t.}$$

$$f'(c) = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$$

$$\because f'(c) = 0$$

$$\therefore f(x_1) = f(x_2)$$

5.9 Monotonicity of functions

Definition Let f be a function defined on an interval I .

- f is **increasing on I** when

$$\forall x_1, x_2 \in I, x_1 < x_2 \implies f(x_1) < f(x_2)$$

- f is **non-decreasing on I** when

$$\forall x_1, x_2 \in I, x_1 < x_2 \implies f(x_1) \leq f(x_2)$$

Theorem Let $a < b$. Let f be a function defined on (a, b) . Then,

$$\forall x \in (a, b), f'(x) > 0 \implies f \text{ is increasing on } (a, b)$$

Theorem Let $a < b$. Let f be a function defined on $[a, b]$. Then,

$$\forall x \in (a, b), f'(x) > 0 \wedge f \text{ is continuous on } [a, b] \implies f \text{ is increasing on } [a, b]$$

Short summary On an open interval

- $f' = 0 \implies f$ constant.
- $f' > 0 \implies f$ increasing.
- $f' < 0 \implies f$ decreasing.

6 Video Playlist 6

Note This chapter focus on *optimization applications*, and there's no video for this topic.

7 Video Playlist 7

7.1 Integral

Integral Let $a < b$, let f be a positive function, then *integral of f from a to b* is denoted as:

$$\int_a^b f(x) dx$$

this is represented as the area of region under function f from $x = a$ to $x = b$.

7.2 Sigma Notation

Sigma Notation The sigma notation, with **index** i , could be represented in the following form:

$$\sum_{i=1}^N a_i = a_1 + a_2 + \cdots + a_N$$

7.3 Supremum and Infimum

Definitions Let $A \subseteq \mathbb{R}$, let $a \in \mathbb{R}$:

- **Upper bound:** a is a upper bound of A means $\forall x \in A, x \leq a$.
- **Least upper bound(l.u.b) / Supremum:** a is the least upper bound or supremum(sup) of A iff a is an upper bound of A and $\forall b \in \{\text{upper bound of } A\}, a \leq b$.

- **Maximum:** if supremum of $A \in A$, it's maximum of A .
- **Bounded above:** A is bounded above if A has (at least) one upper bound.

Definitions (counter-part) Let $A \subseteq \mathbb{R}$, let $a \in \mathbb{R}$:

- **Lower bound:** a is a lower bound of A means $\forall x \in A, x \geq a$.
- **Greatest lower bound(g.l.b) / Infimum:** a is the greatest lower bound (g.l.b) or infimum(inf) of A iff a is a lower bound of A and $\forall b \in \{\text{Lower bound of } A\}, a \geq b$.
- **Minimum:** if infimum of $A \in A$, it's the minimum of A .
- **Bounded below:** A is bounded below if A has (at least) one lower bound.

Theorem: The l.u.b. principle Let $A \subseteq \mathbb{R}$, if A is bounded above and $A \neq \emptyset$, then, A has a least upper bound(supremum).

Theorem: The g.l.b principle Let $A \subseteq \mathbb{R}$, if A is bounded below and $A \neq \emptyset$, then, A has a greatest lower bound(infimum).

7.4 Supremum and Infimum of a function

Definition Supremum of a function f on a domain I is defined as:

$$\sup_{x \in I} f(x) = \sup\{f(x) \mid x \in I\}$$

Theorem Let f be a function defined on domain $I \neq \emptyset$, if f is bounded above, then $\exists \sup_{x \in I} f(x)$. Similarly, if f is bounded below, then $\exists \inf_{x \in I} f(x)$.

Theorem(EVT) Let $a < b$, let f defined on $[a, b]$, if f is continuous on $[a, b]$, then f has a maximum and a minimum on $[a, b]$.

7.5 Definition of Integral (i)

Definition A **partition** of the interval $[a, b]$ is a finite set P , s.t. $\{a, b\} \subseteq P$.

Notation $P = \{x_0, x_1, \dots, x_N\}$ on $[a, b]$. Implicitly, x_i are ordered, such that, $a = x_0 < x_1 < \dots < x_N = b$.

Let f be bounded on $[a, b]$, let $P = \{x_0, x_1, \dots, x_N\}$, let $m_i = \inf_{x \in [x_{i-1}, x_i]} f(x)$, and $M_i = \sup_{x \in [x_{i-1}, x_i]} f(x)$, and $\Delta x_i = x_i - x_{i-1}$.

Definition P-Lower sum of f is defined as:

$$L_P(f) = \sum_{i=1}^N (m_i \Delta x_i)$$

Definition P-Upper sum of f is defined as:

$$U_P(f) = \sum_{i=1}^N (M_i \Delta x_i)$$

Property For all partition P on interval $[a, b]$, the lower sum and upper sum satisfy the following inequality,

$$L_P(f) \leq \int_a^b f(x) dx \leq U_P(f)$$

7.6 Definition of Integral (ii): Properties of $U_P(f)$ and $L_P(f)$

Let f be a bounded function on $[a, b]$, let P and Q be partitions of $[a, b]$, the lower sums and upper sums have the following properties.

1. (Always) $L_P(f) \leq U_P(f)$.
2. If $P \subseteq Q$ (Q is a finer partition), then $L_P(f) \leq L_Q(f) \wedge U_P(f) \geq U_Q(f)$.
3. (Always) $L_P(f) \leq U_Q(f)$

Proof

Let $R = P \cup Q$,

so that, $P \subseteq R \wedge Q \subseteq R$. (R is finer than both P and Q)

$$\begin{aligned} L_P(f) &\leq L_R(f) \leq U_R(f) \leq U_Q(f) \\ \implies L_P(f) &\leq U_Q(f) \end{aligned}$$

■

7.7 Definition of Integral (iii): Upper Integral and Lower Integral

Definition Let f be a bounded function on $[a, b]$, then, lower integral of f from a to b is defined as,

$$I_a^b(f) = \sup\{\text{lower sums of } f\}$$

and the upper integral of f from a to b is defined as,

$$\overline{I_a^b(f)} = \inf\{\text{upper sums of } f\}$$

Then if $I_a^b(f) < \overline{I_a^b(f)}$, then f is **non-integrable** on $[a, b]$.

7.8 An example of integrable function

$$f(x) = \begin{cases} 1 & \text{if } x = 0 \\ 0 & \text{if } x \neq 0 \end{cases} \quad \text{on } [-1, 1]$$

7.9 An example of non-integrable function

$$g(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases} \quad \text{on } [-1, 1]$$

7.10 Integrals as limits

Definition Let $P = \{x_0, x_1, \dots, x_N\}$ be a partition of $[a, b]$, the **norm** of P is defined as:

$$\|P\| = \max\{\Delta x_1, \Delta x_2, \dots, \Delta x_N\}$$

Theorem - Lower Integrals For lower integrals, we have,

$$\underline{I}_a^b(f) = \lim_{\|P\| \rightarrow 0} L_P(f) = \sup\{\text{lower sums of } f\}$$

alternatively, using $\delta - \epsilon$ expression,

$$\forall \epsilon > 0, \exists \delta > 0 \text{ s.t. } \forall P \text{ over } [a, b], \|P\| < \delta \implies |L_P(f) - \underline{I}_a^b(f)| < \epsilon$$

theorem - Upper Integrals For upper integrals, we have,

$$\overline{I}_a^b(f) = \lim_{\|P\| \rightarrow 0} U_P(f)$$

7.11 Riemann Sums

Definition Fix a partition P on $[a, b]$, $m_i = \inf_{x \in [x_{i-1}, x_i]} f(x)$, $M_i = \sup_{x \in [x_{i-1}, x_i]} f(x)$, pick $x_i^* \in [x_{i-1}, x_i]$, so that,

$$\begin{aligned} m_i &\leq f(x_i^*) \leq M_i \\ \implies m_i \Delta x_i &\leq f(x_i^*) \Delta x_i \leq M_i \Delta x_i \\ \implies L_P(f) &= \sum_{i=1}^N (m_i \Delta x_i) \leq \sum_{i=1}^N (f(x_i^*) \Delta x_i) \leq \sum_{i=1}^N (M_i \Delta x_i) = U_P(f) \end{aligned}$$

where the term $\sum_{i=1}^N (f(x_i^*) \Delta x_i)$ is called a **Riemann sum**.

Definition Let f be a bounded function on $[a, b]$, let $P = \{x_0, x_1, \dots, x_N\}$ be a partition on $[a, b]$, for each i , pick **any** point $x_i^* \in [x_{i-1}, x_i]$. then,

$$S_P^*(f) = \sum_{i=1}^N (f(x_i^*) * \Delta x_i)$$

is a **Riemann sum** for f and P . (There are infinitely many Riemann sum).

In general, we have,

$$L_P(f) \leq S_P^*(f) \leq U_P(f)$$

and also,

$$\lim_{\|P\| \rightarrow 0} L_P(f) = \underline{I_a^b(f)}$$

$$\lim_{\|P\| \rightarrow 0} U_P(f) = \overline{I_a^b(f)}$$

and if f is **integrable**, then

$$\lim_{\|P\| \rightarrow 0} L_P(f) = \lim_{\|P\| \rightarrow 0} U_P(f) = \int_a^b f(x) dx$$

By Squeeze Theorem,

$$\lim_{\|P\| \rightarrow 0} S_P^*(f) = \int_a^b f(x) dx$$

7.12 Properties of the integral

Property 1

$$\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$$

Property 2

$$\int_a^b [cf(x)] dx = c \int_a^b f(x) dx$$

Property 3 If f is bounded on $[a, c]$, and f is integrable on $[a, b]$ and integrable on $[b, c]$, then,

$$\int_a^c f(x) dx = \int_a^b f(x) dx + \int_b^c f(x) dx$$

Property 4: Backward Integrals

$$\int_b^a f(x) dx = - \int_a^b f(x) dx$$

Negative function f Integral for negative function is the negative area.

$$\int_a^b f(x) dx$$

8 Video Playlist 8

8.1 Anti-derivatives

Notations

- **Definite integral** $\int_a^b f(x) dx$
- **Indefinite integral** $\int f(x) dx$

Definition Let f be a function defined on an interval, *an anti-derivative* of f is any function F that

$$F' = f$$

Note As a consequence of MVT, if two functions have same derivative on an interval, then they differ by a constant.

8.2 Functions Defined as Integrals

Consider integrable function f , define function F as the definite integral from a , a fixed point in domain of f , to another point x in domain of f , that's,

$$F(x) = \int_a^x f(t) dt$$

Methodology Let I be an interval, let $a \in I$ and let f be a function integrable on I , then for each $x \in I$, compute $F(x) = \int_a^x f(t) dt$ as a number.

8.3 The Fundamental Theorem of Calculus: Part 1

This provides connections between definite integrals and anti-derivatives

Theorem: FTC(part 1)

- Let I be an interval,
- Let $a \in I$,
- Let f be a function on I .

Define $F(x)$ as

$$F(x) = \int_a^x f(t) dt$$

If f is continuous, then F is differentiable and $F' = f$, that's,

$$F'(x) = f(x) \quad \forall x \in I$$

8.4 A Proof of Part 1 of the FTC

Proof.

$$\begin{aligned}
 & \text{Let (fix) } x \in I \\
 & \text{WTS. } F'(x) = f(x) \\
 F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\
 &= \lim_{h \rightarrow 0} \left[\frac{1}{h} (F(x+h) - F(x)) \right] \\
 &= \lim_{h \rightarrow 0} \left[\frac{1}{h} \left(\int_a^{x+h} f(t) dt - \int_a^x f(t) dt \right) \right] \\
 &= \lim_{h \rightarrow 0} \left[\frac{1}{h} \int_x^{x+h} f(t) dt \right]
 \end{aligned}$$

Consider $h > 0$ (for negative h , the proof would be similar)

$$\text{Let } M_h = \sup_{[x, x+h]} (f)$$

$$\text{Let } m_h = \inf_{[x, x+h]} (f)$$

Then we have, by definition of infimum and supremum,

$$m_h \leq \frac{1}{h} \int_x^{x+h} f(t) dt \leq M_h$$

Since f is continuous on $[x, x+h]$, by EVT, it has maximum and minimum on this interval.

$$\exists c_h \in [x, x+h] \text{ s.t. } M_h = f(c_h)$$

$$\exists d_h \in [x, x+h] \text{ s.t. } m_h = f(d_h)$$

$$\because \lim_{h \rightarrow 0} c_h = x \wedge \lim_{h \rightarrow 0} d_h = x$$

$$\therefore \lim_{h \rightarrow 0} M_h = \lim_{h \rightarrow 0, c_h \rightarrow x} f(c_h) = f(x) \text{ (since } f \text{ is continuous.)}$$

$$\text{Similarly, } \lim_{h \rightarrow 0} m_h = \lim_{h \rightarrow 0, d_h \rightarrow x} f(d_h) = f(x)$$

$$\text{By Squeeze Theorem, } \lim_{h \rightarrow 0} \left[\frac{1}{h} \int_x^{x+h} f(t) dt \right] = f(x)$$

$$\therefore F'(x) = f(x) \forall x \in I$$

■

8.5 The Fundamental Theorem of Calculus: Part 2

This provides a quick way to compute definite integrals.

Theorem: FTC(part 2)

- Let $a < b \in \mathbb{R}$,
- let f be continuous on $[a, b]$,

then,

$$\int_a^b f(x) \, dx = G(b) - G(a)$$

where G is any anti-derivative of f .

Notation

$$G(b) - G(a) = G(x)|_{x=a}^{x=b} = G(x)|_a^b$$

8.6 A Proof of Part 2 of the FTC

Proof.

We know that, from the first part of FTC, $G' = f$,

$$\text{WTS. } \int_a^b f(x) \, dx = G(b) - G(a)$$

$$\text{Define } F(x) = \int_a^x f(t) \, dt$$

$$\text{WTS. } F(b) = G(b) - G(a)$$

$$\text{Since } f \text{ is continuous, } F' = f$$

By the consequence of MVT,

$$F' = G' \implies \exists C \in \mathbb{R} \text{ s.t. } F - G = C \forall x \in [a, b]$$

$$\text{at } x = a, F(a) = 0 \implies C = -G(a)$$

$$\implies \forall x \in [a, b] F(x) = G(x) - G(a)$$

$$\text{at } x = b, F(b) = G(b) - G(a)$$

■

8.7 Summary: Definite and indefinite integrals, notation, definitions and theorems.

8.7.1 Definite Integral.

$$\int_a^b f(x) \, dx$$

Theorem (Formal definite) if $\overline{I}_a^b(f) = \underline{I}_a^b(f)$ then $\int_a^b f(x) \, dx = \overline{I}_a^b(f) = \underline{I}_a^b(f)$.

Theorem (FTC: part 2) Choose one anti-derivative $G(x)$ of $f(x)$, then compute the definite integral as $\int_a^b f(x) dx = G(b) - G(a)$.

8.7.2 Indefinite Integral

$$\int f(x) dx \text{ A collection of functions.}$$

Find indefinite integral Find $G(x)$ as one anti-derivative, by the consequence of MVT, then the indefinite integral of f could be constructed as,

$$F(x) = \{G(x) + C \mid C \in \mathbb{R}\}$$

8.7.3 Function Defined by an Integral.

$$F(x) = \int_a^x f(t) dt \text{ This is one function with fixed value of } a.$$

Theorem (FTC: part 1) if f is continuous, then $F'(x) = f(x)$

9 Video Playlist 9

9.1 Integration By Substitution: derivation of the formula

Backwards usage of chain rule.

If $\int f(x) dx = F(x)$ is the anti-derivative of $f(x)$, then

$$F(g(x)) = \int f(g(x))g'(x) dx = F(g(x))$$

9.2 Example 2

9.3 Example 3

9.4 Example 4

Theorem Let $a < b$, let f be a continuous function, let g be a function with continuous derivative in $[a, b]$, assume the range of g on $[a, b]$ is contained in the domain of f . Then,

$$\int_a^b f(g(x))g'(x) dx = \int_{g(a)}^{g(b)} f(u) du$$

9.5 Integration by parts

Backwards product rule

Let f and g be two differentiable function, by product rule of differentiation, we have,

$$\begin{aligned} f'(x)g(x) + f(x)g'(x) &= \frac{d}{dx} f(x)g(x) \\ \implies \int f'(x)g(x) + f(x)g'(x) dx &= f(x)g(x) + C \\ \implies \int f'(x)g(x) dx + \int f(x)g'(x) dx &= f(x)g(x) + C \\ \implies \int f'(x)g(x) dx &= f(x)g(x) - \int f(x)g'(x) dx \end{aligned}$$

The integral constant is implicitly contained in the integral term.

9.6 Examples

Example 1

$$\int x^2 e^x dx$$

Example 2

$$\int e^x \sin x dx$$

Use integration by parts twice.

Example 3

$$\int \arctan x dx$$

Consider the form $1 \times f(x)$ as partition method.

9.7 Integration of products of trigonometric functions

Types

$$\int \sin^n x \cos^m x dx$$

$$\int \sec^n x \tan^m x dx$$

Keys

$$\sin^2(x) + \cos^2(x) = 1$$

$$\sec^2(x) = 1 + \tan^2(x)$$

Summary I Consider the integral in the following form

$$\int \sin^n x \cos^m x \, dx$$

- If **m is odd** then try $u = \sin(x)$, then $du = \cos(x)dx$
- If **n is odd** then try $u = \cos(x)$, then $du = -\sin(x)dx$

10 Video Playlist 10

Note This chapter focus on *volumes*, and there's no video for this topic.

11 Video Playlist 11

11.1 What Is a Sequence

Definition A **sequence** is a function with domain \mathbb{N} .

11.1.1 Conventions

Functions function with domain interval.

- x as variable.
- $f(x)$ as value at x .

Sequence function with domain \mathbb{N} .

- n as variable.
- a_n as value at n .

A sequence is not a set.

11.1.2 Describe sequences

Equation $a_n = \frac{2^n n!}{n+1}$

First few values $\{1, 2, 4, 8, 16, \dots\}$

Words $p_n = n$ -th prime.

Recurrence relation e.g. Fibonacci Sequence.

$$\{F_n\}_{n=0}^{\infty} : F_0 = F_1 = 1, F_n = F_{n-1} + F_{n-2} \, \forall n \geq 2$$

A general definition A sequence is a function with domain $\{n \in \mathbb{Z} \mid n \geq n_0\}$ for some fixed $n_0 \in \mathbb{Z}$.

11.2 The Limit of a Sequence

Example

$$\left\{\frac{n}{n+1}\right\}_{n=0}^{\infty} \quad \lim_{n \rightarrow \infty} \frac{n}{n+1} = 1$$

Definition(Limit) We say that the sequence $\{a_n\}_{n=0}^{\infty}$ converges to the number $L \in \mathbb{R}$ when

$$\forall \epsilon > 0, \exists n_0 \in \mathbb{N} \text{ s.t. } \forall n \in \mathbb{N}, n \geq n_0 \implies |L - a_n| < \epsilon$$

denoted as

$$\lim_{n \rightarrow \infty} a_n = L \text{ or } a_n \rightarrow L$$

Tail: all terms of the sequence after the first few terms.

Every interval centred at L contains a tail of the sequence.

Definition A sequence is **convergent** if it has a limit. This sequence is **divergent** if it does not have a limit.

11.3 Properties of Limits of Sequences

Properties from the limit of functions

- Limit laws: Yes
- Squeeze theorem: Yes
- *L'Hôpital's Rule*: No

11.3.1 Sequence from a function

Let $c \in \mathbb{Z}$ and function f defined on $[c, \infty)$, and define the sequence $\{a_n\}_{n=c}^{\infty}$ as

$$a_n = f(n)$$

We have if $\lim_{n \rightarrow \infty} f(n) = L$ then $\lim_{n \rightarrow \infty} a_n = L$. If $\lim_{n \rightarrow \infty} f(n)$ DNE, then $\lim_{n \rightarrow \infty} a_n$ may or may not exist.

11.3.2 Composite of sequence and function

Theorem If $a_n \rightarrow L$ and f is continuous at L then

$$f(a_n) \rightarrow f(L)$$

11.4 Monotonic and Bounded Sequences

11.4.1 Monotonic Sequences

Definition We say $\{a_n\}_{n=0}^{\infty}$ is **increasing** if

$$\forall n, m \in \mathbb{N}, n < m \implies a_n < a_m$$

Also, we say this sequence is **non-decreasing** if the inequality is in the weak form as

$$\forall n, m \in \mathbb{N}, n < m \implies a_n \leq a_m$$

Definition We say $\{a_n\}_{n=0}^{\infty}$ is **decreasing** if

$$\forall n, m \in \mathbb{N}, n < m \implies a_n > a_m$$

Also, if the inequality is in the weak form as

$$\forall n, m \in \mathbb{N}, n < m \implies a_n \geq a_m$$

we say this sequence is **non-increasing**.

Definition We say a sequence $\{a_n\}_{n=0}^{\infty}$ is **monotonic** if it has any of the four properties above.

Definition $\{a_n\}_{n=0}^{\infty}$ is **eventually decreasing** if

$$\exists n_0 \in \mathbb{N}, \text{ s.t. } \forall n \in \mathbb{N}, n \geq n_0 \implies a_n > a_{n+1}$$

11.4.2 Bounded Sequences

Definition We say a sequence $\{a_n\}_{n=0}^{\infty}$ is **bounded below** if

$$\exists A \in \mathbb{R} \text{ s.t. } \forall n \in \mathbb{N}, A \leq a_n$$

Similarly, the sequence is **bounded above** if

$$\exists B \in \mathbb{R} \text{ s.t. } \forall n \in \mathbb{N}, B \geq a_n$$

Definition We say a sequence is **bounded** if and only if it is both bounded above and below.

Theorem If a sequence is convergent then it is bounded.

Theorem 2A(The monotone convergence theorem for sequence) If a sequence is eventually increasing and bounded above, then it is convergent

Theorem If a sequence is eventually increasing and not bounded above then it divergent to ∞ .

Remark for a sequence:

$$\text{Sequence} \begin{cases} \text{Convergent} \\ \text{Divergent} \begin{cases} \text{to } \infty \\ \text{to } -\infty \\ \text{Oscillating} \end{cases} \end{cases}$$

11.5 Proof: Every convergent sequence is bounded

Theorem Let $\{a_n\}_{n=0}^{\infty}$ be a sequence, if $\{a_n\}_{n=0}^{\infty}$ is convergent then the sequence is bounded. Equivalently,

Proof.

Assume sequence $\{a_n\}_{n=0}^{\infty}$ is convergent.

Let L be the limit.

By the definition of limit, choose $\epsilon = 10$

So that, $\exists n_0 \in \mathbb{N} \text{ s.t. } \forall n \in \mathbb{N}, n \geq n_0 \implies L - 10 \leq a_n \leq L + 10$

Take $A = \min\{a_0, \dots, a_{n_0-1}, L - 10\}$

Take $B = \max\{a_0, \dots, a_{n_0-1}, L + 10\}$

By definition of max and min, let $n \in \mathbb{N}$

case $1n > n_0 \implies A \leq a_n \leq B$

case $2n \geq n_0 \implies L - 10 \leq a_n \leq L + 10$

Since $A \leq L - 10 \wedge B \geq L + 10$

$\implies A \leq a_n \leq B \forall n \in \mathbb{N}$

$\therefore \{a_n\}_{n=0}^{\infty}$ is bounded. ■

11.6 The monotone convergence theorem of sequences

(General) Theorem If a sequence is (eventually) monotonic and bounded then it is convergent.

(Particular Case) Theorem 1 If a sequence is increasing and bounded above the it's convergent.

Proof.

Let $\{a_n\}_{n=0}^{\infty}$ be a sequence that's increasing and bounded above.

Consider $A = \{a_n \mid n \in \mathbb{N}\} \neq \emptyset$

By least upper bound principle, there exists a supremum of set A

Take $L = \sup\{A\}$

Let $\epsilon > 0$

By definition of supremum,

$\exists a_{n_0} \in A$ s.t. $a_{n_0} > L - \epsilon$

Take this value n_0

Since sequence is increasing,

$\forall n \geq n_0$ $a_n > L - \epsilon$

Also, by definition of supremum, $a_n \leq L$

$\implies a_n \leq L + \epsilon$

Therefore, $\forall n \in \mathbb{N}, n \geq n_0 \implies L - \epsilon < a_n < L + \epsilon$

Therefore, $\lim_{n \rightarrow \infty} \{a_n\}_{n=0}^{\infty} = L$

Therefore, $\{a_n\}_{n=0}^{\infty}$ is convergent. ■

11.7 the Big theorem of sequences

Definition (for positive sequences only) Let $\{a_n\}_{n=1}^{\infty}$ and $\{b_n\}_{n=1}^{\infty}$ be positive sequences.

$$a_n \ll b_n \iff \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$$

say $\{a_n\}$ is **much smaller than** $\{b_n\}$.

Theorem for every $a > 0$ and $c > 1$

$$\ln n \ll n^a \ll c^n \ll n! \ll n^n$$

12 Video Playlist 12

12.1 Improper Integral

12.1.1 Improper integral "type 1" (Unbounded domain)

Definition Let $a \in \mathbb{R}$ and f continuous on $[a, \infty]$ the integral of f from a to ∞ , denoted as

$$\int_a^{\infty} f(x) dx = \lim_{b \rightarrow \infty} \int_a^b f(x) dx$$

assuming the limit exists. If the limit exists, the integral is called **convergent**, otherwise, it's called **divergent**.

12.2 The most important family of improper integrals

Let $p \in \mathbb{R}$ consider

$$\int_1^\infty \frac{1}{x^p} dx$$

Summary

$$\int_1^\infty \frac{1}{x^p} dx \text{ is } \begin{cases} \text{convergent} & \text{if } p > 1 \\ \text{divergent to } \infty & \text{if } p \leq 1 \end{cases}$$

12.3 Example

$$\int_0^\infty \sin(x) dx$$

12.4 The most important family of improper integral

Consider

$$I = \int_1^\infty \frac{1}{x^p} dx$$

Summary

1. $p > 1 \iff I$ converges.
2. $p \leq 1 \iff I$ diverges to ∞ .

12.5 Example

Vertical asymptote improper.

$$\int_0^1 \ln(x) dx$$

12.6 Doubly improper integrals

General Strategy Assume A has **multiple** improper.

1. Break A into pieces with **single** improper at their endpoints.
2. If each piece convergent **separately**, then A converges.
3. Else, A diverges, it's *not a number*.

12.7 Basic Comparison Test

Theorem Let $a \in \mathbb{R}$,

Let f and g be *continuous* functions on $[a, \infty)$, and

$$\forall x \geq a, 0 \leq f(x) \leq g(x)$$

we have,

1. $\int_a^\infty g(x) dx < \infty \implies \int_a^\infty f(x) dx < \infty$
2. $\int_a^\infty f(x) dx = \infty \implies \int_a^\infty g(x) dx = \infty$

12.8 Examples

12.9 Limit Comparison Test

Theorem Let $a \in \mathbb{R}$, f and g are *positive* and *continuous* functions on $[a, \infty)$.
And the following limit exists,

$$L = \lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} \in \mathbb{R}$$

Then, $\int_a^\infty f(x) dx$ and $\int_a^\infty g(x) dx$ are **both** convergent or **both** divergent.

12.10 Proof of LCT

Omitted

13 Video Playlist 13

13.1 Infinite Sums

Nothing.

13.2 Definition of series

Definition Series $\sum_{n=1}^\infty a_n$ is defined as

$$\lim_{k \rightarrow \infty} S_k$$

where $S_k = \sum_{n=1}^k a_n$ as finite sum. If the above limit exist, we say series $\sum_{n=1}^\infty a_n$ is convergent (it's a *number*), else series is divergent and it's not a number.

13.3 Example

$$\sum_{n=1}^\infty \frac{1}{n^2 + n} = \sum_{n=1}^\infty \frac{1}{n} - \frac{1}{n+1} = \lim_{k \rightarrow \infty} 1 - \frac{1}{k+1} = 1$$

13.4 Divergent Series Examples

$$S = \sum_{n=1}^{\infty} 1 = \lim_{k \rightarrow \infty} k = \infty$$

$$S = \sum_{n=1}^{\infty} (-1)^n = \begin{cases} 0 & \text{if } k \text{ is odd} \\ 1 & \text{if } k \text{ is even} \end{cases} \quad \text{divergent due to oscillation.}$$

13.5 Geometric Series

Let $x \in \mathbb{R}$

$$S = \sum_{n=0}^{\infty} x^n = \lim_{k \rightarrow \infty} S_k = \lim_{k \rightarrow \infty} \sum_{n=0}^{\infty} x^n$$

Consider

$$S_k = 1 + x + x^2 + \cdots + x^k$$

$$xS_k = x + x^2 + x^3 + \cdots + x^{k+1}$$

$$S_k - xS_k = 1 - x^{k+1}$$

If $x = 1$, the series is simply divergent to ∞ .

$$S_k = \frac{1 - x^{k+1}}{1 - x}, \quad x \neq 1$$

$$S = \lim_{k \rightarrow \infty} \frac{1 - x^{k+1}}{1 - x} = \frac{1 - \lim_{k \rightarrow \infty} (x^{k+1})}{1 - x} = \begin{cases} \frac{1}{1-x} & \iff x \in (-1, 1) \\ \text{Divergent} & \begin{cases} \infty & \iff x > 1 \\ \text{Oscillating} & \iff x \leq -1 \end{cases} \end{cases}$$

13.6 Linearity of series

Simple form of fact

$$\sum_{n=0}^{\infty} a_n + \sum_{n=0}^{\infty} c b_n = \sum_{n=0}^{\infty} a_n + c \sum_{n=0}^{\infty} b_n, \quad \forall c \in \mathbb{R}$$

Theorem If series $\sum_{n=0}^{\infty} a_n$ and $\sum_{n=0}^{\infty} b_n$ are both convergent, then $\sum_{n=0}^{\infty} a_n + b_n$ is also convergent and

$$\sum_{n=0}^{\infty} a_n + b_n = \sum_{n=0}^{\infty} a_n + \sum_{n=0}^{\infty} b_n$$

Proof.

$$\text{Let } \sum_{n=0}^{\infty} a_n = \lim_{k \rightarrow \infty} S_k$$

$$\text{Let } \sum_{n=0}^{\infty} b_n = \lim_{k \rightarrow \infty} T_k$$

$$\text{Let } \sum_{n=0}^{\infty} a_n + b_n = \lim_{k \rightarrow \infty} R_k$$

$$\text{Where } S_k = \sum_{n=0}^k a_n$$

$$T_k = \sum_{n=0}^k b_n$$

$$R_k = \sum_{n=0}^k a_n + b_n$$

$$\text{Since } R_k = S_k + T_k, \forall k \in \mathbb{N}$$

$$\text{By limit laws, } \lim_{k \rightarrow \infty} S_k + \lim_{k \rightarrow \infty} T_k = \lim_{k \rightarrow \infty} R_k$$

$$\implies \sum_{n=0}^{\infty} a_n + \sum_{n=0}^{\infty} b_n = \sum_{n=0}^{\infty} a_n + b_n$$

■

Theorem If series $\sum_{n=0}^{\infty} a_n$ is convergent, then for any $c \in \mathbb{R}$, series $\sum_{n=0}^{\infty} c a_n$ is also convergent and

$$\sum_{n=0}^{\infty} c a_n = c \sum_{n=0}^{\infty} a_n$$

General proof procedure

1. Write series as limit of partial sums.
2. Manipulate partial sums (finite).
3. Manipulate limits.

13.7 the Tail of a series

Fact Consider two series

$$\sum_{n=0}^{\infty} a_n \text{ convergent} \iff \sum_{n=1}^{\infty} a_n \text{ convergent}$$

And $\sum_{n=1}^{\infty} a_n$ is a tail of series $\sum_{n=0}^{\infty} a_n$.

Notation We say $\sum_n^\infty a_n$ is convergent or divergent without specifying the starting index of the series.

Specific form of theorem If $\forall n \in \mathbb{N}$ [Condition(s)] then $\sum_n^\infty a_n$ is convergent or divergent.

General form of theorem If $\exists n_0 \in \mathbb{N}$ s.t. $\forall n \in \mathbb{N}, n \geq n_0 \implies$ [Condition(s)] then $\sum_n^\infty a_n$ is convergent.

13.8 A necessary condition for convergence of series

Fact Series $\sum_{n=0}^\infty a_n$ is convergent if and only if the sequence of its partial sums $\{S_n\}_{n=0}^k$ is convergent.

Theorem If $\sum_{n=0}^\infty a_n$ is convergent then

$$\lim_{n \rightarrow \infty} a_n = 0$$

Note The above theorem is often used as it's contrapositive form

$$\lim_{n \rightarrow \infty} a_n \neq 0 \implies \sum_{n=0}^\infty a_n \text{ is divergent}$$

Proof.

$$\text{Let } S = \sum_{n=0}^\infty a_n \text{ be convergent}$$

$$S = \lim_{k \rightarrow \infty} S_k, \quad S_k = \sum_{n=0}^k a_n$$

$$S = \lim_{k-1 \rightarrow \infty} S_{k-1}$$

$$\implies \lim_{k \rightarrow \infty} S_k - \lim_{k-1 \rightarrow \infty} S_{k-1} = 0$$

By the convergence assumption, those two limits above exist.

$$\implies \lim_{k \rightarrow \infty} S_k - S_{k-1} = 0$$

$$\implies \lim_{k \rightarrow \infty} a_k = 0$$

$$\implies \{a_n\}_{n=1}^\infty \rightarrow 0$$

■

13.9 Positive series

Definition A series $\sum_{n=0}^\infty a_n$ is positive when $\forall n \in \mathbb{N}, a_n > 0$. And a series is positive means it could never diverge to $-\infty$ or *oscillating*.

Notation (For positive series only)

1. $\sum_n^\infty a_n = \infty \iff$ divergent.
2. $\sum_n^\infty a_n < \infty \iff$ convergent.

13.10 The Integral Test

Theorem Let $a \in \mathbb{R}$, let f be a continuous, positive and decreasing function on $[a, \infty)$, then

$$\int_a^\infty f(x) dx < \infty \iff \sum_n^\infty f(n) < \infty$$

That's the improper integral and series have the same convergence/divergence feature. Note as

$$\int_a^\infty f(x) dx \sim \sum_n^\infty f(n)$$

13.11 Examples

p-series consider for what values of p the following series is convergent.

$$\sum_{n=1}^\infty \frac{1}{n^p}$$

Example2

$$\sum_{n=2}^\infty \frac{1}{n \ln n}$$

13.12 Comparison Tests for Series

Works exactly the same as basic and limit comparison tests for improper integrals.

Basic Comparison Test Consider series $\sum_{n=1}^\infty a_n$ and $\sum_{n=1}^\infty b_n$, assume¹

$$\forall n \in \mathbb{N}, 0 \leq a_n \leq b_n$$

then

$$\begin{aligned} \sum_{n=1}^\infty b_n < \infty &\implies \sum_{n=1}^\infty a_n < \infty \\ \sum_{n=1}^\infty a_n = \infty &\implies \sum_{n=1}^\infty b_n = \infty \end{aligned}$$

¹Notice that for large natural number n would be sufficient here also.

Limit Comparison Test Consider series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$, assume existing limit

$$L = \lim_{n \rightarrow \infty} \frac{a_n}{b_n} > 0$$

Then both series convergent or both of them divergent.

14 Video Playlist 14

14.1 Power Series: Example

$$g(x) = \sum_{n=1}^{\infty} \frac{x^n}{n3^n}$$

Domain of $g(x)$ is defined as

$$\{x \in \mathbb{R} \mid g(x) \text{ is convergent}\}$$

The above series convergent when $x \in [-3, 3)$, and $[-3, 3)$ is the **interval of convergence** and 3 is the **radius of convergent**.

14.2 Main Theorem

Definition Let $a \in \mathbb{R}$, a power series centred at a is a function f defined by a equation like

$$f(x) = \sum_{n=0}^{\infty} c_n (x - a)^n$$

Main Theorem Let $f(x)$ be a power series, then

1. Domain of f is an interval centred at a , with radius of convergence R , $0 \leq R \leq \infty$
2. In the *interior* of the interval of convergence, the series is *absolutely convergent*, in the *exterior* of IC, the series is *divergent*, and at the boundaries this theorem is *inconclusive*.
3. In the *interior* of the IC, power series can be treated like polynomial, without change of radius of convergence.

14.3 Taylor Polynomial Definition 1

Definition Let $f(x)$ and $g(x)$ be two functions that are continuous at a , let $n > 0$ and g is a approximation for f near a of order n when

$$\lim_{x \rightarrow a} \frac{f(x) - g(x)}{(x - a)^n} = 0$$

Definition Let $a \in \mathbb{R}$, let f be a continuous function defined at a , let $n \in \mathbb{N}$ then the n^{th} **Taylor Polynomial** for f at a is the polynomial P_n of *smallest* possible degree is an approximating for f near a of order n , that's,

$$\lim_{x \rightarrow a} \frac{f(x) - P_n(x)}{(x - a)^n} = 0$$

14.4 Taylor Polynomial Definition 2

Definition Let $a \in \mathbb{R}$, $n \in \mathbb{N}$ and let f be a C^n function at a , the n^{th} **Taylor Polynomial** for f at a is a polynomial P_n s.t.

$$P_n(a) = f(a), P'_n(a) = f'(a), \dots P_n^{(n)}(a) = f^{(n)}(a)$$

with smallest possible degree.

14.5 Taylor Polynomial Definition 3

Definition Let $a \in \mathbb{R}$, let f be a C^∞ function at a , the **Taylor's series** for f at a is the power series

$$S(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!} (x - a)^k$$

with the fact that

$$\forall k \in \mathbb{N}, S^{(k)}(a) = f^{(k)}(a)$$