ECO326 Advanced Microeconomic Theory A Course in Game Theory

Tianyu Du

April 23, 2019

This work is licensed under a Creative Commons "Attribution-NonCommercial 4.0 International" license.



- GitHub: https://github.com/TianyuDu/Spikey_UofT_Notes
- Website: TianyuDu.com/notes

Contents

1	Strategic Games and Dominant Strategies	3
2	Lecture 2. Jan. 14 2019 Iterated Elimination and Rationalizability 2.1 Iterated Elimination of Strictly Dominated Strategies/Actions	4 4
3	Lecture 3. Jan. 21 2019 Pure Strategy Nash Equilibrium	6
4	Lecture 4. Jan. 28. 2019 Nash Equilibrium: Examples	7
5	Lecture 5 Feb. 7 2019 Mixed Strategies 5.1 Lecture Contents	
6	Lecture 6 Feb. 25 2019 Extensive Form Games 6.1 An Introductory Example	
7	Lecture 7 Mar.4 2019 Examples of Extensive Form Games (Omitted)	20

8	Lecture 8 Mar.11 2019	
	Repeated Games	20
	8.1 Finitely Repeated Games	20
	8.2 Infinitely Repeated Games	20
9	Lecture 9 Mar. 18 2019	
	Bayesian Games	23
	9.1 Introduction	23
	9.2 Formal Definitions	23

1 Lecture 1. Jan. 7 2019

Strategic Games and Dominant Strategies

Definition 1.1 (pg.7). A **preference relation** is a <u>complete reflexive and transitive</u> binary relation.

Definition 1.2 (11.1, lec.1). A (strategic) game consists of

i a finite set of **players** N, with $|N| \ge 2$;

ii for each player $i \in N$, an **action set** $A_i \neq \emptyset$;

iii for each player $i \in N$, a **preference relation** \succeq_i defined on $A := \prod_{i \in N} A_i$.

and can be written as a triple $\langle N, (A_i), (\succeq_i) \rangle$.

Definition 1.3. An action profile is a *n*-tuple of actions $a_i \in A_i$ for each player $i \in N$ and denoted as

$$(a_i)_{i \in \mathcal{N}} \text{ or } (a_i) \tag{1.1}$$

The action profile space is defined as

$$A := \prod_{i \in N} A_i \tag{1.2}$$

And the space of **actions of opponents** is defined as

$$A_{-i} := \prod_{j \neq i} A_j \tag{1.3}$$

Definition 1.4 (Equivalent definition of game). For each player $i \in N$ we can define a **utility** function, $u_i : A \to \mathbb{R}$ such that

$$\forall (a_i), (a_i)' \in A, \ u_i((a_i)) \ge u_i((a_i)') \iff (a_i) \succsim_i (a_i)'$$

$$(1.4)$$

So the game can be defined as a triple $\langle N, (A_i), (u_i) \rangle$.

Definition 1.5. Action $a_i \in A_i$ is strictly dominated by action $\tilde{a}_i \in A_i$ if

$$\forall a_{-i} \in A_{-i}, \ u_i(a_i, a_{-i}) < u_i(\tilde{a}_i, a_{-i}) \tag{1.5}$$

And a_i is **weakly dominated** by \tilde{a}_i if

$$\forall \ a_{-i} \in A_{-i}, \ u_i(a_i, a_{-i}) \le u_i(\tilde{a}_i, a_{-i}) \tag{1.6}$$

and

$$\exists \ a_{-i} \in A_{-i}, \ u_i(a_i, a_{-i}) < u_i(\tilde{a}_i, a_{-i}) \tag{1.7}$$

Corollary 1.1 (Consequence of RCT). It is irrational to play strictly dominated actions. So rational choice theory suggests a player would never play strictly dominated strategies.

Definition 1.6. Action $a_i \in A_i$ is **strictly(weakly) dominant** if it strictly(weakly) dominates all other actions.

Definition 1.7. Action $a_i \in A_i$ is **strictly(weakly) dominated** if there exists another strategy strictly(weakly) dominates a_i .

Definition 1.8. A strategic game $\langle N, (A_i), (\succsim_i) \rangle$ is **finite** if for every $i \in N$, A_i is finite.

2 Lecture 2. Jan. 14 2019 Iterated Elimination and Rationalizability

2.1 Iterated Elimination of Strictly Dominated Strategies/Actions

Definition 2.1 (IESD). Given the initial game,

$$G_0 = \langle N, (A_i^0), (u_i) \rangle \tag{2.1}$$

At stage $k \in \mathbb{N}$,

$$G_k = \langle N, (A_i^k), (u_i) \rangle \tag{2.2}$$

In stage k, for all $i \in N$, find the set of strictly dominated actions, $D_i^k \subseteq A_i^k$.

i) If $\forall i \in N \text{ s.t. } D_i^k = \emptyset$, conclude the profile

$$(A_i^k) (2.3)$$

to be the set of action profiles survive from IESD.

ii) If
$$\exists i \in N \text{ s.t. } D_i^k \neq \emptyset$$
, define
$$\forall i \in N, \ A_i^{k+1} \leftarrow A_i^k \backslash D_i^k$$
 (2.4)

Example 2.1 (Hotelling Model of Politics). Players maximize their votes by choosing where to stand along a natural number line.

- Player $N = \{1, 2\}$
- Action set $A_i = \{1, \dots, M\}$, with $2 \not\mid M$ and M > 3.
- Payoff

$$u_{i}(a_{i}, a_{-i}) = \begin{cases} a_{i} + \frac{1}{2}(a_{-i} - a_{i} - 1) & \text{if } a_{i} < a_{-i} \\ \frac{M}{2} & \text{if } a_{i} = a_{-i} \\ M - [a_{-i} + \frac{1}{2}(a_{i} - a_{-i} - 1)] & \text{if } a_{i} > a_{-i} \end{cases}$$

$$(2.5)$$

Claim i. $a_i = 1$ is strictly dominated by $a_i = 2$.

Proof.

$$u_i(a_i = 1, a_{-i}) = \begin{cases} \frac{M}{2} & \text{if } a_{-i} = 1\\ \frac{a_{-i}}{2} & \text{if } a_{-i} > 1 \end{cases}$$
 (2.6)

$$u_{i}(a_{i} = 2, a_{-i}) = \begin{cases} M - 1 & \text{if } a_{-i} = 1\\ \frac{M}{2} & \text{if } a_{-i} = 2\\ \frac{a_{-i}}{2} + \frac{1}{2} & \text{if } a_{-i} > 2 \end{cases}$$

$$(2.7)$$

Claim ii. $\lfloor \frac{n}{2} \rfloor + 1$ is the only action survives.

Proof. Similarly, we can eliminate all edge-values iteratively.

Definition 2.2. For each $i \in N$, the **best-response function** of this player is a correspondence $Br_i : A_{-i} \rightrightarrows A_i$ defined as

$$Br_i(a_{-i}) := \{ a_i \in A_i : u_i(a_i, a_{-i}) \ge u_i(a_i', a_{-i}) \ \forall a_i' \in A_i \}$$
(2.8)

Definition 2.3. A **belief** of player i (about the actions of the other players) is a <u>probability</u> measure, $\alpha_i \in \Delta(A_{-i})$.

Definition 2.4. a_i is a **best response** to the belief α_i if

$$\forall a_i' \in A_i, \ \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \alpha_i(a_{-i}) \ge \sum_{a_{-i} \in A_{-i}} u_i(a_i', a_{-i}) \alpha_i(a_{-i})$$
(2.9)

more precisely,

$$\forall a_i' \in A_i, \ \mathbb{E}[u_i(a_i, a_{-i}) | \alpha_i] \ge \mathbb{E}[u_i(a_i', a_{-i}) | \alpha_i]$$

$$(2.10)$$

Definition 2.5. $a_i \in A_i$ is a **never best response** if it is not a best response given any belief α_i .

Corollary 2.1 (Iterative Elimination of Never Best Response). Same procedures but D_i^k is the set of never best responses for player i at game G^k .

Example 2.2. For player 1, D is not strictly dominated, but it is a never best response.

Proof. Let α be a probability measure on $\{L, R\}$ such that $\alpha(L) = p \in [0, 1]$.

$$\mathbb{E}[u_1(U)|\alpha] = 10p \tag{2.11}$$

$$\mathbb{E}[u_1(M)|\alpha] = 10 - 10p \tag{2.12}$$

$$\mathbb{E}[u_1(D)|\alpha] = 1 \tag{2.13}$$

Case i

$$p \ge 0.5 \implies \mathbb{E}[u_1(U)|\alpha] \ge 5 > \mathbb{E}[u_1(D)|\alpha]$$
 (2.14)

Case ii

$$p < 0.5 \implies \mathbb{E}[u_1(M)|\alpha] > 5 > \mathbb{E}[u_1(D)|\alpha] \tag{2.15}$$

Therefore, for any belief α , D cannot be a best response. So D is a never best response.

Definition 2.6. An action $a_i \in A_i$ is **rationalizable** if it survives iterative elimination of never best responses (i.e. action a_i is a best response to some mixed strategies of opponents).

Lemma 2.1 (60.1). An action of a player in a *finite* strategic game is a never-best response *if and only if* it is strictly dominated. ¹

¹The concept of *strict domination* will be introduced in section 5.

3 Lecture 3. Jan. 21 2019 Pure Strategy Nash Equilibrium

Definition 3.1. A pure strategy Nash equilibrium is an action profile (a_i) such that

$$\forall i \in N, \ a_i \in Br_i(a_{-i}) \tag{3.1}$$

Remark 3.1. That's, a NE is a situation that if player i knows what other players do, there exists no beneficial deviation from the current strategy.

Remark 3.2 (Interpretations). A Nash equilibrium is

- i) An action profile induces a stable outcome,
- ii) A creditable agreement, such that no player has incentive to break the agreement.

Example 3.1 (Cournot Duopoly). Consider a game with

- i) Player $N = \{1, 2\}$
- ii) Action set $A_i = [0, \infty) \ \forall i \in N$

And revenue R_i defined by

$$R_i = p_i q_i \tag{3.2}$$

where price is linear in quantity supplied,

$$p_i = \alpha - (q_i + q_{-i}), \ \alpha \in \mathbb{R}$$
 (3.3)

and firms face fixed cost $c \in \mathbb{R}$, with the assumption that $\alpha > c$. So the cost function is given by

$$C_i(q_i) = cq_i (3.4)$$

The profit function is given by

$$\Pi_i(q_i, q_{-i}) = (\alpha - (q_i + q_{-i}) - c)q_i \tag{3.5}$$

$$= (\alpha - c - q_{-i})q_i - q_i^2 \tag{3.6}$$

Given $q_{-i} \in A_{-i}$, the best response is given by

$$Br_i(q_{-i}) = \operatorname{argmax}_{q_i \in [0,\infty)} \Pi_i(q_i, q_{-i})$$
(3.7)

$$= \max\{0, \frac{\alpha - c - q_{-i}}{2}\} \ \forall i \in N$$

$$(3.8)$$

Considering the case that both players are producing positive quantities, we can solve q_i^* by

$$Br_i \circ Br_{-i}(q_i) = \frac{\alpha - c - \frac{\alpha - c - q_i}{2}}{2} = q_i$$
(3.9)

$$\implies 2q_i - \frac{q_i}{2} = \frac{\alpha - c}{2} \tag{3.10}$$

$$\implies q_i^* = \frac{\alpha - c}{3} \tag{3.11}$$

Remark 3.3. If fixed cost presents, even if the game is symmetric, the Nash equilibrium could be asymmetric. (e.g. one firm is out of market and the other firm produces the monopoly amount)

Remark 3.4. Nash equilibrium induces an *individual level optimality* instead of the common wealth optimality.

Example 3.2 (Continue Cournot Duopoly). Note that in the Cournot Duopoly game, for each $i \in N$, the Nash equilibrium profit is

$$\Pi_{NE}^* = \left(\alpha - c - \frac{2(\alpha - c)}{3}\right) \frac{\alpha - c}{3} \tag{3.12}$$

$$=\frac{(\alpha-c)^2}{9}\tag{3.13}$$

So the total profit for the two firms is $\frac{2(\alpha-c)^2}{9}$.

Now considering if the two firms form a Cartel, the aggregate quantity produced is

$$Q^* = \operatorname{argmax}_{Q \in \mathbb{R}_{>0}} (\alpha - c - Q)Q \tag{3.14}$$

$$=\frac{\alpha-c}{2}\tag{3.15}$$

$$\implies \Pi_{Cartel}^* = \left(\alpha - c - \frac{\alpha - c}{2}\right) \frac{\alpha - c}{2} \tag{3.16}$$

$$=\frac{(\alpha-c)^2}{4} > \frac{2(\alpha-c)^2}{9} \tag{3.17}$$

The fact $\Pi_{Cartel}^* > 2 \times \Pi_{NE}^*$ suggests the Nash equilibrium action profile did not induce the optimal common wealth outcome. However, the Cartel action profile is not a stable outcome since every player has incentive to increase their production level.

Example 3.3 (Prisoner's Dilemma). The Nash equilibrium (Confess, Confess) is <u>not</u> the best outcome for the two players as a group. The optimal action profile for a group is (Silent, Silent).

Proposition 3.1. No strategy that is eliminated during iterated elimination of *never best response* can be played in a Nash equilibrium.

4 Lecture 4. Jan. 28. 2019 Nash Equilibrium: Examples

Example 4.1 (From last lecture). Consider the payoff matrix Both (A, A) and (B, B) are Nash

$$\begin{array}{c|cccc} & A & B \\ \hline A & (1,1) & (0,0) \\ B & (0,0) & (0,0) \\ \end{array}$$

equilibria. But in the former NE, for each $i \in N$,

$$Br_i(a_{-i} = A) = \{A\}$$
 (4.1)

which is a singleton.

In the second NE, for each $i \in N$,

$$Br_i(a_{-i} = B) = \{A, B\}$$
 (4.2)

We call Nash equilibria of the first type *strict Nash equilibria* and the later one *weak Nash equilibria*. More formal definitions of these two types of Nash equilibria are given below.

Definition 4.1. A strict Nash equilibrium is an action profile (a_i) such that

$$\forall i \in N, \ Br_i(a_{-i}) \text{ is a singleton.}$$
 (4.3)

Definition 4.2. A weak Nash equilibrium is a Nash equilibrium that is not strict.

Example 4.2 (Cournot with n firms). Consider the game

$$\langle N, (\mathbb{R}_{>0}), (\pi_i) \rangle \tag{4.4}$$

Where |N| = n and each firm picks a quantity $q_i \in A_i := \mathbb{R}_{\geq 0}$. Define

$$Q := \sum_{j} q_{j} \tag{4.5}$$

For each $i \in N$, define

$$Q_{-i} := \sum_{j \neq i} q_j \tag{4.6}$$

And the market has linear demand curve

$$P(\lbrace q_i \rbrace_{i \in N}) = \alpha - \sum_j q_j = \alpha - Q \tag{4.7}$$

And firms face fixed cost

$$\forall i \in N, \ C(q_i) = cq_i \text{ where } 0 < c < \alpha$$
 (4.8)

The profit function is

$$\forall i \in N, \ \pi_i(q_i, Q_{-i}) = (\alpha - c - Q_{-i})q_i - q_i^2 \tag{4.9}$$

Question What are the Nash equilibria in this environment? For each firm $i \in N$, the best response correspondence is

$$Br_i(Q_{-i}) = \max\{0, \underset{q_i \in \mathbb{R}_{\geq 0}}{\operatorname{argmax}} \pi_i(q_i, Q_{-i})\}$$
 (4.10)

$$= \max\{0, \frac{\alpha - c - Q_{-i}}{2}\}\tag{4.11}$$

Assume We have a symmetric Nash equilibrium,

$$\forall i \in N, \ q_i^* = q^* = \frac{Q}{n}$$
 (4.12)

$$\implies q^* = Br_i(\frac{n-1}{n}Q) \tag{4.13}$$

$$\implies 2q^* = \alpha - c - (n-1)q^* \tag{4.14}$$

$$\implies q^* = \frac{\alpha - c}{n+1} \tag{4.15}$$

Check Then check the validity of symmetric Nash equilibrium by asserting for every player, if all other players are playing the action suggested by the symmetric Nash equilibrium, then this player should also play it. That's

$$q^* = Br_i(\{q_i\}_{i \neq i}) \tag{4.16}$$

$$= \frac{\alpha - c}{2} - \frac{1}{2} \frac{n - 1}{n + 1} (\alpha - c) \tag{4.17}$$

$$= \frac{1}{2}((\alpha - c) - \frac{n-1}{n+1}(\alpha - c)) \tag{4.18}$$

$$= \frac{1}{2} \frac{2}{n+1} (\alpha - c) \tag{4.19}$$

$$=\frac{\alpha-c}{n+1}=q^*\tag{4.20}$$

Uniqueness We are going to show the symmetric Nash equilibrium is the only possible equilibrium action profile.

Proof. Suppose there exists some non-symmetric Nash equilibrium.

For concreteness, assuming there exists $\epsilon > 0$ such that

$$\exists i, j \in N, \ q_i = q_j + \epsilon \tag{4.21}$$

Define $Q_{-ij} := Q - q_i - q_j$. For firm i,

$$q_i = Br_i(Q_{-i}) = \frac{1}{2}(\alpha - c - Q_{-i})$$
(4.22)

$$= \frac{1}{2}(\alpha - c - Q_{-ij} - q_j) \tag{4.23}$$

$$=\frac{1}{2}(\alpha - c - Q_{-ij} - q_i + \epsilon) \tag{4.24}$$

$$\implies 3q_i = \alpha - c - Q_{-ij} + \epsilon \tag{4.25}$$

$$\implies 3q_i = \alpha - c - Q_{-j} + q_i + \epsilon \tag{4.26}$$

$$\implies 2q_i - \epsilon = \alpha - c - Q_{-i} \tag{4.27}$$

$$\implies q_i - \frac{\epsilon}{2} = Br(Q_{-j}) = q_j \tag{4.28}$$

$$= q_i - \epsilon \tag{4.29}$$

$$\implies \epsilon = 2\epsilon$$
 (4.30)

which contradicts the assumption that $\epsilon > 0$.

Therefore we conclude that the symmetric Nash equilibrium is the only Nash equilibrium in this environment.

Example 4.3 (Cournot duopoly with fixed cost). Consider the game

$$\mathcal{G} = \langle N = (1, 2), (A_i = \mathbb{R}_{>0}), (\pi_i) \rangle \tag{4.31}$$

Where the cost function is defined as

$$C_i(q_i) = \begin{cases} cq_i + f & \text{if } q_i > 0\\ 0 & \text{if } q_i = 0 \end{cases}$$
 (4.32)

where $\alpha > c > 0$.

For each firm $i \in N$, the best response function, conditioned on $q_i > 0$, is

$$q_i = Br_i(q_{-i}) = \frac{\alpha - c - q_{-i}}{2} \tag{4.33}$$

We have to **check the profit** to assert the profit is non-negative while firm i is producing above quantity. Because, otherwise, this firm could always derivate to $q_i = 0$ to avoid loss (earning zero profit).

$$\pi_i(Br_{q_{-i}}, q_{-i}) = (\alpha - q_{-i} - \frac{\alpha - c - q_{-i}}{2} - c)\frac{\alpha - c - q_{-i}}{2} - f \tag{4.34}$$

$$=\frac{(\alpha - c - q_{-i})^2}{4} - f \tag{4.35}$$

$$\pi_i(q_{-i}) \ge 0 \iff \alpha - c - q_{-i} \ge 2\sqrt{f} \tag{4.36}$$

$$\iff q_{-i} \le \alpha - c - 2\sqrt{f}$$
 (4.37)

Therefore we can modify our **best response correspondence** to

$$Br_{i}(q_{-i}) = \begin{cases} \frac{\alpha - c - q_{-i}}{2} & \text{if } q_{-i} \leq \alpha - c - 2\sqrt{f} \\ 0 & \text{if } q_{-i} \geq \alpha - c - 2\sqrt{f} \end{cases}$$
(4.38)

Monopoly: the monopoly amount, conditioned on the firm decides to produce, is $\frac{\alpha-c}{2}$. In the monopoly case, suppose the NE action profile is (the opposite case can be shown by symmetry)

$$\left(\frac{\alpha-c}{2},0\right) \tag{4.39}$$

We have to assert both

$$\begin{cases}
\frac{\alpha - c}{2} \in Br_1(0) \\
0 \in Br_2(\frac{\alpha - c}{2})
\end{cases}$$
(4.40)

$$\begin{cases}
\frac{\alpha - c}{2} \in Br_1(0) \\
0 \in Br_2(\frac{\alpha - c}{2})
\end{cases}$$
(4.40)
$$\Rightarrow \begin{cases}
0 \le \alpha - c - 2\sqrt{f} & \text{for firm 1 to produce positive amount.} \\
\frac{\alpha - c}{2} \ge \alpha - c - 2\sqrt{f} & \text{for firm 2 to produce zero.}
\end{cases}$$

$$\implies 0 \le \alpha - c - 2\sqrt{f} \le \frac{\alpha - c}{2} \tag{4.42}$$

$$\implies -(\alpha - c) \le -2\sqrt{f} \le -\frac{\alpha - c}{2} \tag{4.43}$$

$$\implies \sqrt{f} \in \left[\frac{\alpha - c}{4}, \frac{\alpha - c}{2}\right] \tag{4.44}$$

Positive symmetric equilibrium: we've proven, in the general case, it's impossible for both firms to produce positive but different amounts. Therefore we have to assert

$$\frac{\alpha - c}{3} \in Br_i(\frac{\alpha - c}{3}) \ \forall i \in N$$
 (4.45)

$$\implies \frac{\alpha - c}{3} \le \alpha - c - 2\sqrt{f} \tag{4.46}$$

$$\implies \sqrt{f} \le \frac{1}{3}(\alpha - c) \tag{4.47}$$

Zero symmetric equilibrium: we have to assert

$$0 \in Br_i(0) \ \forall i \in N \tag{4.48}$$

$$\implies 0 \ge \alpha - c - 2\sqrt{f} \tag{4.49}$$

$$\implies \sqrt{f} \ge \frac{\alpha - c}{2} \tag{4.50}$$

Example 4.4 (Discrete price Bertrand duopoly). Consider the following game

$$\mathcal{G} = \langle N = \{1, 2\}, (A_i = \{k\epsilon : k \in \mathbb{Z}_{>0}\}), (\pi_i) \rangle \tag{4.51}$$

The profit function can be derived as

$$\pi_{i}(p_{i}, p_{-i}) = \begin{cases} (\alpha - p_{i})(p_{i} - c) & \text{if } p_{i} < p_{-i} \\ \frac{\alpha - p_{i}}{2}(p_{i} - c) & \text{if } p_{i} = p_{-i} \\ 0 & \text{if } p_{i} > p_{-i} \end{cases}$$

$$(4.52)$$

Claim the only Nash equilibria are (c, c) and $(c + \epsilon, c + \epsilon)$.

Justify (c,c), consider any firm $i \in N$, currently $\pi_i = 0$

$$\uparrow p_i \implies \pi_i \leftarrow \pi_i' = 0 \; \mathsf{X} \tag{4.53}$$

$$\downarrow p_i \implies \pi_i \leftarrow \pi_i' < 0 \; \mathsf{X} \tag{4.54}$$

So no firm has incentive to derivate from this action profile, so (c,c) is a Nash equilibrium by definition

Consider the action profile $(c + \epsilon, c + \epsilon)$, both firms are earning a positive profit $\pi_i > 0$. For any firm $i \in N$,

$$\uparrow p_i \implies \pi_i \leftarrow \pi_i' = 0 \ \mathsf{X} \tag{4.55}$$

$$\downarrow p_i \implies \pi_i \leftarrow \pi_i' = 0 \ \mathsf{X} \tag{4.56}$$

So $(c + \epsilon, c + \epsilon)$ is a Nash equilibrium by definition.

No other Nash equilibrium

Claim: (p_1, p_2) cannot be a Nash equilibrium if any $p_i < c$.

Obviously, set $p_i < c$ would induce negative profit and firm i do better off by setting $p_i \leftarrow c$.

Claim: the symmetric profit (p, p) with $p > c + \epsilon$ cannot be a Nash equilibrium. For both firms, the current profit is

$$\pi_1 = \pi_2 = \frac{1}{2}(\alpha - c - k\epsilon)k\epsilon \tag{4.57}$$

And reducing price by ϵ leads to a profit of

$$\pi_i' = (\alpha - c - k\epsilon + \epsilon)(k - 1)\epsilon > \pi_i \tag{4.58}$$

so such action profile cannot be Nash equilibrium.

Claim (p_1, p_2) with $p_1 \neq p_2$ and $p_1, p_2 > c$ cannot be a Nash equilibrium. The firm charges higher price can always reduce it's price to the price charged by the other firm and gain a positive profit. Claim (c, p_2) with $p_2 \geq c + \epsilon$ cannot be a Nash equilibrium. The firm charging p = c can always increase its price to $c + \epsilon$ to earn positive profit.

Therefore there's no Nash equilibrium other than (c, c) and $(c + \epsilon, c + \epsilon)$.

Example 4.5 (Bertrand duopoly with differentiated products). Consumers are <u>uniformly distributed</u> on a preference line [0, 1].

For a firm $i \in N$, let $x_i \in [0,1]$ measure consumer's preference towards firm i's products. Define

$$x_{-i} := 1 - x_i \sim Unif(0, 1) \tag{4.59}$$

Consumer buy product i if

$$x_i - p_i \ge x_{-i} - p_{-i} \tag{4.60}$$

and purchases product 1 if

$$x_i - p_i \le x_{-i} - p_{-i} \tag{4.61}$$

Then solve the boundary $x^* \in [0,1]$ such that consumers at x^* are indifferent between two products.

$$x + p_i = (1 - x) + p_{-i} (4.62)$$

$$\implies x^* = \frac{1 - p_i + p_{-i}}{2} \tag{4.63}$$

So any consumer with $x_i > x^*$ would choose firm i's product, also because consumers are uniformly distributed on $x_i \in [0, 1]$. So the portion of consumers buying firm i's product is

$$1 - x_i = \frac{1 + p_i - p_{-i}}{2} \tag{4.64}$$

And, clearly, The demand function for firm i is

$$D_{i}(q_{i}, q_{-i}) = \begin{cases} 0 & \text{if } p_{i} \ge p_{-i} + 1\\ \frac{1+p_{i}-p_{-i}}{2} & \text{if } p_{i} \in [p_{-i} - 1, p_{-i} + 1]\\ 1 & \text{if } p_{i} \le p_{-i} - 1 \end{cases}$$

$$(4.65)$$

Consider the case where $|p_i - p_{-i}| \leq 1$,

$$\pi_i = D_i(p_i)(p_i - c) = \frac{1 + p_i - p_{-i}}{2}(p_i - c)$$
(4.66)

Take the first order condition

$$\frac{\partial \pi_i}{\partial p_i} = 0 \tag{4.67}$$

$$\implies p_i = \frac{p_{-i} + c - 1}{2} \tag{4.68}$$

5 Lecture 5 Feb. 7 2019 Mixed Strategies

5.1 Lecture Contents

Assumption 5.1. We assume that the preference relation of each player i satisfies the assumptions of von Neumann and Morgenstern, so that it can be represented by the expected value of some utility function $u_i: A \to \mathbb{R}$.

Definition 5.1. Suppose player i has a *finite* set of pure strategies A_i , then a **mixed strategy** $\sigma_i \in \Delta(A_i)$ is a probability distribution over A_i .

Definition 5.2. The support of σ_i is defined as

$$S(\sigma_i) := \{ a \in A_i : \sigma_i(a) > 0 \}$$

$$(5.1)$$

Definition 5.3. A mixed strategy σ_i is degenerate if

$$\exists a_i \in A_i \ s.t. \ \sigma_i(a_i) = 1 \tag{5.2}$$

and such degenerated mixed strategy is denoted as $e(a_i)$.

Remark 5.1. A pure strategy $a_i \in A_i$ is equivalent to a degenerated mixed strategy $e(a_i)$.

Proposition 5.1. In a finite game, given the independence of randomization, the probability of the action profile $a := (a_i)$ to be realized given mixed strategy profile $\sigma := (\sigma_i)$ is

$$\sigma(a) := \mathbb{P}[a|\sigma] = \prod_{i \in N} \sigma_i(a_i) \tag{5.3}$$

and for player i, the **expected payoff** from mixed strategy profile (σ_i) is

$$U_i(\sigma_i, \sigma_{-i}) := \sum_{a \in A} \underbrace{\prod_{j \in N} \sigma_j(a_j)}_{\mathbb{P}[a|\sigma]} u_i(a) = \mathbb{E}[u_i|\sigma]$$
(5.4)

Proposition 5.2. The expected payoff from mixed strategy profile $(\sigma_i) := (\sigma_i, \sigma_{-i})$ is

$$U_i(\sigma_i, \sigma_{-i}) := \mathbb{E}[u_i(a)|(\sigma_i)] = \sum_{a_i \in A_i} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma_{-i}(a_{-i}) \sigma_i(a_i)$$
 (5.5)

Definition 5.4 (Osborne 2030 lec2). The **mixed extension** of the strategic game $\langle N, (A_i)_{i \in N}, (u_i)_{i \in N} \rangle$ is the following strategic game:

- (i) **Players** N;
- (ii) Action sets $\Delta(A_i)$ for player i;
- (iii) **Preferences** represented by $U_i: \prod_{j\in N} \Delta(A_j) \to \mathbb{R}$, where $U_i(\alpha)$ is the expected value under u_i of the lottery over $\prod_{j\in N} \Delta(A_j)$ induced by mixed strategy profile $\sigma \in \Delta(A)$.

Definition 5.5. A pure strategy $a_i \in A_i$ is **strictly dominated** by a mixed strategy $\sigma_i \in \Delta(A_i)$ if

$$\forall a_{-i} \in A_{-i} \ u_i(a_i, a_{-i}) < U_i(\sigma_i, a_{-i}) \tag{5.6}$$

We say a_i is strictly dominated.

Proposition 5.3. We can also define the strict dominance by replacing $\forall a_{-i} \in A_{-i}$ with $\forall \sigma_{-i} \in \Delta A_{-i}$ since

$$\forall a_{-i} \in A_{-i} \ u_i(a_i, a_{-i}) < U_i(\sigma_i, a_{-i}) \iff \forall \sigma_{-i} \in \Delta(A_{-i}) \ U_i(a_i, \sigma_{-i}) < U_i(\sigma_i, \sigma_{-i})$$
 (5.7)

Proof. Main idea: linearity of expectation operator.

 (\Longrightarrow) Suppose

$$\forall a_{-i} \in A_{-i} \ u_i(a_i, a_{-i}) < U_i(\sigma_i, a_{-i}) \tag{5.8}$$

Let $\sigma_{-i} \in \Delta(A_{-i})$,

$$U_i(a_i, \sigma_{-i}) = \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma_{-i}(a_{-i})$$
(5.9)

$$<\sum_{a_{-i}\in A_{-i}} U_i(\sigma_i, a_{-i})\sigma_{-i}(a_{-i})$$
 (5.10)

$$=U_i(\sigma_i,\sigma_{-i})\tag{5.11}$$

Thus

$$\forall \sigma_{-i} \in \Delta(A_{-i}) \ U_i(a_i, \sigma_{-i}) < U_i(\sigma_i, \sigma_{-i})$$

$$\tag{5.12}$$

 (\Leftarrow) Suppose

$$\forall \sigma_{-i} \in \Delta(A_{-i}) \ U_i(a_i, \sigma_{-i}) < U_i(\sigma_i, \sigma_{-i}) \tag{5.13}$$

Let $a_{-i} \in A_{-i}$, consider σ_{-i} defined as $\sigma_{-i}^*(x) := \mathbb{1}\{x = a_{-i}\}$ so that

$$U_i(a_i, \sigma_{-i}^*) < U_i(\sigma_i, \sigma_{-i}^*) \tag{5.14}$$

$$\implies u_i(a_i, a_{-i}) < U_i(\sigma_i, a_{-i}) \tag{5.15}$$

The equivalence is shown.

Definition 5.6. The **best response** to a mixed strategy $\sigma_{-i} \in \Delta(A_{-i})$ is defined as

$$Br_i(\sigma_{-i}) := \{ \sigma_i \in \Delta(A_i) : \forall \tilde{\sigma}_i \in \Delta(A_i), \ U_i(\sigma_i, \sigma_{-i}) \ge U_i(\tilde{\sigma}_i, \sigma_{-i}) \}$$
 (5.16)

Proposition 5.4. Let $(\sigma_i, \sigma_{-i}) \in \Delta(A)$ then $\sigma_i \in Br_i(\sigma_{-i})$ if and only if

- 1. $\forall a_i, a_k \in \mathcal{S}(\sigma_i), \ a_i \sim_i a_k;$
- 2. and $\forall a_i \in \mathcal{S}(\sigma_i) \ a_k \notin \mathcal{S}(\sigma_i), \ a_i \succsim_i a_k$.

Proof. (\Longrightarrow) case 1: suppose (1) is false, for concreteness, assume

$$\exists a_i, a_k \in \mathcal{S}(\sigma_i) \ s.t. \ U_i(a_i, \sigma_{-i}) > U_i(a_k, \sigma_{-i})$$

$$\tag{5.17}$$

Define

$$U_j := U_i(a_j, \sigma_{-i}) = \sum_{a_{-i} \in A_{-i}} u_i(a_j, a_{-i}) \sigma_{-i}(a_{-i})$$
(5.18)

$$\mathcal{U}_k := U_i(a_k, \sigma_{-i}) = \sum_{a_{-i} \in A_{-i}} u_i(a_k, a_{-i}) \sigma_{-i}(a_{-i})$$
(5.19)

And $\mathcal{U}_j > \mathcal{U}_k$ by assumption. Consider σ'_i defined with

$$\sigma_i'(a) = \begin{cases} \sigma_i(a_j) + \sigma_i(a_k) & \text{if } a = a_j \\ 0 & \text{if } a = a_k \\ \sigma_i(a) & \text{otherwise} \end{cases}$$
 (5.20)

Therefore

$$U_i(\sigma'_i, \sigma_{-i}) := \sum_{a_i \in A_i} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma'_i(a_i) \sigma_{-i}(a_{-i})$$
(5.21)

$$= \sum_{a_i \in A_i \setminus \{a_i, a_k\}} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma'_i(a_i) \sigma_{-i}(a_{-i}) + \sigma'_i(a_j) \mathcal{U}_j + \sigma'_i(a_k) \mathcal{U}_k$$
 (5.22)

$$= \sum_{a_i \in A_i \setminus \{a_i, a_k\}} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma_i(a_i) \sigma_{-i}(a_{-i}) + \sigma_i(a_j) \mathcal{U}_j + \sigma_i(a_k) \mathcal{U}_j$$
 (5.23)

$$= \sum_{a_i \in A_i} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma_i(a_i) \sigma_{-i}(a_{-i}) \equiv U_i(\sigma_i, \sigma_{-i})$$
(5.25)

So that σ'_i strictly dominates σ_i , so $\sigma_i \notin Br_i(\sigma_{-i})$.

case 2: suppose (2) is false, for concreteness, assume

$$\exists a_j \in \mathcal{S}(\sigma_i), \ a_k \notin \mathcal{S}(\sigma_i) \ s.t. \ U_i(a_k, \sigma_{-i}) > U_i(a_j, \sigma_{-i})$$

$$(5.26)$$

Consider mixed strategy σ'_i defined as

$$\sigma'_{i}(a) = \begin{cases} 0 & \text{if } a = \sigma_{j} \\ \sigma_{i}(a_{j}) & \text{if } a = \sigma_{k} \\ \sigma_{i}(a) & \text{otherwise} \end{cases}$$

$$(5.27)$$

By previous definitions, $U_k > U_j$.

Then

$$U_i(\sigma'_i, \sigma_{-i}) := \sum_{a_i \in A_i} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma'_i(a_i) \sigma_{-i}(a_{-i})$$
(5.28)

$$= \sum_{a_i \in A_i \setminus \{a_i, a_k\}} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma'_i(a_i) \sigma_{-i}(a_{-i}) + \sigma'_i(a_j) \mathcal{U}_j + \sigma'_i(a_k) \mathcal{U}_k$$
 (5.29)

$$= \sum_{a_i \in A_i \setminus \{a_j, a_k\}} \sum_{a_{-i} \in A_{-i}} u_i(a_i, a_{-i}) \sigma_i(a_i) \sigma_{-i}(a_{-i}) + 0 \,\mathcal{U}_j + \sigma_i(a_j) \mathcal{U}_k$$
 (5.30)

$$=U_i(\sigma_i,\sigma_{-i})\tag{5.32}$$

Therefore $\sigma_i \notin Br_i(\sigma_{-i})$, modus tollens.

 (\longleftarrow) suppose $\sigma_i \notin Br_i(\sigma_{-i})$, that's

$$\exists \sigma_i' \in \Delta(A_i) \ s.t. \ U_i(\sigma_i', \sigma_{-i}) > U_i(\sigma_i, \sigma_{-i})$$

$$\tag{5.33}$$

Therefore

$$\exists \bar{a}_i, \underline{a}_i \in \mathcal{S}(\sigma_i) \ s.t. \ U_i(\bar{a}_i, \sigma_{-i}) \ge U_i(\sigma_i, \sigma_{-i}) \ge U_i(\underline{a}_i, \sigma_{-i})$$

$$(5.34)$$

$$\wedge \exists \bar{a}_i', \underline{a}_i' \in \mathcal{S}(\sigma_i') \ s.t. \ U_i(\bar{a}_i', \sigma_{-i}) \ge U_i(\sigma_i', \sigma_{-i}) \ge U_i(\underline{a}_i', \sigma_{-i})$$

$$(5.35)$$

$$\implies U_i(\bar{a}_i', \sigma_{-i}) \ge U_i(\sigma_i', \sigma_{-i}) > U_i(a_i, \sigma_{-i}) \ge U_i(\underline{a}_i, \sigma_{-i}) \tag{5.36}$$

$$\implies U_i(\bar{a}_i', \sigma_{-i}) > U_i(\underline{a}_i, \sigma_{-i}) \tag{5.37}$$

$$\implies \bar{a}_i' \succ_i \underline{a}_i \in \mathcal{S}(\sigma_i) \tag{5.38}$$

In both cases, $\overline{a}'_i \in \mathcal{S}(\sigma_i)$ and $\overline{a}'_i \notin \mathcal{S}(\sigma_i)$, the two requirements are falsified, modus tollens.

Remark 5.2. Procedure to solve Nash equilibria for two player finite games,

- (i) Solve for pure strategy Nash equilibria.
- (ii) Eliminate strictly dominated strategies.
- (iii) For player i, solve a mixed strategy scheme σ_j so that player i is indifferent between actions in a non-singleton subset of his/her action set, A_i^* .
- (iv) Player i would play any mixed strategy with support A_i^* .
- (v) Repeat (iii) and (iv) for player j.

5.2 Some Propositions and Proofs

Proposition 5.5 (exer.120.3). A mixed strategy that assigns positive probability of a <u>strictly</u> dominated action is strictly dominated. That's, suppose $\tilde{a}_i \in A_i$ is strictly dominated, then for any $\sigma_i \in \Delta(A_i)$,

$$\tilde{a}_i \in \mathcal{S}(\sigma_i) \implies \sigma_i \text{ strictly dominated}$$
 (5.39)

Note that the converse is **not** true.

Proposition 5.6 (Nash). Every strategic game in which each player has <u>finitely many actions</u> has a mixed strategy Nash equilibrium.

Proposition 5.7. Let $\sigma_{-i} \in \Delta(A_{-i})$, if a pure strategy $a_i \in A_i$ is <u>strictly dominated</u> by another pure strategy $\tilde{a}_i \in A_i$, then

$$\forall \sigma_i \in Br_i(\sigma_{-i}) \ a_i \notin \mathcal{S}(\sigma_i) \tag{5.40}$$

Proof.

Proposition 5.8. Let $\sigma_{-i} \in \Delta(A_{-i})$, if a pure strategy $a_i \in A_i$ is <u>strictly dominated</u> by another mixed strategy $\tilde{\sigma}_i \in \Delta(A_i)$, then

$$\forall \sigma_i \in Br_i(\sigma_{-i}) \ a_i \notin \mathcal{S}(\sigma_i) \tag{5.41}$$

Proof.

Proposition 5.9. For pure strategy $a_i \in A_i$, if there exists $a_{-i} \in A_{-i}$ such that

$$u_i(a_i, a_{-i}) \ge u_i(\tilde{a}_i, a_{-i}) \ \forall \tilde{a}_i \in A_i \tag{5.42}$$

then a_i is **not** strictly dominated (by pure strategies or mixed strategies).

Proposition 5.10. Let $a_i \in A_i$, then the following statements are equivalent

(i)
$$\exists \sigma_i \in \Delta(A_i) \ s.t. \ a_i \in \mathcal{S}(\sigma_i) \land \forall \sigma_{-i} \in \Delta(A_{-i}), \ U_i(a_i, \sigma_{-i}) < U_i(\sigma_i, \sigma_{-i})$$
 (5.43)

(ii)
$$\exists \tilde{\sigma}_i \in \Delta(A_i) \ s.t. \ a_i \notin \mathcal{S}(\tilde{\sigma}_i) \land \forall \sigma_{-i} \in \Delta(A_{-i}), \ U_i(a_i, \sigma_{-i}) < U_i(\tilde{\sigma}_i, \sigma_{-i})$$
 (5.44)

Proof. (\Longrightarrow) Suppose there exists σ_i satisfies (i), Note that $\sigma_i(a_i) \neq 1$, define $\tilde{\sigma}_i$ as

$$\tilde{\sigma}_i(a) = \begin{cases} \frac{\sigma_i(a)}{1 - \sigma_i(a_i)} & \text{if } a \in \mathcal{S}(\sigma_i) \setminus \{a_i\} \\ 0 & \text{otherwise} \end{cases}$$
(5.45)

And $S(\tilde{\sigma}_i) = S(\sigma_i) \setminus \{a_i\}.$ Then, by (i),

$$U_i(a_i, \sigma_{-i}) < \sum_{a_i \in \mathcal{S}(\sigma_i)} U_i(a_j, \sigma_{-i})\sigma_i(a_j)$$
(5.46)

$$\implies (1 - \sigma_i(a_i))U_i(a_i, \sigma_{-i}) < \sum_{a_j \in \mathcal{S}(\sigma_i) \setminus \{a_i\}} U_i(a_j, \sigma_{-i})\sigma_i(a_j)$$
(5.47)

$$\implies U_i(a_i, \sigma_{-i}) < \frac{1}{1 - \sigma_i(a_i)} \sum_{a_j \in \mathcal{S}(\tilde{\sigma}_i)} U_i(a_j, \sigma_{-i}) \sigma_i(a_j)$$
 (5.48)

$$\implies U_i(a_i, \sigma_{-i}) < \sum_{a_j \in \mathcal{S}(\tilde{\sigma}_i)} U_i(a_j, \sigma_{-i}) \frac{\sigma_i(a_j)}{1 - \sigma_i(a_i)}$$

$$(5.49)$$

$$\implies U_i(a_i, \sigma_{-i}) < \sum_{a_j \in \mathcal{S}(\tilde{\sigma}_i)} U_i(a_j, \sigma_{-i}) \tilde{\sigma}_i(a_j) := U_i(\tilde{\sigma}_i, \sigma_{-i})$$
(5.50)

(\iff) Suppose there exists $\tilde{\sigma}_i$ satisfies (ii). Take $\eta \in \mathbb{R}_+$, define σ_i as

$$\sigma_i(a) = \begin{cases} \frac{\tilde{\sigma}_i(a)}{1+\eta} & \text{if } a \in \mathcal{S}(\tilde{\sigma}_i) \\ \frac{\eta}{1+\eta} & \text{if } a = a_i \\ 0 & \text{otherwise} \end{cases}$$
 (5.51)

Note that $S(\sigma_i) = S(\tilde{\sigma}_i) \cup \{a_i\}$. By (ii),

$$U_i(a_i, \sigma_{-i}) < \sum_{a_i \in \mathcal{S}(\tilde{\sigma}_i)} U_i(a_j, \sigma_{-i})\tilde{\sigma}_i(a_j)$$

$$(5.52)$$

$$\implies (1+\eta)U_i(a_i,\sigma_{-i}) < \sum_{a_j \in \mathcal{S}(\tilde{\sigma}_i)} U_i(a_j,\sigma_{-i})\tilde{\sigma}_i(a_j) + U_i(a_i,\sigma_{-i})\eta$$
 (5.53)

$$\implies U_i(a_i, \sigma_{-i}) < \sum_{a_j \in \mathcal{S}(\tilde{\sigma}_i)} U_i(a_j, \sigma_{-i}) \frac{\tilde{\sigma}_i(a_j)}{1+\eta} + \frac{\eta}{1+\eta} U_i(a_i, \sigma_{-i})$$
 (5.54)

$$= \sum_{a_i \in \mathcal{S}(\sigma_i)} U_i(a_j, \sigma_{-i}\sigma_i(a_j)) := U_i(\sigma_i, \sigma_{-i})$$

$$(5.55)$$

Remark 5.3. With above proposition, while we are checking if $a_i \in A_i$ is strictly dominated by some mixed strategies, we don't have to include a_i in the support of potential σ_i .

Proposition 5.11 (Osborne 2030 lec2). Let $\mathcal{G} = \langle N, (A_i), (u_i) \rangle$ and $\mathcal{G}' = \langle N, (\Delta(A_i)), (U_i) \rangle$. Then any Nash equilibrium of \mathcal{G} is a mixed strategy Nash equilibrium (in which each player's strategy is pure) of \mathcal{G}' .

Remark 5.4. Hence mixed strategy may do as well as a pure strategy but can never do strictly better than all pure strategy. Consequently pure strategy remains optimal when mixed strategies are allowed.

Proposition 5.12 (Osborne 2030 lec2). For any mixed strategy Nash equilibrium of \mathcal{G}' in which each player's strategy is pure is a Nash equilibrium.

Remark 5.5. If player optimally chooses a pure strategy when she is allowed to randomize, then when she is prohibited from randomizing the pure strategy remains optimal.

Proposition 5.13 (Osborne 2030 lec2, lemma 33.2). The following statements are equivalent:

- (i) (σ_i^*) is a mixed strategy Nash equilibrium.
- (ii) $\forall i \in N, \ \sigma_i^* \in Br_i(\sigma_{-i}^*).$
- (iii) $\forall i \in N$, every action in support of σ_i^* is a best response to σ_{-i}^* .

6 Lecture 6 Feb. 25 2019 Extensive Form Games

6.1 An Introductory Example




Figure 6.1: Market Entry game

6.2 Extensive Form Game: Assumptions and Terminologies

Assumption 6.1. All games considered in the section are **pure sequential games**, in which, for each sub-history, *exactly* one player gets action.

Assumption 6.2. We assume all players have **perfect information**, they can observe all previous actions. ¹

Definition 6.1. A sequence of actions is called a **history**. For example, $h = (a_1, \ldots, a_t)$ would be a history at period t + 1.

Definition 6.2. A history h belongs to **terminal histories**, T, if the game is over and payoff is generated. That's, there's no action a such that appending a to h would generate another valid history.

Definition 6.3. The sub-histories, S, are histories that are not terminal.

¹This assumption will be relaxed in Bayesian games.

Definition 6.4. Conventionally, the initial history is defined as the empty set.

$$h_0 = \varnothing \in S \tag{6.1}$$

Definition 6.5. Let $h \in S$, then $\tilde{h} := h \oplus a \in S \cup T$ is a **continuation** of $h = (a_1, \dots, a_t)$.

Definition 6.6 (Extensive Form Game). An extensive form game $\Gamma = \langle N, P, H, (u_i) \rangle$ consists

- (i) A finite set of players N;
- (ii) A set of histories $H = S \cup T$ with initial history $h_0 = \emptyset^1$;
- (iii) Player function $P: S \to N$ assigns one player P(h) (in pure sequential games) to act after history h. And an action set A(h) for player P(h) is defined by

$$A(h) := \{ a \in A_{P(h)} : h \oplus \{a\} \in H \}$$
(6.2)

(iv) **Payoff functions** $u_i: T \to \mathbb{R}$ derived from preferences by

$$\forall z_1, z_2 \in T, \ u_i(z_1) \ge u_i(z_2) \iff z_1 \succsim_i z_2 \tag{6.3}$$

Definition 6.7. A strategy s_i for player $i \in N$ is a mapping from $\{h \in S : P(h) = i\}$ to valid actions in A_i for each specific h.

Remark 6.1. A strategy is a <u>complete contingency plan</u> assigns an action to every subhistories such that P(h) = i, even if those subhistories cannot be reached.

Definition 6.8. A **Nash equilibrium** is a <u>strategy profile</u> such that each player's strategy is a best response to strategies of other players.

Definition 6.9. For each $h \in S$, a **subgame** following subhistory h is a 4-tuple $\langle N, P, H', (u_i) \rangle$ where H' is the set of continuations of h. And everything else are inherited from the original game.

Definition 6.10. A subgame perfect Nash equilibrium for Γ is a strategy profile such that there's a Nash equilibrium in every subgame of Γ , including those never reached. This means players are best responding in every subgame.

Lemma 6.1.

$$(s_i) \in \text{SPNE} \implies (s_i) \in \text{NE}$$
 (6.4)

Proof. By the definition of SPNE, take subgame after history $h = \emptyset$.

Proposition 6.1. The set of strategy profiles found by the procedure of backward induction in a finite horizon game is the set of subgame perfect equilibria.

Definition 6.11. An extensive form game Γ is finite if the set H is finite.

Theorem 6.1. Every <u>finite</u> extensive game with perfect information has a subgame perfect equilibrium.

 $[\]overline{^{1}S}$ and T are subhistories and terminal histories sets defined above.

7 Lecture 7 Mar.4 2019 Examples of Extensive Form Games (Omitted)

8 Lecture 8 Mar.11 2019 Repeated Games

8.1 Finitely Repeated Games

Remark 8.1. Comparing extensive form games and repeated games.

- (i) **Extensive Form Game** the structure of game (available action, player, etc.)at each node of history tree varies based on the previous actions (history).
- (ii) Repeated Game Same set of players interact repeatedly, each player can condition her action in period t on other players' actions in periods $\{1, \ldots, t-1\}$.

Theorem 8.1. If a finite repeated game has a unique Nash equilibrium in the game played in each period, then the unique SPNE will have the players playing the NE in each period.

Proof. Backwards induction.

8.2 Infinitely Repeated Games

Definition 8.1 (Infinitely Repeated Games). Let $G = \langle N, (A_i), (\succeq_i) \rangle$ be a strategic game, denote the set of action profile as $A = \times_{i \in N} A_i$. An **infinitely repeated game** of G is an *extensive form* $game \ \Gamma = \langle N, H, P, (\succeq_i^*) \rangle$ where

- $H = \{\varnothing\} \cup (\cup_{t=1}^{\infty} A^t) \cup A^{\infty}.^1$
- $P(h) = N \ \forall h \in H$.
- \succsim_i^* is a preference relation on A^{∞} induced by \succsim_i such that if $(a^t) \in A^{\infty}$, $a, a' \in A$, and $a \succsim_i a'$ then

$$\forall t \in \mathbb{N}, \ (a^1, \dots, a^{t-1}, a, a^{t+1}, \dots) \succsim_i^* (a^1, \dots, a^{t-1}, a', a^{t+1}, \dots)$$
(8.1)

Example 8.1 (Infinitely Repeated Prisoner's Dilemma). Consider the infinitely repeated game induced by the prisoner's dilemma, where C and D denote cooperating and deviating. Suppose y > x > 1. Define payoff in period $t \in \mathbb{N}$ as $u_i(a_i^t, a_{-i}^t)$ for each player $i \in \mathbb{N}$. And both player face

$$\begin{array}{c|c} C & D \\ C & x, x & y, 0 \\ D & y, 0 & 1, 1 \end{array}$$

a time discount factor $\delta \in (0,1)$ measuring how patient they are. For example, the history $h = (CC, CC, \ldots)$ gives each player payoff

$$\sum_{t=1}^{\infty} \delta^{t-1} u_i^t(C, C) = \sum_{t=1}^{\infty} \delta^{t-1} x = \frac{x}{1-\delta}$$
 (8.2)

 $^{{}^{1}}A^{\infty}$ denotes the set of infinite sequences $(a^{t})_{t=1}^{\infty}$ where every $a^{t} \in A$.

Definition 8.2. The **preference** in infinitely repeated games can be defined using a **present** discounted utility function. Let u_i be the utility function induced by (\succeq_i) in game G, then the payoff on history $h = (a^t)_{t=1}^{\infty}$ in the infinitely repeated game generated by G can be defined as

$$U(h) := \sum_{t=1}^{\infty} \delta^{t-1} u(a^t) \text{ for some } \delta \in (0,1)$$

$$(8.3)$$

Lemma 8.1. Two strategic game generate same preferences with discounting in repeated game \iff each player's payoffs in one game are affine transformation of her payoffs in the other game.

Definition 8.3. A strategy can be represented as an automaton consisting of

- (i) A set Q_i of states;
- (ii) An initial state $q_i^0 \in Q_i$;
- (iii) An **output function** $f_i: Q_i \to A_i$;
- (iv) A transition function $\tau_i: Q_i \times A \to Q_i$.

Example 8.2 (Grim Trigger, ∞ -Period Punishment). Grim Trigger in infinitely repeated prisoner dilemma is a strategy defined as

$$s_i^*(\varnothing) = C \tag{8.4}$$

$$s_i^*(h) = \begin{cases} C & \text{if } a_{-i}^{\tau} = C \ \forall \tau \in \{1, \dots, |h|\} \\ D & otherwise \end{cases}$$
 (8.5)

Whether (s_i^*) is a SPNE depends on the parameter set of the game, here $\theta = (x, y, \delta)$.

Proposition 8.1. Grim Trigger is a SPNE in an infinitely repeated Prisoner's Dilemma if and only if

$$\delta \ge \frac{y-x}{y-1} \tag{8.6}$$

Proof. To show (s_i^*) is a SPNE, we will examine all it's alternatives and argue (s_i^*) provides at least much payoff as the supremum of payoffs generated by its alternatives.

Let s_i' be an alternative strategy for player $i \in N$ such that $s_i' \neq s_i$.

Let $k \in \mathbb{N}$ denote the first period D is played in strategy s'_i , let S^k_i denote the set of strategy such that D is played first at period k:

$$S_i^k := \{ s_i \in A_i^\infty : s_i^k = D \land s_i^\tau = C \ \forall \tau < k \}$$

$$(8.7)$$

It's obvious that strategy $\tilde{s}_{i,k} \in \mathcal{S}_i^k$ defined as

$$\tilde{s}_{i,k}^{\tau} = C \mathbb{1}\{\tau < k\} + D \mathbb{1}\{\tau \ge k\} \tag{8.8}$$

satisfies

$$\sum_{t=1}^{\infty} \delta^{t-1} u_i(\tilde{s}_{i,k}, s_{-i}^*) = \max_{\varphi \in \mathcal{S}_i^k} \{ \sum_{t=1}^{\infty} \delta^{t-1} u_i(\varphi, s_{-i}^*) \}$$
 (8.9)

Since k is arbitrary, to show the payoff to player i of playing strategy s_i^* given her opponent's playing s_{-i}^* is bounded below by elements in S_i^k , we only need to show it s_i^* provides at least payoff

as $\tilde{s}_{i,k}$.

Easy to show that, for arbitrary $k \in \mathbb{N}$,

$$U(s_i^*, s_{-i}^*) \ge \mathcal{U}(\tilde{s}_{i,k}, s_{-i}^*) \tag{8.10}$$

$$\iff \sum_{t=1}^{\infty} \delta^{t-1} x \ge \sum_{t=1}^{k-1} \delta^{t-1} x + \delta^{k-1} y + \sum_{t=k+1}^{\infty} \delta^{t-1} 1$$
 (8.11)

$$\iff \delta^{k-1}x + \sum_{t=k+1}^{\infty} \delta^{t-1}x \ge \delta^{k-1}y + \sum_{t=k+1}^{\infty} \delta^{t-1}1$$
(8.12)

$$\iff \delta^{k-1}x + \frac{\delta^k x}{1-\delta} \ge \delta^{k-1}y + \frac{\delta^k}{1-\delta} \tag{8.13}$$

$$\iff x + \frac{\delta x}{1 - \delta} \ge y + \frac{\delta}{1 - \delta}$$
 (8.14)

$$\iff \delta \ge \frac{y-x}{y-1} \tag{8.15}$$

Note that each subgame after period k is another infinitely repeated game where payoff of every play is transformed by multiplying δ^k . By Lemma 8.1, we can use above argument to prove the subgame perfection of Grim trigger strategy when $\delta \geq \frac{y-x}{y-1}$.

Example 8.3 (Tit for Tat, 1-Period Punishment). Tit for tat strategy is defined as

$$s_i^*(\varnothing) = C \tag{8.16}$$

$$s_i^*(h) = \begin{cases} C & \text{if } h[-1] = C \\ D & \text{if } h[-1] = D \end{cases}$$
 (8.17)




Figure 8.1: Three period punishment strategy as an automaton.

Proposition 8.2 (Generalized Limited Punishment, p-Period Punishment). The p-period punishment strategy is defined with $s_i^*(\varnothing) = C$, and if at time period k, $a_{-i}^k = D$, then the player play D for the next p periods as punishment no matter what action the opponent plays during the punishment. And then the player plays C again beginning at time period t = k + p + 1.

Proof. For the deviation not to be profitable, we need

$$\delta^{k-1}y + \sum_{t=k+1}^{k+p} \delta^{t-1}1 \le \delta^{k-1}x + \sum_{t=k+1}^{k+p} \delta^{t-1}x$$
(8.18)

$$\iff \delta^{k-1} y \le \delta^{k-1} x + \frac{(x-1)\delta^k (1-\delta^p)}{1-\delta} \tag{8.19}$$

$$\iff (y-x) \le +\frac{\delta(1-\delta^p)}{1-\delta}(x-1) \tag{8.20}$$

Proposition 8.3. For any value of $k \geq 1$, strategy pair in which each player punishes other for k periods in event of deviation is Nash equilibrium of infinitely repeated game if δ is close enough to 1. The large k is, the smaller lower bound on δ , that's, the mutually desirable outcome (C, C) is sustained by short punishment only if players are relatively patient.

9 Lecture 9 Mar. 18 2019 Bayesian Games

9.1 Introduction

Remark 9.1. Bayesian games allow players to face uncertainty about other players' preferences.

Definition 9.1 (Informal Definition). A **Bayesian game**, G^* , consists a set of players, N, and action set A_i for each player $i \in N$. Also a set of states of world, Ω . Each player has a **prior belief** p_i , which is a probability distribution on Ω . In each play of G^* , one state $\omega \in \Omega$ is realized, and each player i receive a **signal** from his **signal function** $t_i = \tau_i(\omega) \in T_i$, this is referred as the **type** of player i. T_i is the set of all possible types of player i. With $\tau_i(\omega)$ received (player i does not actually observe ω), player i deduces that the actual state realized, ω , is in the set $\tau_i^{-1}(t_i)$. This deduction provides player i's **posterior belief** distribution on Ω . For each $\omega \in \Omega$, the posterior belief of it is the probability of ω , conditioned on $\tau_i(\omega) = t_i$.

$$p_i(\omega|t_i) := \begin{cases} \frac{p_i(\omega)}{p_i(\tau_i^{-1}(t_i))} & \text{if } \omega \in \tau_i^{-1}(t_i) \\ 0 & \text{otherwise} \end{cases}$$
(9.1)

For the conditional probability to make sense, we assume $p_i(\tau_i^{-1}(t_i)) > 0 \ \forall t_i \in T_i$. Since a player in G^* may be uncertain about which pair (a, ω) will be realized, so players' preference relations are defined over *lottery* on $A \times \Omega$.

Definition 9.2 (Informal Definition). A **Bayesian equilibrium** is a strategy profile where each type of each player $(\prod_{i \in N} T_i)$ is best responding given their *posterior beliefs*.

Remark 9.2 (Complication on Belief System). Here we assume individuals have exogenous beliefs. In more complicated models, beliefs are endogenous, and beliefs change based on *other players* (previous) actions, and beliefs can be updated iteratively in multi-period games.

9.2 Formal Definitions

Definition 9.3 (Bayesian Game). A **Bayesian Game**, $G^* = \langle N, \Omega, (A_i), (T_i), (\tau_i), (\tau_i), (t_i), (t_i) \rangle$, consists of

- (i) A finite **player** set N;
- (ii) A set of states Ω . Each $\omega \in \Omega$ represents one realization determines players' types, $\underline{\Omega}$ is assumed to be known by all players;
- (iii) For each player $i \in N$:
 - (a) An **action** set A_i ;
 - (b) a set of types (signals) T_i ;
 - (c) a signal function $\tau_i: \Omega \to T_i$ that associates a type(signal) with each state;

- (d) A probability measure p_i on Ω with $p_i(\tau_i^{-1}(t_i)) > 0 \ \forall t_i \in T_i$, called player i's **prior belief**:
- (e) For each $t_i \in T_i$ we can derive a **posterior belief** (distribution conditioned on $t_i = t_i$) function $\pi_i(\omega|t_i)$ or $\pi_i(t_{-i}|t_i)$, which can be expressed as a probability distribution on Ω ;
- (f) A **preference relation** over probability distribution over $A \times \Omega$. And it's corresponding **payoff** $u_i(a_i, a_{-i}, t_i, t_{-i})$, and here we may assume u_i does *not* depend on t_i .

Remark 9.3. The above definition allows the players to have different *prior beliefs*. These beliefs may be related; *commonly they are identical, coincident with an objective measure*.

Definition 9.4 (Strategy). A (mixed) **strategy**¹ for player i in a Bayesian game is a function defined on T_i

$$\sigma_i: T_i \to \Delta(A_i)$$
 (9.2)

where $\sigma_i(a_i|t_i)$ denotes the chance of player i playing a_i given this player is in type t_i .

Remark 9.4 (Interpretation). For a player i in a Bayesian game, one strategy assigns a probability distribution over his action set for each $t_i \in T_i$.

Definition 9.5. The **expected payoff**, given player i is type t_i and conditioned on player i's own belief π_i , from action a_i is

$$U_i(a_i, \sigma_{-i}|t_i) = \sum_{\omega \in \Omega} u_i(a_i, \sigma_{-i}(\omega)|t_i) \pi_i(\omega|t_i)$$
(9.3)

$$= \int_{\Omega} u_i(a_i, \sigma_{-i}(\omega)|t_i) \ d\pi_i(\omega|t_i) \tag{9.4}$$

Definition 9.6. An action $a_i \in A_i$ is a **best response** to σ_i , given player type t_i , if

$$U_i(a_i, \sigma_{-i}|t_i) \ge U_i(\tilde{a}_i, \sigma_{-i}|t_i) \ \forall \tilde{a}_i \in A_i$$

$$(9.5)$$

The definition for a mixed strategy $\sigma_i \in \Delta(A_i)$ is similar.

Definition 9.7. A Bayesian Nash equilibrium is a strategy profile for each player such that the strategy for each type of each player is a best response given the strategies for the other players deduced from the signal received.

Definition 9.8 (Lottery Generated by a Bayesian Game). The posterior belief of player i, together with an action profile a^* in G^* , generates a lottery $L_i(a^*, t_i)$ over $A \times \Omega$: the probability assigned by $L_i(a^*, t_i)$ to

$$\left((a^*(j, \tau_j(\omega)))_{j \in N}, \omega \right) \tag{9.6}$$

is the posterior belief that the state is ω when he receives the signal t_i and $a^*(j, \tau_j(\omega))$ being the action of player $(j, \tau_j(\omega))$ for each $j \in N$ in the profile a^* .

Definition 9.9 (Formal Definition). A **Nash equilibrium of a Bayesian Game**, $\langle N, \Omega, (A_i), (T_i), (\tau_i), (p_i), (\succsim_i) \rangle$ is a Nash equilibrium of the strategic game defined as follows.

- The set of players is the set of all pairs (i, t_i) for $i \in N$ and $t_i \in T_i$.
- The set of actions of each player (i, t_i) is A_i .

¹This also covers the *pure strategy* case since any pure strategy can be considered as a *degenerate* mixed strategy.

• The preference ordering $\succsim_{i}^{*} (i, t_{i})$ of each player (i, t_{i}) is defined by

$$a^* \succsim_{(i,t_i)}^* b^* \iff L_i(a^*,t_i) \succsim_i L_i(b^*,t_i)$$
 (9.7)

where $L_i(a^*, t_i)$ is the lottery over $A \times \Omega$ that assigns probability $\frac{p_i(\omega)}{p_i(\tau_i^{-1}(\omega))}$ to $(a^*(j, \tau_j(\omega)))_{j \in N}$ if $\omega \in \tau_i^{-1}(t_i)$, zero otherwise.