

# MAT237: Multivariable Calculus

Tianyu Du

Friday 22<sup>nd</sup> March, 2019

## Contents

<b>1</b>	<b>Limits, continuity, and related topics</b>	<b>2</b>
1.1	Open and Closed Sets, Boundary, Interior and Closure . . . . .	2
1.2	Limits and Continuity . . . . .	3
1.3	Limits of Multivariable Functions . . . . .	3
1.3.1	Continuity . . . . .	4
1.3.2	Continuous Functions and Open Sets . . . . .	4
1.4	Sequences and Completeness . . . . .	4
1.5	Compactness . . . . .	6
1.6	the Intermediate Value Theorem . . . . .	6
<b>2</b>	<b>Differentiation and related topics</b>	<b>6</b>
2.1	. . . . .	6
2.2	. . . . .	6
2.3	. . . . .	6
2.4	. . . . .	6
2.5	. . . . .	6
2.6	. . . . .	6
2.7	. . . . .	6
2.8	Optimization . . . . .	6
<b>3</b>	<b>The Implicit and Inverse Function Theorems</b>	<b>7</b>
3.1	The Implicit Function Theorem I . . . . .	7
3.2	Geometric content of the Implicit Function Theorem . . . . .	7
3.3	Transformations, and the Inverse Function Theorem . . . . .	8
<b>4</b>	<b>Integration</b>	<b>9</b>
4.1	Basics . . . . .	9
4.2	Integration on Higher Dimensions . . . . .	9
4.3	Iterated Integrals . . . . .	11
4.4	Change of Variables . . . . .	11
4.5	Further Aspects . . . . .	12
4.5.1	Exchanging Differentiation and Integration . . . . .	12
4.5.2	Improper Integrals . . . . .	12

<b>5</b>	<b>Vector Calculus</b>	<b>13</b>
5.1	Line Integrals . . . . .	13
5.1.1	Arc Length . . . . .	13
5.1.2	Line Integrals of Scalar Functions . . . . .	14
5.1.3	Line Integrals of Vector Fields . . . . .	14
5.1.4	Rectifiable Curves . . . . .	15
5.2	Green's Theorem . . . . .	15
5.2.1	Preliminary Definitions . . . . .	15
5.3	Surface Integrals . . . . .	16
5.3.1	Surface Areas and Surface Integrals . . . . .	16
5.3.2	An invariance property . . . . .	17
5.3.3	Volume and Area . . . . .	18
5.4	Divergence, Gradient and Curl . . . . .	18
5.5	Divergence Theorem . . . . .	18
5.6	Stokes Theorem . . . . .	20
5.6.1	Stokes Theorem in $\mathbb{R}^3$ . . . . .	20
5.6.2	The Generalization . . . . .	21

# 1 Limits, continuity, and related topics

## 1.1 Open and Closed Sets, Boundary, Interior and Closure

**Definition 1.1.1.** Let  $\mathbf{a} \in \mathbb{R}^n$ , and  $r > 0$ . The **open ball with centre  $\mathbf{a}$  and radius  $r$**  is defined as

$$\mathcal{B}(r, \mathbf{a}) := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x} - \mathbf{a}\| < r\}. \quad (1.1.1)$$

**Definition 1.1.2.** The **sphere with centre  $\mathbf{a}$  and radius  $r$**  is defined as

$$\partial\mathcal{B}(r, \mathbf{a}) := \{\mathbf{x} \in \mathbb{R}^n : \|\mathbf{x} - \mathbf{a}\| = r\} \quad (1.1.2)$$

**Definition 1.1.3.** Let  $S \subset \mathbb{R}^n$ ,  $S$  is **bounded** if

$$\exists r > 0 \text{ s.t. } S \subset \mathcal{B}(r, \mathbf{0}) \quad (1.1.3)$$

**Definition 1.1.4.** Let  $S \subset \mathbb{R}^n$ , then the **complement** of  $S$  is defined as

$$S^c := \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x} \notin S\} \quad (1.1.4)$$

**Definition 1.1.5.** Let  $S \subset \mathbb{R}^n$ , the **interior** of  $S$  is defined as

$$S^{int} := \{\mathbf{x} \in \mathbb{R}^n : \exists \varepsilon > 0 \text{ s.t. } \mathcal{B}(\varepsilon, \mathbf{x}) \subset S\} \quad (1.1.5)$$

**Definition 1.1.6.** The **boundary** of  $S$  is defined as

$$\partial S := \{\mathbf{x} \in \mathbb{R}^n : \forall \varepsilon > 0 \mathcal{B}(\varepsilon, \mathbf{x}) \cap S \neq \emptyset \wedge \mathcal{B}(\varepsilon, \mathbf{x}) \cap S^c \neq \emptyset\} \quad (1.1.6)$$

**Theorem 1.1.1.** A point  $\mathbf{x} \in S$  is either a *boundary point* or a *interior point*.

**Definition 1.1.7.** The **closure** of  $S$  is defined as

$$\overline{S} := S^{int} \cup \partial S \quad (1.1.7)$$

**Theorem 1.1.2.** For any  $S \subset \mathbb{R}^n$

$$S^{int} \subset S \subset \overline{S} \quad (1.1.8)$$

**Theorem 1.1.3.** For any  $S \subset \mathbb{R}^n$

$$\partial S = \partial(S^c) \quad (1.1.9)$$

**Definition 1.1.8.** A set  $S \subset \mathbb{R}^n$  is **open** if  $S = S^{int}$ .  $S$  is **closed** if  $S = \overline{S}$ .

**Theorem 1.1.4.**

$$S \text{ is closed} \iff S^c \text{ is open} \quad (1.1.10)$$

*Proof.*

$$S \text{ is closed} \iff \partial S \subset S \iff \partial(S^c) \subset S \quad (1.1.11)$$

$$\iff \text{no point of } S^c \text{ is a boundary point} \iff S^c \text{ is open} \quad (1.1.12)$$

■

## 1.2 Limits and Continuity

### 1.3 Limits of Multivariable Functions

**Definition 1.3.1.** Let  $S \subset \mathbb{R}^n$ ,  $\mathbf{f} : S \rightarrow \mathbb{R}^k$ , and  $\mathbf{a} \in S$ , then

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} \mathbf{f}(\mathbf{x}) = \mathbf{L} \in \mathbb{R}^k \quad (1.3.1)$$

is defined as

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } \forall \mathbf{x} \in S, 0 < \|\mathbf{x} - \mathbf{a}\| < \delta \implies \|\mathbf{f}(\mathbf{x}) - \mathbf{L}\| < \varepsilon \quad (1.3.2)$$

For this definition to be non-trivial, we need  $\mathbf{a}$  not be an isolated point,

$$\forall \delta > 0, \exists \mathbf{x} \in S \text{ s.t. } \|\mathbf{x} - \mathbf{a}\| \in (0, \delta) \quad (1.3.3)$$

**Theorem 1.3.1** (Limit Laws). Let  $S \subset \mathbb{R}^n$  and  $\mathbf{a} \in \mathbb{R}^n$  satisfying (1.3.3). And  $f, g : S \rightarrow \mathbb{R}$ ,  $L, M \in \mathbb{R}$  such that

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} f(\mathbf{x}) = L \quad (1.3.4)$$

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} g(\mathbf{x}) = M \quad (1.3.5)$$

then

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} [f(\mathbf{x}) + g(\mathbf{x})] = L + M \quad (1.3.6)$$

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} [f(\mathbf{x}) \cdot g(\mathbf{x})] = LM \quad (1.3.7)$$

**Theorem 1.3.2** (Squeeze Theorem). Let  $S \subset \mathbb{R}^n$ ,  $\mathbf{a} \in \mathbb{R}^n$  satisfies (1.3.3). Suppose that  $f, g, h : S \rightarrow \mathbb{R}$  and there exists  $p > 0$  and  $L \in \mathbb{R}$  such that

$$\forall \mathbf{x} \in S \cap \mathcal{B}(p, \mathbf{a}) \quad f(\mathbf{x}) \leq g(\mathbf{x}) \leq h(\mathbf{x}) \quad (1.3.8)$$

and

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} f(\mathbf{x}) = \lim_{\mathbf{x} \rightarrow \mathbf{a}} h(\mathbf{x}) = L \quad (1.3.9)$$

then

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} g(\mathbf{x}) = L \quad (1.3.10)$$

**Corollary 1.3.1.** Let  $g, h : S \rightarrow \mathbb{R}$  and

$$|g(\mathbf{x})| \leq h(\mathbf{x}) \quad \forall \mathbf{x} \in S \quad (1.3.11)$$

$$\text{and } \lim_{\mathbf{x} \rightarrow \mathbf{a}} h(\mathbf{x}) = 0 \quad (1.3.12)$$

then

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} g(\mathbf{x}) = 0 \quad (1.3.13)$$

**Theorem 1.3.3.** Assume that  $S \subset \mathbb{R}^n$  and let  $\mathbf{a} \in \mathbb{R}^n$  satisfying (1.3.3). Let  $\mathbf{f} : S \rightarrow \mathbb{R}^k$ , then

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} \mathbf{f}(\mathbf{x}) = \mathbf{L} \iff \lim_{\mathbf{x} \rightarrow \mathbf{a}} f_j(\mathbf{x}) = L_j \quad \forall j \quad (1.3.14)$$

### 1.3.1 Continuity

**Definition 1.3.2.** Let  $S \subset \mathbb{R}^n$  and  $\mathbf{f}: S \rightarrow \mathbb{R}^k$ .  $\mathbf{f}$  is **continuous at**  $\mathbf{a} \in S$  if

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} \mathbf{f}(\mathbf{x}) = \mathbf{f}(\mathbf{a}) \quad (1.3.15)$$

and  $\mathbf{f}$  is **continuous** if  $\mathbf{f}$  is continuous at every point in  $S$ .

**Theorem 1.3.4** (Basic Properties of Continuity). Assume that  $S \subset \mathbb{R}^n$  and  $\mathbf{a} \in S$ ,

- (i) If  $\mathbf{f}: S \rightarrow \mathbb{R}^k$  is continuous at  $\mathbf{a}$ , then every component of  $\mathbf{f}$ ,  $f_j: S \rightarrow \mathbb{R}$ , is continuous at  $\mathbf{a}$ .
- (ii) If  $\mathbf{f}, \mathbf{g}: S \rightarrow \mathbb{R}^k$  are continuous at  $\mathbf{a}$ , then  $\mathbf{f} + \mathbf{g}$  is continuous at  $\mathbf{a}$ .
- (iii) If  $f, g: S \rightarrow \mathbb{R}$  continuous, then  $fg$  is continuous and  $\frac{f}{g}$  is continuous given  $g(\mathbf{a}) \neq 0$ .
- (iv) A composition of continuous functions is continuous.
- (v) The elementary functions of a single variable (trigonometric functions and their inverses, polynomials, exponential and log) are continuous on their domains.

### 1.3.2 Continuous Functions and Open Sets

**Theorem 1.3.5.** Assume that  $\mathbf{f}: \mathbb{R}^n \rightarrow \mathbb{R}^k$ , then the following are equivalent

- (i)  $\mathbf{f}$  is continuous;
- (ii) For every open set  $\mathcal{O} \subset \mathbb{R}^k$ ,  $\mathbf{f}^{-1}(\mathcal{O})$  is also open;
- (iii) For every closed set  $\mathcal{C} \subset \mathbb{R}^k$ ,  $\mathbf{f}^{-1}(\mathcal{C})$  is also closed.

## 1.4 Sequences and Completeness

**Definition 1.4.1.** A sequence  $\{\mathbf{a}_j\}_j$  in  $\mathbb{R}^n$  **converges to the limit**  $\mathbf{L} \in \mathbb{R}^n$  if

$$\forall \varepsilon > 0 \exists J \in \mathbb{N}, \text{ s.t. } \forall j \geq J \implies \|\mathbf{a}_j - \mathbf{L}\| < \varepsilon \quad (1.4.1)$$

**Theorem 1.4.1.**

$$\lim_{j \rightarrow \infty} \mathbf{a}_j = \mathbf{L} \iff \lim_{j \rightarrow \infty} \|\mathbf{a}_j - \mathbf{L}\| = 0 \quad (1.4.2)$$

**Theorem 1.4.2.** Let  $\{a_{jk}\}_j$  be a sequence in  $\mathbb{R}^n$  where  $k \in [n]$ , and let  $\mathbf{L} = (L_1, \dots, L_n) \in \mathbb{R}^n$ , then

$$\lim_{j \rightarrow \infty} \mathbf{a}_j = \mathbf{L} \iff \lim_{j \rightarrow \infty} a_{jk} = L_k \quad \forall k \in [n] \quad (1.4.3)$$

*Proof Idea.*

$$\forall j \in [n], \quad |a_j - L_j| \leq \|\mathbf{a} - \mathbf{L}\| \leq n \max_{k \in [n]} |a_k - L_k| \quad (1.4.4)$$

■

**Axiom 1.1** (the Completeness Axiom). Every *bounded* and *nonempty* set of *real numbers* has a *least upper bound* (**supremum**) and a *greatest lower bound* (**infimum**).

**Theorem 1.4.3** (Monotone Sequence Theorem). Every bounded nondecreasing sequence of real numbers converges to a limit.

*Proof Idea.* Note that such sequence converges to its supremum  $S$ .

Let  $\varepsilon > 0$ , there exists  $j^*$  such that

$$S - \varepsilon < a_{j^*} \leq S \quad (1.4.5)$$

take such  $j^*$  and by the nondecreasing property,

$$\forall j \geq j^* \quad a_j > S - \varepsilon \quad (1.4.6)$$

which implies  $|S - a_j| < \varepsilon$ . ■

**Definition 1.4.2.** A **subsequence** of a sequence  $\{a_j\}_{j \geq j_0}$  in  $\mathbb{R}^n$  is a sequence constructed as  $\{a_{k_j}\}_j$ , such that  $\{k_j\}_j$  is a *strictly increasing* sequence bounded below by  $j_0$ .

**Remark 1.4.1.** Subsequences can be constructed using strictly increasing transformations.

**Proposition 1.4.1.** If  $\{a_j\}_j$  is a sequence in  $\mathbb{R}^n$  converges to  $\mathbf{L}$ , then (i) any subsequence of it converges to the (ii) same limit.

*Proof Idea.* Suppose not and reach a contradiction. ■

**Theorem 1.4.4** (Bounded Sequence Theorem in  $\mathbb{R}$ ). Every bounded sequence in  $\mathbb{R}$  has a convergent subsequence.

*Proof.* Let  $\{a_j\}_j$  be a bounded sequence.

For each  $j \in \mathbb{N}$ , define  $b_{k_j} := \inf_{k > k_j} a_k$ .

Note that  $\{b_j\}$  is non-decreasing and bounded, so it converges to some limit  $\ell$ .

Let  $\{a_{k_j}\}_j$  denote a subsequence of the original sequence, define  $k_0 = j_0$ , and indices are constructed in a recurrent way.

Suppose every index before  $k_j$  has been chosen, we choose  $k_{j+1}$  to be the index such that

$$b_{k_j} \leq a_{k_{j+1}} < b_{k_j} + \frac{1}{j} \quad (1.4.7)$$

by construction,  $\{a_{k_j}\}_j$  is bounded by both  $\{b_{k_j}\}_j$  and  $\{b_{k_j} + \frac{1}{j}\}_j$ , and both bounding sequences converge to  $\ell$ . So  $\{a_{k_j}\}_j$  converges to  $\ell$  by *squeeze theorem*. ■

**Theorem 1.4.5** (Bounded Sequence Theorem). Every bounded sequence in  $\mathbb{R}^n$  has a convergent subsequence.

## 1.5 Compactness

## 1.6 the Intermediate Value Theorem

# 2 Differentiation and related topics

## 2.1

## 2.2

## 2.3

## 2.4

## 2.5

## 2.6

## 2.7

## 2.8 Optimization

**Theorem 2.8.1.** Let  $S \subset \mathbb{R}^n$  be an open set and  $f, g : S \rightarrow \mathbb{R}$  be  $C^1$  functions. If  $\mathbf{x}$  is a *local extremal* satisfying  $g(\mathbf{x}) = 0$ , and  $\nabla g(\mathbf{x}) \neq \mathbf{0}$ , then

$$\exists \lambda \in \mathbb{R} \text{ s.t. } \begin{cases} \nabla f(\mathbf{x}) = \lambda \nabla g(\mathbf{x}) \\ g(\mathbf{x}) = 0 \end{cases} \quad (2.8.1)$$

**Lemma 2.8.1.**  $\nabla g(\mathbf{x})$  is orthogonal to the constraint set  $g^{-1}(0)$ .

**Proposition 2.8.1.** Equations (2.8.1)  $\implies \nabla f(\mathbf{x}) \perp g^{-1}(0)$  at  $\mathbf{x}$ .

**Theorem 2.8.2.** Let  $S \subseteq \mathbb{R}^n$  be an open set, and  $f, \{g_i\}_{i=1}^k : S \rightarrow \mathbb{R}$  be  $C^1$  functions. Define  $\mathbf{g}(\mathbf{x}) : \mathbb{R}^n \rightarrow \mathbb{R}^k \equiv (g_1(\mathbf{x}), g_2(\mathbf{x}), \dots, g_k(\mathbf{x}))$ .

If  $\mathbf{x} \in S$  is a *local extremal* of  $f$  such that  $\mathbf{g}(\mathbf{x}) = \mathbf{0}$ , and  $\{\nabla g_i(\mathbf{x})\}$  are linearly independent (i.e.  $\text{rank}(D\mathbf{g}(\mathbf{x})) = k$ ), then

$$\exists \boldsymbol{\lambda} \in \mathbb{R}^k \text{ s.t. } \begin{cases} \nabla f(\mathbf{x}) = \boldsymbol{\lambda}^T D\mathbf{g}(\mathbf{x}) \\ \mathbf{g}(\mathbf{x}) = \mathbf{0} \end{cases} \quad (2.8.2)$$

**Remark 2.8.1.** Procedure of optimization on *open sets*:

- (i) Find all critical points.
- (ii) Find optimizers among critical points.

**Remark 2.8.2.** Procedure of optimization with *inequality constraints*:

- (i) Find critical points without the constraints.
- (ii) Find critical points on the constraints.
- (iii) Find optimizers among candidates.

### 3 The Implicit and Inverse Function Theorems

#### 3.1 The Implicit Function Theorem I

**Theorem 3.1.1** (Implicit Function Theorem). Let  $S \subseteq \mathbb{R}^{n+k}$  be an open set, and function  $F : S \rightarrow \mathbb{R}^k$  be a  $C^1$  function. Suppose there exists point  $\mathbf{a} \in \mathbb{R}^n, \mathbf{b} \in \mathbb{R}^k$  such that

$$F(\mathbf{a}, \mathbf{b}) = \mathbf{0} \quad (3.1.1)$$

If

$$\det(D_{\mathbf{y}}(F(\mathbf{a}, \mathbf{b}))) \neq 0 \quad (3.1.2)$$

then there exists  $r_0, r_1 > 0$  and a  $C^1$  function  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^k$  such that

$$\forall \mathbf{x} \in \mathcal{B}(r_0, \mathbf{a}), \mathbf{f}(\mathbf{x}) \in \mathcal{B}(r_1, \mathbf{b}) \wedge F(\mathbf{x}, \mathbf{f}(\mathbf{x})) = \mathbf{0} \quad (3.1.3)$$

and define  $\mathbf{y} \equiv \mathbf{f}(\mathbf{x})$ , the derivative of  $\mathbf{f}$  can be found as

$$D\mathbf{f}(\mathbf{x}) = -[D_{\mathbf{y}}F(\mathbf{x}, \mathbf{y})]^{-1}D_{\mathbf{x}}F(\mathbf{x}, \mathbf{y}) \quad (3.1.4)$$

**Remark 3.1.1.** Procedure to prove solvability of non-linear equations

$$\mathbf{F}(\mathbf{x}, \mathbf{y}) = \mathbf{0} \quad (3.1.5)$$

near  $(\mathbf{a}, \mathbf{b})$ .

(i) Verify  $\mathbf{F}(\mathbf{a}, \mathbf{b}) = \mathbf{0}$ .

(ii) Assert

$$\det(D_{\mathbf{y}}\mathbf{F}(\mathbf{a}, \mathbf{b})) \neq 0 \quad (3.1.6)$$

(iii) Approximate solution  $\mathbf{y} = \mathbf{f}(\mathbf{x})$  using

$$\mathbf{f}(\mathbf{x} + \mathbf{h}) \approx \mathbf{a} + D\mathbf{f}(\mathbf{a})\mathbf{h} \quad (3.1.7)$$

$$= \mathbf{a} - [D_{\mathbf{y}}\mathbf{F}(\mathbf{a}, \mathbf{b})]^{-1}D_{\mathbf{x}}\mathbf{F}(\mathbf{a}, \mathbf{b}) \quad (3.1.8)$$

#### 3.2 Geometric content of the Implicit Function Theorem

**Definition 3.2.1.** Let  $S \subseteq \mathbb{R}^n$  and  $\mathbf{a} \in S$ .  $S$  is **singular** at  $\mathbf{a}$  if

$$\forall r > 0 \ S \cap \mathcal{B}(r, \mathbf{a}) \text{ cannot be represented as a } C^1 \text{ graph.} \quad (3.2.1)$$

$S$  is **regular** at  $\mathbf{a}$  if it is not singular there.

**Theorem 3.2.1** ( $k$  dimensional manifold as level set). Let  $U \subseteq \mathbb{R}^n$  and let  $\mathbf{F} : U \rightarrow \mathbb{R}^{n-k}$  be a  $C^1$  function.

$$S \equiv \mathbf{F}^{-1}(\mathbf{0}) \quad (3.2.2)$$

Let  $\mathbf{a} \in U$ , if

$$\text{rank}(D\mathbf{F}(\mathbf{a})) = n - k \quad (3.2.3)$$

then  $\exists r > 0$  such that the *level set* of  $\mathbf{F}$  near  $\mathbf{a}$

$$\mathcal{B}(r, \mathbf{a}) \cap S \quad (3.2.4)$$

can be represented as a  $C^1$  graph.



**Theorem 3.2.2** ( $k$  dimensional manifold as parameterization). Let  $T \subseteq \mathbb{R}^k$  and let  $\mathbf{f} : U \rightarrow \mathbb{R}^n$  be a  $C^1$  function.

$$S \equiv \mathbf{f}(T) \quad (3.2.5)$$

Let  $\mathbf{t} \in T$ , if

$$\text{rank}(\mathbf{f}(\mathbf{t})) = k \quad (3.2.6)$$

then  $\exists r > 0$  such that the *parameterization of  $\mathbf{f}$  near  $\mathbf{t}$*

$$\mathbf{f}(T \cap \mathcal{B}(r, \mathbf{t})) \quad (3.2.7)$$

can be represented as a  $C^1$  graph.

### 3.3 Transformations, and the Inverse Function Theorem

**Example 3.3.1** (Polar coordinate in  $\mathbb{R}^2$ ). Let

$$U \equiv \{(r, \theta) : r > 0 \wedge \theta \in (-\pi, \pi)\} \quad (3.3.1)$$

$$V \equiv \mathbb{R}^2 \setminus \{(x, 0) : x \leq 0\} \quad (3.3.2)$$

Define  $\mathbf{f} : U \rightarrow V$  as

$$\mathbf{f}(r, \theta) \equiv \begin{pmatrix} r \cos(\theta) \\ r \sin(\theta) \end{pmatrix} \quad (3.3.3)$$

**Example 3.3.2** (Spherical coordinate in  $\mathbb{R}^3$ ). Define

$$\mathbf{f}(r, \theta, \varphi) = \begin{pmatrix} r \cos(\theta) \sin(\varphi) \\ r \sin(\theta) \sin(\varphi) \\ r \cos(\varphi) \end{pmatrix} \quad (3.3.4)$$

**Example 3.3.3** (Cylindrical coordinate in  $\mathbb{R}^3$ ). Define

$$\mathbf{f}(r, \theta, z) = \begin{pmatrix} r \cos(\theta) \\ r \sin(\theta) \\ z \end{pmatrix} \quad (3.3.5)$$

**Theorem 3.3.1** (Inverse Function Theorem). Let  $U$  and  $V$  be open subsets in  $\mathbb{R}^n$ , and  $\mathbf{f} : U \rightarrow V$ . Let  $\mathbf{a} \in U$  and define  $\mathbf{b} \equiv \mathbf{f}(\mathbf{a}) \in V$ . If

$$\det(D\mathbf{f}(\mathbf{a})) \neq 0 \quad (3.3.6)$$

then there exists  $M \subseteq U$  and  $N \subseteq V$  such that

- (i)  $\mathbf{a} \in M$  and  $\mathbf{b} \in N$ ,
- (ii)  $\mathbf{f}$  is bijective between  $M$  and  $N$ ,
- (iii)  $\mathbf{f}^{-1} : N \rightarrow M$  is  $C^1$ ,

and **for all  $\mathbf{x} \in M$**  such  $\mathbf{y} \equiv \mathbf{f}(\mathbf{x}) \in N$ ,

$$D\mathbf{f}^{-1}(\mathbf{y}) = [D\mathbf{f}(\mathbf{x})]^{-1} \quad (3.3.7)$$

## 4 Integration

### 4.1 Basics

**Theorem 4.1.1** (**Properties of infimum and supremum**). Let  $A \subseteq \mathbb{R}^n$  and  $A \neq \emptyset$ , and  $f, g : A \rightarrow \mathbb{R}$  are bounded functions. Let  $m$  and  $M$  denote the infimum and supremum respectively, then

- (i)  $m_A f + m_A g \leq m_A(f + g) \leq M_A(f + g) \leq M_A f + M_A g$
- (ii) If  $A' \subseteq A$ , then  $m_A f \leq m_{A'} f \leq M_{A'} f \leq M_A f$
- (iii) If  $f(\mathbf{x}) \leq g(\mathbf{x}) \forall \mathbf{x} \in A$ , then  $m_A f \leq m_A g$  and  $M_A f \leq M_A g$
- (iv)  $|M_A f| \leq M_A |f|$
- (v)  $M_A |f| - m_A |f| \leq M_A f - m_A f$
- (vi)  $\forall c \in \mathbb{R}, M_A(cf) - m_A(cf) = |c|(M_A f - m_A f)$
- (vii)  $M_A f - m_A f = \sup\{f(x) - f(y) : x, y \in A\}$

### 4.2 Integration on Higher Dimensions

**Definition 4.2.1.** A rectangle  $\mathcal{R} \subseteq \mathbb{R}^n$  is defined as

$$\mathcal{R} \equiv \prod_{i=1}^n [a_i, b_i] \quad (4.2.1)$$

where  $a_i, b_i \in \mathbb{R}$  and  $a_i < b_i$ .

**Definition 4.2.2.** A **partition**  $P$  of rectangle  $\mathcal{R} = \prod_{i=1}^n [a_i, b_i]$  is a list of  $n$  **finite** and increasing list of real numbers

$$P = \{L_1, L_2, \dots, L_n\} \quad (4.2.2)$$

where  $L_i = \{e_j\}_{j=0}^{T_i}$  such that

$$a_i = e_0 < e_1 < \dots < e_{T_i} = b_i \quad (4.2.3)$$

and such partition induces a set of rectangles(boxes)  $\mathcal{B}(P) \equiv \{B_j\}_{j=1}^J \subseteq \mathcal{R}$ .

**Definition 4.2.3.** Let  $P$  and  $P'$  be two partitions of  $\mathcal{R}$ . Then  $P'$  is a **refinement** of  $P$  if

$$\forall B_j \in \mathcal{B}(P), B'_j \in \mathcal{B}(P') \quad B'_j \subseteq B_j \vee B'_j \cap B_j = \emptyset \quad (4.2.4)$$

**Definition 4.2.4.** Define the **volume** of rectangle  $\mathcal{R} = \prod_{i=1}^n [a_i, b_i]$  as

$$V^n(\mathcal{R}) \equiv \prod_{i=1}^n (b_i - a_i) \quad (4.2.5)$$

**Definition 4.2.5.** The **lower Riemann sum** of  $f$  with partition  $P$  on  $\mathcal{R}$  is defined as

$$L_P f \equiv \sum_{B_j \in \mathcal{B}(P)} \inf_{\mathbf{x} \in B_j} f(\mathbf{x}) V^n(B_j) \quad (4.2.6)$$

and the **upper Riemann sum** is defined as

$$U_P f \equiv \sum_{B_j \in \mathcal{B}(P)} \sup_{\mathbf{x} \in B_j} f(\mathbf{x}) V^n(B_j) \quad (4.2.7)$$

**Definition 4.2.6.** The **upper integral** and **lower integral** of  $f$  on  $\mathcal{R}$  are defined as

$$\bar{I}_{\mathcal{R}}f \equiv \inf_P U_P f \quad (4.2.8)$$

$$\underline{I}_{\mathcal{R}}f \equiv \sup_P L_P f \quad (4.2.9)$$

**Definition 4.2.7.** A bounded real-valued function  $f$  defined on  $\mathcal{R}$  is **integrable** if

$$\underline{I}_{\mathcal{R}}f = \bar{I}_{\mathcal{R}}f \quad (4.2.10)$$

and the integral is defined as

$$\int \cdots \int_{\mathcal{R}} f \, dV^n \equiv \underline{I}_{\mathcal{R}}f = \bar{I}_{\mathcal{R}}f \quad (4.2.11)$$

**Lemma 4.2.1.** Let  $f$  be a bounded real-valued function defined on  $\mathcal{R}$ ,  $f$  is integrable if and only if  $\forall \epsilon > 0$ , there exists a partition  $P$  of  $\mathcal{R}$  such that

$$U_P f - L_P f < \epsilon \quad (4.2.12)$$

**Theorem 4.2.1.** Let  $f$  and  $g$  be two integrable functions on  $\mathcal{R} \subseteq \mathbb{R}^n$ , let  $c \in \mathbb{R}$ ,

- (i)  $f + g : \mathcal{R} \rightarrow \mathbb{R}$  is integrable and  $\int_{\mathcal{R}}(f + g) = \int_{\mathcal{R}} f + \int_{\mathcal{R}} g$
- (ii)  $c \cdot f$  is integrable and  $\int_{\mathcal{R}} c \cdot f = c \int_{\mathcal{R}} f$
- (iii)  $f(\mathbf{x}) \geq g(\mathbf{x}) \, \forall \mathbf{x} \in \mathcal{R} \implies \int_{\mathcal{R}} f \geq \int_{\mathcal{R}} g$
- (iv)  $|f|$  is integrable and  $|\int_{\mathcal{R}} f| \leq \int_{\mathcal{R}} |f|$

**Definition 4.2.8.** Let  $S \subseteq \mathbb{R}^n$  be a bounded set, and there exists rectangle  $\mathcal{R}$  covers  $S$ , the **indicator function** of  $S$  is  $\chi_S : \mathcal{R} \rightarrow \{0, 1\}$ , defined as

$$\chi_S(\mathbf{x}) \equiv \mathbb{I}(\mathbf{x} \in S) \quad (4.2.13)$$

**Definition 4.2.9.** Let  $S \subseteq \mathbb{R}^n$  be a bounded set, and there exists rectangle  $\mathcal{R}$  covers  $S$ . Let  $f : \mathcal{R} \rightarrow \mathbb{R}$  be a bounded function, then  $f$  is **integrable on  $S$**  if  $\chi_S f$  is integrable on  $\mathcal{R}$ . And

$$\int \cdots \int_S f \, dV^n \equiv \int \cdots \int_{\mathcal{R}} \chi_S f \, dV^n \quad (4.2.14)$$

**Definition 4.2.10.** Let  $Z \subseteq \mathbb{R}^n$ ,  $Z$  has **zero content** if for all  $\epsilon > 0$ , there exists a finite set of rectangles  $\{R_\ell\}_{\ell=1}^L$  covers  $Z$  and

$$\sum_{\ell=1}^L V^n(R_\ell) < \epsilon \quad (4.2.15)$$

**Proposition 4.2.1.** Let  $Z \subseteq \mathbb{R}^n$  has zero content, then

- (i) For any  $Z' \subseteq Z$ ,  $Z'$  has zero content.
- (ii) Finite union of content zero sets has zero content.
- (iii) Let  $f : [a, b] \rightarrow \mathbb{R}$  be an integrable function, it's graph  $\{(x, f(x)) : x \in [a, b]\}$  has zero content.
- (iv) Let  $\mathbf{f} : [a, b] \rightarrow \mathbb{R}^2$  be a  $C^1$  function, the parameterization  $\mathbf{f}([a, b])$  has zero content.

**Theorem 4.2.2.** Let  $\mathcal{R}$  be a rectangle in  $\mathbb{R}^n$  and  $f$  is integrable on  $\mathcal{R}$  if

$$\{\mathbf{x} \in \mathcal{R} : f \text{ is discontinuous at } \mathbf{x}\} \quad (4.2.16)$$

has zero content.

**Proposition 4.2.2** (Folland 4.22). Suppose  $Z \subseteq \mathbb{R}^n$  has zero content. If  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is bounded, then  $f$  is integrable on  $Z$  and  $\int_Z f \, dV^n = 0$ .

### 4.3 Iterated Integrals

**Theorem 4.3.1** (Fubini's Theorem). Let  $\mathcal{R} = [a, b] \times [c, d] \subseteq \mathbb{R}^2$  and  $f : \mathcal{R} \rightarrow \mathbb{R}$  is bounded. Assuming that

- (i)  $f$  is integrable on  $\mathcal{R}$ .
- (ii) for each  $y \in [c, d]$ , the function  $f_y(x) \equiv f(x, y)$  is integrable on  $[a, b]$ .
- (iii) Define  $g(y) \equiv \int_a^b f(x, y) dx$  is integrable on  $[c, d]$ .

Then

$$\iint_{\mathcal{R}} f \, dA = \int_c^d \left( \int_a^b f(x, y) \, dx \right) dy \quad (4.3.1)$$

**Proposition 4.3.1.** Let  $S \subseteq \mathbb{R}^n$  be an unbounded set, and  $f : S \rightarrow \mathbb{R}$ . Then improper integral  $\int \cdots \int_S f \, d^n \mathbf{x}$  is absolutely convergent on  $\mathbb{R}^n$  if and only if  $\int \cdots \int_{\mathbb{R}^n} \chi_S f \, d^n \mathbf{x}$  is absolutely convergent.

### 4.4 Change of Variables

**Theorem 4.4.1** (Change of Variable). Let  $U$  and  $V$  be two open subsets of  $\mathbb{R}^n$ , and let  $\mathbf{G} : U \rightarrow V$  be a  $C^1$  bijection. Let  $T \subset U$  and  $S \subset V$ . Suppose  $\mathbf{G}(T) = S$ , then

$$\int \cdots \int_S f \, d\Omega = \int \cdots \int_T f \circ \mathbf{G} \, |\det D\mathbf{G}| \, d\Theta \quad (4.4.1)$$

**Corollary 4.4.1.** Let  $S$  be a region in  $\mathbb{R}^n$ , suppose  $S$  can be parameterized by  $\mathbf{G} : T \rightarrow S$ . By the change of variable formula, consider the special case  $f(\mathbf{x}) = 1$ ,

$$|S| = \int \cdots \int_S 1 \, d\Omega = \int \cdots \int_T 1 \, |\det D\mathbf{G}(\mathbf{u})| \, d\Theta \quad (4.4.2)$$

**Example 4.4.1** (Polar Coordinate). Define the coordinate transformation mapping from polar to Cartesian,

$$\mathbf{P}(r, \theta) \equiv (x, y) = \begin{pmatrix} r \cos \theta \\ r \sin \theta \end{pmatrix}, \quad \theta \in [0, 2\pi] \quad r \in \mathbb{R}_+ \quad (4.4.3)$$

and  $|\det D\mathbf{P}(r, \theta)| = r$ .

**Example 4.4.2** (Cylindrical Coordinate). Define the coordinate transformation mapping from cylindrical to Cartesian as

$$\mathbf{C}(r, \theta, z) \equiv (x, y, z) = \begin{pmatrix} r \cos \theta \\ r \sin \theta \\ z \end{pmatrix}, \quad \theta \in [0, 2\pi] \quad r \in \mathbb{R}_+ \quad z \in \mathbb{R} \quad (4.4.4)$$

and  $|\det D\mathbf{C}(r, \theta, z)| = r$ .

**Example 4.4.3** (Spherical Coordinate). Define the coordinate transformation mapping from spherical to Cartesian as

$$\mathbf{S}(r, \theta, \varphi) = \begin{pmatrix} r \cos \theta \sin \varphi \\ r \sin \theta \sin \varphi \\ r \cos \varphi \end{pmatrix} \quad (4.4.5)$$

and  $|\det D\mathbf{S}(r, \theta, \varphi)| = r^2 \sin \varphi$

## 4.5 Further Aspects

### 4.5.1 Exchanging Differentiation and Integration

**Theorem 4.5.1** (Exchanging Differentiation and Integration). Let  $f(\mathbf{x}, \mathbf{t}) : S \times T \rightarrow \mathbb{R}$  and define  $F(\mathbf{x}) : S \rightarrow \mathbb{R}$  as

$$F(\mathbf{x}) \equiv \int \cdots \int_T f(\mathbf{x}, \mathbf{t}) \, d\Omega \quad (4.5.1)$$

If

- (i)  $S$  is open and  $T$  is compact and bounded;
- (ii)  $f$  and  $F$  are continuous on their domains;
- (iii) and  $\forall x_j \in \mathbf{x}$ ,  $\frac{\partial f(\mathbf{x}, \mathbf{t})}{\partial x_j}$  is continuous,

then  $F$  is  $C^1$  in  $S$  and for every  $j$ ,

$$\frac{\partial F(\mathbf{x})}{\partial x_j} = \int \cdots \int_T \frac{\partial f(\mathbf{x}, \mathbf{t})}{\partial x_j} \, d\Omega \quad (4.5.2)$$

**Corollary 4.5.1.** By the definition of partial derivative, above theorem is equivalent to

$$\lim_{h \rightarrow 0} \int \cdots \int_T \frac{f(\mathbf{x} + h\mathbf{e}_j, \mathbf{t})}{h} \, d\Omega = \int \cdots \int_T \lim_{h \rightarrow 0} \frac{f(\mathbf{x} + h\mathbf{e}_j, \mathbf{t})}{h} \, d\Omega \quad (4.5.3)$$

### 4.5.2 Improper Integrals

**Definition 4.5.1** (Unbounded Domains). An **improper integral** with unbounded domain  $\int \cdots \int_{\mathbb{R}^n} f \, d\Omega$  is **absolutely convergent** if there exists  $L \in \mathbb{R}$  such that

$$\forall \varepsilon > 0 \, \exists R > 0 \text{ s.t. } \forall S \subseteq \mathbb{R}^n \, B(R, \mathbf{0}) \subset S \implies \left| \int \cdots \int_S f \, d\Omega - L \right| < \varepsilon \quad (4.5.4)$$

**Theorem 4.5.2.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a continuous function, and that

$$\lim_{R \rightarrow \infty} \int \cdots \int_{B(R, \mathbf{0})} |f| \, d\Omega \text{ exists} \quad (4.5.5)$$

then  $\int \cdots \int_{\mathbb{R}^n} f \, d\Omega$  is absolutely convergent.

**Corollary 4.5.2** (Equivalence). Above improper integral  $\int \cdots \int_{\mathbb{R}^n} f \, d\Omega$  is absolutely convergent if set

$$\left\{ \int \cdots \int_{B(R, \mathbf{0})} |f| \, d\Omega : R \in \mathbb{R}_{++} \right\} \quad (4.5.6)$$

is bounded.

**Corollary 4.5.3.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a continuous function, if

$$\exists p > n, \, C > 0 \text{ s.t. } |f(\mathbf{x})| \leq \frac{1}{\|\mathbf{x}\|^p} \, \forall \mathbf{x} \in \mathbb{R}^n \quad (4.5.7)$$

then  $\int \cdots \int_{\mathbb{R}^n} f \, d\Omega$  is absolutely convergent.

**Definition 4.5.2** (Unbounded Function). Let  $S \subset \mathbb{R}^n$ ,  $\mathbf{a} \in \mathbb{R}^n$ . Consider a function  $f : S \setminus \{\mathbf{a}\} \rightarrow \mathbb{R}$ . Then the improper integral  $\int \cdots \int_S f d\Omega$  is absolutely convergent if

$$\exists L \in \mathbb{R} \text{ s.t. } \forall \varepsilon > 0 \exists r > 0 \text{ s.t. } \forall U \subset S \text{ s.t. } \mathbf{a} \in U^{\text{int}} \wedge U \subset B(r, \mathbf{a}), \left| \int \cdots \int_{S \setminus U} f d\Omega - L \right| < \varepsilon \quad (4.5.8)$$

**Theorem 4.5.3.** Let  $f : S \setminus \{\mathbf{a}\} \rightarrow \mathbb{R}$ , if

$$\lim_{r \rightarrow 0} \int \cdots \int_{S \setminus B(r, \mathbf{a})} |f| d\Omega \text{ exists} \quad (4.5.9)$$

then  $\int \cdots \int_S f d\Omega$  is absolutely convergent.

**Corollary 4.5.4** (Equivalence). If the set

$$\left\{ \iint_{S \setminus B(r, \mathbf{a})} |f| d\Omega : r \in \mathbb{R}_{++} \right\} \quad (4.5.10)$$

is bounded, then  $\int \cdots \int_S f d\Omega$  is absolutely convergent.

**Corollary 4.5.5.** Let  $f : S \setminus \{\mathbf{a}\} \rightarrow \mathbb{R}$ , if

$$\exists p < n, C > 0 \text{ s.t. } |f(\mathbf{x})| \leq \frac{C}{\|\mathbf{x} - \mathbf{a}\|^p} \quad \forall \mathbf{x} \in S \setminus \{\mathbf{a}\} \quad (4.5.11)$$

then the improper integral  $\int \cdots \int_S f d\Omega$  is absolutely convergent.

## 5 Vector Calculus

### 5.1 Line Integrals

#### 5.1.1 Arc Length

**Definition 5.1.1.** Let  $C$  be a smooth curve in  $\mathbb{R}^n$  parameterized by  $C^1$  function  $\mathbf{g}$  such that  $\mathbf{g}'(t) \neq \mathbf{0}$  for every appropriate  $t$ .

$$C \equiv \{\mathbf{g}(t) : t \in [a, b]\} \quad (5.1.1)$$

and the **arc length** of  $C$  is defined as

$$\int_C d^n \mathbf{x} \equiv \int_C ds \equiv \int_a^b \|\mathbf{g}'(t)\| dt \quad (5.1.2)$$

**Proposition 5.1.1.** The arc length of a curve  $C$  is an intrinsic property of the geometric object  $C$  and should not depend on the particular parameterization we use.

*Proof.* Let  $\varphi : [c, d] \rightarrow [a, b]$  be a bijection, so that  $\mathbf{h} \equiv \mathbf{g} \circ \varphi$  is also a valid parameterization of  $C$  such that

$$C \equiv \{\mathbf{h}(u) : u \in [c, d]\} \quad (5.1.3)$$

The arc length of  $C$  can be computed using

$$\int_C ds = \int_c^d \|\mathbf{h}'(u)\| du \quad (5.1.4)$$

$$= \int_c^d \|\mathbf{g}'(\varphi(u))\| \times \|\varphi'(u)\| du \quad (5.1.5)$$

$$= \int_a^b \|\mathbf{g}'(t)\| dt \text{ by change of variable formula.} \quad (5.1.6)$$

■

**Remark 5.1.1** (Interpretations). Suppose  $\mathbf{g}$  is a parameterization of  $C$ .

- (i)  $\int_a^b \mathbf{g}'(t) dt = \mathbf{g}(b) - \mathbf{g}(a)$  measures the distance between two endpoints of  $C$ .
- (ii) Choosing a parameterization is effectively choosing an **orientation** for the curve  $C$ .

**Definition 5.1.2.** A function  $\mathbf{g} : [a, b] \rightarrow \mathbb{R}^n$  is called **piecewise smooth** if

- (i) it's *continuous*, and
- (ii) it's derivate exists and is continuous except at finitely many points  $t_j$ , at which the one-sided limits exists.

### 5.1.2 Line Integrals of Scalar Functions

**Definition 5.1.3.** Let smooth curve  $C \subseteq \mathbb{R}^n$ ,  $f : C \rightarrow \mathbb{R}$  and  $\mathbf{g}$  be a parameterization of  $C$ , then

$$\int_C f ds = \int_a^b f(\mathbf{g}(t)) \|\mathbf{g}'(t)\| dt \quad (5.1.7)$$

**Remark 5.1.2.** The line integrals of scalar functions are also independent from the choices of parameterizations.

**Definition 5.1.4.**

$$\text{Average of } f \text{ over } C \equiv \frac{\int_C f ds}{\int_C ds} \quad (5.1.8)$$

### 5.1.3 Line Integrals of Vector Fields

**Definition 5.1.5.** Let smooth  $C \in \mathbb{R}^n$  with parameterization  $\mathbf{g}$  and  $\mathbf{F} : C \rightarrow \mathbb{R}^n$  defined on it, the **line integral** of  $\mathbf{F}$  over  $C$  is defined as

$$\int_C \mathbf{F} \cdot d\mathbf{x} = \int_a^b \mathbf{F}(\mathbf{g}(t)) \cdot \mathbf{g}'(t) dt \quad (5.1.9)$$

**Proposition 5.1.2.** The line integral  $\int_C \mathbf{F} \cdot d\mathbf{x}$  is independent of the parameterization *as long as the orientation is unchanged*.

**Theorem 5.1.1** (The Fundamental Theorem of Line Integral). Let  $f : C \rightarrow \mathbb{R}$  defined on smooth curve  $C$  parameterized by  $\mathbf{g} : [a, b] \rightarrow \mathbb{R}^n$ , then

$$\int_C \nabla f(\mathbf{x}) \cdot d^n \mathbf{x} = f(\mathbf{g}(b)) - f(\mathbf{g}(a)) \quad (5.1.10)$$

*Proof.*

$$\int_C \nabla f(\mathbf{x}) \cdot d^n \mathbf{x} = \int_a^b \frac{\partial f(\mathbf{g}(t))}{\partial \mathbf{g}(t)} \cdot \mathbf{g}'(t) dt \quad (5.1.11)$$

$$= \int_a^b \frac{\partial f(\mathbf{g}(t))}{\partial t} dt = f(\mathbf{g}(b)) - f(\mathbf{g}(a)) \quad (5.1.12)$$

■

### 5.1.4 Rectifiable Curves

**Remark 5.1.3.** Let  $C$  be a curve in  $\mathbb{R}^n$  parameterized by injection  $\mathbf{g} : [a, b] \rightarrow \mathbb{R}^n$  such that  $\mathbf{g}'(t) \neq \mathbf{0}$ . Let  $P$  be a partition of  $[a, b]$ . Denote

$$L_P(C) \equiv \sum_j \|\mathbf{g}(t_j) - \mathbf{g}(t_{j-1})\| \quad (5.1.13)$$

**Definition 5.1.6.** A curve  $C$  is **rectifiable** if the set  $\{L_P(C) : P\}$  is bounded. And the arc length of  $C$  is defined as

$$L(C) \equiv \sup\{L_P(C) : P\} \quad (5.1.14)$$

**Theorem 5.1.2.** The supremum found above,  $L(C)$  is the precisely the arc length of  $C$ :

$$L(C) = \int_a^b \|\mathbf{g}'(t)\| dt \quad (5.1.15)$$

## 5.2 Green's Theorem

### 5.2.1 Preliminary Definitions

**Definition 5.2.1.** A **simple closed curve** is a curve with parameterization  $\mathbf{g} : [a, b] \rightarrow \mathbb{R}^n$  where

- (i)  $\mathbf{g}$  is continuous;
- (ii)  $\mathbf{g}(a) = \mathbf{g}(b)$ ;
- (iii)  $\mathbf{g}$  is injective with its domain restricted to  $(a, b)$ .

**Definition 5.2.2.** A *simple closed curve* is **piecewise smooth** if it has a parameterization  $\mathbf{g}$  such that

- (i)  $\mathbf{g}$  is continuously differentiable with  $\mathbf{g}'(t) \neq \mathbf{0}$  except finitely many breakpoints;
- (ii)  $\mathbf{g}'(t)$  is *one side continuous* at breakpoints of the curve.

**Definition 5.2.3.** A **regular region**  $S \subseteq \mathbb{R}^n$  is a set satisfying both

- (i)  $S$  is compact;
- (ii)  $\overline{S^{int}} = S$ .

**Definition 5.2.4.** Let  $S \subseteq \mathbb{R}^2$ ,  $S$  has **piecewise smooth boundary** if  $\partial S$  consists of one or more *disjoint, piecewise smooth, simple closed curve*.

**Definition 5.2.5.** Let  $S \subseteq \mathbb{R}^2$ , then **positive orientation** on  $\partial S$  is the orientation on each of the closed curves that make up the boundary such that the region is on the *left* with respect to the positive direction on the curve.

**Theorem 5.2.1** (Green's Theorem). Suppose  $S \subseteq \mathbb{R}^2$  is a regular region with piecewise smooth region  $\partial S$ . Suppose  $\mathbf{F}$  is a  $C^1$  vector field defined on  $\overline{S}$ , then

$$\int_{\partial S} \mathbf{F} \cdot d\mathbf{x} = \iint_S \left( \frac{\partial F_2}{\partial x_1} - \frac{\partial F_1}{\partial x_2} \right) dA \quad (5.2.1)$$



**Corollary 5.2.1.** Suppose  $S$  is a regular region in  $\mathbb{R}^2$  with piecewise smooth boundary  $\partial S$ , and let  $\mathbf{n}(\mathbf{x})$  be the *unit outward normal* vector to  $\partial S$  at  $\mathbf{x} \in \partial S$ . Suppose also that  $\mathbf{F}$  is a vector field defined on  $\overline{S}$ , then

$$\int_{\partial S} \mathbf{F} \cdot \mathbf{n} \, ds = \iint_S \left( \frac{\partial F_1}{\partial x_1} + \frac{\partial F_2}{\partial x_2} \right) dA \quad (5.2.2)$$

*Proof.* Let  $\mathbf{g}(t)$  be a parameterization of boundary  $\partial S$ . Then the tangent vector would be  $\mathbf{g}'(t)$  and we can conclude the *outer normal vector*  $\mathbf{n}$  is  $\frac{(g'_2(t), -g'_1(t))}{\|(g'_2(t), -g'_1(t))\|}$ . Then

$$\int_{\partial S} \mathbf{F} \cdot \mathbf{n} \, ds = \int_T \mathbf{F} \circ \mathbf{g} \cdot \frac{(g'_2(t), -g'_1(t))}{\|(g'_2(t), -g'_1(t))\|} \|\mathbf{g}'(t)\| \, dt \quad (5.2.3)$$

$$= \int_T F_1 g'_2(t) - F_2 g'_1(t) \, dt \quad (5.2.4)$$

$$= \int_T \begin{pmatrix} -F_2 \\ F_1 \end{pmatrix} \cdot \begin{pmatrix} g'_1(t) \\ g'_2(t) \end{pmatrix} \, dt \quad (5.2.5)$$

$$= \int_{\partial S} \begin{pmatrix} -F_2 \\ F_1 \end{pmatrix} \cdot d^2 \mathbf{x} \quad (5.2.6)$$

$$= \iint_S \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} \, dA \text{ By Green's Theorem} \quad (5.2.7)$$

■

## 5.3 Surface Integrals

### 5.3.1 Surface Areas and Surface Integrals

**Definition 5.3.1.** Suppose  $S$  is a surface in  $\mathbb{R}^3$  and parameterized by

$$\mathbf{G}(\mathbf{u}) : R \rightarrow S \quad (5.3.1)$$

where  $\text{rank}(D\mathbf{G}(\mathbf{u})) = 2$  for every  $\mathbf{u} \in R \setminus Z$  where  $Z$  is a probably empty set with zero content. If  $\|\frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v}\|$  is integrable, then

$$\text{Area}(S) \equiv \iint_R \left\| \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right\| d\Theta \quad (5.3.2)$$

**Definition 5.3.2.** Let  $f : S \rightarrow \mathbb{R}$  be a real-valued continuous function defined on a super set of  $S$ , the **integral of a real-valued function on a surface** is defined as

$$\iint_S f(\mathbf{x}) \, dA \equiv \iint_R f(\mathbf{G}(\mathbf{u})) \left\| \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right\| d\Theta \quad (5.3.3)$$

**Definition 5.3.3.** Let  $\mathbf{F} : S \rightarrow \mathbb{R}^3$  be a continuous vector field defined on a super set of  $S$ , the **integral of vector field on a surface** is defined as

$$\iint_S \mathbf{F}(\mathbf{x}) \cdot \mathbf{n} \, dA \equiv \iint_R \mathbf{F}(\mathbf{G}(\mathbf{u})) \cdot \left( \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right) d\Theta \quad (5.3.4)$$

**Remark 5.3.1.** Surface integrals of **real-valued functions** are independent of the choice of parametrization.

**Remark 5.3.2.** But the choice of parameterization can change the sign of surface integrals of vector fields. We need to choose the direction of the normal,  $\mathbf{n}$ .

**Definition 5.3.4.** Let  $S \subseteq \mathbb{R}^3$  be a two dimensional sub-manifold, and  $f$  is a real-valued function defined on a super set of  $S$ . Define the **average of  $f$  over  $S$**  as

$$\text{aver of } f \text{ over } S \equiv \frac{\iint_S f \, dA}{\iint_S 1 \, dA} \quad (5.3.5)$$

**Remark 5.3.3.** A note on the relation between integrals of a vector field and a real-valued function. The surface of vector field  $\mathbf{F}$  on  $S$  is defined by *reducing  $\mathbf{F}$  to a real-valued function  $\mathbf{F} \cdot \mathbf{n}$*  and then follow the definition of conventional real-valued function on  $S$ . Define  $h \equiv \mathbf{F} \cdot \mathbf{n}$ ,

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, dA = \iint_S h \, dA \quad (5.3.6)$$

$$\equiv \iint_R h(\mathbf{G}(\mathbf{u})) \left\| \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right\| d\Theta \quad (5.3.7)$$

$$= \iint_R \mathbf{F}(\mathbf{G}(\mathbf{u})) \cdot \mathbf{n}(\mathbf{G}(\mathbf{u})) \left\| \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right\| d\Theta \quad (5.3.8)$$

$$= \iint_R \mathbf{F}(\mathbf{G}(\mathbf{u})) \cdot \frac{\frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v}}{\left\| \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right\|} \left\| \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right\| d\Theta \quad (5.3.9)$$

$$= \iint_R \mathbf{F}(\mathbf{G}(\mathbf{u})) \cdot \left( \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right) d\Theta \quad (5.3.10)$$

### 5.3.2 An invariance property

**Remark 5.3.4.** As mentioned above, given  $\mathbf{n}(\mathbf{x})$  fixed, we can define the surface integral of vector field as the surface integral of a real-valued function defined as  $h(\mathbf{x}) \equiv \mathbf{F}(\mathbf{x}) \cdot \mathbf{n}(\mathbf{x})$ . And as argued before, one  $\mathbf{n}$  is fixed (i.e. orientation is fixed), the value of integral is deterministic. Therefore we can conclude the integral of a vector field  $\mathbf{F}$  over a surface  $S$  depends on the orientation of  $S$  but otherwise independent of the parameterization.

**Remark 5.3.5.** Let  $S \subseteq \mathbb{R}^3$  be a two dimensional sub-manifold parameterized by  $\mathbf{G} : R \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}^3$  such that  $\text{rank}(\mathbf{G}(\mathbf{u})) = 2$  for all but zero-content sets on its domain.

Let  $\varphi : W \subseteq \mathbb{R}^2 \rightarrow R$  be a bijection such that  $\mathbf{H} \equiv \mathbf{G} \circ \varphi : W \rightarrow \mathbb{R}^3$  is another parameterization of  $S$ .

Now consider the integral of vector field  $\mathbf{F}$  under parameterization  $\mathbf{H}$ ,

$$\iint_S \mathbf{F} \cdot \mathbf{u} \, dA = \iint_W \mathbf{F}(\mathbf{H}) \cdot \left( \frac{\partial \mathbf{H}}{\partial s} \times \frac{\partial \mathbf{H}}{\partial t} \right) d\Theta \quad (5.3.11)$$

$$= \iint_W \mathbf{F} \circ \mathbf{G} \circ \varphi \cdot \left( \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right) \text{det } D\varphi \, d\Theta \quad (5.3.12)$$

$$= \pm \iint_R \mathbf{F} \circ \mathbf{G} \cdot \left( \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right) d\Theta \text{ (change of variable)} \quad (5.3.13)$$

**Theorem 5.3.1** (Invariance). Let  $\mathbf{G} : R \rightarrow \mathbb{R}^3$  and  $\mathbf{H} \equiv \mathbf{G} \circ \varphi : W \rightarrow \mathbb{R}^3$  be two parameterizations of  $S$ , then

$$\iint_R f \circ \mathbf{G} \left\| \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right\| d\Theta = \iint_W f \circ \mathbf{H} \left\| \frac{\partial \mathbf{H}}{\partial s} \times \frac{\partial \mathbf{H}}{\partial t} \right\| d\Theta \quad (5.3.14)$$

and

$$\iint_R \mathbf{F} \circ \mathbf{G} \cdot \left( \frac{\partial \mathbf{G}}{\partial u} \times \frac{\partial \mathbf{G}}{\partial v} \right) d\Theta = \pm \iint_W \mathbf{F} \circ \mathbf{H} \cdot \left( \frac{\partial \mathbf{H}}{\partial u} \times \frac{\partial \mathbf{H}}{\partial v} \right) d\Theta \quad (5.3.15)$$

### 5.3.3 Volume and Area

**Theorem 5.3.2.** Let  $R$  be an arbitrary regular region in  $\mathbb{R}^3$ , and let  $S$  be the boundary surface of  $R$ , define

$$S_h \equiv \{\mathbf{x} + \delta \mathbf{n} : \mathbf{x} \in S \wedge \delta \in [0, h]\} \quad (5.3.16)$$

where  $S_h$  can be interpreted as *a shell of region  $R$  with thickness  $h$* . Then the surface area of  $S$  is

$$\text{area}(S) = \lim_{h \rightarrow 0} \frac{|S_h|}{h} \quad (5.3.17)$$

## 5.4 Divergence, Gradient and Curl

**Definition 5.4.1.** Let  $U \subseteq \mathbb{R}^n$  be an open set, and define real-valued function  $f : U \rightarrow \mathbb{R}$  and vector field  $\mathbf{F} : U \rightarrow \mathbb{R}^n$ . Then we define

1. The **gradient** of  $f$  as  $\nabla f$ ;
2. The **divergence** of  $\mathbf{F}$  as  $\nabla \cdot \mathbf{F}$ ;
3. The **curl** of  $\mathbf{F}$  as  $\nabla \times \mathbf{F}$ .

**Definition 5.4.2.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a  $C^1$  real-valued function, define the **Laplacian** of  $f$  as a mapping from *real-valued functional space* to *real-valued functional space* defined as

$$\text{div}(\text{grad})f \equiv \sum_j \partial_j^2 f = \Delta f = \nabla^2 f \quad (5.4.1)$$

**Theorem 5.4.1.** For every  $C^2$  real valued function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ ,

$$\text{curl}(\text{grad}f) = \mathbf{0} \quad (5.4.2)$$

For every  $C^2$  vector field defined in  $\mathbb{R}^3$  or a subset of it,

$$\text{div}(\text{curl}\mathbf{F}) = 0 \quad (5.4.3)$$

Note that the domain of  $f$  and  $\mathbf{F}$  must be  $\mathbb{R}^3$  or a subset of it, otherwise the curl operation is not well-defined.

**Theorem 5.4.2** (Product rules).

$$\text{grad}(fg) = f \text{ grad}g + \text{grad}f \cdot g \quad (5.4.4)$$

$$\text{div}(f\mathbf{G}) = f \text{ div}\mathbf{G} + \text{grad}f \cdot \mathbf{G} \quad (5.4.5)$$

$$\text{curl}(f\mathbf{G}) = f \text{ curl}\mathbf{G} + \text{grad}f \times \mathbf{G} \quad (5.4.6)$$

## 5.5 Divergence Theorem

**Remark 5.5.1.** vector field integral on boundary (2-dimensional sub-manifold) of region in  $\mathbb{R}^3$  ( $\mathbf{F} \cdot \mathbf{n} \, dA$  2-form) and scalar valued function ( $\text{div}(\mathbf{F}) \, dV$  3-form) in a region (3-dimensional sub-manifold).

**Theorem 5.5.1** (Divergence Theorem). Let  $R \subseteq \mathbb{R}^3$  be a *regular region* with *piece-wise smooth* boundary  $\partial S$ . And  $\mathbf{n}$  is the *outer normal vector* on  $\partial S$ , then,

$$\iint_{\partial S} \mathbf{F} \cdot \mathbf{n} \, dA = \iiint_S \operatorname{div}(\mathbf{F}) \, dV \quad (5.5.1)$$

*Proof.*

**Definition 5.5.1.** A region  $R \subseteq \mathbb{R}^3$  is said to be *xy-simple* if and only if it can be expressed as the following form

$$R = \{(x, y, z) \in \mathbb{R}^3 : (x, y) \in W, \varphi_1(x, y) \leq z \leq \varphi_2(x, y)\} \quad (5.5.2)$$

Suppose  $S$  is simple in terms of all combinations of  $x, y, z$ .

Then

$$\iint_{\partial S} \mathbf{F} \cdot \mathbf{n} \, dA = \iint_{\partial S} F_1 n_1 + F_2 n_2 + F_3 n_3 \, dA \quad (5.5.3)$$

Consider  $\iint_{\partial S} F_3 n_3 \, dA$ , since  $R$  is *xy-simple*,

$$\iint_{\partial S} F_3 n_3 \, dA = \iint_{\partial S} F_3 \mathbf{k} \cdot \mathbf{n} \, dA \quad (5.5.4)$$

Note that except the bottom and top sides, which are parameterized by  $\mathbf{G}_1(x, y) = (x, y, \varphi_1(x, y))$  and  $\mathbf{G}_2(x, y) = (x, y, \varphi_2(x, y))$ , the outer normal vector of those region has form  $(\cdot, \cdot, 0)$ , and therefore  $\mathbf{n} \cdot \mathbf{k} = 0$  for every  $\mathbf{x}$  on those regions, and contribute nothing to the integral.

Therefore, to evaluate  $\iint_{\partial S} F_3 \mathbf{k} \cdot \mathbf{n} \, dA$ , we only need to consider the upper and bottom surfaces. Also note that  $\mathbf{n}$  has opposite  $z$  component on those two surfaces.

Moreover, the undirected  $\mathbf{n}$  on those two surfaces is

$$\tilde{\mathbf{n}} = \begin{pmatrix} 1 \\ 0 \\ \partial_x \varphi_i \end{pmatrix} \times \begin{pmatrix} 0 \\ 1 \\ \partial_y \varphi_i \end{pmatrix} = \begin{pmatrix} -\partial_x \varphi_i \\ \partial_x \varphi_i - \partial_y \varphi_i \\ 1 \end{pmatrix} \quad (5.5.5)$$

$$\implies \iint_{\partial S} F_3 \mathbf{k} \cdot \mathbf{n} \, dA = \iint_{\partial S} F_3 \, dA \quad (5.5.6)$$

$$= \iint_{\text{upper } \partial S} F_3 \, dA - \iint_{\text{lower } \partial S} F_3 \, dA \quad (5.5.7)$$

$$= \iint_W F_3(x, y, \varphi_2(x, y)) \, dx dy - \iint_W F_3(x, y, \varphi_1(x, y)) \, dx dy \quad (5.5.8)$$

$$= \iint_W \int_{\varphi_1(x, y)}^{\varphi_2(x, y)} \partial_3 F_3 \, dz dx dy = \iiint_S \partial_3 F_3 \, dV \quad (5.5.9)$$

We can prove the equalities involving the other two components, and the proof is completed by the fact that any open set in  $\mathbb{R}^n$  can be written as a countable union of *almost disjoint* cubes, which are simple and the boundary of  $S$  has zero content.  $\blacksquare$

**Proposition 5.5.1** (Geometric Interpretation of Divergence). Let  $S \subset \mathbb{R}^3$ ,  $\mathbf{F} : S \rightarrow \mathbb{R}^3$ ,  $\mathbf{a} \in S$ ,

$$\operatorname{div}(\mathbf{F})(\mathbf{a}) = \lim_{r \rightarrow 0} \frac{3}{4\pi r^2} \iiint_{\mathcal{B}(\mathbf{a}, r)} \operatorname{div}(\mathbf{F})(\mathbf{x}) \, d\mathbf{x} \quad (5.5.10)$$

$$= \lim_{r \rightarrow 0} \frac{3}{4\pi r^2} \underbrace{\iint_{\partial \mathcal{B}(\mathbf{a}, r)} \mathbf{F} \cdot \mathbf{n} \, d\mathbf{x}}_{\text{flux through boundary}} \quad (5.5.11)$$

thus  $\operatorname{div}(\mathbf{F})(\mathbf{a}) > 0$  if and only if at point  $\mathbf{a}$ , matters are flowing away from this point.

**Corollary 5.5.1** (Green's Formula). Suppose  $R \subset \mathbb{R}^3$  and  $f, g : R \rightarrow \mathbb{R}$  are  $C^2$  functions, then

$$\iint_{\partial S} f \nabla g \cdot \mathbf{n} \, dA = \iiint_S \nabla f \cdot \nabla g + f \nabla^2 g \, dV \quad (5.5.12)$$

$$\iint_{\partial S} (f \nabla g - g \nabla f) \, dA = \iiint_S (f \nabla^2 g - g \nabla^2 f) \, dV \quad (5.5.13)$$

*Proof.*

$$\iint_{\partial S} f \nabla g \cdot \mathbf{n} \, dA = \iiint_S \operatorname{div}(f \nabla g) \, dV \quad (5.5.14)$$

$$= \iiint_S f \operatorname{div}(\nabla g) + \nabla f \cdot \nabla g \, dV = \iiint_S f \nabla^2 g + \nabla f \cdot \nabla g \, dV \quad (5.5.15)$$

The second formula can be proved directly using divergence theorem the first formula. ■

## 5.6 Stokes Theorem

### 5.6.1 Stokes Theorem in $\mathbb{R}^3$

**Theorem 5.6.1** (Stokes Theorem, Special Case). Let  $S$  be a 2 dimensional sub-manifold in  $\mathbb{R}^3$ , and let  $\mathbf{F}$  be a vector field defined on some neighbour of  $S$ , then

$$\int_{\partial S} \mathbf{F} \cdot d\mathbf{x} = \iint_S \operatorname{curl}(\mathbf{F}) \cdot \mathbf{n} \, dA \quad (5.6.1)$$

**Remark 5.6.1.** In above theorem,  $\omega \equiv \mathbf{F} \cdot d\mathbf{x}$  is a 1-form in  $\mathbb{R}^3$  and  $d\omega \equiv \operatorname{curl}(\mathbf{F}) \cdot \mathbf{n} \, dA$  is a 2-form in  $\mathbb{R}^3$ .

**Corollary 5.6.1.** Let  $S$  be a closed surface in  $\mathbb{R}^3$ , that's,  $\partial S = \emptyset$ , and let  $\mathbf{n}$  denote the outer normal vector, and  $\mathbf{F}$  is a  $C^1$  vector field. Then

$$\iint_S \operatorname{curl}(\mathbf{F}) \cdot \mathbf{n} \, dA = 0 \quad (5.6.2)$$

*Proof.* We can construct a *small* simple closed curve  $C$  on  $S$  and divide  $S$  into two regions sharing the same boundary. And note that given orientation fixed on  $S$ , the orientation on  $\partial S_1$  and  $\partial S_2$  are opposite. Then

$$\iint_S \operatorname{curl}(\mathbf{F}) \cdot \mathbf{n} \, dA = \iint_{S_1} \operatorname{curl}(\mathbf{F}) \cdot \mathbf{n} \, dA + \iint_{S_2} \operatorname{curl}(\mathbf{F}) \cdot \mathbf{n} \, dA \quad (5.6.3)$$

$$= \int_{\partial S_1} \mathbf{F} \cdot d\mathbf{x} - \int_{\partial S_2} \mathbf{F} \cdot d\mathbf{x} = \int_C \mathbf{F} \cdot d\mathbf{x} - \int_C \mathbf{F} \cdot d\mathbf{x} = 0 \quad (5.6.4)$$

■

**Proposition 5.6.1** (Geometric Interpretation of Curl). Let  $R \subset \mathbb{R}^3$  be a 2 dimensional sub-manifold with  $\mathbf{n}$  as outer normal vector on it, and  $\mathbf{a} \in R$ ,

$$\operatorname{curl}(F)(\mathbf{a}) \cdot \mathbf{n}(\mathbf{a}) = \lim_{r \rightarrow 0} \frac{1}{2\pi r^2} \iint_{\mathcal{D}(\mathbf{a}, r)} \operatorname{curl}(F)(\mathbf{x}) \cdot \mathbf{n}(\mathbf{x}) \, dA \quad (5.6.5)$$

$$= \lim_{r \rightarrow 0} \frac{1}{2\pi r^2} \int_{\partial \mathcal{D}(\mathbf{a}, r)} \mathbf{F} \cdot d\mathbf{x} \quad (5.6.6)$$

$$(5.6.7)$$

If we think of  $\mathbf{F}$  as a force field, then  $\int_{\partial\mathcal{D}(\mathbf{a},r)} \mathbf{F} \cdot d\mathbf{x}$  represents the work done by  $\mathbf{F}$  on a particle moves around  $\partial\mathcal{D}(\mathbf{a},r)$ . Thus  $\text{curl}(\mathbf{F}) \cdot \mathbf{u}$  represents the **tendency of the force  $\mathbf{F}$  to push the particle around  $\partial\mathcal{D}(\mathbf{a},r)$  in a direction compatible with  $\mathbf{n}$ .**

### 5.6.2 The Generalization

**Proposition 5.6.2** (Properties of Exterior Products). Let  $\alpha_1, \alpha_2$  and  $\beta$  be 1-forms on  $\mathbb{R}^n$  and  $f_1, f_2$  are continuous functions defined on  $\mathbb{R}^n$ ,

#### 1. Distributive

$$(f_1\alpha_1 + f_2\alpha_2) \wedge \beta = f_1(\alpha_1 \wedge \beta) + f_2(\alpha_2 \wedge \beta) \quad (5.6.8)$$

$$\beta \wedge (f_1\alpha_1 + f_2\alpha_2) = f_1(\beta \wedge \alpha_1) + f_2(\beta \wedge \alpha_2) \quad (5.6.9)$$

#### 2. Anti-commutative

$$\beta \wedge \alpha = -\alpha \wedge \beta \quad (5.6.10)$$

**Theorem 5.6.2** (Divergence Theorem in  $\mathbb{R}^n$ ). Let  $R$  be a regular region in  $\mathbb{R}^n$  bounded by a piecewise smooth hyper-surface  $\partial R$ . Note here  $R$  is a  $n$  dimensional sub-manifold and  $\partial R$  is a  $n-1$  dimension sub-manifold. Then

$$\int \cdots \int_{\partial R} \mathbf{F} \cdot \mathbf{n} dV^{n-1} = \iiint_R \text{div}(\mathbf{F}) dV^n \quad (5.6.11)$$

where if  $\partial R$  is parameterized by  $\mathbf{G}(u_1, \dots, u_{n-1})$ , then

$$\mathbf{n} dV^{n-1} = \det \begin{pmatrix} \mathbf{e}_1 & \cdots & \mathbf{e}_n \\ \partial_1 G_1 & \cdots & \partial_1 G_n \\ \vdots & & \vdots \\ \partial_{n-1} G_1 & \cdots & \partial_{n-1} G_n \end{pmatrix} \quad (5.6.12)$$

**Definition 5.6.1.** A 0-form on  $\mathbb{R}^n$  is a real valued function  $f$ .

**Remark 5.6.2.** While writing the basis elements  $dx_i \wedge dx_j$  with the variables in *cyclic order*. That's  $dx$  before  $dy$  before  $dz$  before  $dx$  in  $\mathbb{R}^3$  case.

**Definition 5.6.2.** A  $k$ -form in  $\mathbb{R}^n$  takes the expression of linear combination of  $C(n, k)$  basis elements  $\{\beta_i\}_i$ .

**Example 5.6.1.** A 2-form  $\omega$  in  $\mathbb{R}^3$  can be expressed using a 3-element basis

$$\omega = \sum_{1 \leq i < j \leq 3} C_{ij}(\mathbf{x}) \beta_{ij} \quad (5.6.13)$$

$$\beta_{ij} \in \{dx \wedge dy, dy \wedge dz, dx \wedge dz\} \quad (5.6.14)$$

**Definition 5.6.3.** Let  $\omega = \sum_{j=1}^{C(n,k)} f_j \beta_j$  be a  $k$ -form in  $\mathbb{R}^n$ , then it's **exterior derivative** is defined to be the  $(k+1)$ -form in  $\mathbb{R}^n$  defined as

$$d\omega \equiv \sum_j df_j \wedge \beta_j \quad (5.6.15)$$

where  $df_j$  can be computed using *total derivative*.

**Example 5.6.2.** In  $\mathbb{R}^3$ , the *exterior derivative* for a 0-form  $f$  is its **gradient**, which is a 1-form. And the exterior derivate of a 1-form in  $\mathbb{R}^3$  is its curl

$$\omega := F_1 dx + F_2 dy + F_3 dz \quad (5.6.16)$$

$$\implies d\omega = dF_1 \wedge dx + dF_2 \wedge dy + dF_3 \wedge dz \quad (5.6.17)$$

$$= (\partial_1 F_1 dx + \partial_2 F_1 dy + \partial_3 F_1 dz) \wedge dx \quad (5.6.18)$$

$$+ (\partial_1 F_2 dx + \partial_2 F_2 dy + \partial_3 F_2 dz) \wedge dy \quad (5.6.19)$$

$$+ (\partial_1 F_3 dx + \partial_2 F_3 dy + \partial_3 F_3 dz) \wedge dz \quad (5.6.20)$$

$$= (\partial_1 F_2 - \partial_2 F_1) dx \wedge dy + (\partial_2 F_3 - \partial_3 F_2) dy \wedge dz + (\partial_3 F_1 - \partial_1 F_3) dz \wedge dx \quad (5.6.21)$$

$$= \text{curl}(\mathbf{F}) \quad (5.6.22)$$

The exterior derivate of a 2-form in  $\mathbb{R}^3$  is its divergence

$$\omega := A dy \wedge dz + B dz \wedge dx + C dx \wedge dy \quad (5.6.23)$$

$$\implies d\omega = (\partial_1 A dx + \partial_2 A dy + \partial_3 A dz) \wedge dy \wedge dz \quad (5.6.24)$$

$$+ (\partial_1 B dx + \partial_2 B dy + \partial_3 B dz) \wedge dz \wedge dx \quad (5.6.25)$$

$$+ (\partial_1 C dx + \partial_2 C dy + \partial_3 C dz) \wedge dx \wedge dy \quad (5.6.26)$$

$$= (\partial_1 A + \partial_2 B + \partial_3 C) dx \wedge dy \wedge dz \quad (5.6.27)$$

$$= \text{div}(\mathbf{F}) \quad (5.6.28)$$

**Theorem 5.6.3** (Stokes Theorem, 5.77). Let  $M$  be a smooth, oriented  $k$  dimensional sub-manifold of  $\mathbb{R}^n$  with a piecewise smooth boundary  $\partial M$ , and let  $\partial M$  carry the orientation that is (in a suitable sense) compatible with the one on  $M$ . If  $\omega$  is a  $(k-1)$ -form of class  $C^1$  on an open set containing  $M$ , then

$$\int \cdots \int_{\partial M} \omega = \iint \cdots \int_M d\omega \quad (5.6.29)$$

**Theorem 5.6.4.** The *boundary* of a (smoothly bounded) region  $M$  in a  $k$  dimensional manifold is a  $(k-1)$  dimensional manifold with no boundary.

That's let  $M$  be a  $k$  dimensional manifold with piecewise smooth boundary  $\partial M$ , then

$$\partial(\partial M) = \emptyset \quad (5.6.30)$$

**Theorem 5.6.5.** For any  $k$ -form  $\omega$  on  $\mathbb{R}^n$ ,

$$d(d\omega) = 0 \quad (5.6.31)$$

*Proof.* Let  $M$  be a  $k$  dimensional sub-manifold in  $\mathbb{R}^n$  with piecewise smooth boundary, and  $\omega$  is a  $(k-2)$ -form on  $\mathbb{R}^n$ , so  $d(d\omega)$  is a  $k$ -form on  $\mathbb{R}^n$ . And

$$\iiint \cdots \int_M d(d\omega) = \iint \cdots \int_{\partial M} d\omega \quad (5.6.32)$$

$$= \int \cdots \int_{\partial(\partial M)} \omega = \int \cdots \int_{\emptyset} \omega = 0 \quad (5.6.33)$$

■