

MAT237: Lecture Notes

Advanced Calculus

Tianyu Du

September 6, 2018

Contents

1	Lecture 1 September 6 2018	2
1.1	The Geometry of Euclidean Space	2
1.2	Subspaces of \mathbb{R}^n	3
1.3	Cross Product	3
1.4	Functions of Several Variables	4

1 Lecture 1 September 6 2018

1.1 The Geometry of Euclidean Space

Example 1.1. Consider $(1, 2) \in \mathbb{R}^2$ as a point or a vector.

Remark 1.1. All vectors in this course are considered as column vectors. Reasoning: suppose a linear function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then the transformation can be implemented as

$$f(\vec{x}) = A\vec{x}, \quad A \in mn$$

if \vec{x} is a column vector.

Let $\vec{a}, \vec{b} \in \mathbb{R}^n$, the **dot product** $\cdot : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is defined as,

$$ab = \sum_i a_i b_i$$

Let $\vec{a} \in \mathbb{R}^n$, the **Euclidean norm** $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined as

$$\|\vec{a}\| = \sqrt{ab}$$

Interpretation the Euclidean norm of \vec{a} , $\|\vec{a}\|$ is the length of \vec{a} , or the distance of \vec{a} from the origin. And $\|\vec{a} - \vec{b}\|$ is the distance from \vec{a} to \vec{b} .

Two vectors $\vec{a}, \vec{b} \in \mathbb{R}^n$ is **orthogonal** if and only if

$$ab = 0$$

Theorem 1.1. (Cauchy Schwarz inequality)

$$|ab| \leq \|\vec{a}\| \|\vec{b}\|$$

Theorem 1.2. (Triangle inequality)

$$\|\vec{a} + \vec{b}\| \leq \|\vec{a}\| + \|\vec{b}\|$$

Theorem 1.3.

$$ab = \|\vec{a}\| \|\vec{b}\| \cos \theta$$

where θ is the angle between \vec{a} and \vec{b}

If $\vec{u} \in \mathbb{R}^n$ is a **unit vector** if

$$\|\vec{u}\| = 1$$

The **projection** of \vec{a} onto the line through \vec{u} is defined as

$$(ua)\vec{u}$$

1.2 Subspaces of \mathbb{R}^n

A subspace V of \mathbb{R}^n is a subset of \mathbb{R}^n such that

$$\vec{a}, \vec{b} \in V \wedge c_1, c_2 \in \mathbb{R} \implies c_1\vec{a} + c_2\vec{b} \in V$$

Example 1.2. Suppose

$$A = \begin{pmatrix} 1 & 3 \\ 2 & 7 \\ -1 & 0 \end{pmatrix}$$

And consider

$$V = \{A\vec{x} : \vec{x} \in \mathbb{R}^n\}$$

V is a subspace with dimension 2.

Theorem 1.4. Let $A \in mn$ with $m > n$ and columns are independent then $V = \{A\vec{x} : \vec{x} \in \mathbb{R}^n\}$ is a n -dimensional subspace of \mathbb{R}^n .

Example 1.3. Consider

$$A = \begin{pmatrix} 3 & 1 & 0 \\ 1 & 9 & -2 \end{pmatrix}$$

and

$$V = \{\vec{x} \in \mathbb{R}^3 : A\vec{x} = \vec{0}\}$$

Then V is a 1-dimensional subspace of \mathbb{R}^3 .

Theorem 1.5. $A \in mn$ and $m < n$ and rows are linearly independent, then $\{\vec{x} \in \mathbb{R}^n : A\vec{x} = \vec{0}\}$ is a $(n - m)$ dimensional subspace.

1.3 Cross Product

(*Only available in \mathbb{R}^3*) is a way to multiplying two vectors in \mathbb{R}^3 to get another vector in \mathbb{R}^3 . Let $\vec{a}, \vec{b} \in \mathbb{R}^3$ then the cross product $\times : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is defined as

$$\vec{a} \times \vec{b} := \det \begin{pmatrix} \vec{i} & \vec{j} & \vec{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{pmatrix}$$

$$\text{where } \vec{i} = (1, 0, 0), \vec{j} = (0, 1, 0), \vec{k} = (0, 0, 1)$$

Remark 1.2. $\vec{a} \times \vec{b}$ is the vector such that

1. orthogonal to both \vec{a} and \vec{b} .
2. it's length is $||\vec{a}|| ||\vec{b}|| \sin \theta$.

Let $\vec{a}, \vec{b} \in \mathbb{R}^3$, then

1. $\vec{a} \times \vec{b} = \vec{b} \times \vec{a}$
2. $\vec{a} \times \vec{a} = \vec{0}$
3. $(c_1 \vec{a}_1 + c_2 \vec{a}_2) \times \vec{b} = c_1(\vec{a}_1 \times \vec{b}) + c_2(\vec{a}_2 \times \vec{b})$
4. $(\vec{a} \times \vec{b}) \times \vec{c} \neq \vec{a} \times (\vec{b} \times \vec{c})$

1.4 Functions of Several Variables

Remark 1.3. Idea of differential calculus: more general functions can then be approximated by linear functions.

Consider function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, the graph of f is

$$\{(x, y, z) : z = f(x, y)\} \subseteq \mathbb{R}^3$$