

CSC412/2506 Winter 2020: Probabilistic Learning and Reasoning

Tianyu Du

February 8, 2020

Contents

1	Introduction	2
2	Probabilistic Models	2
3	Directed Graphical Models	2
4	Exact Inference	2
4.1	Variable Elimination	2
4.2	Intermediate Factors	2
4.3	Sum-Product Inference	3
4.4	Complexity of Variable Elimination Ordering	3
5	Message passing, Hidden Markov Models, and Sampling	4
6	Sampling and Monte Carlo Methods	4

1 Introduction

2 Probabilistic Models

3 Directed Graphical Models

4 Exact Inference

Notation 4.1. Let X denote the set of all random variables in the model, and

1. X_E = The observed evidence;
2. X_F = The unobserved variable we want to infer;
3. $X_R = X - \{X_F, X_E\}$ = Remaining variables, extraneous to query.

The model defines the joint distribution of all random variables:

$$p(X_E, X_F, X_R) \quad (4.1)$$

Definition 4.1. The joint distribution over evidence and subject of inference is

$$p(X_F, X_E) = \sum_{X_R} p(X_F, X_E, X_R) \quad (4.2)$$

Definition 4.2. The conditional probability distribution for inference given evidence is


$$p(X_F|X_E) = \frac{p(X_F, X_E)}{p(X_E)} = \frac{p(X_F, X_E)}{\sum_{X_F} p(X_F, X_E)} \quad (4.3)$$

Definition 4.3. The distribution of evidence can be computed as

$$p(X_E) = \sum_{X_F, X_R} p(X_F, X_E, X_R) \quad (4.4)$$

4.1 Variable Elimination

4.2 Intermediate Factors



$$p(A, B, C) = \sum_X p(X)p(A|X)p(B|A)p(C|B, X) \quad (4.5)$$

$$= p(B|A) \underbrace{\sum_X p(X)p(A|X)p(C|B, X)}_{\text{unnormalized}} \quad (4.6)$$

Definition 4.4. A **factor** ϕ describes the local relation between random variables, meanwhile, $\int d\phi$ is not necessarily one.

Remark 4.1. Let $X_\ell \subseteq X$ be a group of local random variables, then $p(X_\ell)$ is automatically a factor $\phi(X_\ell)$.

$$p(A, B, C) = \sum_X \underbrace{p(X)p(A|X)p(B|A)p(C|B, X)}_{\text{from graphical representation}} \quad (4.7)$$

$$= \sum_X \underbrace{\phi(X)\phi(A, X)\phi(A, B)\phi(X, B, C)}_{\text{factor representation}} \quad (4.8)$$

$$= \phi(A, B) \sum_X \phi(X)\phi(A, X)\phi(X, B, C) \quad (4.9)$$

$$= \phi(A, B) \underbrace{\tau(A, B, C)}_{\text{another factor}} \quad (4.10)$$

4.3 Sum-Product Inference

Theorem 4.1. Consider a graphical model with random variables $X = Y \cup Z$. For an random variable Y in a directed or undirected model, $P(Y)$ can be computed using the **sum-product**

$$\tau(Y) = \sum_z \prod_{\phi \in \Phi} \phi(\text{Scope}[\phi] \cap Z, \text{Scope}[\phi] \cap Y) \quad (4.11)$$

where Φ is a set of factors.

Remark 4.2. For directed models,

$$\Phi = \{\phi_{x_i}\}_{i=1}^N = \{p(x_i | \text{parents}(x_i))\}_{i=1}^N \quad (4.12)$$

4.4 Complexity of Variable Elimination Ordering

Theorem 4.2. The complexity of the variable elimination algorithm is

$$\mathcal{O}(mk^{N_{max}}) \quad (4.13)$$

where

- (i) m is the number of initial factors $|\Phi|$;
- (ii) k is the number of states each random variable takes, assumed to be equal;
- (iii) N_i is the number of random variables within each summation;
- (iv) $N_{max} = \max_i N_i$.

5 Message passing, Hidden Markov Models, and Sampling

6 Sampling and Monte Carlo Methods