

Notes on Measure Theory

Tianyu Du

June 8, 2020

1 Sigma Algebra

Definition 1.1. For a set X , a set $\mathcal{A} \subseteq \mathcal{P}(X)$ is a **σ -algebra** if it satisfies the following properties:

1. $\emptyset, X \in \mathcal{A}$;
2. for all $A \in \mathcal{A}$, $A^c \in \mathcal{A}$ as well;
3. for a sequence in \mathcal{A} , $\{A_i\}_{i \in \mathbb{N}}$, the union $\bigcup_{i \in \mathbb{N}} A_i \in \mathcal{A}$ as well.

An element $A \in \mathcal{A}$ is called a **\mathcal{A} -measurable set**.

Remark 1.1. It's easy to show that the largest σ -algebra of set X is the power set $\mathcal{P}(X)$, and the smallest σ -algebra is $\{\emptyset, X\}$.

Theorem 1.1. Let $\{\mathcal{A}_i\}_{i \in I}$ be the collection of all σ -algebra on X . Then, $\bigcap_{i \in I} \mathcal{A}_i$ is also a σ -algebra on X .

Proof. ■

Definition 1.2. For $\mathcal{M} \subseteq \mathcal{P}(X)$ (\mathcal{A} is not necessarily a σ -algebra), the smallest σ -algebra (by taking intersections) containing \mathcal{M} is defined as the **σ -algebra generated by \mathcal{M}** . The generated σ -algebra is simply the intersection of all σ -algebras that are supersets of \mathcal{M} .

$$\sigma(\mathcal{M}) = \bigcap_{\mathcal{A} \supseteq \mathcal{M} \text{ s.t. } \mathcal{A} \text{ is } \sigma\text{-algebra}} \mathcal{A} \quad (1)$$

Definition 1.3. Let (X, τ) be a topological space, then the **Borel algebra** is σ -algebra generated by the collection of open sets τ .

$$\mathcal{B}(X) := \sigma(\tau) \quad (2)$$

Remark 1.2. We do not use the entire power set for analysis because it's too large to construct a sensible measure on (see Theorem 1.2).

Theorem 1.2. There is no measure μ on $(\mathbb{R}, \mathcal{P}(\mathbb{R}))$ satisfying the following two conditions: (i) $\mu((a, b]) = b - a$ for every $a < b$ and (ii) $\mu(x + A) = \mu(A)$ for all $a \in \mathbb{R}$ and $A \in \mathcal{P}(\mathbb{R})$.

Proof. Suppose, for contradiction, there exists such a measure μ , then $\mu((0, 1]) = 1 < \infty$.

Claim: the only measure on $(\mathbb{R}, \mathcal{P}(\mathbb{R}))$ satisfying $\mu((0, 1]) < \infty$ and $\mu(x + A) = \mu(A)$ is the zero measure.

To prove the claim, let $I := (0, 1]$ and define the following equivalence relation on I :

$$x \sim y \iff x - y \in \mathbb{Q} \quad (3)$$

the corresponding equivalence class of x on I can be written as

$$[x] = \{x + r : r \in \mathbb{Q} \wedge x + r \in I\} \quad (4)$$

The collection of all such equivalence classes, \mathcal{A} , is a disjoint decomposition of I . (for every $x \in I$, $[x]$ must in \mathcal{A} and $x \in [x]$ trivially. If there exists different $[x] \neq [y]$ but $[x] \cap [y] \neq \emptyset$, take $z \in [x] \cap [y]$, by the transitivity of equivalence relation, $x \sim z \sim y$. Therefore, $[x] = [y]$, contradiction.)

For each $[x] \in \mathcal{A}$, take exactly one $a_x \in [x]$ and define set $A := \{a_x : [x] \in \mathcal{A}\}$. As a result, set A satisfies the following two properties:

1. $\forall x \in I, \exists a_x \in A$ s.t. $a_x \in [x]$.
2. $\forall x, y \in A, x \sim y \implies x = y$.

Since $\mathbb{Q} \cup (-1, 1]$ is countable, let $(r_n)_{n \in \mathbb{N}}$ be an enumeration of all elements in it.

For each $n \in \mathbb{N}$, define $A_n := r_n + A$.

Note that for any m, n such that $A_m \cap A_n \neq \emptyset$, take $x \in A_m \cap A_n$. By definition,

$$x = r_n + a_n \quad (5)$$

$$x = r_m + a_m \quad (6)$$

where $a_n, a_m \in A$ and $r_n, r_m \in \mathbb{Q}$. Consequently,

$$a_n - a_m = r_m - r_n \in \mathbb{Q} \quad (7)$$

Therefore, $a_n \sim a_m$. By the second property of A , $a_n = a_m$. Thus, $r_m = r_n$ and $m = n$.

Take the contraposition of what we just proved, $m \neq n \implies A_m \cap A_n = \emptyset$.

Let $z \in (0, 1]$, there exists some $a \in A$ such that $z \in [a]$. That is, $z = a + r$ for some $r \in \mathbb{Q} \cap (-1, 1]$. There must exist some $m \in \mathbb{N}$ such that $r_m = r$, and consequently, $z \in A_m$.

Therefore, $(0, 1] \subseteq \bigcup_{n \in \mathbb{N}} A_n \subseteq (-1, 2]$ (the second relation is obvious). Moreover,

$$\mu((0, 1]) \leq \mu\left(\bigcup_{n \in \mathbb{N}} A_n\right) \leq \mu((-1, 2]) = \mu((-1, 0]) + \mu((0, 1]) + \mu((1, 2]) = 3\mu((0, 1]) \quad (8)$$

Note that we just proved $\bigcup_{n \in \mathbb{N}} A_n$ is a disjoint union, hence,

$$\mu((0, 1]) \leq \sum_{n=1}^{\infty} \mu(A_n) \leq 3\mu((0, 1]) \quad (9)$$

$$\implies ((0, 1])\mu \leq \sum_{n=1}^{\infty} \mu(A + r_n) \leq 3\mu((0, 1]) \quad (10)$$

$$\implies \mu((0, 1]) \leq \sum_{n=1}^{\infty} \mu(A) \leq 3\mu((0, 1]) \quad (11)$$

Since $\mu((0, 1])$ is finite, the only value $\mu(A)$ can take is zero, and $\mu(I) = 0$ as well. Consequently, for any set $S \in \mathcal{P}(\mathbb{R})$, if $S \subseteq I$, then $\mu(S) \leq \mu(I)$ and $\mu(S) = 0$. Otherwise, let $l = \lfloor \inf(S) \rfloor$ and $u = \lceil \sup(S) \rceil$.

$$I \subseteq S \subseteq \bigcup_{n=l}^u (n, n+1] \quad (12)$$

Therefore,

$$0 \leq \mu(S) \leq \sum_{n=l}^u \mu(n + (0, 1]) = \sum_{n=l}^u \mu((0, 1]) = 0 \quad (13)$$

It's shown that $\mu(S) = 0$ for every $S \subseteq \mathcal{P}(\mathbb{R})$.

This leads to a contradiction to the first property required ($\mu((a, b]) = b - a$). ■

2 Measurable Spaces and Measurable Maps

Definition 2.1. Let (X_1, \mathcal{A}_1) and (X_2, \mathcal{A}_2) be two measurable spaces. A function $f : X_1 \rightarrow X_2$ is a **measurable map** with respect to \mathcal{A}_1 and \mathcal{A}_2 if

$$f^{-1}(A_2) \in \mathcal{A}_1 \quad \forall A_2 \in \mathcal{A}_2 \quad (14)$$

That is, the pre-image of every set in \mathcal{A}_2 is an element in \mathcal{A}_1 as well.

Theorem 2.1. Let (Ω, \mathcal{A}) be a measurable space, then the indicator function for any $A \in \mathcal{A}$, $\mathcal{X}_A : \Omega \rightarrow \mathbb{R}$, is measurable with respect to \mathcal{A} and $\mathcal{B}(\mathbb{R})$.

$$\mathcal{X}_A(\omega) := \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{otherwise} \end{cases} \quad (15)$$

Proof. ■

Theorem 2.2. The composition of measurable maps is measurable.

Proof. ■

Theorem 2.3. For measurable spaces (X, \mathcal{A}) and $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ and measurable maps $f, g : \Omega \rightarrow \mathbb{R}$, $f + g$, $f - g$ and $|f|$ are measurable.

Proof. ■

3 Lebesgue Measures