GSD RESEARCH MISSION PACKAGE

Highly Nonlinear Systems, Clouds, and the Open Frontier

Frank Merle, Fluid Dynamics, Atmospheric Physics, Numerical Weather Prediction, and the Erdős Problems

A Deep Research Study Across Six Cross-Cutting Modules

> *"By engaging with the chaos, Merle discovered simplicity. Much of his work focuses on special structures, called solitons, that persist amid the mayhem of nonlinear systems."*
>
> — on the 2026 Breakthrough Prize in Mathematics

Vision → Research → Mission

Full Pipeline Execution

Date: April 20, 2026 | Status: Ready for GSD Orchestrator
Activation Profile: Fleet | Estimated: 14 context windows across 5 sessions
Six modules | Cross-cutting math, physics, atmospheric science, and AI

Contents

1 Mission Vision **6**
1.1 Vision . . . 6
1.2 Problem Statement . . . 7
1.3 Core Concept . . . 8
1.3.1 Domain Map . . . 8
1.4 Research Modules . . . 8
1.4.1 M1: Frank Merle and Highly Nonlinear Evolution Equations . . . 8
1.4.2 M2: Fluid Dynamics — Mathematics and Physics . . . 9
1.4.3 M3: WMO Cloud Atlas — Complete Taxonomy . . . 9
1.4.4 M4: Cloud Microphysics and Atmospheric Thermodynamics . . . 9
1.4.5 M5: Numerical Weather Prediction and AI Forecasting . . . 10
1.4.6 M6: Open Problems in Mathematics and Physics . . . 10
1.5 Chipset Configuration . . . 10
1.6 Scope Boundaries . . . 11
1.6.1 In Scope (v1.0) . . . 11
1.6.2 Out of Scope (Future Considerations) . . . 11
1.7 Success Criteria . . . 12
1.8 Relationship to Other Documents . . . 13
1.9 The Through-Line . . . 13

2 How to Use This Document **14**

3 Module 1 Reference: Frank Merle and Highly Nonlinear PDEs **15**
3.1 Biographical Foundation . . . 15
3.2 Awards and Recognition . . . 15
3.3 Contribution Stream 1: L^2-Critical Nonlinear Schrödinger Blow-Up . . . 16
3.4 Contribution Stream 2: Soliton Resolution via Channels of Energy . . . 16
3.5 Contribution Stream 3: KdV Soliton Collision and Blow-Up . . . 17
3.6 Contribution Stream 4: Compressible Euler/Navier–Stokes Implosion (2019–2022) . . . 17
3.7 Bibliography for Module 1 . . . 18

4 Module 2 Reference: Fluid Dynamics — Mathematics and Physics **18**
4.1 The Navier–Stokes Equations . . . 18
4.2 The Reynolds Number . . . 19
4.3 Vorticity and Vortex Stretching . . . 19
4.4 The Leray–Hopf Weak Solution Framework . . . 19
4.5 Kolmogorov 1941 Theory of Turbulence (K41) . . . 19
4.6 The Clay Millennium Prize Problem for Navier–Stokes . . . 20
4.7 Bibliography for Module 2 . . . 21

5 Module 3 Reference: WMO Cloud Atlas — Complete Taxonomy **21**
5.1 Historical Foundation . . . 21
5.2 The Ten Genera . . . 21
5.3 The Fifteen Species . . . 23

5.4 The Nine Varieties . . . 24
5.5 Supplementary Features and Accessory Clouds . . . 24
5.6 Special Clouds . . . 25
5.7 Upper-Atmospheric Clouds . . . 26
5.8 Bibliography for Module 3 . . . 26

6 Module 4 Reference: Cloud Microphysics and Atmospheric Thermodynamics 26
6.1 Saturation Vapor Pressure and Clausius–Clapeyron . . . 27
6.2 Adiabatic Ascent and the Lifting Condensation Level . . . 27
6.3 Köhler Theory of Cloud Droplet Activation . . . 27
6.4 The Wegener–Bergeron–Findeisen (WBF) Process . . . 28
6.5 Collision–Coalescence (Warm Rain) . . . 28
6.6 Ice Nucleation Modes . . . 28
6.7 Aerosol Indirect Effects . . . 29
6.8 Bibliography for Module 4 . . . 29

7 Module 5 Reference: Numerical Weather Prediction and AI Forecasting 30
7.1 Historical Foundations . . . 30
7.2 The Primitive Equations . . . 30
7.3 Numerical Methods . . . 31
7.4 Data Assimilation . . . 32
7.5 Predictability and the Lorenz Limit . . . 33
7.6 Operational NWP Systems (2026) . . . 34
7.7 The AI Weather Models (2022–2026) . . . 34
7.8 Bibliography for Module 5 . . . 36

8 Module 6 Reference: Open Problems — Erdős, Millennium, and Physics 36
8.1 The Erdős Problems Database . . . 36
8.2 Erdős Problem #1196 — Solved by AI, April 2026 . . . 37
8.3 Other AI Contributions to Erdős Problems (2025–2026) . . . 38
8.4 The Seven Clay Millennium Prize Problems . . . 38
8.5 Other Major Open Problems in Mathematics . . . 39
8.6 Major Open Problems in Physics (2026) . . . 40
8.7 Bibliography for Module 6 . . . 41

9 Cross-Module Synthesis (Reference) 41

10 Safety and Sensitivity Considerations (All Modules) 42

11 Milestone Specification 43
11.1 Mission Objective . . . 43
11.2 Architecture Overview . . . 43
11.3 System Layers . . . 43
11.4 Deliverables . . . 44
11.5 Component Breakdown . . . 45
11.6 Model Assignment Rationale . . . 45

12 Wave Execution Plan 45

12.1 Wave Summary . . . 46
12.2 Wave 0: Foundation . . . 46
12.3 Wave 1A: Math & Open Problems Track (parallel with 1B) . . . 46
12.4 Wave 1B: Atmospheric Track (parallel with 1A) . . . 47
12.5 Wave 2: Synthesis . . . 47
12.6 Wave 3: Publish + Verify . . . 47
12.7 Cache Optimization Strategy . . . 47
12.8 Token Budget . . . 48
12.9 Dependency Graph . . . 48

13 Test Plan **48**
13.1 Test Categories . . . 48
13.2 Safety-Critical Tests . . . 49
13.3 Core Functional Tests (sample) . . . 50
13.4 Integration Tests . . . 51
13.5 Edge Case Tests . . . 51
13.6 Verification Matrix . . . 52

14 Mission Crew Manifest (Fleet Profile) **53**

15 Execution Summary **54**

16 Closing Through-Line **54**

STAGE 1 — VISION DOCUMENT

1 Mission Vision

Date: April 20, 2026

Status: Vision Compiled / Research Integrated

Depends on: None — standalone exploratory research mission

Context: A six-module deep-research mission spanning the mathematics of nonlinear evolution equations (with focus on Frank Merle's recently-honored body of work), the physics and mathematics of fluid dynamics, the comprehensive WMO classification of clouds, the science and mathematics of numerical weather prediction, and the present landscape of open problems in mathematics — including the just-resolved Erdős Problem #1196 — and physics.

1.1 Vision

On April 18, 2026, Frank Merle was awarded the Breakthrough Prize in Mathematics. His citation *"for breakthroughs in nonlinear evolution equations, with regards to their stability, singularity formation, or resolution into solitons"* sits at the precise intersection of every topic in this mission. To understand Merle's work is to understand the mathematics of fluid singularity. To understand fluid singularity is to understand why *water and air* — the two substances we live inside — still defeat the analytical tools of the most powerful mathematical century in human history. And to look honestly at what we cannot yet predict about the atmosphere, while admiring the precise Latin taxonomy we have built to name every variation of cloud, is to be reminded that we live at a strange threshold: we have named the mountains we cannot yet climb.

Five days earlier, on April 13, 2026, the Erdős Problem #1196 — the Erdős–Sárközy–Szemerédi conjecture on primitive sets — fell to a single 80-minute reasoning trace from GPT-5.4 Pro. Jared Lichtman, who had worked on the problem for seven years, had built the strongest prior bound; the AI's proof produced a strikingly compact result in a way that no human researcher had managed. Two of the most consequential mathematical events of the year, separated by less than a week, both pointed at the same fact: *the boundary between what we can solve and what we cannot is moving in front of us, and we should look carefully at where it sits today.*

This mission catalogs that boundary across six modules. It begins with Merle as the human protagonist of a story about taming nonlinear chaos through solitons. It steps into the Navier–Stokes equations — still a Millennium Prize problem — where Merle's recent work with Raphaël, Rodnianski, and Szeftel produced the first proof that the compressible Euler equations admit finite-time imploding singularities. It catalogs every cloud the World Meteorological Organization has named, and the physics by which each one forms — because clouds are precisely what nonlinear fluid equations *produce*, in a moist, rotating, radiating atmosphere. It documents the mathematics by which we forecast that atmosphere, including the recent AI revolution that has made GraphCast, Pangu-Weather, and FengWu credible challengers to the European

Centre for Medium-Range Weather Forecasts' Integrated Forecast System. And it closes with the open problems — the seven Millennium Problems, the Erdős Problems database now under active assault by AI systems, and the major unsolved questions in physics from quantum gravity to dark matter.

The through-line is the *spaces between* — between order and chaos, between continuous fluid and discrete observation, between human proof and AI proof, between the named and the unnamed. The Amiga Principle is everywhere: the right specialized tool, applied at the right level of architectural abstraction, produces outsized outcomes. Merle did not crush the nonlinear Schrödinger equation by force; he found the soliton structure inside it. Kolmogorov did not solve turbulence; he identified the scale hierarchy that lets us reason about it. The WMO did not predict every cloud; it built the genus–species–variety taxonomy that lets every observer in every nation file what they see into a coherent global picture. This mission honors that lineage by structuring its own research the same way: specialized modules, precise vocabulary, and architectural leverage where brute-force breadth would have failed.

1.2 Problem Statement

1. **Nonlinear evolution equations resist analytical understanding.** The Navier–Stokes equations are 180 years old, govern the motion of every fluid that matters to human life, and remain a Clay Millennium Prize Problem. Frank Merle's body of work — from the L^2-critical nonlinear Schrödinger blow-up theory of the 1990s to the 2022 *Annals of Mathematics* paper proving compressible-Euler implosion — has fundamentally shifted what is provable, but no comprehensive synthesis exists that connects his contributions to the broader landscape of fluid singularity, soliton resolution, and turbulence theory.

2. **Cloud taxonomy is fragmented across audiences.** The WMO International Cloud Atlas catalogs ten genera, fifteen species, nine varieties, and a dozen-plus supplementary features and accessory clouds, plus upper-atmospheric and special clouds. No single accessible reference co-locates the formal taxonomy with the underlying microphysics (Köhler theory, Wegener–Bergeron–Findeisen process, collision–coalescence) and thermodynamics (lifting condensation level, saturation vapor pressure, Clausius–Clapeyron) that make each genus appear when and where it does.

3. **Weather forecasting is undergoing an AI revolution that needs honest framing.** GraphCast, Pangu-Weather, FengWu, FuXi, AIFS, and GenCast have all matched or exceeded the European Centre's Integrated Forecast System on most standard metrics, while running in under a minute on a single TPU rather than hours on a supercomputer. They have weaknesses (extreme precipitation, novel regimes, training-data biases) that practicing forecasters must understand, but the existing literature splits between breathless promotion and reflexive skepticism. A grounded reference matters now.

4. **Erdős Problem #1196 has just been solved by AI, and the broader Erdős Problems database is becoming a live frontier.** Terence Tao's `erdosproblems.com` site now tracks AI contributions in real time. #1196 was solved by GPT-5.4 Pro on April 13, 2026 after a seven-year human attack. #1202, #152, #218, #281, and

#333 have all seen substantive AI contributions in 2026. Anyone working at the boundary of AI and mathematics needs a map of what is open, what has fallen, and what the methodological controversies are.

5. **Open problems in mathematics and physics need a unified modern reference.** The seven Millennium Prize Problems were posed in 2000; only the Poincaré Conjecture has been solved. Wikipedia's *List of unsolved problems in physics* is comprehensive but unstructured. Anyone trying to orient themselves to the current mathematical and physical frontier in 2026 needs a curated, source-backed index that connects the Millennium Problems to the Erdős Problems to the major open questions in physics, with enough technical specificity to be useful and enough breadth to be navigable.

1.3 Core Concept

One sentence: Catalog → Connect → Synthesize.

The mission produces six self-contained research modules that catalog their respective domains rigorously, connect across modules through explicit cross-references (Merle's compressible-Euler work touches the Navier–Stokes Millennium Problem; the cloud taxonomy interfaces with the NWP parameterization problem; the open-problems module catalogs both the Erdős frontier where AI is succeeding and the fluid frontier where it is not), and synthesize through a final integration document that draws the through-line connecting Merle's solitons, Kolmogorov's energy cascade, WMO's genera, and the open frontier.

1.3.1 Domain Map

```
SIX-MODULE RESEARCH ARCHITECTURE
+---------------------------+ +---------------------------+ |
M1: FRANK MERLE & | | M2: FLUID DYNAMICS: | | HIGHLY NONLINEAR
PDEs |---| MATH & PHYSICS | | (biographical, technical)| |
(NSE, Euler, turbulence) | +---------------------------+
+---------------------------+ | | | +---------------------+ | |
v v +---------------------------+ +---------------------------+
| M3: WMO CLOUD ATLAS: | | M4: CLOUD MICROPHYSICS | | COMPLETE
TAXONOMY |---| & THERMODYNAMICS | | (genera/species/var.)
| | (Kohler, WBF, droplet) | +---------------------------+
+---------------------------+ | | +-----------+------------------+ v
+---------------------------+ | M5: NUMERICAL WEATHER | | PREDICTION
+ AI | | (NWP, DA, EnKF, ML) | +---------------------------+ | v
+---------------------------+ | M6: OPEN PROBLEMS: | | Erdos #1196,
Millennium, | | unsolved math & physics | +---------------------------+
| v +---------------------------+ | SYNTHESIS / THROUGH- | | LINE
INTEGRATION DOC | +---------------------------+
```

1.4 Research Modules

1.4.1 M1: Frank Merle and Highly Nonlinear Evolution Equations

Biographical context (Marseille, ENS, Berestycki, Cergy-Pontoise, IHES Chair), career arc, and a precise technical synthesis of the four major contribution streams:

(1) L^2-critical nonlinear Schrödinger blow-up dynamics with Pierre Raphaël; (2) the Kenig–Merle channels-of-energy + concentration–compactness method for the soliton resolution conjecture; (3) the Martel–Merle–Raphaël work on KdV soliton collision and blow-up; and (4) the Merle–Raphaël–Rodnianski–Szeftel proof of finite-time singularity formation in compressible Euler and Navier–Stokes (2019, 2022) which won the 2023 Bôcher Prize and the 2023 Clay Research Award. Closes with the 2026 Breakthrough Prize award.

1.4.2 M2: Fluid Dynamics — Mathematics and Physics

Derivation and properties of the incompressible Navier–Stokes equations $\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\rho^{-1}\nabla p + \nu\Delta\mathbf{v} + \mathbf{f}$ and $\nabla \cdot \mathbf{v} = 0$, the inviscid Euler equations, the Reynolds number and the laminar/turbulent transition, vorticity and vortex stretching, the Leray–Hopf weak solution framework, and Kolmogorov's 1941 statistical theory of turbulence (K41) with the famous $E(k) \sim \epsilon^{2/3}k^{-5/3}$ inertial-range spectrum. The Clay Millennium Prize formulation of the Navier–Stokes existence and smoothness problem closes the module.

1.4.3 M3: WMO Cloud Atlas — Complete Taxonomy

Every cloud the WMO International Cloud Atlas formally recognizes:

- **Ten genera**, organized by altitude (high, middle, low/multi-level), with full Latin name, abbreviation, mean base altitude in temperate latitudes, typical composition (water/ice/mixed), and characteristic appearance.
- **Fifteen species**, describing shape and internal structure (fibratus, uncinus, spissatus, castellanus, floccus, stratiformis, nebulosus, lenticularis, volutus, fractus, humilis, mediocris, congestus, calvus, capillatus), with the genera each species most frequently occurs with.
- **Nine varieties**, describing transparency and arrangement (intortus, vertebratus, undulatus, radiatus, lacunosus, duplicatus, translucidus, perlucidus, opacus).
- **Supplementary features and accessory clouds** (incus, mamma, virga, praecipitatio, arcus, tuba, pileus, velum, pannus, fluctus, asperitas, cavum, murus, cauda).
- **Special clouds and upper-atmospheric clouds** (homogenitus and homomutatus from human activity; nacreous polar stratospheric clouds; noctilucent mesospheric clouds).

1.4.4 M4: Cloud Microphysics and Atmospheric Thermodynamics

The physics that makes each genus form. Saturation vapor pressure and the Clausius–Clapeyron relation. Adiabatic ascent, lifting condensation level, and the dry/moist adiabatic lapse rates. Köhler theory of cloud-droplet activation on cloud condensation nuclei, with the curvature (Kelvin) and solute (Raoult) terms in tension. The Wegener–Bergeron–Findeisen process by which ice crystals grow at the expense of supercooled water droplets. Collision–coalescence in warm rain. Ice nucleation modes (homogeneous and heterogeneous), riming, and aggregation. Mixed-phase cloud thermodynamics. The aerosol indirect effects (Twomey, Albrecht).

1.4.5 M5: Numerical Weather Prediction and AI Forecasting

Bjerknes' 1904 vision; Richardson's 1922 hand calculation; the Charney 1950 ENIAC barotropic-vorticity-equation forecast; the rise of primitive-equation models in the 1960s. The mathematical structure of NWP: the seven primitive equations (three momentum, mass continuity, thermodynamic energy, state, moisture). Discretization on staggered grids (Arakawa A–E), spectral methods, finite-volume methods, and the Courant–Friedrichs–Lewy stability condition. Sub-grid-scale parameterization (convection, radiation, boundary layer, microphysics, land surface). Data assimilation: optimal interpolation, 3D-Var, 4D-Var, the Ensemble Kalman Filter, and hybrid 4DEnVar. Ensemble forecasting (Lorenz's chaos, Epstein's 1969 stochastic-Monte-Carlo proposal, today's operational EPS systems). The 2022–2026 AI revolution: Pangu-Weather (3D Earth-Specific Transformer), GraphCast (Graph Neural Network), FengWu, FuXi, AIFS (ECMWF), GenCast (probabilistic), with their performance, weaknesses, and open questions.

1.4.6 M6: Open Problems in Mathematics and Physics

The Erdős Problems database curated by Terence Tao. Detailed treatment of **Erdős Problem #1196**: the Erdős–Sárközy–Szemerédi conjecture on primitive sets, with the precise statement, prior bounds (Erdős 1934, Lichtman 2023), and the April 13, 2026 GPT-5.4 Pro proof. Survey of notable AI contributions to other Erdős problems (#1202, #120, #152, #218, #281, #333). The seven Clay Millennium Prize Problems with current status. Curated catalog of major open problems in mathematics (Riemann hypothesis, Birch–Swinnerton-Dyer, Hodge conjecture, P vs NP, Yang–Mills mass gap, twin prime conjecture, Goldbach, Collatz). Curated catalog of major open problems in physics (quantum gravity, dark matter, dark energy, matter–antimatter asymmetry, neutrino masses and Dirac/Majorana nature, strong CP problem, hierarchy problem, cosmological constant problem, fine-tuning, measurement problem in QM, theory of everything).

1.5 Chipset Configuration


```
name: nonlinear-systems-clouds-open-problems version: 1.0.0 description:
"Six-module deep research mission across nonlinear PDEs, fluid dynamics,
cloud taxonomy, NWP, and open problems."
skills: merle-pde: domain: mathematical-physics description: "Frank
Merle biography + nonlinear PDE blow-up theory" fluid-math: domain:
mathematical-physics description: "Navier-Stokes, Euler, turbulence,
Kolmogorov K41" cloud-atlas: domain: atmospheric-science description:
"Complete WMO cloud taxonomy with physics per type" cloud-microphysics:
domain: atmospheric-science description: "Kohler theory, WBF process,
droplet/ice growth" nwp-forecast: domain: atmospheric-science description:
"NWP equations, data assimilation, AI weather models" open-problems:
domain: mathematics-physics description: "Erdos #1196, Millennium
Problems, unsolved physics"
agents: topology: "leader-worker" agents: - name: "FLIGHT" (Mission
orchestrator) role: "go/no-go gates, wave decomposition" - name: "PLAN"
role: "wave optimizer, dependency manager" - name: "SCOUT" role: "source
gathering, gap-fill research" - name: "EXEC-A" role: "Track A executor
(M1, M2, M6)" - name: "EXEC-B" role: "Track B executor (M3, M4, M5)"
```

```
- name: "CRAFT-MATH" role: "PDE / number theory technical specialist"
- name: "CRAFT-FLUID" role: "fluid dynamics + atmospheric physics
specialist" - name: "CRAFT-METEO" role: "WMO taxonomy + NWP specialist"
- name: "CRAFT-AI" role: "AI weather models + AI mathematics specialist"
- name: "VERIFY" role: "source validation, equation-form checking" -
name: "INTEG" role: "cross-module synthesis, through-line construction" -
name: "CAPCOM" role: "human approval at wave boundaries" - name: "BUDGET"
role: "token budget tracking" - name: "SURGEON" role: "research-quality
degradation detection" - name: "LOG" role: "audit trail, commit messages"
- name: "TOPO" role: "agent topology management" - name: "ARCH" role:
"cross-cutting structural decisions"
evaluation: gates: pre_deploy: - check: "source_quality"
threshold: 100 (zero non-authoritative sources) action: "block"
- check: "equation_form_check" command: "verify all PDE
statements against canonical references" action: "block" - check:
"cloud_taxonomy_completeness" threshold: 100 (all 10 genera, 15 species,
9 varieties) action: "block"
```

1.6 Scope Boundaries

1.6.1 In Scope (v1.0)

- Frank Merle's published research career through the 2026 Breakthrough Prize
- The incompressible and compressible Navier–Stokes and Euler equations
- Kolmogorov K41 turbulence theory and modern refinements
- Every cloud type formally cataloged in the WMO International Cloud Atlas (2017 edition and post-2017 additions including *asperitas*, *volutus*, *cavum*, *murus*, *cauda*, *fluctus*, homogenitus, homomutatus)
- All major numerical weather prediction techniques in operational use through 2026, including the AI weather models that emerged 2022–2025
- The Erdős Problems database, with detailed treatment of #1196 and representative coverage of other AI-touched problems
- The seven Clay Millennium Prize Problems with current (April 2026) status
- Major curated open problems in mathematics and physics

1.6.2 Out of Scope (Future Considerations)

- Computational fluid dynamics implementations and codes (separate mission)
- Climate-change projections and policy implications (separate domain)
- Detailed weather-modification engineering (cloud seeding, geoengineering)
- AI safety analysis of mathematics-solving systems (separate mission)
- Full proof reconstruction for any specific Erdős problem (reference only)
- Quantum field theory derivations underlying open physics problems

1.7 Success Criteria

1. Frank Merle's biographical and professional record is documented from authoritative sources (IHES, CY Cergy Paris University, Breakthrough Prize Foundation, peer-reviewed publication record).
2. All four of Merle's major contribution streams are described with correct technical content and primary-source citations to the relevant papers in *Inventiones Mathematicae*, *Annals of Mathematics*, and *Communications in Mathematical Physics*.
3. The Navier–Stokes and Euler equations are stated in their canonical mathematical form with all symbols defined, and the Millennium Prize formulation matches the Clay Mathematics Institute statement.
4. All ten WMO cloud genera are individually profiled with mean altitude, composition, characteristic appearance, and primary-source attribution to the WMO International Cloud Atlas.
5. All fifteen species and nine varieties are tabled with formal definitions and the genera with which each most frequently occurs.
6. Cloud microphysics covers Köhler theory, the Wegener–Bergeron–Findeisen process, collision–coalescence, and ice nucleation with the relevant physical equations.
7. NWP is documented from Bjerknes (1904) through the 2026 AI weather models, with primary-source citations to ECMWF, NOAA, and peer-reviewed papers introducing each AI model.
8. Erdős Problem #1196 is stated precisely with its history, prior bounds, and the April 13, 2026 GPT-5.4 Pro resolution documented.
9. The seven Millennium Prize Problems are listed with current status (April 2026) and primary-source attribution to Clay Mathematics Institute.
10. Every major claim is attributed to a specific primary source. Zero entertainment-media citations.
11. The synthesis document explicitly connects Merle → Navier–Stokes → turbulence → atmospheric flow → cloud formation → NWP predictability → open problems.
12. PDF compiles cleanly with three XeLaTeX passes, including a complete table of contents, working hyperlinks, and accurate page numbers throughout.

1.8 Relationship to Other Documents

Document	Relationship
The Space Between	Provides the mathematical-philosophical spine; this mission's treatment of nonlinear PDEs, turbulence, and chaos extends and grounds chapters on dynamical systems and the mathematics of complexity.
The Quark and the Compiler	Shares the framing of computing history and mathematical evolution; the AI-solves-Erdős-#1196 thread directly extends the essay's argument about the evolution of computational reasoning.
The Hundred Voices	Provides a literary frame for the long arc from the "cosmic purple" physics of the early universe to the named taxonomies of the present moment.
GSD Mathematical Foundations Conversation	Establishes the unit-circle / mathematical-bedrock framework into which this mission's PDE and turbulence content fits.
GSD Foundational Knowledge Pack	This mission is a candidate sub-pack focused on *nonlinear systems and prediction*.

1.9 The Through-Line

The Amiga Principle taught us that architectural elegance and specialized execution paths produce outsized outcomes. This mission's six modules are that principle in research form: rather than a single sprawling survey, six precise specialists — the Merle biographer, the fluid mathematician, the cloud taxonomist, the microphysicist, the NWP modeler, the open-problems cataloger — working in parallel, then synthesized into a single coherent through-line. The *spaces between* are everywhere here: between the linear and the nonlinear, between the well-posed and the singular, between the named cloud and the un-named transitional form, between the human proof and the AI proof, between the problem solved and the problem adjacent to it that remains. We have learned to live at thresholds. This mission is a careful inventory of one such threshold, taken in April 2026, on a particularly meaningful week.

STAGE 2 — RESEARCH REFERENCE

2 How to Use This Document

This Research Reference is the source-grounded technical companion to the Vision Document above. Where the Vision says *"six modules covering Merle, fluids, clouds, NWP, and open problems,"* this Reference provides the actual statements, equations, biographical facts, taxonomic catalogs, and citations that the mission agents must use. An execution agent should be able to produce its assigned module using only this Reference and the corresponding component spec.

Key source organizations. Every claim in this Reference is attributed to one or more of the following authoritative sources:

- **Institut des Hautes Études Scientifiques (IHES)** — biographical and award records for Merle.
- **Breakthrough Prize Foundation** — 2026 prize citation.
- **American Mathematical Society** — Bôcher Prize records.
- **Clay Mathematics Institute** — Millennium Prize Problem statements; Clay Research Award.
- **World Meteorological Organization (WMO)** — International Cloud Atlas, the legal Annex I to the Technical Regulations (WMO-No. 49).
- **NOAA (National Oceanic and Atmospheric Administration)** — NWS forecast model documentation; cloud reference.
- **NCEP (National Centers for Environmental Prediction)** — operational forecast systems documentation.
- **ECMWF (European Centre for Medium-Range Weather Forecasts)** — Integrated Forecast System; AIFS; data assimilation literature.
- **Peer-reviewed journals** — *Inventiones Mathematicae*, *Annals of Mathematics*, *Communications in Mathematical Physics*, *Science*, *Nature*, *Journal of Fluid Mechanics*, *Atmospheric Chemistry and Physics*, *Monthly Weather Review*, *npj Climate and Atmospheric Science*.
- **erdosproblems.com** (curated by Terence Tao) — Erdős Problems database with formal statements, current status, and AI-contribution tracking.
- **Wikipedia** — used for standardized equation forms and high-level historical timelines, with primary-source verification of all substantive claims.

3 Module 1 Reference: Frank Merle and Highly Nonlinear PDEs

3.1 Biographical Foundation

Frank Merle was born on November 22, 1962, in Marseille, France. He attended the École Normale Supérieure (ENS) and received his Ph.D. in 1987 from the Université Paris VI (Pierre et Marie Curie, now Sorbonne Université) under the direction of Henri Berestycki. His thesis, *Contributions à l'étude de certaines équations aux dérivées partielles non linéaires de la physique mathématique*, was defended on November 18, 1987 before an examination jury that included Haïm Brezis, Claude Bardos, Jean Ginibre, and Pierre-Louis Lions.

After his Ph.D., Merle became a researcher for the CNRS at ENS. In 1989–1990 he was an assistant professor at the Courant Institute of Mathematical Sciences at New York University. Since 1991 he has been a professor at the Université de Cergy-Pontoise (now CY Cergy Paris Université). From 1998 to 2003 he was a member of the Institut Universitaire de France. He was a visiting scholar at the Institute for Advanced Study (Princeton) in 1996, 2001, and 2003–2004, and held visiting professorships at Stanford, Rutgers, Chicago, MSRI Berkeley, Leiden, and the University of Tokyo.

Since 2009 Merle has held the joint CY Cergy Paris Université–IHES Chair in Analysis at the Institut des Hautes Études Scientifiques in Bures-sur-Yvette, with funding from the Simons Foundation. He leads research within the CNRS–CY Cergy joint laboratory for analysis, geometry, and modeling (UMR 8088).

3.2 Awards and Recognition

Year	Award / Recognition
1997	Prix de l'Institut Poincaré en Physique Théorique
1998	Invited speaker, International Congress of Mathematicians, Berlin
2000	Prix Charles-Louis de Saulse de Freycinet, Académie des Sciences
2005	Bôcher Memorial Prize (American Mathematical Society)
2005	CNRS Silver Medal
2014	Plenary speaker, International Congress of Mathematicians, Seoul
2018	Prix Ampère de l'Électricité (Académie des Sciences)
2020	Member, Academia Europaea (mathematics section)
2023	Bôcher Memorial Prize (with Raphaël, Rodnianski, Szeftel)
2023	Clay Research Award (with Raphaël, Rodnianski, Szeftel)
2026	**Breakthrough Prize in Mathematics ($3M)**

The 2026 Breakthrough Prize citation reads:

> *"For breakthroughs in nonlinear evolution equations, with regards to their stability, singularity formation, or resolution into solitons."*

— Breakthrough Prize Foundation, April 18, 2026

3.3 Contribution Stream 1: L^2-Critical Nonlinear Schrödinger Blow-Up

The L^2-critical (or mass-critical) nonlinear Schrödinger equation in dimension $d = 2$ is

$$i\partial_t u + \Delta u + |u|^2 u = 0, \qquad u(0,x) = u_0(x) \in H^1(\mathbb{R}^2),$$

where $u : \mathbb{R}_t \times \mathbb{R}^2_x \to \mathbb{C}$.

Critical mass is the L^2 norm of the ground state Q, the unique radial positive solution of $\Delta Q - Q + Q^3 = 0$. Below critical mass, solutions exist globally; at and above critical mass, finite-time blow-up can occur.

Merle's first major contribution (1990, *Communications in Mathematical Physics* 129, 223–240) proved the existence of solutions blowing up at exactly k prescribed points for the critical Schrödinger equation. The work with Pierre Raphaël in the early 2000s (*Geometric and Functional Analysis* 13, 591–642 (2003); *Inventiones Mathematicae* 156, 565–672 (2004); *Communications in Mathematical Physics* 253, 675–704 (2005)) established the universal log-log blow-up rate

$$\|\nabla u(t)\|^2_{L^2} \sim \frac{\log|\log(T-t)|}{T-t}$$

for a class of initial data slightly above critical mass, and proved quantization of the blow-up mass — the singular part accumulates a quantized amount of L^2 mass at the blow-up point.

3.4 Contribution Stream 2: Soliton Resolution via Channels of Energy

The **soliton resolution conjecture** states that a generic global solution to a nonlinear dispersive equation will, as $t \to \infty$, decompose asymptotically into a finite sum of decoupled solitons (modulated traveling waves) plus a radiation term that disperses to zero in L^p for $p > 2$. As Scientific American summarized for the 2026 prize: "a soliton is a special solution to nonlinear equations, such as fluid equations, that doesn't send energy away to infinity — it keeps all its energy contained and keeps the same shape."

Merle's collaboration with Carlos Kenig at Chicago, joined later by Thomas Duyckaerts, developed the *channels of energy* method coupled with the Bahouri–Gérard concentration–compactness profile decomposition. The key insight is a lower bound on the energy of an outgoing wave outside the light cone, which forces blow-up solutions to either contain a soliton or disperse cleanly. Duyckaerts–Kenig–Merle (2011, *Journal of the European Mathematical Society*) used this to resolve the conjecture for the energy-critical wave equation in the radial case, three-dimensional setting. Subsequent work by Duyckaerts, Jia, Kenig, Merle, and others has extended the result substantially.

3.5 Contribution Stream 3: KdV Soliton Collision and Blow-Up

With Yvan Martel and (later) Pierre Raphaël, Merle revealed the structure of singularities in the L^2-critical generalized Korteweg–de Vries (gKdV) equation

$$\partial_t u + \partial_x(\partial_{xx} u + u^5) = 0, \qquad u : \mathbb{R}_t \times \mathbb{R}_x \to \mathbb{R}.$$

Martel–Merle (2002, *Annals of Mathematics* 155, 235–280) proved stability of the blow-up profile and a sharp lower bound on the blow-up rate. Subsequent work demonstrated that solitons in this and similar equations can collide nearly elastically — the classical integrable KdV (u^2 nonlinearity instead of u^5) preserves solitons exactly through collision; the supercritical case approximates this only inelastically, emitting a small dispersive remainder.

3.6 Contribution Stream 4: Compressible Euler/Navier–Stokes Implosion (2019–2022)

This is the work that produced the 2023 Bôcher Prize, the 2023 Clay Research Award, and — as the most-cited recent contribution — a major piece of the 2026 Breakthrough Prize citation.

The team — Frank Merle (CY Cergy Paris–IHES), Pierre Raphaël (Schlumberger Chair at IHES, Cambridge), Igor Rodnianski (Princeton), and Jérémie Szeftel (CNRS, Sorbonne) — proved finite-time singularity formation in the compressible Euler equations and the compressible Navier–Stokes equations. The key paper, *On the implosion of a compressible fluid II: Singularity formation*, appeared in the *Annals of Mathematics* (2022).

The compressible Euler equations for an ideal isentropic fluid with density ρ, velocity $\mathbf{u}$, and pressure $p = K\rho^\gamma$ (adiabatic constant $\gamma > 1$) are

$$\begin{aligned} \partial_t \rho + \nabla \cdot (\rho \mathbf{u}) &= 0 \quad \text{(continuity)} \\ \partial_t(\rho \mathbf{u}) + \nabla \cdot (\rho \mathbf{u} \otimes \mathbf{u}) + \nabla p &= 0 \quad \text{(momentum).} \end{aligned}$$

The team constructed self-similar solutions where the density and velocity diverge in finite time at a point, with a precisely characterized self-similar profile. The construction extends to the compressible Navier–Stokes equations (with viscosity) for certain parameter ranges, demonstrating that viscosity does not always prevent singularity formation when compressibility allows imploding profiles.

This work also established blow-up for the energy-supercritical defocusing nonlinear Schrödinger equation $i\partial_t u + \Delta u - |u|^{p-1}u = 0$ in some regimes, a result long believed (but not proved) to be impossible because the defocusing sign of the nonlinearity was thought to suppress singularity. The construction exploited an unexpected connection between the supercritical NLS and compressible fluid dynamics.

3.7 Bibliography for Module 1

- Wikipedia, "Frank Merle (mathematician)" — biographical record.
- Institut des Hautes Études Scientifiques, archive of Merle-related events: https://www.ihes.fr/en/tag/merle/.
- Breakthrough Prize Foundation, "Breakthrough Prize Announces 2026 Laureates," April 18, 2026: https://breakthroughprize.org/News/98.
- Scientific American, "Master of chaos wins $3-million math prize for 'blowing up' equations," April 19, 2026.
- Merle, F.; Raphaël, P. (2003). "Sharp upper bound on the blow up rate for critical nonlinear Schrödinger equation." *Geometric and Functional Analysis* 13, 591–642.
- Merle, F.; Raphaël, P. (2004). "On universality of blow-up profile for L^2 critical nonlinear Schrödinger equation." *Inventiones Mathematicae* 156, 565–672.
- Martel, Y.; Merle, F. (2002). "Stability of blow-up profile and lower bounds for blow-up rate for the critical generalized KdV equation." *Annals of Mathematics* 155, 235–280.
- Merle, F.; Raphaël, P.; Rodnianski, I.; Szeftel, J. (2022). "On the implosion of a compressible fluid II: Singularity formation." *Annals of Mathematics*.
- Duyckaerts, T.; Kenig, C.; Merle, F. (2011). "Universality of blow-up profile for small radial Type II blow-up solutions of energy-critical wave equation." *Journal of the European Mathematical Society*.

4 Module 2 Reference: Fluid Dynamics — Mathematics and Physics

4.1 The Navier–Stokes Equations

For a Newtonian fluid of constant density ρ and constant kinematic viscosity $\nu = \mu/\rho$, the incompressible Navier–Stokes equations in $\mathbb{R}^3$ are

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\frac{1}{\rho}\nabla p + \nu \Delta \mathbf{v} + \mathbf{f}(\mathbf{x}, t),$$
$$\nabla \cdot \mathbf{v} = 0.$$

Here $\mathbf{v}(\mathbf{x}, t)$ is the velocity field, $p(\mathbf{x}, t)$ is the pressure, $\mathbf{f}$ is an external body force per unit mass, and $\Delta = \nabla^2$ is the Laplacian. The first equation is conservation of momentum (Newton's second law for a fluid parcel); the second is conservation of mass for an incompressible flow. The convective acceleration term $(\mathbf{v} \cdot \nabla)\mathbf{v}$ is the source of nonlinearity that defeats analytical treatment in three dimensions.

If $\nu = 0$, the equations reduce to the **Euler equations** for an inviscid fluid:

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\frac{1}{\rho}\nabla p + \mathbf{f}, \qquad \nabla \cdot \mathbf{v} = 0.$$

4.2 The Reynolds Number

For a flow with characteristic velocity V and characteristic length L, the Reynolds number

$$\mathrm{Re} = \frac{VL}{\nu}$$

is the dimensionless ratio of inertial to viscous forces. Transition from laminar to turbulent flow occurs at a critical Re that depends on geometry: for pipe flow, transition begins around $\mathrm{Re} \approx 2300$; for atmospheric boundary-layer flow, $\mathrm{Re} \sim 10^5$ or higher and turbulence is fully developed.

4.3 Vorticity and Vortex Stretching

The **vorticity** $\boldsymbol{\omega} = \nabla \times \mathbf{v}$ is twice the local angular velocity of fluid parcels. Taking the curl of the incompressible Navier–Stokes equations gives the vorticity equation

$$\partial_t \boldsymbol{\omega} + (\mathbf{v} \cdot \nabla)\boldsymbol{\omega} = (\boldsymbol{\omega} \cdot \nabla)\mathbf{v} + \nu \Delta \boldsymbol{\omega}.$$

The term $(\boldsymbol{\omega} \cdot \nabla)\mathbf{v}$ is **vortex stretching**: when a fluid parcel is stretched in the direction of its vorticity, conservation of angular momentum amplifies the vorticity. Vortex stretching is the three-dimensional mechanism behind the energy cascade and is absent in two dimensions, which is why two-dimensional turbulence has an *inverse* energy cascade rather than the direct cascade of three dimensions.

4.4 The Leray–Hopf Weak Solution Framework

Jean Leray (1934) introduced the modern mathematical framework for the Navier–Stokes equations, proving the existence of **global weak solutions** in three dimensions for finite-energy initial data. A Leray–Hopf weak solution satisfies the Navier–Stokes equations in a distributional sense and obeys the energy inequality

$$\frac{1}{2}\|\mathbf{v}(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla \mathbf{v}(s)\|_{L^2}^2 \, ds \le \frac{1}{2}\|\mathbf{v}_0\|_{L^2}^2.$$

Eberhard Hopf (1951) extended Leray's results. The crucial open question is **regularity**: are these weak solutions actually smooth? In two dimensions, weak solutions are smooth; in three dimensions, this remains a Millennium Prize Problem.

4.5 Kolmogorov 1941 Theory of Turbulence (K41)

Andrey Kolmogorov's 1941 papers (*Doklady Akademii Nauk SSSR* 30, 299 and 32, 16) established the modern statistical theory of turbulence. The key hypotheses are:

1. **Local isotropy.** At sufficiently high Reynolds number, the small-scale turbulent motions are statistically isotropic, regardless of the anisotropy of the large-scale forcing.

2. **First similarity hypothesis.** The statistics of small-scale motions are universally determined by the kinematic viscosity ν and the mean energy dissipation rate per unit mass ε.

3. **Second similarity hypothesis (inertial subrange).** For length scales r such that $\eta \ll r \ll L$ (where η is the Kolmogorov dissipation scale and L is the integral scale), the statistics depend only on ε, not on ν.

The Kolmogorov microscale is

$$\eta = \left(\frac{\nu^3}{\varepsilon}\right)^{1/4}.$$

For typical atmospheric turbulence, $\eta \approx 1$ mm; for tornado flows, η can be smaller still.

The famous **five-thirds law** for the energy spectrum in the inertial subrange is

$$E(k) = C_K\, \varepsilon^{2/3}\, k^{-5/3}, \quad \eta^{-1} \gg k \gg L^{-1},$$

where $C_K \approx 1.5$ is the empirical Kolmogorov constant. The $-5/3$ exponent is one of the most robust experimental laws in fluid mechanics, observed across more than ten orders of magnitude in Re from laboratory wind tunnels to atmospheric boundary layers to galactic plasmas.

The cascade picture is captured by Lewis Fry Richardson's verse:

> *"Big whirls have little whirls that feed on their velocity, / And little whirls have lesser whirls and so on to viscosity."*

Energy is injected at large scales by mean-flow shear, transferred conservatively through the inertial subrange by nonlinear vortex stretching, and dissipated into heat by viscosity at scales of order η.

4.6 The Clay Millennium Prize Problem for Navier–Stokes

The Clay Mathematics Institute's official statement asks for proof or counterexample of the following:

> **Navier–Stokes Existence and Smoothness.** For every smooth, divergence-free, finite-energy initial velocity field $\mathbf{v}_0 : \mathbb{R}^3 \to \mathbb{R}^3$, there exists a smooth, globally defined velocity field $\mathbf{v}(\mathbf{x}, t)$ and pressure field $p(\mathbf{x}, t)$ that solve the incompressible Navier–Stokes equations on $\mathbb{R}^3 \times [0, \infty)$ with bounded kinetic energy.

Status as of April 2026: unsolved. The Merle–Raphaël–Rodnianski–Szeftel result on *compressible* Navier–Stokes implosion does not directly resolve the incompressible Millennium Problem, but represents the most important fluid-singularity proof of the past decade and demonstrates that intuitions about viscosity universally preventing singularity must be abandoned in compressible regimes.

4.7 Bibliography for Module 2

- Clay Mathematics Institute, "Navier–Stokes Equation," Millennium Problems: https://www.claymath.org/millennium/navier-stokes-equation/.
- Wikipedia, "Navier–Stokes existence and smoothness"; "Turbulence"; "Energy cascade"; "Vorticity".
- Leray, J. (1934). "Sur le mouvement d'un liquide visqueux emplissant l'espace." *Acta Mathematica* 63, 193–248.
- Kolmogorov, A.N. (1941). "The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers." *Doklady Akademii Nauk SSSR* 30, 299–303.
- Frisch, U. (1995). *Turbulence: The Legacy of A.N. Kolmogorov.* Cambridge University Press.
- Sreenivasan, K.R. (1984, 1995, 2009). Empirical work on dissipation anomaly and Kolmogorov constant.

5 Module 3 Reference: WMO Cloud Atlas — Complete Taxonomy

5.1 Historical Foundation

The present international cloud classification dates to 1803, when the amateur English meteorologist **Luke Howard** published *Essay on the Modification of Clouds*, introducing the four core Latin terms *cirrus* (curl, fiber), *stratus* (layer), *cumulus* (heap), and *nimbus* (rain-bearing). Howard's scheme survived because it was *descriptive of process and form, not just appearance* — it grouped clouds by the way air moves and water condenses, not by the shapes a viewer happened to perceive.

The World Meteorological Organization (WMO) maintains the **International Cloud Atlas** as the legal global standard. The text of the Atlas constitutes Annex I to the Technical Regulations (WMO-No. 49) and has the status of standard practices and procedures binding on member states. The 2017 edition added new official classifications, including the variety *volutus* (the roll cloud) and the supplementary feature *asperitas* (proposed by the Cloud Appreciation Society in 2009 and formally accepted in 2017), among others.

5.2 The Ten Genera

Each observed cloud is a member of *one and only one* genus. Altitude bands below are temperate-latitude averages; clouds are generally lower in polar regions and higher in the tropics.

Genus	Abbr.	Level	Base (km)	Composition / Character
Cirrus	Ci	High	5–13	Ice. White, delicate filaments, patches, narrow bands; fibrous (hair-like) appearance and silky sheen.
Cirrocumulus	Cc	High	5–13	Ice (and supercooled water at lower limit). Thin patch/sheet of small cloudlets, no shading.
Cirrostratus	Cs	High	5–13	Ice. Transparent whitish veil; produces halo phenomena around sun/moon.
Altocumulus	Ac	Middle	2–7	Water (often supercooled), some ice. White/grey patches, sheets, layers of laminae or rolls.
Altostratus	As	Middle	2–7	Water, ice, mixed. Greyish/bluish striated, fibrous, or uniform sheet covering all/most of sky; sun visible as through ground glass.
Nimbostratus	Ns	Multi	2–7 base, often extends down	Water, mixed phase. Grey, often dark, diffuse cloud layer; continuous precipitation; thick enough to blot out sun.
Stratocumulu	Sc	Low	0–2	Water (occasionally ice in cold). Grey/whitish patches/sheets/layers of rounded masses, rolls, with dark parts.
Stratus	St	Low	0–2	Water droplets. Generally grey, uniform base; fog when at the surface; may yield drizzle.
Cumulus	Cu	Low (vertical)	0.5–2 base	Water (ice in tops of large Cu). Detached, dense, sharp-outlined heap clouds; flat base, cauliflower top.
Cumulonimbu	Cb	All levels	0.5–2 base, top to tropopause	Water at base, ice at top; often full mixed phase. Heavy, dense vertical tower with anvil top; thunderstorm cloud.

The three *cirro-* genera (Ci, Cc, Cs) are entirely above 5 km in temperate latitudes and are composed almost exclusively of ice crystals because temperatures aloft are well below freezing. The three *alto-* genera (Ac, As, Ns) sit at intermediate heights and are typically composed of liquid water droplets, including supercooled droplets below freezing. The four low-level genera (Sc, St, Cu, Cb) generally have liquid bases; Cb is the exceptional vertical cloud whose top can exceed 12 km and tap mid-latitude ice processes throughout its column.

5.3 The Fifteen Species

Most genera are subdivided into species based on shape and internal structure. A cloud bears the name of *at most one* species.

Species	Most-frequent genera	Definition
fibratus	Ci, Cs	Detached fibres or filaments, nearly straight or irregularly curved, without hooks or tufts.
uncinus	Ci	Hook-shaped or comma-shaped, terminating in a tuft, the upper part not in the form of a rounded protuberance.
spissatus	Ci	Sufficiently dense to appear greyish when viewed toward the sun.
castellanus	Ci, Cc, Ac, Sc	Turreted cloudlets in the upper part, common base, crenellated battlement appearance, indicates instability.
floccus	Ci, Cc, Ac	Each cloudlet is a small tuft of cumulus form, often with virga falling from it.
stratiformis	Ac, Sc, Cc	Spread out in a horizontal layer or sheet of large extent.
nebulosus	Cs, St	Veil-like cloud showing no distinct details.
lenticularis	Cc, Ac, Sc	Lens-shaped or almond-shaped, often very elongated, with sharp outlines; characteristic of orographic flow.
volutus	Sc, Ac (rare Ci)	Long, typically low, horizontal, detached, tube-shaped cloud mass that appears to roll slowly about a horizontal axis.
fractus	St, Cu	Irregular shreds with ragged appearance; common in pre-frontal precipitation.
humilis	Cu	Flattened cumulus of slight vertical extent (height $\approx$ width); “fair-weather cumulus.”
mediocris	Cu	Moderate vertical extent, with small protuberances and sproutings on the upper part.
congestus	Cu	Marked vertical extent, with bulging protuberances often resembling a cauliflower; can produce showers.
calvus	Cb	At least some upper protuberances are losing their cumuliform appearance and beginning to form a smooth ice mass; no distinct anvil yet.
capillatus	Cb	Distinct cirriform (fibrous, striated) parts in the upper portion; usually accompanied by precipitation, anvil (incus), mamma.

The four cumulus species (humilis, mediocris, congestus, the transition to cumulonimbus calvus then capillatus) form a continuous developmental sequence: a fair-weather

Cu humilis can grow to mediocris, then congestus ("towering cumulus"), then glaciate at the top to become a young Cb calvus, then mature into Cb capillatus with its characteristic anvil. This progression is the visible signature of an unstable atmosphere with sufficient moisture and lift.

5.4 The Nine Varieties

Varieties describe transparency or arrangement; a single cloud can bear multiple varieties simultaneously (e.g., *Altocumulus stratiformis duplicatus translucidus* — a layered Ac in two superposed sheets, both translucent enough to see the sun's outline).

Variety	Most-frequent genera	Definition
intortus	Ci	Filaments very irregularly curved, often seemingly entangled in capricious manner.
vertebratus	Ci	Elements arranged like vertebrae, ribs, or a fish skeleton.
undulatus	Cc, Cs, Ac, As, Sc, St	Patches/sheets/layers showing undulations; sometimes a double system of undulations.
radiatus	Ci, Ac, As, Sc, Cu	Broad parallel bands or arrangement in parallel bands which, by perspective, appear to converge to a horizon point.
lacunosus	Cc, Ac (rare Sc)	Thin patches with regularly distributed round holes giving honeycomb/lace appearance.
duplicatus	Ci, Cs, Ac, As, Sc	Patches or layers at slightly different levels, sometimes partly merged.
translucidus	Ac, As, Sc, St	Greater part is translucent enough to reveal sun or moon position.
perlucidus	Ac, Sc	Distinct (sometimes very small) spaces between elements, allowing sun, moon, or blue sky to be seen.
opacus	Ac, As, Sc, St	Greater part is sufficiently opaque to mask the sun or moon completely.

5.5 Supplementary Features and Accessory Clouds

Supplementary features are attached to or partly merged with a parent cloud. Accessory clouds are smaller separate clouds that occur near or attached to a larger cloud. Selected entries:

Feature	Type	Definition / Physics
incus	Supplemen	Anvil-shaped upper portion of a Cb capillatus, formed when the rising thermal hits the tropopause and spreads horizontally.
mamma	Supplemen	Hanging udder-like protuberances on the under surface; cold pockets descending into clear air through evaporative cooling, common under Cb anvils.
virga	Supplemen	Precipitation falling from the cloud base but evaporating before reaching the ground (visible streaks).
praecipitatio	Supplemen	Precipitation reaching the ground.
arcus	Supplemen	Dense horizontal roll/shelf at the leading edge of a Cb gust front; the visible boundary of the cold outflow.
tuba	Supplemen	Cloud column or inverted cone protruding from a base; if reaching the ground = tornado.
fluctus	Supplemen	Kelvin–Helmholtz instability waves on the top of a cloud layer (curls).
asperitas	Supplemen	Wave-like undulations on the underside, more chaotic and less horizontally organized than undulatus; formally adopted 2017.
cavum	Supplemen	Hole punch in a layer of supercooled liquid cloud; ice-crystal-induced glaciation evacuates the surrounding water.
murus	Supplemen	Wall cloud; localized lowering attached to the rain-free base of a supercell Cb; a precursor to tornado formation.
cauda	Supplemen	Tail cloud; horizontal cloud that connects a wall cloud (murus) to the precipitation core.
pileus	Accessory	Small cap/hood above a rising cumulus; condensation in the orographic-like updraft of the rising thermal.
velum	Accessory	Thin sheet/veil pierced by the rising parent Cu/Cb; the tower has punched through a stable layer.
pannus	Accessory	Ragged shreds in a layer below the parent cloud, typically below precipitating Ns or Cb.
flumen	Accessory	“Beaver tail” inflow cloud; horizontal cloud band feeding moist air into a supercell Cb’s updraft.

5.6 Special Clouds

Homogenitus. Clouds formed as a consequence of human activity: contrails (cirrus homogenitus), cumulus homogenitus from industrial heat sources, stratus homogenitus from cooling-tower exhaust, and others.

Homomutatus. Clouds that form when a homogenitus cloud (typically a contrail) persists long enough that the wind shear distorts it into a different cloud form (often cirrus uncinus or fibratus).

Cataractagenitus. Clouds generated by the local turbulence and moisture release of a major waterfall.

Flammagenitus. Pyrocumulus and pyrocumulonimbus that form above wildfires and volcanic eruptions; can become severe with their own lightning ("dry" thunderstorms).

Silvagenitus. Stratus formed above forest canopies through evapotranspiration.

5.7 Upper-Atmospheric Clouds

Above the troposphere, two formal classes are recognized:

Nacreous (polar stratospheric clouds, PSCs). Form at altitudes 15–25 km in the polar winter stratosphere when temperatures drop below approximately $-78\,^{\circ}$C. Composed of nitric acid trihydrate (Type Ia), supercooled ternary solutions (Type Ib), and pure ice (Type II). Iridescent in the right twilight geometry. Catalyze the heterogeneous chlorine chemistry that destroys stratospheric ozone in the Antarctic spring.

Noctilucent (polar mesospheric clouds, PMCs). The highest clouds in Earth's atmosphere, at $\sim$76–85 km altitude in the summer polar mesosphere. Composed of micron-scale ice crystals nucleated on meteoric dust at temperatures near $-130\,^{\circ}$C. Visible only during deep twilight (sun 6–16° below horizon) when sunlight still illuminates the high-altitude particles against a dark sky.

5.8 Bibliography for Module 3

- World Meteorological Organization, *International Cloud Atlas*: https://cloudatlas.wmo.int/ (Annex I to Technical Regulations, WMO-No. 49).
- NOAA JetStream, "Ten Basic Clouds": https://www.noaa.gov/jetstream/clouds/ten-basic-clouds.
- NOAA JetStream, "The Four Core Types of Clouds."
- SKYbrary Aviation Safety, "Cloud Classification: Genera and Species."
- Wikipedia, "List of cloud types"; "Polar stratospheric cloud"; "Noctilucent cloud."
- Encyclopædia Britannica, "Climate — Cloud types" (altitude bands).
- Howard, L. (1803). *Essay on the Modification of Clouds.*

6 Module 4 Reference: Cloud Microphysics and Atmospheric Thermodynamics

6.1 Saturation Vapor Pressure and Clausius–Clapeyron

The saturation vapor pressure e_s over a flat surface of pure water satisfies the Clausius–Clapeyron relation

$$\frac{de_s}{dT} = \frac{L_v\, e_s}{R_v T^2},$$

where $L_v \approx 2.5 \times 10^6$ J/kg is the latent heat of vaporization and $R_v = 461.5$ J/(kg·K) is the specific gas constant for water vapor.

Integrating with the August–Roche–Magnus approximation gives the practical formula

$$e_s(T)\ (\text{hPa}) \approx 6.112 \exp\left(\frac{17.625\, T}{T + 243.04}\right), \quad T \text{ in } ^\circ\text{C}.$$

Crucially, the saturation vapor pressure over *ice* is lower than over *water* at the same subzero temperature. This difference is the thermodynamic engine of the Wegener–Bergeron–Findeisen process below.

6.2 Adiabatic Ascent and the Lifting Condensation Level

A parcel of unsaturated air rising adiabatically cools at the **dry adiabatic lapse rate**

$$\Gamma_d = \frac{g}{c_p} \approx 9.8\ ^\circ\text{C/km},$$

where g is gravity and c_p the specific heat of air at constant pressure. Condensation begins at the **lifting condensation level (LCL)**, where the parcel temperature equals the dew point. Above the LCL, the parcel cools at the smaller **moist adiabatic lapse rate** $\Gamma_m \approx 4$–7 °C/km, because latent heat release from condensation partially offsets adiabatic cooling.

The LCL is the thermodynamic answer to "why does a cumulus have a flat base?" Every parcel rising from the surface reaches saturation at the same height, so the cloud bases align horizontally regardless of the irregular topography of the rising thermals.

6.3 Köhler Theory of Cloud Droplet Activation

Pure-water condensation does not occur until extreme supersaturations (relative humidity >120% in the laboratory). In the atmosphere, **cloud condensation nuclei (CCN)** — hygroscopic aerosol particles such as sea salt, sulfate, and certain mineral dusts — enable droplet formation at much lower supersaturations.

Köhler's equation (Hilding Köhler, 1936) gives the equilibrium saturation ratio $S = e/e_s$ for a droplet of radius r formed on a soluble nucleus of dissolved solute mass m_s:

$$\ln S(r) \;=\; \underbrace{\frac{2\sigma M_w}{\rho_w R T r}}_{\text{Kelvin (curvature) term}} \;-\; \underbrace{\frac{i\, m_s\, M_w}{M_s\left(\frac{4}{3}\pi r^3 \rho_w - m_s\right)}}_{\text{Raoult (solute) term}},$$

where σ is surface tension, M_w and M_s are molar masses of water and solute, ρ_w

is water density, R the gas constant, and i the van't Hoff factor.

The **Kelvin term** raises the equilibrium vapor pressure for small curved droplets (smaller drops require higher humidity to be stable). The **Raoult term** lowers it (dissolved solute reduces water activity). The two terms produce a curve with a single maximum at the **critical supersaturation** S^* at the **critical radius** r^*. A droplet smaller than r^* at supersaturation below S^* is in stable equilibrium (haze); a droplet pushed above r^* becomes **activated** and grows freely as a cloud droplet.

The implication: a cloud's microstructure is set by the size and chemistry distribution of its CCN. Polluted continental air with abundant CCN produces many small droplets (Twomey indirect effect: brighter, longer-lived clouds); clean maritime air produces fewer larger droplets that more readily collide and coalesce into rain.

6.4 The Wegener–Bergeron–Findeisen (WBF) Process

In a mixed-phase cloud (temperatures between roughly 0 and -38 °C where supercooled liquid and ice can coexist), the difference between saturation vapor pressure over water e_s^w and over ice e_s^i at the same temperature drives a one-way mass transfer. The maximum saturation difference $e_s^w - e_s^i$ occurs near -12 °C.

At equilibrium between liquid droplets and the surrounding vapor, the vapor pressure $e \approx e_s^w$. But $e_s^w > e_s^i$, so the same vapor is *supersaturated* with respect to ice. Ice crystals therefore grow by deposition; this growth depletes the vapor below e_s^w; liquid droplets then evaporate to maintain their saturation. Net result: ice grows at the expense of supercooled liquid.

This mechanism, theorized by Alfred Wegener (1911), elaborated by Tor Bergeron (1933), and refined experimentally by Walter Findeisen (late 1930s), is the dominant precipitation mechanism in mid- and high-latitude clouds. Together with collision–coalescence (next), it explains why most rain we experience originated as snow.

6.5 Collision–Coalescence (Warm Rain)

In purely warm (above-freezing) clouds, especially over tropical oceans, precipitation forms by collision and coalescence of liquid droplets. A typical cloud droplet is $\sim$10 μm radius; a typical raindrop is $\sim$1 mm. Spanning the factor of $\sim$100 in radius (factor of 10^6 in volume) requires accelerated growth.

The fall velocity of a droplet scales roughly as r^2 in the Stokes regime (small drops) and as $\sqrt{r}$ for larger drops. Differential fall speeds allow larger "collector" drops to overtake smaller ones; the **collection efficiency** — the product of geometric collision cross-section, hydrodynamic deflection corrections, and coalescence probability — governs the rate.

6.6 Ice Nucleation Modes

Homogeneous nucleation. Pure water below approximately -38 °C freezes spontaneously without need for a nucleus. This sets the upper limit of supercooling for

cloud droplets and explains why cirrus clouds at very high altitudes are ice-only.

Heterogeneous nucleation. Above $-38\,^{\circ}$C, ice formation requires an **ice nucleating particle (INP)**: mineral dust (especially K-feldspar), biological particles (Pseudomonas syringae, fungal spores), soot, and certain organics. INPs are far rarer than CCN, by typical factors of 10^4 or more; this scarcity is precisely what makes the WBF process efficient — a few ice crystals dispersed among many supercooled droplets.

Heterogeneous modes include: *deposition* (vapor $\to$ ice on dry INP), *condensation freezing* (vapor $\to$ liquid $\to$ ice on hygroscopic INP), *immersion freezing* (INP already in droplet triggers freezing on cooling), and *contact freezing* (INP collides with droplet from outside).

6.7 Aerosol Indirect Effects

Twomey effect (first indirect). More CCN $\to$ more droplets of smaller mean size for the same liquid water content $\to$ greater total droplet surface area $\to$ greater cloud albedo. Polluted clouds reflect more sunlight back to space.

Albrecht effect (second indirect). Smaller droplets coalesce less efficiently $\to$ less precipitation $\to$ longer cloud lifetime $\to$ more time-integrated radiative forcing. Polluted clouds also persist longer.

These indirect effects constitute one of the largest uncertainties in present-day climate sensitivity estimates.

6.8 Bibliography for Module 4

- Lohmann, U.; Lüönd, F.; Mahrt, F. (2016). *An Introduction to Clouds: From the Microscale to Climate.* Cambridge University Press.
- Pruppacher, H.R.; Klett, J.D. (2010). *Microphysics of Clouds and Precipitation* (2nd ed.). Springer.
- Rogers, R.R.; Yau, M.K. (1989). *A Short Course in Cloud Physics.* Pergamon.
- Wikipedia, “Wegener–Bergeron–Findeisen process”; “Köhler theory.”
- Wegener, A. (1911). *Thermodynamik der Atmosphäre.*
- Bergeron, T. (1933). “On the physics of clouds and precipitation,” Proc. 5th Assembly of the Intl. Union of Geodesy & Geophysics, Lisbon.
- Twomey, S. (1977). “The influence of pollution on the shortwave albedo of clouds.” *Journal of the Atmospheric Sciences* 34, 1149–1152.
- Albrecht, B.A. (1989). “Aerosols, cloud microphysics, and fractional cloudiness.” *Science* 245, 1227–1230.

7 Module 5 Reference: Numerical Weather Prediction and AI Forecasting

7.1 Historical Foundations

Bjerknes (1904). The Norwegian physicist Vilhelm Bjerknes published *Das Problem der Wettervorhersage* ("The Problem of Weather Forecasting"), arguing that weather prediction is an initial-value problem in mathematical physics: given the present state of the atmosphere and the governing equations, the future state is in principle determined.

Richardson (1922). Lewis Fry Richardson made the first numerical weather forecast, by hand, using a hydrostatic version of Bjerknes's primitive equations on a grid covering Central Europe. His 6-hour forecast for surface pressure change came out at 145 hPa — about two orders of magnitude too large. The cause was an imbalance between pressure and wind-velocity initial conditions; the diagnosis identified the need for what would later become *data assimilation*.

Charney, Fjørtoft, von Neumann (1950). Using the ENIAC, the team at the Institute for Advanced Study (Princeton) computed the first *successful* numerical forecast — a 24-hour forecast over North America using the barotropic potential vorticity equation. This single result opened the modern era of NWP.

Operational primitive-equation models (1966 onward). West Germany and the United States began running operational primitive-equation forecasts in 1966, joined by the United Kingdom (1972) and Australia (1977). A six-layer hemispheric primitive-equation model came online in 1966; the first global model in 1974.

Chaos and ensemble forecasting (1990s onward). Following Lorenz's identification of the predictability limit (next subsection), operational centers began running **ensemble prediction systems (EPS)**: many forecasts from slightly perturbed initial conditions and physics, providing probabilistic forecasts. ECMWF's EPS launched in 1992; NCEP's GEFS in 1992 also.

AI revolution (2022–2026). A new family of machine-learning weather models trained on the ECMWF ERA5 reanalysis matched and then exceeded the deterministic skill of the world's best NWP systems on standard metrics, running in under a minute on a single TPU.

7.2 The Primitive Equations

The atmospheric primitive equations are a coupled nonlinear PDE system on the rotating Earth, comprising:

1. **Three momentum equations** (Newton's second law including Coriolis force, pressure gradient, gravity, and friction):

$$\frac{D\mathbf{v}}{Dt} = -2\mathbf{\Omega} \times \mathbf{v} - \frac{1}{\rho}\nabla p + \mathbf{g} + \mathbf{F}_{\text{fric}},$$

where $D/Dt = \partial_t + \mathbf{v} \cdot \nabla$ is the material derivative and $\mathbf{\Omega}$ is Earth's rotation vector.

2. **Mass continuity:**
$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0.$$

3. **Thermodynamic energy** (first law of thermodynamics):
$$c_p \frac{DT}{Dt} - \frac{1}{\rho}\frac{Dp}{Dt} = Q,$$
with diabatic heating Q from radiation, latent heat release, and surface fluxes.

4. **Equation of state** (ideal gas):
$$p = \rho R T.$$

5. **Moisture continuity** (one or more species: water vapor, cloud water, cloud ice, rain, snow, graupel):
$$\frac{Dq_v}{Dt} = S_{q_v},$$
with sources/sinks S from microphysical processes.

In practice the equations are written in pressure or hybrid sigma-pressure vertical coordinates and the hydrostatic approximation $\partial p/\partial z = -\rho g$ is invoked at scales larger than $\sim$10 km; non-hydrostatic models retain full vertical momentum at convection-permitting resolutions.

7.3 Numerical Methods

Discretization. Spatial discretization uses finite differences on **Arakawa A–E grids** (with the Arakawa C-grid being most common for its momentum–mass conservation properties), spectral transforms (spherical harmonics in global models), or finite-volume methods. Time integration uses explicit, semi-implicit, or split-explicit schemes.

The Courant–Friedrichs–Lewy (CFL) condition. For an explicit scheme to be stable, the time step Δt must satisfy
$$\Delta t \leq \frac{\Delta x}{c},$$
where c is the fastest wave speed (sound waves in compressible models, gravity waves in shallow-water models). For a 1 km grid with sound speed 340 m/s, $\Delta t < 3$ s. This drives the use of semi-implicit schemes that treat the fastest waves implicitly.

Sub-grid-scale parameterization. Processes smaller than the grid must be *parameterized* — represented statistically in terms of resolved variables. Major parameterization categories:

- **Convection** (Kuo, Arakawa–Schubert, Tiedtke, Kain–Fritsch, Zhang–McFarlane): cumulus and Cb representation in coarse-grid models.
- **Cloud microphysics** (single-moment, double-moment, bin schemes): the WBF and collision–coalescence physics from Module 4 implemented as bulk transfer rates.
- **Radiation** (RRTMG, FLG): shortwave and longwave radiative transfer through clear and cloudy atmosphere.
- **Boundary layer** (YSU, Mellor–Yamada–Janjic, ACM2): turbulent fluxes of heat, moisture, and momentum near the surface.
- **Land surface** (Noah, Noah-MP, CLM): soil moisture, vegetation, snowpack, and surface energy balance.
- **Gravity wave drag**: orographic and non-orographic gravity wave breaking momentum deposition.

7.4 Data Assimilation

The atmosphere is observed by an inhomogeneous patchwork of sources: radiosondes (twice-daily soundings), surface stations, ships and buoys, commercial aircraft, weather radar, ground GPS, and — dominantly — satellites. Globally, more than 200 million observations per day are processed at major centers.

The data assimilation problem. Combine the observations $\mathbf{y}^o$, the previous short-range forecast (*background*) $\mathbf{x}^b$, and their error statistics to produce the best estimate of the current atmospheric state (*analysis*) $\mathbf{x}^a$.

Optimal interpolation (1960s–80s). Linear regression onto observation departures with empirically prescribed background error covariance.

3D-Var. Minimize the cost function

$$J(\mathbf{x}) = \tfrac{1}{2}(\mathbf{x}-\mathbf{x}^b)^T\mathbf{B}^{-1}(\mathbf{x}-\mathbf{x}^b) + \tfrac{1}{2}(\mathbf{y}^o - H(\mathbf{x}))^T\mathbf{R}^{-1}(\mathbf{y}^o - H(\mathbf{x})),$$

where $\mathbf{B}$ is background error covariance, $\mathbf{R}$ is observation error covariance, and H is the (possibly nonlinear) observation operator mapping model state to observation space.

4D-Var. Same cost function generalized over a time window using the model's tangent linear and adjoint to propagate the analysis increment through time. Operational at ECMWF since 1997 and Météo-France since 2000.

Ensemble Kalman Filter (EnKF). Use a Monte Carlo ensemble of forecasts to estimate the flow-dependent background error covariance $\mathbf{B}$, then apply the Kalman update. Introduced by Evensen (1994); operational at multiple centers.

Hybrid 4DEnVar. Combine the static **B** of variational methods with the flow-dependent **B** of the EnKF. NCEP implemented this operationally for the Global Forecast System (GFS) in 2016.

7.5 Predictability and the Lorenz Limit

Edward Lorenz's 1963 paper *Deterministic Nonperiodic Flow* (*J. Atmos. Sci.* 20, 130–141) introduced the three-equation "Lorenz system"

$$\begin{aligned}\dot{x} &= \sigma(y - x),\\ \dot{y} &= x(\rho - z) - y,\\ \dot{z} &= xy - \beta z,\end{aligned}$$

with classical parameters $\sigma = 10$, $\beta = 8/3$, $\rho = 28$.

For these parameters the system possesses a strange attractor (the "Lorenz butterfly") on which trajectories diverge exponentially. This is the canonical demonstration of **sensitive dependence on initial conditions** — the butterfly effect, popularized by Lorenz's 1972 talk "Predictability: Does the Flap of a Butterfly's Wings in Brazil Set Off a Tornado in Texas?"

The practical consequence: even with a perfect model and arbitrarily good (but finite-precision) initial conditions, atmospheric forecasts lose skill exponentially, with a doubling time of roughly 1–2 days for synoptic-scale features. This caps useful deterministic forecasts at about 14 days. Beyond that, only statistical (climate) forecasting is meaningful.

7.6 Operational NWP Systems (2026)

System	Center	Notes
IFS / HRES	ECMWF (Reading, UK)	Long the gold standard for medium-range deterministic. 9 km horizontal grid; 4D-Var with hybrid EnVar.
IFS / ENS	ECMWF	51-member ensemble; 18 km grid; out to 15 days.
GFS	NCEP (USA)	Global Forecast System; 13 km grid; hybrid 4DEnVar.
GEFS	NCEP	Global Ensemble Forecast System; 31 members.
UM	UK Met Office	Unified Model; same code for global and limited-area, NWP and climate.
ICON	DWD (Germany)	Icosahedral grid, non-hydrostatic.
ARPEGE / AROME	Météo-France	Global ARPEGE plus regional convection-resolving AROME.
GDPS / CMC	Environment Canada	Global Deterministic Prediction System.
JMA GSM	Japan Meteorological Agency	Spectral global model.
GRAPES	China Meteorological Admin.	Global/Regional Assimilation and Prediction System.

7.7 The AI Weather Models (2022–2026)

A new generation of machine-learning weather prediction (MLWP) systems, trained on the ECMWF ERA5 reanalysis (43 years of hourly global atmospheric state at 0.25° resolution), now matches or exceeds the IFS deterministic skill on most standard metrics, while running in under a minute on a single accelerator.

Model	Developer	Architecture and Notable Properties
FourCastNet (FCN)	NVIDIA	2022; Adaptive Fourier Neural Operator (AFNO); first MLWP at ∼IFS scale.
Pangu-Weather	Huawei	2022; 3D Earth-Specific Transformer (3DEST); ∼256M parameters; first AI model to outperform IFS-HRES on all variables and lead times to one week.
GraphCast	Google DeepMind	2023 (*Science*); Graph Neural Network on icosahedral multimesh; outperformed HRES on 90% of 1380 verification targets at 0.25°.
FourCastNet v2 (FCN2)	NVIDIA	Spherical Harmonics Neural Operator; more stable autoregressive rollouts.
FengWu	Shanghai AI Lab et al.	Multi-modal cross-modal fusion transformer with uncertainty loss; best-performer in regional inter-comparisons.
FuXi	Fudan University	Cascade architecture optimized per lead-time band.
AIFS	ECMWF	ECMWF's own production AI model; graph neural network; running operationally alongside IFS.
GenCast	Google DeepMind	2024–25; first competitive *probabilistic* AI ensemble; matches or exceeds ECMWF ENS on most metrics.

Strengths. Speed (sub-minute 10-day forecasts), competitive medium-range deterministic skill, accurate cyclone tracking, energy efficiency.

Weaknesses (as of 2026).

- **Extreme precipitation underestimation**: GraphCast and Pangu-Weather underestimate 99th-percentile precipitation events by 20–35%, versus 10–15% for IFS-HRES (Hamill et al., 2024).
- **Smoothing**: forecast fields are spectrally too smooth at long lead times, producing low-bias of extremes generally.
- **Out-of-distribution failures**: novel regimes (unprecedented Arctic ice loss, regime shifts) fall outside training data.
- **Convective precipitation**: high-resolution NWP with explicit convection still beats AI on 1–6 hour rainfall.
- **No clear path to "physics from data only"**: AI models *depend* on NWP-produced ERA5 reanalysis for training; without traditional NWP and the global observation network, AI weather has no training signal.

Operational integration. As of 2026, AI weather models run alongside traditional NWP at major centers. ECMWF runs AIFS in production; NOAA's National Hurricane

Center references GraphCast and Pangu-Weather in internal forecast discussions. The dominant workflow is *multi-model comparison* — AI models are one more voice in the discussion alongside deterministic and ensemble NWP, with model disagreement itself treated as a signal of forecast uncertainty.

7.8 Bibliography for Module 5

- Bjerknes, V. (1904). "Das Problem der Wettervorhersage betrachtet vom Standpunkt der Mechanik und Physik." *Meteorologische Zeitschrift* 21, 1–7.
- Richardson, L.F. (1922). *Weather Prediction by Numerical Process.* Cambridge University Press.
- Charney, J.G.; Fjørtoft, R.; von Neumann, J. (1950). "Numerical integration of the barotropic vorticity equation." *Tellus* 2, 237–254.
- Lorenz, E.N. (1963). "Deterministic nonperiodic flow." *Journal of the Atmospheric Sciences* 20, 130–141.
- Bauer, P.; Thorpe, A.; Brunet, G. (2015). "The quiet revolution of numerical weather prediction." *Nature* 525, 47–55.
- Pu, Z.; Kalnay, E. (2019). "Numerical weather prediction basics: models, numerical methods, and data assimilation," in *Handbook of Hydrometeorological Ensemble Forecasting*, Springer.
- Bi, K. et al. (2023). "Pangu-Weather: A 3D high-resolution model for fast and accurate global weather forecast." arXiv:2211.02556.
- Lam, R. et al. (2023). "Learning skillful medium-range global weather forecasting" (GraphCast). *Science* 382, 1416–1421.
- Price, I. et al. (2025). GenCast probabilistic ensemble.
- Charlton-Perez, A. et al. (2024). "Do AI models produce better weather forecasts than physics-based models? A quantitative evaluation case study of Storm Ciarán." *npj Climate and Atmospheric Science*.
- Wikipedia, "Numerical weather prediction"; "Data assimilation"; "Lorenz system."

8 Module 6 Reference: Open Problems — Erdős, Millennium, and Physics

8.1 The Erdős Problems Database

Paul Erdős (1913–1996), one of the most prolific mathematicians in history, posed thousands of open problems across number theory, combinatorics, graph theory, and set theory. Many were stated as bets ranging from $10 to $10,000 and persist as open challenges decades after his death.

The site https://www.erdosproblems.com, curated by Fields Medalist **Terence Tao** and a community of contributors, maintains a structured database of these problems with formal statements, known partial results, historical context, and cross-references. The companion GitHub repository `teorth/erdosproblems` also tracks AI contributions in a wiki at `wiki/AI-contributions-to-Erdős-problems`.

8.2 Erdős Problem #1196 — Solved by AI, April 2026

The conjecture. A set $A \subset \mathbb{N}$ is called **primitive** if no element of A divides any other. The **Erdős–Sárközy–Szemerédi conjecture**, listed as Erdős Problem #1196, asks:

> For any x, if $A \subset [x, \infty)$ is a primitive set of integers, is it true that
>
> $$\sum_{a \in A} \frac{1}{a \log a} \;<\; 1 + o(1),$$
>
> where the $o(1)$ term tends to zero as $x \to \infty$?

The example $A = \{p : p \text{ prime}\}$ achieves $\sum 1/(p \log p) \approx 1.636$ at $x = 2$ but the relevant comparison is the behavior as $x \to \infty$. Erdős himself proved (1934) that the infinite sum $\sum_{a \in A} 1/(a \log a)$ converges for any primitive A, the famous *Erdős primitive sum*.

Prior bounds. Jared Lichtman proved in 2023 that

$$\sum_{a \in A} \frac{1}{a \log a} \;<\; e^{\gamma} \cdot \frac{\pi}{4} + o(1) \approx 1.399 + o(1),$$

where $\gamma \approx 0.5772$ is the Euler–Mascheroni constant. Lichtman also showed (2020) that for A_k, the set of integers with exactly k prime factors, $\sum_{a \in A_k} 1/(a \log a) \geq 1 + O(k^{-1/2+o(1)})$, making the conjectural bound of 1 tight up to lower-order terms. Gorodetsky, Lichtman, and Wong (2024) refined this to $\sum = 1 - (c + o(1))k^2 2^{-k}$ with $c \approx 0.0656$.

Resolution by GPT-5.4 Pro (April 13, 2026). OpenAI's GPT-5.4 Pro, prompted by Eric Price, produced a one-shot proof after approximately 80 minutes of reasoning. The proof establishes the stronger statement

$$\sum_{\substack{a \in A \\ a > x}} \frac{1}{a \log a} \;\leq\; 1 + O\left(\frac{1}{\log x}\right)$$

for any primitive set $A \subset \mathbb{N}$, immediately implying the conjecture. As Lichtman commented:

> *"I care deeply about this problem, and I've been thinking about it for the past 7 years. I'd frequently talk to Maynard about it in our meetings, and consulted over the years with several experts... The problem was not a question of low-visibility per se. Rather, it seems like a proof which becomes strikingly compact post-hoc, but the construction is quite special among many similar variations."*

Status. Marked **formally solved** on `erdosproblems.com` (page last edited April 15, 2026); a Lean formalization exists in the Google DeepMind `formal-conjectures` repository.

8.3 Other AI Contributions to Erdős Problems (2025–2026)

The `teorth/erdosproblems` wiki tracks the following recent significant AI contributions:

Erd. #	AI System	Date	Outcome
1196	GPT-5.4 Pro	13 Apr 2026	Full solution; literature presence comparable, found afterwards.
1202	GPT-5.4 Pro	1 Apr 2026	Full solution.
120	DeepMind prover agent	14 Mar 2026	Solution to variant problem (Lean).
152	DeepMind prover agent	3 Apr 2026	Full solution (Lean); literature found later (Erdős, Sárközy, Sós 1994).
218	GPT	28 Jan 2026	Conditional solution.
281	GPT-5.2 Pro	17 Jan 2026	Full solution.
333	Claude Opus 4.5, GPT-5.2 Pro	—	Listed contribution.

The wiki includes important caveats: it is not a benchmark; some problems were initially stated incorrectly; absence of past progress may reflect obscurity rather than difficulty; and an AI proof does not automatically qualify as a publishable journal paper. A central methodological concern is *survivorship bias* — only the successful AI runs are reported, so the underlying success rate is unknown.

8.4 The Seven Clay Millennium Prize Problems

In May 2000, the Clay Mathematics Institute announced seven problems with $1 million prizes. As of April 2026, only one is solved.

Problem	Status (Apr 2026)	Brief Statement
Poincaré Conjecture	**Solved** (Perelman, 2003)	Every simply connected closed 3-manifold is homeomorphic to the 3-sphere.
Riemann Hypothesis	Open	All non-trivial zeros of the Riemann zeta function lie on the critical line $\Re(s) = 1/2$.
P vs NP	Open	Whether problems whose solutions can be verified in polynomial time can also be *solved* in polynomial time.
Navier–Stokes Existence & Smoothness	Open	For every smooth, finite-energy, divergence-free $\mathbf{v}_0 \in \mathbb{R}^3$, do smooth global solutions exist?
Yang–Mills Existence & Mass Gap	Open	Construct a quantum non-Abelian gauge theory on $\mathbb{R}^4$ and prove a positive mass gap.
Birch & Swinnerton-Dyer Conjecture	Open	The rank of the group of rational points on an elliptic curve equals the order of vanishing of its L-function at $s = 1$.
Hodge Conjecture	Open	Every Hodge class on a non-singular complex projective variety is a rational linear combination of cohomology classes of algebraic cycles.

Grigori Perelman (2003) proved the Poincaré conjecture using Hamilton's Ricci flow with surgery; he was awarded but *declined* both the Fields Medal (2006) and the $1M Millennium Prize (2010), citing his belief that Hamilton's contribution was equally fundamental.

8.5 Other Major Open Problems in Mathematics

- **Twin Prime Conjecture.** There exist infinitely many primes p such that $p+2$ is also prime. Zhang (2013) showed infinitely many prime pairs differ by at most $7 \cdot 10^7$; Maynard, Tao, and the Polymath project reduced this gap dramatically (now under 250).

- **Goldbach Conjecture.** Every even integer greater than 2 is the sum of two primes. Verified computationally to $4 \cdot 10^{18}$; Helfgott (2013) proved the related ternary Goldbach conjecture.

- **Collatz Conjecture (3n+1).** For every positive integer n, the iteration $n \mapsto n/2$ if even, $3n + 1$ if odd, eventually reaches 1. Tao (2019) proved a probabilistic "almost all" version.

- **abc Conjecture.** For any $\epsilon > 0$, only finitely many coprime triples (a, b, c) with $a + b = c$ satisfy $c > \mathrm{rad}(abc)^{1+\epsilon}$. Mochizuki claimed proof via inter-universal Teichmüller theory (2012, published 2021); the international community has not reached consensus on the proof's validity.

- **Generalized Riemann Hypothesis (GRH)**, the Langlands program, the Hodge–Newton inequality, the Kakeya conjecture, the Sato–Tate conjecture (proved 2008–2011 in various cases).

8.6 Major Open Problems in Physics (2026)

Quantum Gravity. Reconcile general relativity with quantum mechanics. Candidates include string theory, loop quantum gravity, causal dynamical triangulations, asymptotic safety, and emergent / entropic gravity proposals. The April 2026 publication by Sharmila, Vermeulen, and Datta (*Nature Communications*) unified detection schemes for spacetime fluctuations across LIGO, QUEST, and GQuEST tabletop experiments.

Dark Matter. Constitutes ∼27% of the universe's mass-energy content; gravitationally evident but not directly detected. Candidate identities include WIMPs (weakly interacting massive particles), axions, sterile neutrinos, primordial black holes, or modifications of gravity (MOND, TeVeS).

Dark Energy. Constitutes ∼68%; drives accelerating cosmic expansion. The cosmological-constant interpretation suffers from a ∼120-order-of-magnitude discrepancy between QFT vacuum energy estimates and observed value.

Hubble Tension. Local measurements of H_0 (∼73 km/s/Mpc from Type Ia supernovae) and CMB-based inferences (∼67 km/s/Mpc from Planck under ΛCDM) disagree at $\geq 5\sigma$; the discrepancy resists straightforward systematic explanations.

Matter–Antimatter Asymmetry (Baryogenesis). The observed universe is overwhelmingly matter; mechanisms (Sakharov conditions, leptogenesis, electroweak baryogenesis) remain unconfirmed.

Neutrino Mass and Nature. Neutrinos have nonzero mass (oscillation experiments), but absolute mass scale is unknown. Whether neutrinos are Dirac or Majorana fermions is open; neutrinoless double-beta-decay experiments (KATRIN, KamLAND-Zen, LEGEND) test this.

Strong CP Problem. QCD permits a CP-violating θ-term; experiment constrains $|\theta| \lesssim 10^{-10}$. Why so small? Peccei–Quinn symmetry and the axion are the leading candidate explanation.

Hierarchy Problem. Why is the electroweak scale (∼246 GeV) so much smaller than the Planck scale ($\sim 10^{19}$ GeV)? Supersymmetry was the leading proposed answer; null results from the LHC have weakened this. Composite Higgs and extra-dimensional models continue.

Measurement Problem in Quantum Mechanics. The relationship between unitary Schrödinger evolution and apparent wave-function collapse remains unresolved at the foundational level. Interpretations include Copenhagen, Many-Worlds, decoherence, GRW, Bohmian mechanics, QBism, and relational QM.

Color Confinement. An analytical proof of confinement in non-Abelian gauge theory — and more broadly the construction of quantum Yang–Mills with mass gap — is itself one of the seven Millennium Problems.

Theory of Everything. Whether a single coherent framework can unify all fundamental forces and matter remains the deepest open question in physics. String theory, M-theory, and various quantum-gravity programs offer partial paths.

8.7 Bibliography for Module 6

- Tao, T. (curator). *Erdős Problems Database.* https://www.erdosproblems.com/.
- `teorth/erdosproblems` GitHub wiki, "AI contributions to Erdős problems."
- Erdős, P. (1934). "A theorem of Sylvester and Schur." *Journal of the London Mathematical Society* 9, 282–288.
- Lichtman, J.D. (2023). "A proof of the Erdős primitive set conjecture." *Forum of Mathematics, Pi*.
- Gorodetsky, O.; Lichtman, J.D.; Wong, M.W. (2024). Refined estimates on Erdős primitive sums.
- Clay Mathematics Institute, "Millennium Problems": https://www.claymath.org/millennium-problems/.
- Wikipedia, "Millennium Prize Problems"; "List of unsolved problems in physics"; "List of unsolved problems in mathematics."
- Sharmila, B.; Vermeulen, S.M.; Datta, A. (2026). "Signatures of correlation of spacetime fluctuations in laser interferometers." *Nature Communications* 17, 67313.
- Helfgott, H.A. (2013). "The ternary Goldbach conjecture is true." arXiv:1312.7748.
- Tao, T. (2019). "Almost all orbits of the Collatz map attain almost bounded values." arXiv:1909.03562.

9 Cross-Module Synthesis (Reference)

The six modules form a closed loop of mathematical and physical understanding from abstract PDEs through cloud taxonomy to the open frontier:

1. **M1 → M2:** Merle's compressible-Euler implosion proof (M1) sits inside the broader fluid-singularity question (M2) of which the incompressible Navier–Stokes Millennium Problem is the most famous instance.

2. **M2 → M5:** The same Navier–Stokes equations whose regularity is unproven (M2) are integrated numerically every six hours by operational NWP centers worldwide (M5) — a striking case of "engineering precedes mathematics."
3. **M2 → M3 → M4:** Atmospheric flow (M2) produces clouds (M3) through microphysical processes (M4) that NWP must parameterize (M5).
4. **M5 → M2:** Lorenz's chaos (M5) emerged from a simplified model of atmospheric convection (M2), demonstrating that the predictability limit is intrinsic to the equations themselves.
5. **M6 ↔ All:** The Navier–Stokes Millennium Problem (M6) is the open question at the heart of M2; the Erdős frontier (M6) shows where AI is succeeding while the fluid frontier (M2, M5) shows where it is not yet; the open physics problems (M6) include the quantum-gravity questions that bound any *theory of everything* that would also have to explain why the atmospheric Navier–Stokes equations generate the clouds we see.

10 Safety and Sensitivity Considerations (All Modules)

Concern	Mitigation	Boundary
Mathematical claim accuracy	Every theorem statement and equation form verified against the original paper or a standard textbook. No paraphrased "proofs" presented as proven.	ABSOLUTE
Attribution of AI contributions	GPT-5.4 Pro #1196 result presented with full source attribution (Lichtman, Price, erdosproblems.com); methodological caveats (survivorship bias, no peer review yet) explicitly noted.	GATE
Climate / weather projections	Only data and projections from authoritative agencies (NOAA, ECMWF, IPCC, WMO) cited; no policy advocacy.	GATE
Open problem status	Status as of April 2026 explicitly dated; statements of "open" or "solved" attributed to authoritative source (Clay Math Institute, erdosproblems.com).	GATE
WMO classification	Latin names spelled correctly, including post-2017 additions; cited to the official Cloud Atlas Annex I.	ABSOLUTE
Polar stratospheric / noctilucent clouds	Discussed as natural phenomena; ozone-depletion mechanism described scientifically without geoengineering or weather-modification advocacy.	ANNOTATE

STAGE 3 — MISSION PACKAGE

11 Milestone Specification

11.1 Mission Objective

Produce a six-module deep-research deliverable covering Frank Merle's body of work in nonlinear evolution equations, the mathematics and physics of fluid dynamics, the complete WMO cloud taxonomy, cloud microphysics, numerical weather prediction (including the 2022–2026 AI weather models), and the present landscape of open problems in mathematics (with deep treatment of Erdős Problem #1196) and physics. The deliverable is a single publication-quality reference document, source-grounded throughout, with explicit cross-module synthesis.

11.2 Architecture Overview

```
WAVE EXECUTION ARCHITECTURE
Wave 0 (Foundation) Sequential, Haiku scaffolding +-----------------------------------
| shared schemas, source index, color/typography | +-----------------------------------
| v Wave 1 (Parallel research) Two parallel tracks +-------------------+
+-------------------------+ | Track A: M1+M2+M6 | | Track
B: M3+M4+M5 | | (math/PDE/open) | | (atmospheric/NWP) | |
Sonnet+Opus mix | | Sonnet+Opus mix | +-------------------+
+-------------------------+  /  / v v Wave 2 (Synthesis) Sequential,
Opus +-------------------------------------------------+ | Cross-module
integration; through-line | +-------------------------------------------------+
| v Wave 3 (Publish + Verify) Sequential, Sonnet +-----------------------------------
| LaTeX assembly, 3-pass XeLaTeX, verification | +-----------------------------------
```

11.3 System Layers

1. **Foundation Layer (Wave 0)** — shared schemas (citation format, glossary, color tokens, typography), source index, document skeleton.
2. **Research Layer (Wave 1)** — six parallel module surveys with verification of every cited source.
3. **Synthesis Layer (Wave 2)** — cross-module integration, the through-line narrative, the M1→M6 closing arc.
4. **Publication Layer (Wave 3)** — LaTeX assembly with three-pass XeLaTeX compilation, hyperlink verification, page-number / TOC consistency.
5. **Verification Layer (Wave 3, parallel)** — success-criteria mapping, safety-test execution, source-quality audit.

11.4 Deliverables

#	Deliverable	Acceptance Criteria	Component Spec
D1	Vision Document	8–12 success criteria; Fleet chipset; through-line	01-vision.tex
D2	M1 Reference (Merle)	4 contribution streams; awards; equation forms	02-m1-merle.tex
D3	M2 Reference (Fluids)	NSE/Euler stated; K41 derivation; CMI quote	03-m2-fluids.tex
D4	M3 Reference (Cloud Atlas)	10 genera + 15 species + 9 varieties + features	04-m3-clouds.tex
D5	M4 Reference (Microphysics)	Köhler eq; WBF; collision-coalescence	05-m4-microphys.tex
D6	M5 Reference (NWP)	Primitive eqs; DA; Lorenz; AI models 2022–2026	06-m5-nwp.tex
D7	M6 Reference (Open Problems)	Erdős #1196 details; 7 Millennium; physics	07-m6-open.tex
D8	Cross-Module Synthesis	Explicit M1→M6 connections	08-synthesis.tex
D9	Mission Package (Stage 3)	Waves, components, tests, crew	09-mission.tex
D10	Compiled PDF	3 XeLaTeX passes; clean compile; LastPage works	build artifact
D11	index.html landing	Download links; color match to LaTeX scheme	index.html

11.5 Component Breakdown

Component	Spec Document	Dependencie	Model	Est. Tokens
Vision	01-vision.tex	none	Opus	18 K
M1: Merle	02-m1-merle.tex	Vision	Opus	22 K
M2: Fluids	03-m2-fluids.tex	Vision, M1	Opus	24 K
M3: Cloud Atlas	04-m3-clouds.tex	Vision	Sonnet	28 K
M4: Microphys.	05-m4-microphys.tex	Vision, M3	Sonnet	22 K
M5: NWP	06-m5-nwp.tex	Vision, M2, M3, M4	Sonnet	26 K
M6: Open Probs.	07-m6-open.tex	Vision, M2	Opus	24 K
Synthesis	08-synthesis.tex	all M1–M6	Opus	14 K
Mission Spec	09-mission.tex	Vision (read-only)	Sonnet	16 K
LaTeX assembly	build pipeline	D1–D9	Haiku	6 K
Verification	test plan	D10	Sonnet	12 K
Total				**∼212 K**

11.6 Model Assignment Rationale

Model	% of Tokens	Why
Opus	∼48%	Vision narrative; M1 (mathematical biography needs nuance); M2 (PDE technical content); M6 (open-problem judgment); cross-module synthesis. Judgment-heavy and accuracy-critical.
Sonnet	∼49%	M3 (taxonomic compilation, large but structured); M4 (technical compilation); M5 (large structured survey); mission spec; verification. Structured assembly.
Haiku	∼3%	LaTeX shell, color tokens, citation format scaffolding. Pure templating.

The split is approximately 48 / 49 / 3 Opus / Sonnet / Haiku, slightly more Opus-heavy than the typical 30/60/10 because of the judgment density in M1 (prize citations and biographical accuracy), M2 (PDE statements that must be mathematically correct), and M6 (open-problem status, which is fast-changing and requires careful attribution).

12 Wave Execution Plan

12.1 Wave Summary

Wave	Purpose	Mode	Components	Model
0	Foundation	Sequential	shared schemas, source index, skeleton	Haiku
1A	Math/Open track	Parallel	M1 (Merle), M2 (Fluids), M6 (Open)	Opus
1B	Atmospheric track	Parallel	M3 (Atlas), M4 (Microphys.), M5 (NWP)	Sonnet+Opus
2	Synthesis	Sequential	cross-module through-line	Opus
3	Publish + Verify	Sequential	LaTeX, compile, test, index.html	Sonnet+Haiku

12.2 Wave 0: Foundation

Task	Description	Model
W0.1 LaTeX skeleton	Preamble, packages, color scheme, custom environments.	Haiku
W0.2 Source index	Cross-checked list of authoritative sources for each module.	Sonnet
W0.3 Citation format	Verify consistent citation style across all modules.	Haiku
W0.4 Glossary	Shared terminology table (PDE terms, WMO Latin, NWP acronyms).	Sonnet

12.3 Wave 1A: Math & Open Problems Track (parallel with 1B)

Task	Description	Model
W1A.1 M1: Merle	Biography, awards table, four contribution streams with equation-form verification.	Opus
W1A.2 M2: Fluids	NSE/Euler/vorticity equations; Reynolds; Leray–Hopf; K41 with $-5/3$ derivation; Millennium Problem statement.	Opus
W1A.3 M6: Open Probs	Erdős #1196 detailed, AI-contribution table, 7 Millennium Problems, math & physics open questions.	Opus

12.4 Wave 1B: Atmospheric Track (parallel with 1A)

Task	Description	Model
W1B.1 M3: Cloud Atlas	10 genera + 15 species + 9 varieties + supplementary features + special + upper-atmospheric.	Sonnet
W1B.2 M4: Microphys.	Clausius-Clapeyron, LCL, Köhler theory, WBF, collision-coalescence, ice nucleation, indirect effects.	Sonnet
W1B.3 M5: NWP	History (Bjerknes-Richardson-Charney), primitive equations, DA, Lorenz, operational systems, AI models 2022-2026.	Opus

12.5 Wave 2: Synthesis

Task	Description	Model
W2.1 Cross-references	Build the M1→M2→M5→M6 connection chain explicitly.	Opus
W2.2 Through-line	Write the synthesis narrative connecting Merle's solitons, Kolmogorov's cascade, WMO's taxonomy, NWP's chaos limit, and the open frontier.	Opus
W2.3 Closing quote box	Domain-relevant closing quote with through-line echo.	Opus

12.6 Wave 3: Publish + Verify

Task	Description	Model
W3.1 LaTeX assembly	Concatenate D1–D9 into mission.tex.	Haiku
W3.2 XeLaTeX pass 1	First compile; resolves `\section` numbering, populates .toc.	Haiku
W3.3 XeLaTeX pass 2	Second compile; resolves TOC, cross-references, lastpage.	Haiku
W3.4 XeLaTeX pass 3	Third compile; final hyperlink and lastpage resolution.	Haiku
W3.5 Verification	Execute test plan; map success criteria to outcomes.	Sonnet
W3.6 index.html	Build landing page with download links and color-matched header.	Sonnet

12.7 Cache Optimization Strategy

- **Wave 0 outputs cached** into the prompt for all Wave 1 agents (LaTeX skeleton, source index, glossary, citation format).
- **Wave 1 outputs cached** for Wave 2 (synthesis reads all six module references from cache rather than re-loading).

- **Five-minute cache TTL exploited**: parallel Wave 1A and Wave 1B launched within the same five-minute window so both share the cached Wave 0 foundation.
- **Citation database persisted**: built once in Wave 0, referenced by all module agents.

12.8 Token Budget

Wave	Opus	Sonnet	Haiku	Wave Total
0 (Foundation)	0	4 K	6 K	10 K
1A (Math/Open)	70 K	0	0	70 K
1B (Atmospheric)	26 K	50 K	0	76 K
2 (Synthesis)	14 K	0	0	14 K
3 (Publish + Verify)	0	28 K	8 K	36 K
Total	**110 K**	**82 K**	**14 K**	**~206 K**

12.9 Dependency Graph

```
[Wave 0] shared foundation | +---------------+---------------+ | | [Wave
1A] [Wave 1B] math/open track atmospheric track +-----+-----+-----+
+-----+-----+-----+ | M1 | M2 | M6 | | M3 | M4 | M5 | +-----+-----+-----+
+-----+-----+-----+   / / / /   / / / /   / / /  / / / /  /  / / / v
v v v v +---------------------------------+ | [Wave 2: Synthesis] | |
cross-module through-line | +---------------------------------+ | v
+---------------------------------+ | [Wave 3: Publish+Verify] | | LaTeX,
3 passes, tests, html | +---------------------------------+
```

13 Test Plan

13.1 Test Categories

Category	Count	Priority	Action	Notes
Safety-critical	7	Mandatory	BLOCK	Source quality, equation accuracy, attribution.
Core functional	18	Required	BLOCK	Module-by-module content acceptance.
Integration	8	Required	BLOCK	Cross-module references resolve.
Edge cases	6	Best-effort	LOG	Robustness, formatting, boundary cases.
Total	**39**			2–4 tests per success criterion.

13.2 Safety-Critical Tests

ID	Verifies	Expected
SC-SRC	Source quality	Every citation traces to government agency, peer-reviewed journal, professional org, or curated authoritative database (erdosproblems.com, claymath.org). Zero entertainment media.
SC-NUM	Numerical attribution	Every figure (H_0 values, percentages, dates, prize amounts) attributed to specific source.
SC-EQN	Equation form correctness	NSE, Euler, Köhler, Kolmogorov $E(k)$, and Lorenz system equations match canonical references.
SC-DATE	Time-sensitive claim dating	Erdős #1196 status, Breakthrough Prize 2026, Millennium Problem status all dated with "as of April 2026."
SC-ATTR	AI contribution honesty	GPT-5.4 Pro #1196 result attributed to OpenAI/Price/Lichtman; survivorship-bias caveat present.
SC-WMO	WMO Latin spelling and authority	Every cloud Latin name spelled correctly; cited to WMO Cloud Atlas Annex I.
SC-ADV	No policy advocacy	Climate, AI-safety, weather-modification topics presented as evidence without policy recommendations.

13.3 Core Functional Tests (sample)

ID	Verifies	Expected
CF-M1.1	Merle biographical completeness	Birth date, ENS, PhD year+advisor, Cergy chair, IHES chair, all awards through 2026 Breakthrough Prize.
CF-M1.2	Four contribution streams	Each stream named, dated, with primary citation.
CF-M2.1	NSE statement correctness	Both momentum and continuity equations stated with all symbols defined.
CF-M2.2	K41 derivation	Energy spectrum $E(k) \sim \varepsilon^{2/3}k^{-5/3}$ stated with inertial subrange bounds.
CF-M3.1	10 genera complete	All ten genera present with abbreviation, level, base altitude, composition.
CF-M3.2	15 species complete	All fifteen species in the table; matched to most-frequent genera.
CF-M3.3	9 varieties complete	All nine varieties; matched to most-frequent genera.
CF-M3.4	Supplementary features	At least 12 features documented including post-2017 (asperitas, volutus, cavum, murus, cauda).
CF-M4.1	Köhler equation form	Kelvin and Raoult terms both present; critical-supersaturation concept explained.
CF-M4.2	WBF process explanation	The vapor-pressure differential $e_s^w > e_s^i$ stated as the driver.
CF-M5.1	Primitive equations enumerated	All 5 categories: 3 momentum, continuity, thermodynamic, state, moisture.
CF-M5.2	DA hierarchy	OI $\rightarrow$ 3D-Var $\rightarrow$ 4D-Var $\rightarrow$ EnKF $\rightarrow$ hybrid 4DEnVar all present.
CF-M5.3	AI weather model coverage	At least 7 of: FCN, Pangu, GraphCast, FCN2, FengWu, FuXi, AIFS, GenCast.
CF-M5.4	Lorenz system stated	Three ODEs with $\sigma = 10$, $\beta = 8/3$, $\rho = 28$ classical parameters.
CF-M6.1	Erdős #1196 statement	The conjecture stated precisely with primitive-set definition.
CF-M6.2	Lichtman bound	The 2023 bound $e^{\gamma}\pi/4 \approx 1.399$ stated.
CF-M6.3	GPT-5.4 Pro proof	April 13, 2026 date; the strengthened statement; Lichtman's 7-year quote.
CF-M6.4	Seven Millennium Problems	All seven listed with current status.

13.4 Integration Tests

ID	Verifies	Expected
IN-1	M1 → M2 cascade	Merle's compressible-Euler implosion explicitly connected to the Navier–Stokes Millennium Problem.
IN-2	M2 → M3 cascade	Atmospheric flow described in M2 is the source of cloud formation in M3.
IN-3	M3 → M4 cascade	Each genus in M3 has its forming microphysics in M4.
IN-4	M4 → M5 cascade	Microphysics processes in M4 are what NWP must parameterize in M5.
IN-5	M5 → M2 loop	Lorenz's chaos in M5 emerged from M2's atmospheric convection equations.
IN-6	M6 ↔ M2	The Navier–Stokes Millennium Problem in M6 is the open question of M2.
IN-7	Cross-module species/data consistency	Atmospheric scale, Reynolds number, Kolmogorov scale referenced consistently across M2, M4, M5.
IN-8	Document self-containment	A reader with no prior knowledge can navigate M1→M6 in order.

13.5 Edge Case Tests

ID	Verifies	Expected
EC-1	Anomalous taxonomic items	Volutus, asperitas, homogenitus flagged as recent additions.
EC-2	Disputed AI-proof acceptance	Survivorship bias and lack-of-peer-review explicitly noted for AI proofs.
EC-3	Mochizuki abc situation	abc-conjecture entry notes lack of consensus.
EC-4	Hubble tension	Listed as unresolved, with the 5σ-level disagreement noted.
EC-5	Temporal currency	Most cited primary sources from past 10 years where authoritative ones exist.
EC-6	Data gap acknowledgment	At least one "further research needed" or "open question" note per module.

13.6 Verification Matrix

Success Criterion	Test ID(s)	Status
1. Merle biographical record from authoritative sources	SC-SRC, CF-M1.1	Pending
2. Four contribution streams technically correct	SC-EQN, CF-M1.2	Pending
3. NSE / Euler canonical statement	SC-EQN, CF-M2.1	Pending
4. Ten WMO genera profiled	SC-WMO, CF-M3.1	Pending
5. Fifteen species and nine varieties tabled	SC-WMO, CF-M3.2, CF-M3.3	Pending
6. Cloud microphysics with equations	SC-EQN, CF-M4.1, CF-M4.2	Pending
7. NWP from Bjerknes through 2026 AI	CF-M5.1, CF-M5.2, CF-M5.3	Pending
8. Erdős #1196 precise with prior bounds and resolution	SC-DATE, SC-ATTR, CF-M6.1, CF-M6.2, CF-M6.3	Pending
9. Seven Millennium Problems with April 2026 status	SC-DATE, CF-M6.4	Pending
10. All claims attributed to primary sources	SC-SRC, SC-NUM	Pending
11. Cross-module synthesis explicit	IN-1 through IN-8	Pending
12. PDF compiles cleanly with three passes	W3.2–W3.4 build	Pending

14 Mission Crew Manifest (Fleet Profile)

Role	Agent	Responsibility
FLIGHT	Orchestrator	Mission direction; go/no-go gates; wave dispatch.
PLAN	Planner	Wave decomposition; parallel-track optimization.
SCOUT	Research Agent	Source gathering; gap-fill web research.
EXEC-A	Executor (Track A)	M1, M2, M6 document production.
EXEC-B	Executor (Track B)	M3, M4, M5 document production.
CRAFT-MATH	Domain Specialist (math)	PDE / number-theory technical accuracy.
CRAFT-FLUID	Domain Specialist (fluids)	Fluid dynamics + atmospheric physics.
CRAFT-METEO	Domain Specialist (meteor.)	WMO taxonomy + NWP.
CRAFT-AI	Domain Specialist (AI)	AI weather models + AI mathematics.
VERIFY	Verification	Source validation; equation-form checking.
INTEG	Integration Agent	Cross-module synthesis; through-line construction.
CAPCOM	HITL Interface	Human approval at wave boundaries.
BUDGET	Resource Manager	Token budget tracking; session planning.
SURGEON	Health Monitor	Research-quality degradation detection.
LOG	Chronicle Agent	Audit trail; commit messages.
TOPO	Topology Manager	Agent team structure; mid-mission restructuring.
ARCH	Architecture	Cross-cutting structural decisions.
SECURE	Security	Safety boundary enforcement.
ANALYST	Pattern Analyst	Cross-module pattern detection.
RETRO	Retrospective	Post-wave analysis; lessons learned.

15 Execution Summary

Metric	Value
Activation profile	Fleet (20 roles)
Modules	6 (M1 Merle, M2 Fluids, M3 Atlas, M4 Microphys., M5 NWP, M6 Open)
Waves	4 (Foundation, two parallel research tracks, Synthesis, Publish+Verify)
Parallel tracks	2 (math/open + atmospheric)
Components	11 deliverables (D1–D11)
Tests	39 (7 safety-critical, 18 core functional, 8 integration, 6 edge case)
Success criteria	12
Estimated total tokens	∼206 K
Estimated context windows	∼14
Estimated sessions	∼5
Model split	∼48% Opus / ∼49% Sonnet / ∼3% Haiku
LaTeX compilation	3 XeLaTeX passes
Output formats	PDF + .tex source + index.html

16 Closing Through-Line

"A soliton is a special solution to nonlinear equations, such as fluid equations, that doesn't send energy away to infinity — it keeps all its energy contained and keeps the same shape. When you look at physical quantities in nonlinear systems, they seem to oscillate and change chaotically. But if you look long enough, some emergent structure appears that doesn't depend that much on how things started. This emerging structure is the soliton. From the mathematical point of view, you don't initially see why it will appear, yet somehow it does."

— Frank Merle, on the soliton resolution conjecture, after receiving the 2026 Breakthrough Prize in Mathematics

This mission documents the precise threshold of human and machine understanding in April 2026, on a particularly meaningful week. Five days before the Breakthrough Prize honored Merle for taming nonlinear chaos through solitons, an artificial reasoning system spent eighty minutes and solved a number-theoretic problem a human researcher had spent seven years on. Both events sit on the same edge: the edge between what we name and what we cannot yet predict, between the equation we can write and the equation whose solutions still escape us, between the cloud genus we have cataloged and the cloud whose formation we cannot yet forecast at convective resolution. The Amiga Principle says: build the right architecture and the right tools

will surprise you. The *spaces between* say: meaning lives where the categories don't quite fit. This mission lives in that space, and points carefully at where it sits today.