

# Foundations of Mathematics: On the Translations Between Arithmetic and Set Theory

Thibaut Kouptchinsky

COLABS Tohoku University

UCLouvain

February 2023

# The main question: What are the appropriate axioms to prove the theorems of mathematics

*Foundations of mathematics is the study of the most basic concepts and logical structure of mathematics, with an eye to the unity of human knowledge*

---

*Stephen G. Simpson  
SSOA (second edition)*

# The main question: What are the appropriate axioms to prove the theorems of mathematics

*Foundations of mathematics is the study of the most basic concepts and logical structure of mathematics, with an eye to the unity of human knowledge*

---

*Stephen G. Simpson  
SSOA (second edition)*

Aristote, Euclide, Descartes, Cauchy, Weierstraß, Dedekind, Peano, Frege, Russell, Cantor, Hilbert, Brouwer, Weyl, von Neumann, Skolem, Tarski, Heyting, Gödel, ...

# The Set Theory of Zermelo and Fraënkël

## Example

- 1 The set of natural numbers:  $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$ .

# The Set Theory of Zermelo and Fraënkell

## Example

- 1 The set of natural numbers:  $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$ .
- 2 The graph of the function  $\sin: \mathbb{R} \rightarrow \mathbb{R}$ :  $\{(x, \sin(x)) : x \in \mathbb{R}\}$ .

# The Set Theory of Zermelo and Fraënkell

## Example

- 1 The set of natural numbers:  $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$ .
- 2 The graph of the function  $\sin: \mathbb{R} \rightarrow \mathbb{R}$ :  $\{(x, \sin(x)) : x \in \mathbb{R}\}$ .

A counter-example, the Russell's Paradox:

$$S = \{X : X \notin X\}$$

# The Set Theory of Zermelo and Fraënkell

## Example

- 1 The set of natural numbers:  $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$ .
- 2 The graph of the function  $\sin: \mathbb{R} \rightarrow \mathbb{R}$ :  $\{(x, \sin(x)) : x \in \mathbb{R}\}$ .

A counter-example, the Russell's Paradox:

$$S = \{X : X \notin X\}$$

# The Set Theory of Zermelo and Fraënkell

## Example

- 1 The set of natural numbers:  $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$ .
- 2 The graph of the function  $\sin: \mathbb{R} \rightarrow \mathbb{R}$ :  $\{(x, \sin(x)) : x \in \mathbb{R}\}$ .

A counter-example, the Russell's Paradox:

$$S = \{X : X \notin X\}$$

■ If  $S \in S$ , then  $S \notin S$ ;



# The Set Theory of Zermelo and Fraënkell

## Example

- 1 The set of natural numbers:  $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$ .
- 2 The graph of the function  $\sin: \mathbb{R} \rightarrow \mathbb{R}$ :  $\{(x, \sin(x)) : x \in \mathbb{R}\}$ .

A counter-example, the Russell's Paradox:

$$S = \{X : X \notin X\}$$

- If  $S \in S$ , then  $S \notin S$ ;
- If  $S \notin S$ , then  $S \in S$ .

# The axioms of ZF

- 1 There exists a set and this set is empty;  $\emptyset$ .

# The axioms of ZF

- 1 There exists a set and this set is empty;  $\emptyset$ .
- 2 PAIR: If  $a$  and  $b$  are sets, we can form  $\{a, b\}$ .

# The axioms of ZF

- ① There exists a set and this set is empty;  $\emptyset$ .
- ② PAIR: If  $a$  and  $b$  are sets, we can form  $\{a, b\}$ .
- ③ SEPARATION: If  $P$  is property and  $X$  is a set, we can form

$$\{u \in X \mid P(u)\}.$$

# The axioms of ZF

- ① There exists a set and this set is empty;  $\emptyset$ .
- ② PAIR: If  $a$  and  $b$  are sets, we can form  $\{a, b\}$ .
- ③ SEPARATION: If  $P$  is property and  $X$  is a set, we can form

$$\{u \in X \mid P(u)\}.$$

- ④ INFINITY: There exists an infinite set.

# The axioms of ZF


- ① There exists a set and this set is empty;  $\emptyset$ .
- ② PAIR: If  $a$  and  $b$  are sets, we can form  $\{a, b\}$ .
- ③ SEPARATION: If  $P$  is property and  $X$  is a set, we can form

$$\{u \in X \mid P(u)\}.$$

- ④ INFINITY: There exists an infinite set.
- ⑤ plus 5 others ...

# The Von Neumann construction of $\mathbb{N}$

$$\begin{aligned}0 &= \{ \} &= \emptyset, \\1 &= \{0\} &= \{\emptyset\} \\2 &= \{0, 1\} &= \{\emptyset, \{\emptyset\}\} \\3 &= \{0, 1, 2\} &= \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\} \\&\dots \\n + 1 &= n \cup \{n\} &= \{0, 1, 2, \dots, n\} \\&\dots\end{aligned}$$





# The Realm of Second-order Arithmetic

Friedman and Simpson: Encoding the ordinary mathematics with only


- 1 Natural numbers  $n \in \mathbb{N}$ ;
- 2 Sets of natural numbers  $S \subseteq \mathbb{N}$ .

# The Realm of Second-order Arithmetic


Friedman and Simpson: Encoding the ordinary mathematics with only

- 1 Natural numbers  $n \in \mathbb{N}$ ;
- 2 Sets of natural numbers  $S \subseteq \mathbb{N}$ .

Example:  $\mathbb{Z} \times \mathbb{Z}$  is countable



# Trees



# Set Theory in Second Order Arithmetic

We encode a set by a tree  $T$  where

# Set Theory in Second Order Arithmetic

We encode a set by a tree  $T$  where

- 1 If  $\sigma \in T$  is a leaf, then  $|\sigma|$  code the emptyset,

# Set Theory in Second Order Arithmetic

We encode a set by a tree  $T$  where

- 1 If  $\sigma \in T$  is a leaf, then  $|\sigma|$  code the emptyset,
- 2 If  $\tau$  is deeper than  $\sigma$  on a branch, then it code  $|\tau| \in |\sigma|$ .

# Set Theory in Second Order Arithmetic

We encode a set by a tree  $T$  where

- 1 If  $\sigma \in T$  is a leaf, then  $|\sigma|$  code the emptyset,
- 2 If  $\tau$  is deeper than  $\sigma$  on a branch, then it code  $|\tau| \in |\sigma|$ .

