

Determinacy on the edge of second order arithmetic

Thibaut Kouptchinsky

Logic Seminar Tohoku University

UCLouvain

Janvier 12, 2023

Program

- 1 A tool of descriptive set theory
- 2 The theorem of Wolfe as a warm up
- 3 Determinacy of Π_3^0 Differences

What is determinacy?

Consider a set A and a payoff set $X \subseteq A^\omega$.

I: a_0 a_2 a_{2n} \dots $(a_i)_{i < \omega} \stackrel{?}{\in} X$

II: a_1 a_3 a_{2n+1}

Player I wins if yes. Otherwise player II wins.

Axiom of determinacy (AD): “All these games are determined”.
(False in ZF + C.)

The Borel and projective hierarchy

Motivations and applications

Theorem (Mycielski-Swierczkowski; Mazur, Banach; Davis)

ZF + AD proves that every set of real numbers is Lebesgue measurable (M1), has the Baire property (M2), and has the perfect set property (M3).

Motivations and applications

Theorem (Mycielski-Swierczkowski; Mazur, Banach; Davis)

ZF + AD proves that every set of real numbers is Lebesgue measurable (M1), has the Baire property (M2), and has the perfect set property (M3).

- Study these properties for projective Σ^1_n sets in ω^ω (Blackwell, 1967).

Motivations and applications

Theorem (Mycielski-Swierczkowski; Mazur, Banach; Davis)

ZF + AD proves that every set of real numbers is Lebesgue measurable (M1), has the Baire property (M2), and has the perfect set property (M3).

- Study these properties for projective Σ^1_n sets in ω^ω (Blackwell, 1967).
- Are Σ^1_2 , Σ^1_3 , etc sets Lebesgue measurable?

Motivations and applications

Theorem (Mycielski-Swierczkowski; Mazur, Banach; Davis)

ZF + AD proves that every set of real numbers is Lebesgue measurable (M1), has the Baire property (M2), and has the perfect set property (M3).

- Study these properties for projective Σ^1_n sets in ω^ω (Blackwell, 1967).
- Are Σ^1_2 , Σ^1_3 , etc sets Lebesgue measurable?
- Applications in measure theory, descriptive set theory, harmonic analysis, ergodic theory, dynamical systems etc.

Borel Determinacy

- First best result (1964): $\text{Det}(\Sigma_3^0)$ by Davis.

Borel Determinacy

- First best result (1964): $\text{Det}(\Sigma_3^0)$ by Davis.
- The proof can be carried out in $\text{ZC}^- + \Sigma_1\text{Replacement}$ (Martin).

Borel Determinacy

- First best result (1964): $\text{Det}(\Sigma^0_3)$ by Davis.
- The proof can be carried out in $\text{ZC}^- + \Sigma_1\text{Replacement}$ (Martin).
- Friedman (1968): Borel determinacy requires existence of V_{ω_1} .

Borel Determinacy

- First best result (1964): $\text{Det}(\Sigma_3^0)$ by Davis.
- The proof can be carried out in $\text{ZC}^- + \Sigma_1\text{Replacement}$ (Martin).
- Friedman (1968): Borel determinacy requires existence of V_{ω_1} .

Theorem (Martin, ZFC)

All Borel games are determined.

The theorem of Wolfe

Theorem ($ZC^- + \Sigma_1\text{-REPLACEMENT}$)

All Σ_2^0 games are determined.

Definition (Player I's strategies)

Given $G(T, X)$, a strategy S_I is a subtree of T such that

Definition (Player I's strategies)

Given $G(T, X)$, a strategy S_I is a subtree of T such that

- 1 Every even-length node has one unique child,

Definition (Player I's strategies)

Given $G(T, X)$, a strategy S_I is a subtree of T such that

- 1 Every even-length node has one unique child,
- 2 Every child of an odd-length node that lies in T lies also in the strategy,

Definition (Player I's strategies)

Given $G(T, X)$, a strategy S_I is a subtree of T such that

- 1 Every even-length node has one unique child,
- 2 Every child of an odd-length node that lies in T lies also in the strategy,

A strategy S is said to be winning if $[S] \subseteq X$.

Definition (Player I's strategies)


Given $G(T, X)$, a strategy S_I is a subtree of T such that

- 1 Every even-length node has one unique child,
- 2 Every child of an odd-length node that lies in T lies also in the strategy,


A strategy S is said to be winning if $[S] \subseteq X$.

In a quasistrategy the player's response has not to be unique.

Example: A binary game



Strategy and quasistrategy



A technical lemma

Lemma

Let $B \subseteq A \subseteq [T]$ with B being closed. If player I has no winning strategy in the game $G(T, A)$, then there is a strategy τ for II such that every $x \in [\tau]$ has a finite initial segment p verifying

$[T_p] \cap B = \emptyset$ and I has still no winning strategy in $G(T_p, A)$

Proof of the theorem of Wolfe

$$A = \bigcup_{i < \omega} A_i, \quad A_i \text{ closed in } [T];$$

Proof of the theorem of Wolfe

$$A = \bigcup_{i < \omega} A_i, \quad A_i \text{ closed in } [T];$$

■ Suppose $G(T, A)$ is not a win for I;

Proof of the theorem of Wolfe

$$A = \bigcup_{i < \omega} A_i, \quad A_i \text{ closed in } [T];$$

- Suppose $G(T, A)$ is not a win for I;
- Apply the lemma for $B = A_0$ to get $\tau_0 \subset T$ and p_0 ;

Proof of the theorem of Wolfe

$$A = \bigcup_{i < \omega} A_i, \quad A_i \text{ closed in } [T];$$

- Suppose $G(T, A)$ is not a win for I;
- Apply the lemma for $B = A_0$ to get $\tau_0 \subset T$ and p_0 ;
- Apply the lemma for $B = A_{n+1}$ to get $\tau_{n+1} \subset T_{p_n}$ and p_{n+1} ;

Proof of the theorem of Wolfe

$$A = \bigcup_{i < \omega} A_i, \quad A_i \text{ closed in } [T];$$

- Suppose $G(T, A)$ is not a win for I;
- Apply the lemma for $B = A_0$ to get $\tau_0 \subset T$ and p_0 ;
- Apply the lemma for $B = A_{n+1}$ to get $\tau_{n+1} \subset T_{p_n}$ and p_{n+1} ;
- ...

Proof of the theorem of Wolfe

$$A = \bigcup_{i < \omega} A_i, \quad A_i \text{ closed in } [T];$$

- Suppose $G(T, A)$ is not a win for I;
- Apply the lemma for $B = A_0$ to get $\tau_0 \subset T$ and p_0 ;
- Apply the lemma for $B = A_{n+1}$ to get $\tau_{n+1} \subset T_{p_n}$ and p_{n+1} ;
- ...
- Our τ avoid all the A_i and hence is winning for II.

Theorem (Montalbán and Shore)

$$\forall n \in \omega, \Pi_{n+2}^1\text{-CA}_0 \vdash \text{Det}(n\text{-}\Pi_3^0)$$

Theorem (Montalbán and Shore)

$$\forall n \in \omega, \Pi_{n+2}^1\text{-CA}_0 \vdash \text{Det}(n\text{-}\Pi_3^0)$$

However, $\Pi_{n+2}^1\text{-CA}_0$ is not the right set of axioms for $\text{Det}(n\text{-}\Pi_3^0)$.

Theorem (Montalbán and Shore)

$\forall n \in \omega, \Pi_{n+2}^1\text{-CA}_0 \vdash \text{Det}(n\text{-}\Pi_3^0)$

However, $\Pi_{n+2}^1\text{-CA}_0$ is not the right set of axioms for $\text{Det}(n\text{-}\Pi_3^0)$.

Theorem (MedSalem and Tanaka)

Borel determinacy does not imply $\Delta_2^1\text{-CA}_0$.

Theorem (Montalbán and Shore)

$$\forall n \in \omega, \Pi_{n+2}^1\text{-CA}_0 \vdash \text{Det}(n\text{-}\Pi_3^0)$$

However, $\Pi_{n+2}^1\text{-CA}_0$ is not the right set of axioms for $\text{Det}(n\text{-}\Pi_3^0)$.

Theorem (MedSalem and Tanaka)

Borel determinacy does not imply $\Delta_2^1\text{-CA}_0$.

$\text{Det}(n\text{-}\Pi_3^0)$ is the Π_3^1 sentence

$$\forall X \exists Y \forall Z (X \in n\text{-}\Pi_3^0) \rightarrow \begin{cases} Y \in S_I \wedge Z \in S_{II} \rightarrow Y \oplus Z \in X; \\ Y \in S_{II} \wedge Z \in S_I \rightarrow Z \oplus Y \in X. \end{cases}$$

Consider $\emptyset = A_m \subseteq \cdots \subseteq A_1 \subseteq A_0$, Π_0^3 sets.

$$A_i = \bigcap_{k < \omega} A_{i,k} \quad \text{and} \quad A_{i,k} = \bigcup_{j < \omega} A_{i,k,j}.$$

Definition

Assume $0 \leq k < \omega$.

Definition

Assume $0 \leq k < \omega$.

1 Σ^1_k -DC₀:

$$\forall n \forall X \exists Y \eta(n, X, Y) \rightarrow \exists Z \forall n \eta(n, (Z)^n, (Z)_n).$$

Definition

Assume $0 \leq k < \omega$.

1 $\Sigma_k^1\text{-DC}_0$:

$$\forall n \forall X \exists Y \eta(n, X, Y) \rightarrow \exists Z \forall n \eta(n, (Z)^n, (Z)_n).$$

2 Strong $\Sigma_k^1\text{-DC}_0$:

$$\exists Z \forall n \forall Y (\eta(n, (Z)^n, Y) \rightarrow \eta(n, (Z)^n, (Z)_n)).$$

Definition

Assume $0 \leq k < \omega$.

1 $\Sigma_k^1\text{-DC}_0$:

$$\forall n \forall X \exists Y \eta(n, X, Y) \rightarrow \exists Z \forall n \eta(n, (Z)^n, (Z)_n).$$

2 Strong $\Sigma_k^1\text{-DC}_0$:

$$\exists Z \forall n \forall Y (\eta(n, (Z)^n, Y) \rightarrow \eta(n, (Z)^n, (Z)_n)).$$

Strong $\Sigma_{m+2}^1\text{-DC}_0$ is Π_4^1 conservative over $\Pi_{m+2}^1\text{-CA}_0$.

Definition

We define $\Sigma^1_{|s|+2}$ relations $P^s(S)$ by induction on $|s| \leq m$:

Definition

We define $\Sigma^1_{|s|+2}$ relations $P^s(S)$ by induction on $|s| \leq m$:

■ When $|s| = 0$, $P^\emptyset(S)$ iff

$G(A, S)$ is a win for I if $l := m - |s|$ is even .

Definition

We define $\Sigma_{|s|+2}^1$ relations $P^s(S)$ by induction on $|s| \leq m$:

- When $|s| = 0$, $P^\emptyset(S)$ iff

$G(A, S)$ is a win for I if $l := m - |s|$ is even .

- For $|s| = n + 1$ and l is even, $P^s(S)$ iff there is a quasistrategy U for I in S such that

$$[U] \subseteq A \cup A_{l,s(n)} \quad \text{and} \quad P^{s[n]}(U) \text{ fails.}$$

Definition

We define $\Sigma^1_{|s|+2}$ relations $P^s(S)$ by induction on $|s| \leq m$:

- When $|s| = 0$, $P^\emptyset(S)$ iff

$G(A, S)$ is a win for I if $l := m - |s|$ is even .

- For $|s| = n + 1$ and l is even, $P^s(S)$ iff there is a quasistrategy U for I in S such that

$$[U] \subseteq A \cup A_{l,s(n)} \quad \text{and} \quad P^{s[n]}(U) \text{ fails.}$$

A quasistrategy U witnesses $P^s(S)$ if U is as required in the appropriate clause, the latter being a $\Pi^1_{|s|+1}$ sentence.

Definition

A quasistrategy U for I locally witnesses $P^s(S)$ if $|s| = n + 1$ and l is even if: $\exists D \subseteq S \forall d \in D$, there is a quasistrategy R^d for II in S_d such that:

Definition

A quasistrategy U for I locally witnesses $P^s(S)$ if $|s| = n + 1$ and l is even if: $\exists D \subseteq S \forall d \in D$, there is a quasistrategy R^d for II in S_d such that:

- 1 $\forall d \in D \cap U, U_d \cap R^d$ witnesses $P^s(R^d)$.

Definition

A quasistrategy U for I locally witnesses $P^s(S)$ if $|s| = n + 1$ and l is even if: $\exists D \subseteq S \forall d \in D$, there is a quasistrategy R^d for II in S_d such that:

- 1 $\forall d \in D \cap U, U_d \cap R^d$ witnesses $P^s(R^d)$.
- 2 $[U] \setminus \bigcup_{d \in D} [R^d] \subseteq A$.

Definition

A quasistrategy U for I locally witnesses $P^s(S)$ if $|s| = n + 1$ and l is even if: $\exists D \subseteq S \forall d \in D$, there is a quasistrategy R^d for II in S_d such that:

- 1 $\forall d \in D \cap U, U_d \cap R^d$ witnesses $P^s(R^d)$.
- 2 $[U] \setminus \bigcup_{d \in D} [R^d] \subseteq A$.
- 3 $\forall p \in S \exists^{\leq 1} d \in D, d \subseteq p \wedge p \in R^d$.

Definition

A quasistrategy U for I locally witnesses $P^s(S)$ if $|s| = n + 1$ and l is even if: $\exists D \subseteq S \forall d \in D$, there is a quasistrategy R^d for II in S_d such that:

- 1 $\forall d \in D \cap U, U_d \cap R^d$ witnesses $P^s(R^d)$.
- 2 $[U] \setminus \bigcup_{d \in D} [R^d] \subseteq A$.
- 3 $\forall p \in S \exists \leq^1 d \in D, d \subseteq p \wedge p \in R^d$.

We observe that “ U locally witnesses $P^s(S)$ ” is a $\Sigma^1_{|s|+2}$ sentence.

Lemma (1)

If U locally witnesses $P^s(S)$, then U witnesses $P^s(S)$.

Proof of lemma 1

■ First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.

Proof of lemma 1

- First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.
- $|s| = n + 1 \leq m, n = 0$:

Proof of lemma 1

- First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.
- $|s| = n + 1 \leq m, n = 0$:
 - Suppose $P^\emptyset(U)$ holds, there is τ winning for II.

Proof of lemma 1

- First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.
- $|s| = n + 1 \leq m, n = 0$:
 - Suppose $P^\emptyset(U)$ holds, there is τ winning for II.
 - 2 and 3 implies $\exists d \in D \cap \tau \forall x \supset d \ x \in [\tau] \rightarrow x \in [R^d]$.

Proof of lemma 1

- First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.
- $|s| = n + 1 \leq m, n = 0$:
 - Suppose $P^\emptyset(U)$ holds, there is τ winning for II.
 - 2 and 3 implies $\exists d \in D \cap \tau \forall x \supset d \ x \in [\tau] \rightarrow x \in [R^d]$.
 - So τ_d is winning for II in $G(U_d \cap R^d, A)$, contradiction.

Proof of lemma 1

- First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.
- $|s| = n + 1 \leq m, n = 0$:
 - Suppose $P^\emptyset(U)$ holds, there is τ winning for II.
 - 2 and 3 implies $\exists d \in D \cap \tau \forall x \supset d \ x \in [\tau] \rightarrow x \in [R^d]$.
 - So τ_d is winning for II in $G(U_d \cap R^d, A)$, contradiction.
- By induction hypothesis we have to build for $n > 1, \hat{U}, \hat{D}$ and $\{\hat{R}^d : d \in \hat{D}\}$ locally witnessing $P^{s[n-1]}(\hat{S})$.

Proof of lemma 1

- First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.
- $|s| = n + 1 \leq m$, $n = 0$:
 - Suppose $P^\emptyset(U)$ holds, there is τ winning for II.
 - 2 and 3 implies $\exists d \in D \cap \tau \forall x \supset d \ x \in [\tau] \rightarrow x \in [R^d]$.
 - So τ_d is winning for II in $G(U_d \cap R^d, A)$, contradiction.
- By induction hypothesis we have to build for $n > 1$, \hat{U} , \hat{D} and $\{\hat{R}^d : d \in \hat{D}\}$ locally witnessing $P^{s[n-1]}(\hat{S})$.
- We want to define \hat{U} such that \hat{U} falls out of the R^d and hence ends up in A (we suppose $m - n$ odd).

Proof of lemma 1

- First clause “ $[U] \subseteq A \cup A_{l,s(n)}$ ” follows from 1 and 2.
- $|s| = n + 1 \leq m, n = 0$:
 - Suppose $P^\emptyset(U)$ holds, there is τ winning for II.
 - 2 and 3 implies $\exists d \in D \cap \tau \forall x \supset d \ x \in [\tau] \rightarrow x \in [R^d]$.
 - So τ_d is winning for II in $G(U_d \cap R^d, A)$, contradiction.
- By induction hypothesis we have to build for $n > 1$, \hat{U} , \hat{D} and $\{\hat{R}^d : d \in \hat{D}\}$ locally witnessing $P^{s[n-1]}(\hat{S})$.
- We want to define \hat{U} such that \hat{U} falls out of the R^d and hence ends up in A (we suppose $m - n$ odd).
- We use $\Sigma_{|s|}^1\text{-AC}_0$ to pick up witnesses for $P^{s[n-1]}(\hat{R}^d)$ which will be our $\hat{U}_d \cap \hat{R}^d$.

Definition

We say that $P^s(S)$ fails everywhere if $P^s(S_p)$ fails for every $p \in S$.
This is a $\Pi^1_{|S|+2}$ sentence.

Definition

We say that $P^s(S)$ fails everywhere if $P^s(S_p)$ fails for every $p \in S$.
This is a $\Pi_{|S|+2}^1$ sentence.

Lemma (2)

If $P^s(S)$ fails, then there is a quasistrategy W for I if l is odd in S such that $P^s(W)$ fails everywhere.

Proof of lemma 2

- $|s| = 0$, then II does not have winning strategy and we take I 's non-losing quasistrategy as W (Using $\Pi_2^1\text{-CA}_0$).

Proof of lemma 2

- $|s| = 0$, then II does not have winning strategy and we take I's non-losing quasistrategy as W (Using Π_2^1 -CA₀).
- $|s| = n + 1$ (and l even), we use $\Pi_{|s|+2}^1$ -CA₀ to set

$$d \in D \leftrightarrow d \in S \wedge P^s(S_d) \wedge \neg P^s(S_{d[|d|-1]}). \quad (d \text{ is minimal})$$

Proof of lemma 2

- $|s| = 0$, then II does not have winning strategy and we take I 's non-losing quasistrategy as W (Using $\Pi_2^1\text{-CA}_0$).
- $|s| = n + 1$ (and l even), we use $\Pi_{|s|+2}^1\text{-CA}_0$ to set

$$d \in D \leftrightarrow d \in S \wedge P^s(S_d) \wedge \neg P^s(S_{d[|d|-1]}). \quad (d \text{ is minimal})$$

- We will use $\Sigma_{|s|}^1\text{-AC}_0$ to choose a witness U^d .

Proof of lemma 2

- $|s| = 0$, then II does not have winning strategy and we take I 's non-losing quasistrategy as W (Using $\Pi_2^1\text{-CA}_0$).
- $|s| = n + 1$ (and l even), we use $\Pi_{|s|+2}^1\text{-CA}_0$ to set

$$d \in D \leftrightarrow d \in S \wedge P^s(S_d) \wedge \neg P^s(S_{d[|d|-1]}). \quad (d \text{ is minimal})$$

- We will use $\Sigma_{|s|}^1\text{-AC}_0$ to choose a witness U^d .
- We now consider the game $G(S, B)$ where

$$B = \{x \in [S] \mid \exists d \in D \ d \subseteq x\}.$$

Proof of lemma 2

- $|s| = 0$, then II does not have winning strategy and we take I's non-losing quasistrategy as W (Using Π^1_2 -CA₀).
- $|s| = n + 1$ (and l even), we use $\Pi^1_{|s|+2}$ -CA₀ to set

$$d \in D \leftrightarrow d \in S \wedge P^s(S_d) \wedge \neg P^s(S_{d \upharpoonright |d|-1}). \quad (d \text{ is minimal})$$

- We will use $\Sigma^1_{|s|}$ -AC₀ to choose a witness U^d .
- We now consider the game $G(S, B)$ where

$$B = \{x \in [S] \mid \exists d \in D \ d \subseteq x\}.$$

- Using preceding lemma, $G(S, B)$ is not a win for I.

Proof of lemma 2

- $|s| = 0$, then II does not have winning strategy and we take I's non-losing quasistrategy as W (Using Π_2^1 -CA₀).
- $|s| = n + 1$ (and l even), we use $\Pi_{|s|+2}^1$ -CA₀ to set

$$d \in D \leftrightarrow d \in S \wedge P^s(S_d) \wedge \neg P^s(S_{d \upharpoonright |d|-1}). \quad (d \text{ is minimal})$$

- We will use $\Sigma_{|s|}^1$ -AC₀ to choose a witness U^d .
- We now consider the game $G(S, B)$ where

$$B = \{x \in [S] \mid \exists d \in D \ d \subseteq x\}.$$

- Using preceding lemma, $G(S, B)$ is not a win for I.
- Again we define W as non-losing II's quasistrategy (and use preceding lemma to show it is as required).

Definition

For $|s| = n + 1$, W strongly witnesses $P^s(S)$ if, for all $p \in W$, W_p witnesses $P^s(S_p)$, that is, W witnesses $P^s(S)$ and $P^{s[n]}(W)$ fails everywhere. This is a $\Pi_{|s|+1}^1$ sentence.

Lemma (3)

If $P^s(S)$, then there is a W that strongly witnesses it.

Proof of lemma 3

Take U such that

$$[U] \subseteq A \cup A_{l,s(n)} \quad \text{and} \quad P^{s[n]}(U) \text{ fails.}$$

Proof of lemma 3

Take U such that

$$[U] \subseteq A \cup A_{l,s(n)} \quad \text{and} \quad P^{s[n]}(U) \text{ fails.}$$

Apply preceding lemma to get a W such $P^{s[n]}(W)$ fails everywhere.

Lemma (4)

If $|s| = n + 1$, then at least one of $P^s(S)$ and $P^{s[n]}(S)$ holds.

Proof of lemma 4

- Reverse induction on $n < m$, $m - n$ odd, suppose $P^s(S)$ fails.

Proof of lemma 4

- Reverse induction on $n < m$, $m - n$ odd, suppose $P^s(S)$ fails.
- Using strong Σ^1_{m+2} -DC₀, we define by induction a quasistrategy U for II in S along with $D \subseteq S$ and R^d for $d \in D$ showing that

U (locally) witnesses $P^{s[n]}(S)$ if $(n > 0) \vee n = 0$.

Proof of lemma 4

- Reverse induction on $n < m$, $m - n$ odd, suppose $P^s(S)$ fails.
- Using strong Σ_{m+2}^1 -DC₀, we define by induction a quasistrategy U for II in S along with $D \subseteq S$ and R^d for $d \in D$ showing that

U (locally) witnesses $P^{s[n]}(S)$ if $(n > 0) \vee n = 0$.

- The method to ensure $x \in [U] \setminus_{d \in D} R^d$ implies $x \in \bar{A}$ is as follow:

Proof of lemma 4

- Reverse induction on $n < m$, $m - n$ odd, suppose $P^s(S)$ fails.
- Using strong Σ^1_{m+2} -DC₀, we define by induction a quasistrategy U for II in S along with $D \subseteq S$ and R^d for $d \in D$ showing that

U (locally) witnesses $P^{s[n]}(S)$ if $(n > 0) \vee n = 0$.

- The method to ensure $x \in [U] \setminus_{d \in D} R^d$ implies $x \in \bar{A}$ is as follow:
 - 1 We define U such that $x \in \bar{A} \cup A_{m-n-2,j}$, $\forall j < \omega$ such that $x \notin \bar{A} \cup A_{m-n-2}, \cdot$

Proof of lemma 4

- Reverse induction on $n < m$, $m - n$ odd, suppose $P^s(S)$ fails.
- Using strong $\Sigma_{m+2}^1\text{-DC}_0$, we define by induction a quasistrategy U for II in S along with $D \subseteq S$ and R^d for $d \in D$ showing that

U (locally) witnesses $P^{s[n]}(S)$ if $(n > 0) \ n = 0$.

- The method to ensure $x \in [U] \setminus_{d \in D} R^d$ implies $x \in \bar{A}$ is as follow:
 - 1 We define U such that $x \in \bar{A} \cup A_{m-n-2,j}$, $\forall j < \omega$ such that $x \notin \bar{A} \cup A_{m-n-2}$.
 - 2 We make sure $x \notin A_{m-n-1,s(n),j}$, $\forall j < \omega$ such that $x \notin A_{m-n-1}$.

Proof of lemma 4

- Reverse induction on $n < m$, $m - n$ odd, suppose $P^s(S)$ fails.
- Using strong $\Sigma_{m+2}^1\text{-DC}_0$, we define by induction a quasistrategy U for II in S along with $D \subseteq S$ and R^d for $d \in D$ showing that

U (locally) witnesses $P^{s[n]}(S)$ if $(n > 0) \vee n = 0$.

- The method to ensure $x \in [U] \setminus_{d \in D} R^d$ implies $x \in \bar{A}$ is as follow:
 - 1 We define U such that $x \in \bar{A} \cup A_{m-n-2,j}, \forall j < \omega$ such that $x \notin \bar{A} \cup A_{m-n-2},$
 - 2 We make sure $x \notin A_{m-n-1,s(n),j}, \forall j < \omega$ such that $x \notin A_{m-n-1}.$
 - 3 We then use $\bar{A} \cup A_{m-n-2} \setminus A_{m-n-1} \subseteq \bar{A}.$

Proof of the theorem

■ Suppose m is odd and $G(A, T)$ is not a win for II; $P^\emptyset(T)$ fails.

Proof of the theorem

- Suppose m is odd and $G(A, T)$ is not a win for II; $P^\emptyset(T)$ fails.
- Take W^\emptyset such that $P^\emptyset(W^\emptyset)$ fails everywhere.

Proof of the theorem

- Suppose m is odd and $G(A, T)$ is not a win for II; $P^\emptyset(T)$ fails.
- Take W^\emptyset such that $P^\emptyset(W^\emptyset)$ fails everywhere.
- We define a quasistrategy U for I in W^\emptyset by induction on $|p|$ for $p \in U$. In the same time, we use strong Σ_3^1 -DC₀ to define for $|p| = j + 1$ a quasistrategy W^p for I such that

W^p strongly witnesses $P^{\langle j \rangle}(W_p^{p[j]})$.

Proof of the theorem

- Suppose then $p \in U$, $|p| = j + 1$ and W^p has been defined.
The child q of p in U are those of p in W^p .

Proof of the theorem

- Suppose then $p \in U$, $|p| = j + 1$ and W^p has been defined.
The child q of p in U are those of p in W^p .
- $P^\emptyset(W^p)$ fails everywhere and so,

$P^\emptyset(W^p_q)$ fails for each child q of p in U .

Proof of the theorem

- Suppose then $p \in U$, $|p| = j + 1$ and W^p has been defined.
The child q of p in U are those of p in W^p .
- $P^\emptyset(W^p)$ fails everywhere and so,

$P^\emptyset(W^p_q)$ fails for each child q of p in U .

- By preceding lemma $P^{(j)}(W^p_q)$ and we choose a W^q that strongly witnesses it.

Consider any play $x \in [U]$.

Consider any play $x \in [U]$.

■ For all j ,

$\forall j \ x \in [W^{x[j+1]}]$, which witnesses $P^{\langle j \rangle}$.

Consider any play $x \in [U]$.

■ For all j ,

$\forall j \ x \in [W^{x[j+1]}]$, which witnesses $P^{\langle j \rangle}$.

■ Then $\forall j \ x \in A \cup A_{m-1,j}$,

Consider any play $x \in [U]$.

■ For all j ,

$\forall j \ x \in [W^{x[j+1]}]$, which witnesses $P^{\langle j \rangle}$.

■ Then $\forall j \ x \in A \cup A_{m-1,j}$,

■ But $\bigcap_{j < \omega} A_{m-1,j} = A_{m-1} \subseteq A$.

Thank you for your attention!