

Foundations of Mathematics: On the Translations Between Arithmetic and Set Theory

Thibaut Kouptchinsky

COLABS Tohoku University

UCLouvain

February 2023

;

The main question: What are the appropriate axioms to prove the theorems of mathematics

Foundations of mathematics is the study of the most basic concepts and logical structure of mathematics, with an eye to the unity of human knowledge

*Stephen G. Simpson
SSOA (second edition)*

The main question: What are the appropriate axioms to prove the theorems of mathematics

Foundations of mathematics is the study of the most basic concepts and logical structure of mathematics, with an eye to the unity of human knowledge

*Stephen G. Simpson
SSOA (second edition)*

Aristote, Euclide, Descartes, Cauchy, Weierstraß, Dedekind, Peano, Frege, Russell, Cantor, Hilbert, Brouwer, Weyl, von Neumann, Skolem, Tarski, Heyting, Gödel, ...

The Set Theory of Zermelo and Fraënkël

Example

- 1 The set of natural numbers: $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$.

The Set Theory of Zermelo and Fraënkell

Example

- 1 The set of natural numbers: $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$.
- 2 The graph of the function $\sin: \mathbb{R} \rightarrow \mathbb{R}$: $\{(x, \sin(x)) : x \in \mathbb{R}\}$.

The Set Theory of Zermelo and Fraënkël

Example

- 1 The set of natural numbers: $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$.
- 2 The graph of the function $\sin: \mathbb{R} \rightarrow \mathbb{R}$: $\{(x, \sin(x)) : x \in \mathbb{R}\}$.
- 3 The set of the subsets of the complex number $\mathcal{P}(\mathbb{C}) = \{A : A \subseteq \mathbb{C}\}$.

The Set Theory of Zermelo and Fraënkell

Example

- 1 The set of natural numbers: $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$.
- 2 The graph of the function $\sin: \mathbb{R} \rightarrow \mathbb{R}$: $\{(x, \sin(x)) : x \in \mathbb{R}\}$.
- 3 The set of the subsets of the complex number $\mathcal{P}(\mathbb{C}) = \{A : A \subseteq \mathbb{C}\}$.

A counter-example, the Russel's Paradox:

$$S = \{X : X \notin X\}$$

The Set Theory of Zermelo and Fraënkell

Example

- ① The set of natural numbers: $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$.
- ② The graph of the function $\sin: \mathbb{R} \rightarrow \mathbb{R}$: $\{(x, \sin(x)) : x \in \mathbb{R}\}$.
- ③ The set of the subsets of the complex number $\mathcal{P}(\mathbb{C}) = \{A : A \subseteq \mathbb{C}\}$.

A counter-example, the Russel's Paradox:

$$S = \{X : X \notin X\}$$

■ If $S \in S$, then $S \notin S$;

The Set Theory of Zermelo and Fraënkël

Example

- 1 The set of natural numbers: $\mathbb{N} = \{0, 1, 2, 3, \dots, 2^{82,589,933} - 1, \dots\}$.
- 2 The graph of the function $\sin: \mathbb{R} \rightarrow \mathbb{R}$: $\{(x, \sin(x)) : x \in \mathbb{R}\}$.
- 3 The set of the subsets of the complex number $\mathcal{P}(\mathbb{C}) = \{A : A \subseteq \mathbb{C}\}$.

A counter-example, the Russel's Paradox:

$$S = \{X : X \notin X\}$$

- If $S \in S$, then $S \notin S$;
- If $S \notin S$, then $S \in S$.

The axioms of ZF

- 1 There exists a set and this set is empty; \emptyset .

The axioms of ZF

- 1 There exists a set and this set is empty; \emptyset .
- 2 EXTENSIONALITY Two sets are equal if they have the same elements.

The axioms of ZF

- ① There exists a set and this set is empty; \emptyset .
- ② EXTENSIONALITY Two sets are equal if they have the same elements.
- ③ PAIR If a and b are sets, we can form $\{a, b\}$.

The axioms of ZF

- ① There exists a set and this set is empty; \emptyset .
- ② EXTENSIONALITY Two sets are equal if they have the same elements.
- ③ PAIR If a and b are sets, we can form $\{a, b\}$.
- ④ SEPARATION If P is property and X is a set, we can form

$$\{u \in X \mid P(u)\}.$$

The axioms of ZF

- ① There exists a set and this set is empty; \emptyset .
- ② EXTENSIONALITY Two sets are equal if they have the same elements.
- ③ PAIR If a and b are sets, we can form $\{a, b\}$.
- ④ SEPARATION If P is property and X is a set, we can form

$$\{u \in X \mid P(u)\}.$$

- ⑤ INFINITY There exists an infinite set.

The axioms of ZF

- ① There exists a set and this set is empty; \emptyset .
- ② EXTENSIONALITY Two sets are equal if they have the same elements.
- ③ PAIR If a and b are sets, we can form $\{a, b\}$.
- ④ SEPARATION If P is property and X is a set, we can form

$$\{u \in X \mid P(u)\}.$$

- ⑤ INFINITY There exists an infinite set.
- ⑥ POWER SET If X is a set, we can form $\mathcal{P}(X)$

The axioms of ZF


- ① There exists a set and this set is empty; \emptyset .
- ② EXTENSIONALITY Two sets are equal if they have the same elements.
- ③ PAIR If a and b are sets, we can form $\{a, b\}$.
- ④ SEPARATION If P is property and X is a set, we can form

$$\{u \in X \mid P(u)\}.$$

- ⑤ INFINITY There exists an infinite set.
- ⑥ POWER SET If X is a set, we can form $\mathcal{P}(X)$
- ⑦ plus 3 others ...

The Von Neumann construction of \mathbb{N}

$$\begin{aligned}0 &= \{ \} &= \emptyset, \\1 &= \{0\} &= \{\emptyset\} \\2 &= \{0, 1\} &= \{\emptyset, \{\emptyset\}\} \\3 &= \{0, 1, 2\} &= \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\} \\&\dots \\n + 1 &= n \cup \{n\} &= \{0, 1, 2, \dots, n\} \\&\dots\end{aligned}$$



The Realm of Second-order Arithmetic

Friedman and Simpson: Encoding the ordinary mathematics with only


- 1 Natural numbers $n \in \mathbb{N}$;
- 2 Sets of natural numbers $S \subseteq \mathbb{N}$.

The Realm of Second-order Arithmetic

Friedman and Simpson: Encoding the ordinary mathematics with only

- 1 Natural numbers $n \in \mathbb{N}$;
- 2 Sets of natural numbers $S \subset \mathbb{N}$.

Example: $\mathbb{Z} \times \mathbb{Z}$ is countable



Trees


Definition

A tree is a subset of finite sequences of natural numbers ($T \subseteq \mathbb{N}^{<\infty}$) such that if $\sigma = \langle m_0, m_1, \dots, m_i, \dots, m_k \rangle$ is in T , then every initial segment of σ (for example $\tau = \langle m_0, m_1, \dots, m_i \rangle$) is also in T .


Trees

Definition

A tree is a subset of finite sequences of natural numbers ($T \subseteq \mathbb{N}^{<\infty}$) such that if $\sigma = \langle m_0, m_1, \dots, m_i, \dots, m_k \rangle$ is in T , then every initial segment of σ (for example $\tau = \langle m_0, m_1, \dots, m_i \rangle$) is also in T .



Another example of tree



Set Theory in Second Order Arithmetic

We encode a set by a tree T where

Set Theory in Second Order Arithmetic

We encode a set by a tree T where

- 1 If $\sigma \in T$ is an end node (a leaf), then $|\sigma|$ code the emptyset,

Set Theory in Second Order Arithmetic

We encode a set by a tree T where

- 1 If $\sigma \in T$ is an end node (a leaf), then $|\sigma|$ code the emptyset,
- 2 If τ is a direct child node of σ , then it code $|\tau| \in |\sigma|$.

Set Theory in Second Order Arithmetic

We encode a set by a tree T where

- ① If $\sigma \in T$ is an end node (a leaf), then $|\sigma|$ code the emptyset,
- ② If τ is a direct child node of σ , then it code $|\tau| \in |\sigma|$.

