

Efficient Task Offloading in MEC via UAV-UGV Collaboration

Tianchi Su and Qian Ma

School of Intelligent Systems Engineering, Sun Yat-sen University, Shenzhen, China
sutch@mail2.sysu.edu.cn, maqian25@mail.sysu.edu.cn

1 Appendix A

First, we define the quadratic Lyapunov function:

$$L(t) = \frac{1}{2} \left(\sum_{j \in \mathcal{A}} Q_j^t{}^2 + \sum_{k \in \mathcal{G}} Q_k^t{}^2 \right), \quad (1)$$

and Lyapunov drift,

$$\Delta(\mathbf{Q}(t)) = \mathbb{E}\{L(t+1) - L(t) | \mathbf{Q}(t)\},$$

where $\mathbf{Q}(t) = \{Q_j^t, Q_k^t\}_{j \in \mathcal{A}, k \in \mathcal{G}}$. According to [1], for any $\lambda > 0$, there exist a strategies π^t satisfied,

$$\begin{aligned} \mathbb{E}\{T_{sum}^t(\pi^t) | \mathbf{Q}(t)\} &\leq T_{sum}^{opt} + \lambda, \\ \mathbb{E}\{E_j^t(\pi^t) - e_j^t - \bar{E}_A\} &\leq \lambda, \forall j \in \mathcal{A}, \\ \mathbb{E}\{E_k^t(\pi^t) - \bar{E}_G\} &\leq \lambda, \forall k \in \mathcal{G}. \end{aligned}$$

Based on (1), we have

$$\begin{aligned} \Delta(\mathbf{Q}(t)) + V \sum_{t=1}^T T_{sum}^t &\leq (B_A + B_G)T + V \sum_{t=1}^T T_{sum}^t \\ &+ \leq \mathbb{E}\left\{\sum_{j \in \mathcal{A}} Q_j^t (E_j^t - e_j^t - \bar{E}_A) | \mathbf{Q}(t)\right\} + \mathbb{E}\left\{\sum_{k \in \mathcal{G}} Q_k^t (E_k^t - \bar{E}_G) | \mathbf{Q}(t)\right\}, \end{aligned}$$

taking $\lambda \rightarrow 0$, we have for any $\tilde{t} > 0$,

$$\frac{1}{\tilde{t}} \sum_{t=1}^{\tilde{t}} \mathbb{E}\{T_{sum}^t | \mathbf{Q}(t)\} \leq T_{sum}^{opt} + \frac{B_A + B_G}{V}.$$

2 Appendix B

We let $f^*(\mathbb{M}_G, \Gamma)$ denote the objective function value achieved under optimal strategies, and $\hat{f}(\mathbb{M}_G, \Gamma)$ be the approximated objective function value obtained

via the SAA method. Consider a larger sample set \mathcal{N}' of size $|\mathcal{N}'| \gg |\mathcal{N}|$, where the mean SAA estimate is given by:

$$\hat{f}_{\mathcal{N}'}(\mathbb{M}_G, \Gamma) = \frac{1}{|\mathcal{N}'|} \sum_{n \in \mathcal{N}'} f^n(\mathbb{M}_G, \Gamma), \quad (2)$$

with variance:

$$\sigma_{\mathcal{N}'}^2 = \frac{1}{|\mathcal{N}'|(|\mathcal{N}'| - 1)} \sum_{n \in \mathcal{N}'} \left(f^n(\mathbb{M}_G, \Gamma) - \hat{f}_{\mathcal{N}'}(\mathbb{M}_G, \Gamma) \right)^2. \quad (3)$$

Additionally, consider a collection $\mathcal{C} = \{\mathcal{N}_1, \mathcal{N}_2, \dots, \mathcal{N}_C\}$ of i.i.d. SAA replications, each with sample size $|\mathcal{N}|$. The mean SAA estimate across these replications is:

$$\bar{f}_{|\mathcal{N}|, \mathcal{C}}(\mathbb{M}_G, \Gamma) = \frac{1}{|\mathcal{C}|} \sum_{c \in \mathcal{C}} \hat{f}_{|\mathcal{N}|}^{(c)}(\mathbb{M}_G, \Gamma), \quad (4)$$

with the corresponding variance:

$$\sigma_{|\mathcal{N}|, \mathcal{C}}^2 = \frac{1}{|\mathcal{C}|(|\mathcal{C}| - 1)} \sum_{c \in \mathcal{C}} \left(\hat{f}_{|\mathcal{N}|}^{(c)}(\mathbb{M}_G, \Gamma) - \bar{f}_{|\mathcal{N}|, \mathcal{C}}(\mathbb{M}_G, \Gamma) \right)^2. \quad (5)$$

Since $\mathbb{E}[\hat{f}_{\mathcal{N}}] = \mathbb{E}[\bar{f}_{|\mathcal{N}|, \mathcal{C}}]$ and $\mathbb{E}[\hat{f}_{\mathcal{N}}] \leq f^*(\mathbb{M}_G, \Gamma)$ (indicating a negative bias) as established in [2], we analyze the optimality gap $\hat{f}_{\mathcal{N}'}(\mathbb{M}_G, \Gamma) - f^*(\mathbb{M}_G, \Gamma)$. Given that

$$\mathbb{E} \left[\hat{f}_{\mathcal{N}'}(\mathbb{M}_G, \Gamma) - \bar{f}_{|\mathcal{N}|, \mathcal{C}}(\mathbb{M}_G, \Gamma) \right] \geq \hat{f}(\mathbb{M}_G, \Gamma) - f^*(\mathbb{M}_G, \Gamma), \quad (6)$$

and invoking the Central Limit Theorem as in [2], the optimality gap with confidence level ρ for a sample size $|\mathcal{N}|$ is:

$$\Delta_{|\mathcal{N}|, \rho} = (\bar{f}_{\mathcal{N}'}(\mathbb{M}_G, \Gamma) - \bar{f}_{|\mathcal{N}|, \mathcal{C}}(\mathbb{M}_G, \Gamma)) + z_\rho \left(\sigma_{\mathcal{N}'}^2 + \sigma_{|\mathcal{N}|, \mathcal{C}}^2 \right)^{\frac{1}{2}}, \quad (7)$$

where $z_\rho = \phi^{-1}(1 - \rho)$. This result establishes an upper bound on the deviation from optimality under the given confidence level.

References

1. Neely, M.: Stochastic Network Optimization with Application to Communication and Queueing Systems, vol. 3 (01 2010)
2. Kleywegt, A.J., Shapiro, A., Homem-de Mello, T.: The sample average approximation method for stochastic discrete optimization. SIAM Journal on Optimization **12**(2), 479–502 (2002)