



OPEN ACCESS

EDITED AND REVIEWED BY
Liren Qian,
Chinese PLA General Hospital, China

*CORRESPONDENCE
Lu-Hong Xu
✉ xulhong@mail.sysu.edu.cn
Jian-Pei Fang
✉ fangjpei@mail.sysu.edu.cn

†These authors have contributed
equally to this work

RECEIVED 29 August 2026
REVISED 29 August 2026
ACCEPTED 07 September 2026
PUBLISHED 18 September 2026

CITATION
Liu D-D, Hong W-C, Qiu K-Y,
Li X-Y, Liu Y, Zhu L-W, Lai W-X,
Chen H-, Yang H-Q, Xu L-H
and Fang J-P (2026) Correction:
Umbilical cord blood: a promising
source for allogeneic CAR-T cells.
Front. Oncol. 16:1982369.
doi: 10.3389/fonc.2026.1982369

COPYRIGHT
© 2026 Liu, Hong, Qiu, Li, Liu, Zhu, Lai,
Chen, Yang, Xu and Fang. This is an
open-access article distributed under the
terms of the [Creative Commons
Attribution License \(CC BY\)](#). The use,
distribution or reproduction in other
forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Correction: Umbilical cord blood: a promising source for allogeneic CAR-T cells

Dian-Dian Liu^{1,2†}, Wei-Cong Hong^{1,2†}, Kun-Yin Qiu^{1,2}, Xin-Yu Li^{1,2},
Yong Liu^{1,2}, Li-Wen Zhu^{1,2}, Wei-Xin Lai^{1,2}, Han- Chen^{1,2},
Hua-Qing Yang^{1,2}, Lu-Hong Xu^{1,2*} and Jian-Pei Fang^{1,2*}

¹Department of Pediatrics, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, China,

²Key Laboratory of Malignant Tumor Gene Regulation and Target Therapy of Guangdong Higher
Education Institutes, Sun Yat-sen University, Guangzhou, China

KEYWORDS

chimeric antigen receptor T cells, umbilical cord blood, peripheral blood, acute
lymphoblastic leukemia, T-lymphocyte subsets, NCG-immunodeficient mice

A Correction on

Umbilical cord blood: a promising source for allogeneic CAR-T cells

By Liu D-D, Hong W-C, Qiu K-Y, Li X-Y, Liu Y, Zhu L-W, Lai W-X, Chen H, Yang H-Q, Xu L-H and
Fang J-P (2022). *Front. Oncol.* 12:944248. doi: 10.3389/fonc.2022.944248

There was a mistake in [Figure 4B](#) as published the panel corresponding to the CB
CAR-T group at day 21 was inadvertently misused. The corrected [Figure 4](#) appears below.

In the Statistical analysis section, “Kruskal–Wallis Friedman tests” has been corrected to
“Kruskal–Wallis test,” “one-way analysis of variance” has been corrected to “one-way
ANOVA,” and “significant” has been corrected to “significant.”

The correction has been made to the section **Materials and methods**, *Statistical analysis*:
“SPSS 24.0 and GraphPad Prism version 8.0 (GraphPad Software, San Diego, CA, USA)
were used to perform one-way analysis of variance, t-tests (with Welch’s correction for
unequal variances), and Kruskal–Wallis test, depending on the data distribution (normal
distribution or not). p-values <0.05 were treated as statistically significant (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).”

In the **Results** section, *Antitumor efficacy of cord blood CAR-T, peripheral blood CAR-T,
and peripheral blood (patient) CAR-T in vivo*, Paragraph 2, line 4 to 6. the numbers of
surviving mice in the CB CAR-T and PB CAR-T groups were reversed and have
been corrected.

“At the end of the experiment, there were three surviving mice in the CB CAR-T group
and four in the PB CAR-T group ([Figure 4B](#)).” has been corrected to “At the end of the
experiment, there were four surviving mice in the CB CAR-T group and three in the PB
CAR-T group ([Figure 4B](#)).”

The original version of this article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organiza-

tions, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 4

CB CAR-T and PB CAR-T had better antitumor efficacy in vivo. (A) Flow chart of the in vivo experiment. (B) Bioluminescence imaging of NALM6-GL-burdened mice treated with the three groups of CAR-T cells and control T cells. (C) Survival analysis of mice treated with the three groups of CAR-T cells and control T cells. (D) Proportions of CAR-T cells in the peripheral blood of mice on days 14 and 28. (E) Proportions of tumor cells in the peripheral blood of mice on days 14 and 28. * $p < 0.05$, ** $p < 0.01$.