



OPEN

Correction: Estuaries of the Tinto, Odiel and Piedras rivers as source of new species of *Pseudomonas* with biofertilizer potential under stress conditions

Noris J. Flores-Duarte^{ID}, Ignacio D. Rodríguez-Llorente^{ID}, Eloisa Pajuelo^{ID},
Susana Redondo-Gómez^{ID}, Enrique Mateos-Naranjo^{ID} & Salvadora Navarro-Torre^{ID}

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-025-31256-y>, published online 09 December 2025

In the original version of this Article, the name of the type strain of *Pseudomonas onubensis* sp. nov and the deposit number from CECT were incorrect, where

“The type strain is L1^T (= MIS C2^T = CECT 31157^T = LMG 34132^T) and was isolated from the rhizosphere of *Medicago* spp. plants.”

now reads:

“The type strain is L1^T (= MIS 1^T = CECT 31092^T = LMG 34132^T) and was isolated from the rhizosphere of *Medicago* spp. plants.”

Description of *Pseudomonas onubensis* sp. nov.

Pseudomonas onubensis (o.nu.ben'sis. L. fem. adj. *onubensis*, of or belonging to Onuba, the ancient Latin name of Huelva, a city in Spain, where the type strain was isolated).

Cells are individual and motile Gram-negative rods. Colonies are light-yellow, convex, irregular, smooth, and wet with undulate edges on TSA plates (pH = 7) at 30 °C for 24 h (optimal conditions). The growth ranges are pH 6–9, 5–40 °C, and 0–0.5 M NaCl.

Catalase- and oxidase-positive. According to the API 20NE test, it can assimilate D-glucose, L-arabinose, D-mannose, D-mannitol, N-acetyl-glucosamine, potassium gluconate, capric acid, malic acid, and trisodium citrate. Nitrate reduction, indole formation, D-glucose fermentation, aesculin and gelatin hydrolysis, and assimilation of D-maltose, adipic acid, and phenylacetic acid are negative. Strong enzymatic activity of arginine dihydrolase and no activity of urease or β -galactosidase are observed.

The genome of the type strain L1^T has 6,653,686 bp assembled into 383 contigs, with a mean coverage of 121.3 × and an N50 value of 50,308 bp. The G + C content is 59.9%.

The type strain is L1^T (= MIS 1^T = CECT 31092^T = LMG 34132^T) and was isolated from the rhizosphere of *Medicago* spp. plants. The GenBank accession number for the 16S rRNA gene sequence is OM397092. The GenBank/EMBL/DDBJ accession number for the draft genome is CBFHAZ01.

The original Article has been corrected.

Published online: 24 September 2026

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2026