

AI-Driven Nutrient Optimization Systems for Precision Agriculture: A Data-Integrated Approach to Reducing Nutrient Runoff and Enhancing Soil Health in Agricultural Systems

Obebe Olufisayo Andrew¹, Akinola Ebenezer Abayomi²

^{1,2}Kentucky State University, Frankfort, USA.

Abstract

Nutrients play a fundamental role in plant growth, soil fertility, and overall agricultural productivity. However, inefficient fertilizer application in conventional farming systems often leads to nutrient imbalance, reduced soil quality, and increased environmental degradation. The emergence of precision agriculture and artificial intelligence (AI) has introduced new opportunities for improving nutrient management through data-driven decision-making. This study investigates an AI-driven nutrient optimization system designed to enhance fertilizer efficiency while reducing nutrient runoff and improving soil health. Findings indicate that AI-based systems significantly improve nutrient use efficiency by enabling site-specific fertilizer recommendations and adaptive management strategies. The integration of multi-source agricultural data enhances prediction accuracy for soil nutrient status and crop demand, while also supporting early identification of nutrient loss risks such as runoff and leaching. The study concludes that AI-driven nutrient optimization provides a sustainable and scalable approach for modern agriculture, supporting both productivity enhancement and environmental protection through intelligent, data-integrated farming systems.

Keywords: Artificial intelligence, Digital agriculture, Nutrient, Soil health, Data integration, Machine learning, Sustainable agriculture.

1. Introduction

Nutrients are essential chemical elements required for plant growth, crop development, and productive agricultural systems. Primary macronutrients such as nitrogen, phosphorus, and potassium support root growth, photosynthesis, and biomass production, while micronutrients help maintain plant health and physiological balance (Noulas et al., 2023).

Balanced nutrient supply is therefore critical to crop yield, food production, and long-term soil fertility. Nutrient management refers to the planned application and monitoring of nutrient inputs to meet crop demand efficiently while protecting environmental resources. Its importance has increased as agriculture faces the challenge of raising productivity while sustaining soil quality and ecosystem health (Khambalkar et al., 2025). Efficient fertilizer use is now recognized as vital for food security, waste reduction, and environmental protection (Penuelas et al., 2023). Proper nutrient management also improves soil structure and biological activity (Mamatha et al., 2024).

Conventional fertilizer practices often rely on uniform application methods that ignore variations in soil conditions, crop demand, and climate. This can cause over-application in some areas and nutrient deficiency in others, lowering efficiency and increasing production costs (Cai et al., 2025). Poor nutrient allocation may also reduce profitability through unnecessary input expenditure and lower yields (Rodrigues, 2022). Environmental concerns linked to inefficient nutrient use have grown significantly. Excess nitrogen and phosphorus often move beyond the crop root zone through runoff and leaching, causing eutrophication, water pollution, and ecosystem decline (Mahmud et al., 2021). Nutrient losses can also release greenhouse gases such as nitrous oxide, increasing climate-related risks (Wang et al., 2024). Artificial intelligence has created new opportunities for precision agriculture and nutrient optimization. Machine learning, deep learning, and predictive analytics can process complex farm data, detect patterns, predict crop responses, and generate accurate fertilizer recommendations (Liakos et al., 2018). Kamilaris and Prenafeta-Boldú (2018) noted that deep learning is especially effective for analyzing imagery, sensor streams, and other large agricultural datasets. This makes AI highly suitable for adaptive nutrient decision-making.

Digital agriculture has also expanded access to data from soil sensors, weather stations, satellite imagery, drones, machinery, and geospatial platforms. These big data systems enable real-time collection, storage, and analysis of farm information, improving management efficiency and decision quality (Wolfert et al., 2017). Wider availability of such technologies has accelerated precision agriculture adoption across many farming systems (Lee et al., 2021).

Conventional nutrient management methods based only on soil tests or fixed fertilizer schedules are often inadequate for modern farming conditions. Effective decisions now require simultaneous consideration of soil variability, climate uncertainty, crop demand, production cost, and environmental risk. AI-enabled systems can integrate these variables into a smarter and more responsive optimization framework (Ennaji et al., 2023). This study aims to develop an AI-driven nutrient optimization system for precision agriculture using a data-integrated framework that improves soil health while reducing nutrient runoff.

2. Foundations of AI-Driven Nutrient Management Systems

AI-driven nutrient management systems are built on the integration of agronomic knowledge, digital technologies, and data-based decision support tools. These systems aim to improve fertilizer efficiency, crop productivity, and environmental sustainability through accurate nutrient planning and site-specific management (Penuelas et al., 2023). The growing demand for sustainable food production has made advanced nutrient management an important component of modern agriculture.

Precision agriculture provides the operational foundation for these systems by using sensors, geospatial tools, and variable-rate technologies to match nutrient application with field conditions. This approach reduces waste, improves input efficiency, and supports better farm management decisions (Rodrigues, 2022). Site-specific nutrient management technologies have also expanded the ability to monitor soil variability and crop nutrient demand across agricultural landscapes (Vullaganti et al., 2025). Artificial intelligence strengthens nutrient management by analyzing complex datasets and generating predictive recommendations. Machine learning models can evaluate soil properties, weather trends, and crop responses to recommend suitable fertilizer strategies (Liakos et al., 2018). Deep learning techniques are also valuable for processing sensor signals, imagery, and other large agricultural datasets (Kamilaris & Prenafeta-Boldú, 2018). Recent studies show that AI applications in nutrient management can optimize chemical fertilizer use while reducing environmental pressure. These systems support efficient nutrient

allocation, lower losses, and improved sustainability outcomes (Ennaji et al., 2023; Cai et al., 2025). As digital farming continues to expand, AI-driven nutrient management systems are becoming a core foundation for resilient and efficient agricultural production.

2.1 Concept of Nutrient Optimization in Precision Agriculture

Nutrient optimization in precision agriculture refers to the strategic management of plant nutrients to supply crops with the right nutrient source, at the right rate, right time, and right place in order to maximize productivity while minimizing waste and environmental harm (Miao, 2023). It integrates agronomic knowledge with digital technologies such as sensors, geographic information systems (GIS), artificial intelligence, and data analytics to improve fertilizer decisions based on field-specific conditions rather than generalized recommendations (Mishra & Mishra, 2024). This approach has become increasingly important as farming systems seek to improve efficiency, profitability, and sustainability.

Macronutrients such as nitrogen (N), phosphorus (P), and potassium (K) are required in large quantities and are fundamental for photosynthesis, root development, energy transfer, and yield formation (Gürbüz, 2025). Micronutrients, including zinc, iron, boron, and manganese, are needed in smaller amounts but remain essential for enzyme activation, metabolic regulation, and crop quality (Shrivastav et al., 2020). Deficiency or imbalance of either macro- or micronutrients can significantly reduce crop performance and soil productivity.

A major objective of nutrient optimization is improving nutrient use efficiency (NUE), which measures how effectively applied nutrients are converted into crop output. Low NUE remains a widespread challenge across major global cropping systems, leading to economic losses and environmental pollution (Liu et al., 2025). Practices such as balanced fertilization, crop-responsive nutrient scheduling, and integrated nutrient management help increase efficiency and resilience under changing climatic conditions (Govindasamy et al., 2023; Pierre & Regmi, 2026).

Modern agriculture is gradually shifting from conventional blanket fertilizer application, where identical rates are applied uniformly across fields, to site-specific nutrient management. This transition recognizes spatial variability in soil fertility, moisture, and crop demand within farms. By using precision technologies, farmers can apply nutrients variably according to actual field needs, thereby reducing over-application, lowering costs, and enhancing environmental sustainability (Akhil et al., 2025; Hasan et al., 2026).

2.2 Soil Health and Nutrient Dynamics in Agricultural Systems

Soil health refers to the continued capacity of soil to function as a living ecosystem that sustains plants, animals, and humans while supporting environmental quality and agricultural productivity. In sustainable agriculture, healthy soils are essential because they regulate water, cycle nutrients, store carbon, and maintain biodiversity needed for long-term crop production (Lehmann et al., 2020). Soil health assessment has evolved beyond crop yield measurements to include physical, chemical, and biological indicators that determine the resilience and productivity of agricultural soils (Karlen et al., 2019). Sustainable farming systems, therefore, depend on maintaining soil functionality rather than focusing only on short-term nutrient inputs (Bünemann et al., 2018).

Soil organic matter is one of the most important components of healthy soils because it improves nutrient retention, aggregation, water holding capacity, and cation exchange processes. It also serves as a major energy source for soil microorganisms that drive decomposition and nutrient mineralization (Audette et al., 2021). Higher soil organic matter levels can enhance rhizosphere nutrient cycling by influencing microbial metabolic activity and bacterial ecological strategies (Liang et al., 2026). Microbial

communities play a central role in converting organic residues into plant-available nutrients such as nitrogen, phosphorus, and sulfur, thereby sustaining soil fertility and crop growth (Dincă et al., 2022). The use of organic fertilizers has also been shown to increase microbial diversity and stabilize nutrient cycling processes in agricultural soils (Chen et al., 2025).

Soil pH strongly affects nutrient solubility and plant uptake because highly acidic or alkaline conditions can reduce the availability of essential nutrients and increase toxicity risks. Maintaining optimum pH is therefore necessary for balanced fertilization and efficient nutrient absorption (Hartemink & Barrow, 2023). Soil texture also influences nutrient dynamics because clay-rich soils generally retain more nutrients and water than sandy soils, which are more prone to leaching losses (Zheng et al., 2021). In addition, soil structure determines aeration, root penetration, and water infiltration, while soil moisture controls microbial activity and nutrient transport within the soil profile (Dewangan et al., 2024).

Improper fertilizer management can degrade soil productivity over time. Excessive fertilizer use may lead to soil acidification, salinity buildup, nutrient imbalances, loss of beneficial microbes, and contamination of surrounding ecosystems (Yanardağ, 2024). Conversely, under-fertilization can deplete nutrient reserves, reduce soil biological activity, and lower crop yields over successive seasons. Therefore, long-term agricultural sustainability depends on balanced nutrient replenishment strategies that maintain soil fertility without causing environmental damage (Bhattacharyya & Furtak, 2022).

2.3 Artificial Intelligence in Precision Agriculture

Artificial intelligence (AI) in precision agriculture refers to the use of computational systems that mimic human decision-making to improve farming efficiency, productivity, and sustainability. Core AI methodologies include machine learning, deep learning, and predictive analytics, which enable computers to identify patterns, learn from historical data, and generate recommendations for farm management decisions (Liakos et al., 2018). Machine learning models are increasingly applied to complex agricultural datasets such as weather records, sensor outputs, and crop observations to support real-time management decisions (Wang & Wang, 2026). In soil science, machine learning tools are also used to analyze soil variability, classify soil properties, and estimate nutrient conditions with greater precision than conventional methods (Padarian et al., 2020).

In remote sensing, one of the most important applications of AI in agriculture for crop monitoring, drones and smart sensors generate large volumes of field data that AI systems can analyze to detect crop stress, diseases, moisture deficiencies, and growth anomalies at early stages (Talaviya et al., 2020). This allows farmers to respond quickly to changing field conditions and reduce production risks. AI also supports automated monitoring systems that continuously track crop performance throughout the growing season, improving farm responsiveness and operational efficiency (Mishra & Mishra, 2024).

AI has also made major contributions to crop yield prediction through the use of advanced learning algorithms. Deep learning and reinforcement learning models can combine climate, soil, management, and historical yield data to forecast crop output with improved accuracy (Elavarasan & Vincent, 2020). Hybrid neural network models such as convolutional neural networks-recurrent neural networks (CNN-RNN) frameworks have proven effective in modeling temporal and spatial relationships affecting agricultural yield performance (Khaki et al., 2020). Recent developments integrating explainable artificial intelligence further improve trust and transparency in predictive systems used for farm planning (Mohan et al., 2025).

In nutrient management, AI-driven recommendation systems help determine appropriate fertilizer types, rates, timing, and placement according to site-specific crop and soil needs. These systems process data

from soil tests, sensors, satellite imagery, and management records to optimize nutrient applications while reducing waste and environmental losses (Miao, 2023). Predictive analytics also enables farmers to anticipate nutrient deficiencies and make proactive fertilizer adjustments before yield losses occur (Jabed & Murad, 2024).

Decision support systems (DSS) powered by AI are becoming central tools for fertilizer optimization in precision agriculture. These platforms integrate agronomic knowledge with real-time data analytics to generate actionable recommendations for farmers and agronomists. AI-based DSS can improve nutrient use efficiency, lower production costs, and enhance sustainability outcomes through evidence-based fertilizer decisions (Mishra & Mishra, 2024). Despite these benefits, AI adoption in agriculture faces several limitations and challenges. Many farming regions still lack reliable digital infrastructure, quality datasets, and the technical expertise required to implement advanced systems. High investment costs, limited digital literacy, and uncertainty about economic returns can also slow adoption among farmers (Ansari et al., 2024). In addition, model accuracy may decline when algorithms trained in one region are transferred to different soils, climates, or crop systems, highlighting the need for localized calibration and continuous system improvement (Liakos et al., 2018). *Figure 1* illustrates the distribution of machine learning techniques applied in agricultural systems, highlighting their usage across different agricultural domains such as crop production, soil management, water systems, and farmer decision support (Cravero et al., 2026).




Figure 1: Machine Learning Techniques Employed in the Development of Agricultural Systems (Cravero et al., 2026).

2.4 Data Integration in Smart Nutrient Management

Data integration is central to smart nutrient management because effective fertilizer decisions depend on combining multiple agricultural data sources into a unified system. Important sources include field

sensors, Internet of Things (IoT) devices, drones, satellite imagery, geographic information systems (GIS), and farm management records that provide information on soil conditions, crop growth, weather patterns, and management history (Soussi et al., 2024). These technologies improve the accuracy and timeliness of nutrient recommendations by generating continuous farm-level data (Kour & Arora, 2020).

Data fusion techniques are used to combine soil, crop, and climate datasets into meaningful decision outputs. Machine learning models can integrate heterogeneous data streams to identify nutrient deficiencies, predict crop demand, and optimize fertilizer application across variable field conditions (Wang & Wang, 2026). In soil management, data-driven models are increasingly used to analyze soil variability and nutrient status for precision interventions (Padarian et al., 2020). AI-based systems also support crop yield forecasting by linking environmental and management data for proactive nutrient planning (Elavarasan & Vincent, 2020).

Despite these advantages, several challenges affect integrated agricultural data systems; Poor sensor calibration, missing values, inconsistent data formats, and weak interoperability between devices can reduce decision accuracy and system reliability (Fizza et al., 2021). Limited digital infrastructure, adoption costs, and compatibility issues across platforms also constrain implementation, particularly in developing farming regions (Ansari et al., 2024). Real-time interoperable systems are therefore needed to ensure seamless communication among smart farm technologies (El-Dosuky, 2025).

Real-time analytics plays a vital role in nutrient decision-making by enabling immediate responses to changing soil and crop conditions. AI systems can process incoming field data continuously and recommend timely fertilizer adjustments that improve nutrient use efficiency and reduce losses (Mishra & Mishra, 2024). Such intelligent platforms are becoming essential tools for future on-farm planning, productivity enhancement, and sustainable nutrient stewardship (Hasan et al., 2026). *Table 1* presents examples of IoT-based smart agriculture platforms used for data collection, monitoring, and decision support in precision farming systems. These platforms integrate sensor technologies, cloud systems, and analytics tools to enhance agricultural productivity and resource management (Soussi et al., 2024).

Table 1: Examples of Platforms Available for Smart Agriculture (Soussi et al., 2024)

IoT Platform	Application
Phytoprove (2019)	Develops tools for early, rapid, and non-destructive determination of nitrogen and water supply status of plants.
Telemetry2u	Provides integrated hardware and software system for measuring temperature and irrigation control.
Sensoterra	Provides soil moisture monitoring using integrated sensor systems.
CropX (2013)	Enables measurement of electrical conductivity, soil moisture, and temperature for precision nutrient management.
ONFARM	Provides dashboard-based farm data visualization for real-time decision support.
Phytech	Uses data-driven insights to optimize irrigation and improve plant health and resource efficiency.
Semios (2020)	Provides real-time monitoring of pests, diseases, and crop health for yield optimization.

2.5 Nutrient Runoff and Environmental Sustainability

Nutrient runoff and leaching are major environmental challenges associated with modern agricultural production. They occur when excess nutrients, especially nitrogen and phosphorus, are transported from farmland through surface runoff, erosion, irrigation drainage, or downward water movement beyond the root zone. These losses are often caused by excessive fertilizer application, poor timing of nutrient placement, heavy rainfall events, unsuitable irrigation practices, and limited crop nutrient uptake (Juncal et al., 2023). In many farming systems, inefficient nutrient management increases the likelihood of nutrients escaping into surrounding ecosystems instead of being utilized by crops (Del Rossi et al., 2023). The environmental consequences of nutrient runoff are significant. Nutrient enrichment of rivers, lakes, and coastal waters can trigger eutrophication, leading to algal blooms, oxygen depletion, and degradation of aquatic ecosystems. Excessive nitrogen and phosphorus use for major crops has also been linked to biodiversity loss in freshwater environments and wider ecological disruption (Jwaideh et al., 2022). In addition, agricultural soils are an important source of greenhouse gas emissions such as nitrous oxide, which contributes substantially to climate change (Basheer et al., 2024). Nitrogen losses through volatilization and gaseous emissions, therefore, represent both environmental and agronomic inefficiency (Pan et al., 2022). These concerns have increased policy and regulatory attention toward nutrient management in agriculture. Governments and environmental agencies are adopting stricter standards on fertilizer use, water quality protection, and emission reduction targets. Economic analyses show that nutrient pollution imposes substantial social and environmental costs, creating a strong case for improved regulations, incentives, and sustainable nutrient stewardship policies (Del Rossi et al., 2023). Predictive tools are increasingly important for protecting the environment while maintaining farm productivity. Decision-support systems, nutrient budgeting models, runoff forecasting tools, and emission risk assessments help farmers anticipate losses before they occur and adjust management practices accordingly. Such tools support nutrient neutrality strategies by improving fertilizer timing, placement, and application rates based on environmental risk conditions (Juncal et al., 2023). Advanced predictive approaches are therefore essential for climate-smart and environmentally sustainable agriculture (Pan et al., 2022).

3. AI-Driven Nutrient Optimization System Design

AI-driven nutrient optimization systems are designed to combine sensing technologies, data analytics, and intelligent decision tools to improve fertilizer management in precision agriculture. These systems typically integrate soil sensors, climate data, satellite imagery, and farm records into a unified digital platform that supports accurate nutrient recommendations and efficient resource use (Soussi et al., 2024). A core component of system design is data fusion, where multiple data streams are combined using machine learning models to provide a complete understanding of field conditions. Sensor fusion techniques help interpret soil moisture, nutrient levels, crop health, and weather variability for timely nutrient planning and adaptive farm management (Wang & Wang, 2026). AI systems can therefore generate more precise recommendations than conventional single-source approaches. The soil intelligence layer focuses on collecting and analyzing soil information such as pH, organic matter, nutrient content, and moisture status. Machine learning tools are increasingly used to classify soils, predict nutrient deficiencies, and map field variability for site-specific interventions (Padarian et al., 2020). This improves the accuracy of fertilizer placement and reduces unnecessary application.

The nutrient optimization layer converts field data into practical fertilizer decisions, AI models evaluate crop demand, soil nutrient supply, and environmental conditions to recommend suitable fertilizer type, rate, timing, and placement. Precision nutrient management systems can significantly improve nutrient use efficiency while lowering production costs and minimizing environmental losses (Miao, 2023). Decision support systems are also central to modern nutrient optimization platforms. These tools provide farmers with dashboards, mobile alerts, and scenario analysis for better on-farm decisions. AI-supported systems enhance farmer decision-making by translating complex data into clear management actions (Kaushik et al., 2025). Data-driven farm management platforms are increasingly important for improving productivity and sustainability (Mishra & Mishra, 2024). Real-time capability is another critical design feature, Modern precision agriculture systems continuously monitor changing field conditions and update recommendations instantly, allowing rapid responses to crop stress, rainfall events, or nutrient deficiencies. Real-time architectures are becoming essential for scalable and sustainable smart farming systems (Bhat et al., 2026).

3.1 Systems-Level Framework Architecture

A systems-level framework architecture for AI-driven nutrient optimization is designed as an integrated digital platform that connects data collection, analysis, optimization, and advisory functions for efficient fertilizer management (Soussi et al., 2024). Such frameworks support precision agriculture by coordinating multiple technologies into one intelligent decision environment (Bhat et al., 2026). The first layer is the sensing layer, where soil sensors, weather stations, drones, satellite systems, and smart farm devices collect real-time data on nutrient levels, moisture, crop health, and environmental conditions (Soussi et al., 2024). Continuous sensing improves field visibility and creates the raw data needed for nutrient decisions (Miller et al., 2025).

The second layer is the analytics layer, where machine learning models process raw datasets into useful insights such as soil classification, nutrient deficiency detection, and crop demand forecasting (Padarian et al., 2020). Data analytics also improves accuracy by combining multiple data sources into a clearer representation of field variability (Wang & Wang, 2026). The third layer is the optimization layer, where AI algorithms recommend the best fertilizer source, rate, timing, and placement according to crop and soil requirements (Miao, 2023). This layer aims to maximize nutrient use efficiency while reducing runoff, leaching, and unnecessary input costs (Mishra & Mishra, 2024). The fourth layer is the advisory layer, where recommendations are delivered through dashboards, mobile applications, or automated farm systems for practical use by farmers and agronomists (Kaushik et al., 2025). Clear decision interfaces help transform technical outputs into actionable farm operations (Mishra & Mishra, 2024).

Data flow within the framework begins with field-level sensing and moves through transmission, analytics, optimization, and final decision outputs in real time (Bhat et al., 2026). Feedback loops allow the system to update recommendations whenever field conditions change (Wang & Wang, 2026). Interoperability is essential because these systems must connect across sensors, machinery, cloud platforms, and farm software from different providers (Soussi et al., 2024). Standardized data exchange improves compatibility, scalability, and long-term adoption of digital nutrient management systems (Bhat et al., 2026).

3.2 Soil Intelligence and Analytics Layer

The soil intelligence and analytics layer is responsible for collecting, processing, and interpreting soil data to support precise nutrient management decisions. It forms a core component of AI-driven nutrient optimization systems by transforming raw soil information into actionable insights for farmers and

agronomists (Padarian et al., 2020). Smart sensing systems now enable continuous digital monitoring of soil conditions across agricultural fields (Soussi et al., 2024).

This layer gathers soil physical data such as texture, bulk density, moisture, and temperature using in-field sensors and remote sensing tools (Miller et al., 2025). It also captures chemical data including pH, nitrogen, phosphorus, potassium, salinity, and organic carbon, needed for fertilizer planning (Hossen et al., 2021). Biological indicators such as microbial activity and soil health status are increasingly monitored through AI-enabled soil restoration systems (Pandey et al., 2025). Whereas, before analysis, raw soil datasets require preprocessing to improve quality and consistency. Common steps include noise removal, handling missing values, normalization, scaling, and integration of multi-source data from sensors, satellites, and farm records (Wang & Wang, 2026). Effective soil data sharing frameworks also improve interoperability and access across institutions and digital platforms (Padarian & McBratney, 2020).

Artificial intelligence models are then applied for soil classification and nutrient deficiency detection. Machine learning algorithms can classify soils according to fertility zones, texture groups, and management capability classes with higher speed and accuracy than conventional methods (Padarian et al., 2020). Deep learning systems are also being used to detect nutrient problems through soil image analysis and nutrient profiling tools (Pandey et al., 2025). Soil properties often differ significantly within the same farm, making uniform fertilizer application inefficient. In line with this, machine learning models using satellite, weather, clay, and yield data can map field heterogeneity and predict spatial nutrient patterns for site-specific nutrient management (Kammerlander et al., 2025). Such analytics improve fertilizer targeting, reduce waste, and enhance environmental sustainability (Soussi et al., 2024).

3.3 Nutrient Optimization Layer

The nutrient optimization layer is the decision engine of an AI-driven nutrient management system, where field data are converted into fertilizer recommendations that improve productivity and sustainability. This layer uses computational models to determine the right nutrient source, rate, timing, and placement for specific crop conditions (Miao, 2023). Data-driven nutrient planning is increasingly recognized as essential for efficient farm management under changing environmental conditions (Mishra & Mishra, 2024). Machine learning-based fertilizer recommendation systems analyze historical and real-time farm data to generate accurate nutrient prescriptions. These systems can learn relationships between soil fertility, crop response, and management practices to improve fertilizer efficiency over time (Wang & Wang, 2026). AI-based farm decision tools are also being developed to guide crop and nutrient management through automated recommendations (Kaushik et al., 2025).

Additionally, effective nutrient optimization requires integration of soil status, crop nutrient demand, and climatic variables. Soil nutrient levels, moisture conditions, crop growth stage, temperature, and rainfall patterns all influence nutrient availability and crop uptake (Elavarasan & Vincent, 2020). Combining these variables allows the system to make more adaptive and location-specific fertilizer decisions (Mishra & Mishra, 2024).

Multi-objective optimization is important because fertilizer decisions must balance yield improvement, production cost, and environmental protection. Studies show that low nutrient use efficiency remains widespread across major crops, highlighting the need for smarter allocation strategies (Liu et al., 2025). Improving nitrogen use efficiency is especially critical for sustaining crop productivity while reducing nutrient losses under climate stress (Govindasamy et al., 2023). The nutrient optimization layer also supports variable-rate and adaptive nutrient application strategies. Instead of uniform fertilizer application, nutrients are supplied according to within-field variability and changing crop needs during the season

(Akhil et al., 2025). Precision nutrient technologies can therefore enhance agronomic efficiency while lowering environmental risks associated with fertilizer misuse (Pierre & Regmi, 2026).

3.4 Environmental Risk Prediction Layer

The environmental risk prediction layer focuses on forecasting nutrient losses from agricultural systems, particularly runoff, leaching, and gaseous emissions. It uses AI and hydrological models to estimate how nutrients move through soil and water systems under different management and environmental conditions (Juncal et al., 2023). Such predictive approaches are increasingly important for supporting nutrient neutrality and sustainable land management strategies (Shao et al., 2021). Risk prediction models incorporate key environmental drivers such as rainfall intensity, terrain slope, soil properties, and land management practices. These variables strongly influence nutrient transport pathways, especially under extreme weather conditions that increase runoff and erosion risk (Yang et al., 2025). Machine learning approaches further improve prediction accuracy by capturing complex interactions between hydrological and agricultural factors (Elsayed et al., 2024).

Nutrient leaching and runoff risks are also influenced by fertilizer application type and timing. Studies show that different fertilization strategies significantly affect nitrogen and phosphorus losses in agricultural watersheds (Reisinger et al., 2025). Similar findings confirm that nutrient loss patterns vary across soil types, crop systems, and seasonal biomass conditions (Assa et al., 2024).

AI-based models are increasingly used to simulate nutrient concentrations in surface water and predict pollution loads in agricultural catchments. These models integrate spatial and temporal data to estimate how nutrients move from fields into surrounding water bodies (Lee et al., 2024). Machine learning approaches have demonstrated strong performance in identifying key drivers of nutrient runoff, particularly precipitation and land-use intensity (Elsayed et al., 2024). Threshold-based alert systems are an important component of environmental risk prediction. These systems trigger warnings when nutrient levels or environmental conditions exceed safe limits, enabling timely intervention to reduce pollution risks (Juncal et al., 2023). Such systems support compliance with environmental regulations and promote sustainable fertilizer use practices (Shao et al., 2021).

3.5 Decision Support and Advisory Layer

The decision support and advisory layer transforms AI-driven nutrient model outputs into practical guidance for farmers, ensuring that complex analytics are usable at the field level. It typically operates through farmer-oriented dashboards and mobile-based platforms that present real-time recommendations on fertilizer use, crop management, and soil conditions (Soussi et al., 2024). These systems improve accessibility by delivering actionable insights directly to end users in simplified formats (Miao, 2023). Explainable AI (XAI) plays a key role in this layer by improving transparency and trust in nutrient recommendations. XAI techniques help users understand why specific fertilizer or management decisions are suggested, making AI systems more interpretable and reliable for agricultural decision-making (Mishra & Mishra, 2024). This transparency is essential for increasing the adoption of AI-based advisory tools in farming systems (Kaushik et al., 2025).

Scenario analysis is another important function of the advisory layer, allowing farmers to evaluate different management options before implementation. AI-based decision support systems can simulate outcomes such as yield response, fertilizer efficiency, and environmental impact under different input strategies (Elavarasan & Vincent, 2020). Such scenario-based tools support proactive planning and improved resource allocation in agriculture (Hasan et al., 2026). Human expertise remains essential in AI-driven advisory systems, particularly through the integration of agronomists and extension officers into decision

workflows. These hybrid systems combine machine intelligence with expert validation to improve the accuracy and contextual relevance of recommendations (Mishra & Mishra, 2024). Studies show that human-in-the-loop designs enhance trust, usability, and adoption of agricultural AI systems (Soussi et al., 2024).

4. Implementation Strategies and Operational Considerations

Successful implementation of AI-driven nutrient optimization systems requires robust digital infrastructure capable of collecting, storing, and processing large volumes of agricultural data. Smart sensors, IoT networks, cloud platforms, and edge computing systems are essential for enabling continuous data exchange and real-time decision support in farming environments (Soussi et al., 2024). Real-time system architectures further improve responsiveness and operational efficiency under changing field conditions (Bhat et al., 2026). Reliable model deployment also depends on high-quality datasets and strong analytical frameworks. Machine learning systems require accurate soil, climate, crop, and management data for training, calibration, and validation before field application (Cravero et al., 2026). Soil parameter prediction models demonstrate the importance of integrating satellite, weather, and yield data to strengthen model accuracy and robustness (Kammerlander et al., 2025).

Interoperability is another major operational consideration because agricultural technologies often operate on different hardware and software platforms. Seamless integration among sensors, machinery, farm software, and advisory systems improves usability and scalability of AI solutions (Wang & Wang, 2026). Standardized data-sharing systems are therefore necessary for efficient smart farming ecosystems (Soussi et al., 2024). Human capacity and user adoption are equally important for successful implementation. Farmers need practical interfaces such as dashboards and mobile tools that convert complex analytics into clear management actions (Kaushik et al., 2025). Data-driven advisory systems become more effective when combined with farmer knowledge, extension support, and agronomic expertise (Mishra & Mishra, 2024). Ultimately, long-term sustainability depends on economic viability, system maintenance, and adaptability across crops and regions. Precision nutrient management systems must remain cost-effective while continuously updating models as new seasonal data become available (Miao, 2023). The integration of IoT and AI technologies is increasingly viewed as essential for scalable and future-ready agricultural systems (Miller et al., 2025).

4.1 Data Infrastructure and Governance

Effective implementation of AI-driven nutrient optimization systems depends on a strong data infrastructure capable of storing, processing, and exchanging large volumes of agricultural information. Big data architectures such as cloud platforms, edge computing systems, and agricultural data warehouses are increasingly used to manage sensor, satellite, climate, and farm management datasets in real time (Ngo et al., 2019). Modern intelligent agricultural ecosystems also rely on scalable digital infrastructures that support continuous analytics and automated decision-making (Cravero et al., 2026). Cloud computing systems improve accessibility by allowing farmers, agronomists, and institutions to access shared data and decision tools remotely. These systems enable rapid processing of high-volume datasets generated from IoT devices and precision agriculture technologies (Soussi et al., 2024). Real-time cloud-edge architectures also improve system responsiveness under changing field conditions (Bhat et al., 2026).

Data ownership, privacy, and cybersecurity remain major governance concerns in digital agriculture. Farmers often generate valuable operational data, yet uncertainty over ownership rights, access control, and commercial use can limit willingness to share information (Rouzky et al., 2025). Secure governance

frameworks are therefore necessary to protect sensitive farm data from misuse, unauthorized access, and cyber threats (Wang & Wang, 2026). Standardization and interoperability are essential because agricultural technologies often use incompatible data formats and disconnected platforms. Shared standards improve communication between sensors, machinery, farm software, and analytics tools across different providers (San Emeterio de la Parte et al., 2023). Open-data and open-source approaches can further enhance transparency, innovation, and system integration in precision agriculture ecosystems (Jeppesen et al., 2022).

Governance models for shared agricultural data ecosystems should balance innovation, collaboration, and user protection. Federated platforms and cooperative data-sharing systems are increasingly proposed to allow multiple stakeholders to benefit from common datasets while retaining local control of sensitive information (Pincheira et al., 2025). Such governance approaches are important for building trust and supporting long-term adoption of AI-enabled nutrient management systems (Rouzky et al., 2025)

4.2 Model Development and Validation

Effective nutrient optimization systems depend on selecting AI and optimization models that match the nature of agricultural data and decision objectives. Machine learning techniques such as random forests, support vector machines, neural networks, and reinforcement learning are widely used because they can model complex relationships between soil, crop, and climate variables (Liakos et al., 2018). Model choice should also consider computational efficiency, interpretability, and scalability for real farm environments (Cravero et al., 2026).

Model development requires structured training, testing, calibration, and validation procedures to ensure accurate predictions. Historical datasets are commonly divided into training and testing subsets, while cross-validation methods are used to reduce overfitting and improve generalization performance (Padarian et al., 2020). Calibration with field observations further strengthens reliability of nutrient recommendation and soil prediction models (Kammerlander et al., 2025). According to Mishra and Mishra (2024), managing uncertainty and bias is essential because agricultural datasets may contain noise, missing values, seasonal anomalies, or sampling imbalance. Poor-quality data can reduce model accuracy and create biased recommendations that do not reflect field realities (Mishra & Mishra, 2024). Therefore, robust models require preprocessing, sensitivity analysis, and explainable AI methods that help users understand prediction confidence and decision logic (Mohan et al., 2025).

Continuous learning is increasingly important in dynamic farming systems where weather, soil fertility, and management practices change over time. Updating models with seasonal observations and historical farm records allows systems to improve prediction performance across successive production cycles (Wang & Wang, 2026). This adaptive learning approach supports more resilient and location-specific nutrient management decisions.

4.3 Policy, Regulatory, and Institutional Support

Effective implementation of AI-driven nutrient optimization systems requires supportive agricultural policies that encourage adoption of precision farming technologies. Public policies that promote digital agriculture, smart machinery, and data-driven input management can improve productivity while reducing waste and environmental pressure (Cravero et al., 2026). Financial support measures such as subsidies and innovation grants are also important for helping farmers overcome high initial investment costs (Sanyaolu & Sadowski, 2024).

Environmental regulations play a central role in guiding fertilizer use and nutrient management practices. Regulatory frameworks increasingly emphasize nutrient-use efficiency, reduced runoff, and compliance

with sustainability targets related to soil and water protection (Juncal et al., 2023). Precision nutrient technologies can help farmers meet these standards through more accurate timing, placement, and rates of fertilizer application (Maaz et al., 2025).

Extension services and public institutions are essential for translating policy goals into farm-level adoption. Advisory agencies, research institutes, and agricultural ministries often provide training, technical guidance, and demonstration programs that help farmers understand and apply precision agriculture tools (Nkosi et al., 2025). Strengthening extension capacity is particularly important in developing regions where digital skills and technical awareness may be limited (Onyango et al., 2021). Well-designed incentive structures can accelerate adoption of sustainable agricultural technologies. Evidence shows that payments, tax incentives, cost-sharing schemes, and sustainability rewards can encourage farmers to invest in environmentally beneficial innovations (Piñeiro et al., 2020). When aligned with nutrient stewardship goals, such incentives support wider use of AI-enabled nutrient optimization systems and long-term agricultural sustainability.

4.4 Stakeholder Engagement and Adoption

Stakeholder engagement is a central requirement for the successful deployment of digital and AI-driven agricultural technologies because system effectiveness depends on collaboration among farmers, agronomists, researchers, and agri-tech companies. Precision agriculture adoption is strongly influenced by how well technologies align with farmer needs, contextual realities, and institutional support systems (Soussi et al., 2024). Therefore, integrated smart agricultural ecosystems depend on coordinated participation across all actors in the value chain (Cravero et al., 2026).

Farmers play the primary role as end-users, while agronomists and extension agents act as intermediaries who translate technical outputs into practical field decisions. Research shows that adoption is higher when farmers receive continuous technical support and when innovation systems connect them directly with experts and service providers (Mishra & Mishra, 2024). Agri-tech companies and researchers contribute by developing tools that integrate sensor data, machine learning models, and decision support systems for real-world use (Wang & Wang, 2026).

However, adoption is often constrained by barriers such as high cost, limited digital literacy, inadequate infrastructure, and lack of trust in automated systems. Studies indicate that financial limitations and insufficient training significantly reduce farmers' willingness to adopt precision technologies (Cravero et al., 2026). In addition, weak digital skills and limited access to infrastructure continue to slow down technology diffusion in many farming communities (Miller et al., 2025). Capacity building and training initiatives are essential to overcome these constraints and improve adoption outcomes. Continuous farmer education, demonstration farms, and extension-led training programs help bridge the gap between innovation and practice (Kaushik et al., 2025). Evidence also shows that digital literacy and hands-on training significantly improve farmers' confidence in using smart agricultural systems (Soussi et al., 2024). Participatory design and a co-creation approach further enhance adoption by involving farmers directly in the development of agricultural technologies. Co-design ensures that tools are user-friendly, context-specific, and aligned with real farm challenges (Bhat et al., 2026). This collaborative model strengthens trust, improves usability, and increases long-term sustainability of precision agriculture solutions (Mishra & Mishra, 2024).

4.5 Scalability and Long-Term Sustainability

Scalability of AI-driven nutrient optimization systems depends on their ability to function across diverse crops, farm sizes, and agro-ecological zones. Modern precision agriculture frameworks are increasingly

designed as integrated ecosystems that combine sensing, analytics, and decision systems to support large-scale deployment across heterogeneous environments (Cravero et al., 2026). These systems rely on flexible digital architectures that can adapt to varying climatic, soil, and management conditions (Soussi et al., 2024).

Economic feasibility is a key determinant of long-term adoption, particularly for smallholder farmers who face cost and infrastructure constraints. Studies show that while AI-based precision agriculture improves efficiency and productivity, its adoption is strongly influenced by affordability and access to technology (Bhat et al., 2026). Scalable and low-cost digital solutions are therefore essential to ensure inclusivity across different farming systems (Wang & Wang, 2026). System maintenance and continuous updating are necessary to maintain model accuracy over time. Agricultural AI systems must be regularly retrained using new seasonal and historical farm data to account for changing climate conditions, soil variability, and management practices (Mishra & Mishra, 2024). Continuous learning frameworks help ensure that predictive performance remains reliable across production cycles (Kammerlander et al., 2025).

Long-term sustainability of nutrient optimization systems is closely linked to their contribution to regenerative and climate-smart agriculture. AI-enabled precision farming can improve resource efficiency, reduce environmental degradation, and support soil restoration practices when properly implemented (Miao, 2023). Evidence suggests that integrating AI with sustainable nutrient management strategies enhances both productivity and environmental resilience in agricultural systems.

5. Conclusion

AI-driven nutrient optimization systems represent a major advancement in modern agriculture by integrating data analytics, machine learning, and precision sensing into a unified decision-support framework. These systems enable more accurate nutrient management by combining soil, crop, and environmental information for improved fertilizer recommendations and resource efficiency.

The transition from conventional fertilization practices to data-driven precision agriculture enhances productivity while reducing environmental impacts such as nutrient runoff, leaching, and soil degradation. By optimizing nutrient application at the right time and place, these systems improve soil health and long-term agricultural sustainability. Integrated data systems and real-time analytics strengthen decision-making by providing continuous monitoring of field conditions and enabling adaptive nutrient management. This improves responsiveness to climate variability and supports more efficient use of agricultural inputs.

Effective implementation of AI-based nutrient management requires collaboration among farmers, researchers, policymakers, and technology developers. Strong institutional support, digital infrastructure, and capacity-building initiatives are essential for scaling these technologies across diverse agricultural systems. Overall, AI-enabled nutrient stewardship offers a sustainable pathway for improving global food production while protecting environmental resources, supporting climate-smart agriculture, and enhancing long-term soil fertility management.

References

3. Akhil, K., Renukaswamy, N. S., Sowjanya, J., & Aruna, K. S. (2025). Site-specific nutrient management: Revolutionizing fertilization for future farming. *INTERNATIONAL JOURNAL*, 9(3), 544-549.

4. Anas, M., Liao, F., Verma, K. K., Sarwar, M. A., Mahmood, A., Chen, Z. L., Li, Q., Zeng, X. P., Liu, Y., & Li, Y. R. (2020). Fate of nitrogen in agriculture and environment: agronomic, eco-physiological and molecular approaches to improve nitrogen use efficiency. *Biological research*, 53(1), 47. <https://doi.org/10.1186/s40659-020-00312-4>
5. Ansari, M. N., Gupta, S., Kumar, D., & Hussain, B. (2024). Adoption challenges and incentives for IoT and AI in agriculture. In *Agriculture 4.0* (pp. 355-372). CRC Press.
6. Assa, B. G., Bhowmick, A., & Cholo, B. E. (2024). Assessing Nitrate Leaching and Runoff Coefficients in the Dynamic Interplay of Seasonal Crop Biomass: A Study of Surface and Groundwater Nitrate Contamination in the Bilate Cropland Watershed. *Environmental Advances*, 16, 100528.
7. Audette, Y., Congreves, K. A., Schneider, K., Zaro, G. C., Nunes, A. L., Zhang, H., & Voroney, R. P. (2021). The effect of agroecosystem management on the distribution of C functional groups in soil organic matter: A review. *Biology and Fertility of Soils*, 57(7), 881-894.
8. Basheer, S., Wang, X., Farooque, A. A., Nawaz, R. A., Pang, T., & Neokye, E. O. (2024). A review of greenhouse gas emissions from agricultural soil. *Sustainability*, 16(11), 4789.
9. Bhat, M. H., da Silva Rickiel, F., Sameer, B., Aeshna, S., & Moore, K. J. (2026). Precision Agriculture Through a Real-Time Systems Perspective: A Narrative Review. *Agronomy*, 16(5), 552.
10. Bhattacharyya, S. S., & Furtak, K. (2022). Soil–Plant–Microbe interactions determine soil biological fertility by altering rhizospheric nutrient cycling and biocrust formation. *Sustainability*, 15(1), 625.
11. Bünemann, E. K., Bongiorno, G., Bai, Z., Creamer, R. E., De Deyn, G., De Goede, R., ... & Brussaard, L. (2018). Soil quality—A critical review. *Soil biology and biochemistry*, 120, 105-125.
12. Cai, B., Shi, F., Geremew, B. A., Addis, A. K., Abate, M. C., Dessie, W., & Bayu, T. (2025). Precision agriculture techniques for optimizing chemical fertilizer use and environmental sustainability: a systematic review. *Frontiers in Agronomy*, 7, 1665444.
13. Chen, X., Ma, X., Liu, Z., Gu, H., Fang, H., Shen, Z., ... & Liu, J. (2025). Organic fertilizers increase microbial community diversity and stability slowing down the transformation process of nutrient cycling. *Environmental Microbiome*, 20(1), 130.
14. Coble, K. H., Mishra, A. K., Ferrell, S., & Griffin, T. (2018). Big data in agriculture: A challenge for the future. *Applied Economic Perspectives and Policy*, 40(1), 79-96.
15. Cravero, A., Sepúlveda, S., Gutiérrez, F., & Muñoz, L. (2026). From Precision Agriculture to Intelligent Agricultural Ecosystems: A Systematic Review of Machine Learning and Big Data Applications. *Agronomy*, 16(5), 516.
16. Del Rossi, G., Hoque, M. M., Ji, Y., & Kling, C. L. (2023). The economics of nutrient pollution from agriculture. *Annual Review of Resource Economics*, 15(1), 105-130.
17. Dewangan, S. K., Kashyap, N., Tigga, D., & Turkey, N. (2024). The role of physico-chemical properties in soil functionality: a literature review. *EPRA International Journal of Research & Development*, 9(12), 167-173.
18. Dincă, L. C., Grenni, P., Onet, C., & Onet, A. (2022). Fertilization and soil microbial community: a review. *Applied Sciences*, 12(3), 1198.
19. Elavarasan, D., & Vincent, P. D. (2020). Crop yield prediction using deep reinforcement learning model for sustainable agrarian applications. *IEEE access*, 8, 86886-86901.
20. El-Dosuky, M. (2025). Real-time Framework for Interoperable Semantic-driven Internet-of-Things in Smart Agriculture. *arXiv preprint arXiv:2510.05187*.

21. Elsayed, A., Rixon, S., Levison, J., Binns, A., & Goel, P. (2024). Machine learning models for prediction of nutrient concentrations in surface water in an agricultural watershed. *Journal of Environmental Management*, 372, 123305.
22. Ennaji, O., Vergütz, L., & El Allali, A. (2023). Machine learning in nutrient management: A review. *Artificial Intelligence in Agriculture*, 9, 1-11.
23. Fizza, K., Jayaraman, P. P., Banerjee, A., Georgakopoulos, D., & Ranjan, R. (2021). Evaluating sensor data quality in Internet of Things smart agriculture applications. *Ieee Micro*, 42(1), 51-60.
24. Govindasamy, P., Muthusamy, S. K., Bagavathiannan, M., Mowrer, J., Jagannadham, P. T. K., Maity, A., ... & Tiwari, G. (2023). Nitrogen use efficiency—a key to enhance crop productivity under a changing climate. *Frontiers in Plant Science*, 14, 1121073.
25. Gürbüz, M. A. (2025). Identification of Multiple Extraction Methods to be Used in Extraction of Macro and Micronutrients of Neutral and Alkaline Soils by a Multi-Criteria Decision-Making Technique (TOPSIS). *Journal of Soil Science and Plant Nutrition*, 25(2), 4669-4686.
26. Hartemink, A. E., & Barrow, N. J. (2023). Soil pH-nutrient relationships: the diagram. *Plant and Soil*, 486(1), 209-215.
27. Hasan, T., Mudgal, G., Ramasamy, V., Kumar, A., & Kugonza, D. R. (2026). Precision Futures: Leveraging Modern Technology for Intelligent On-Farm Planning and Production Efficiency. In *Smart Sustainable Materials Advancing Re-manufacturing and Technology: Smart and Sustainable Development in Agricultural Technology: Ecological and Practical Applications (Volume IV)* (pp. 247-275). Cham: Springer Nature Switzerland.
28. Hossen, M. A., Diwakar, P. K., & Ragi, S. (2021). Total nitrogen estimation in agricultural soils via aerial multispectral imaging and LIBS. *Scientific Reports*, 11(1), 12693.
29. Howari, F. (2025). AI-Remote Sensing for Soil Variability Mapping and Precision Agrochemical Management: A Comprehensive Review of Methods, Limitations, and Climate-Smart Applications. *Agrochemicals*, 5(1), 1.
30. Jabed, M. A., & Murad, M. A. A. (2024). Crop yield prediction in agriculture: A comprehensive review of machine learning and deep learning approaches, with insights for future research and sustainability. *Heliyon*, 10(24).
31. Jeppesen, J. H., Jacobsen, R. H., Jørgensen, R. N., & Toftegaard, T. S. (2022). Towards data-driven precision agriculture using open data and open source software. *arXiv preprint arXiv:2204.05582*.
32. Johnston, A. M., & Bruulsema, T. W. (2014). 4R nutrient stewardship for improved nutrient use efficiency. *Procedia Engineering*, 83, 365-370.
33. Juncal, M. J. L., Masino, P., Bertone, E., & Stewart, R. A. (2023). Towards nutrient neutrality: A review of agricultural runoff mitigation strategies and the development of a decision-making framework. *Science of the Total Environment*, 874, 162408.
34. Jwaideh, M. A., Sutanudjaja, E. H., & Dalin, C. (2022). Global impacts of nitrogen and phosphorus fertiliser use for major crops on aquatic biodiversity. *The International Journal of Life Cycle Assessment*, 27(8), 1058-1080.
35. Kamilaris, A., & Prenafeta-Boldú, F. X. (2018). Deep learning in agriculture: A survey. *Computers and electronics in agriculture*, 147, 70-90.
36. Kammerlander, C., Kolb, V., Luegmair, M., Scheermann, L., Schmailzl, M., Seufert, M., ... & Schön, T. (2025). Machine learning models for soil parameter prediction based on satellite, weather, clay and yield data. *arXiv preprint arXiv:2503.22276*.

37. Karlen, D. L., Veum, K. S., Sudduth, K. A., Obrycki, J. F., & Nunes, M. R. (2019). Soil health assessment: Past accomplishments, current activities, and future opportunities. *Soil and tillage research*, 195, 104365.
38. Kaushik, P., Yadav, S., Kumar, R., Kiran, K., Balakrishnan, M., & Abraham, K. (2025). AI-Driven Agriculture System for Crop Recommendation and Farmer Decision-Making Support. In *2025 Modern Electronics Devices and Intelligent Communication Systems (MEDCOM)* (pp. 1013-1017). IEEE.
39. Khaki, S., Wang, L., & Archontoulis, S. V. (2020). A CNN-RNN framework for crop yield prediction. *Frontiers in plant science*, 10, 1750.
40. Khambalkar, P. A., Agrawal, S., Dhaliwal, S. S., Yadav, S. S., Sadawarti, M. J., Singh, A., ... & Afreen, N. (2025). Sustainable nutrient management balancing soil health and food security for future generations. *Applied Food Research*, 5(2), 101087.
41. Kour, V. P., & Arora, S. (2020). Recent developments of the internet of things in agriculture: a survey. *Ieee Access*, 8, 129924-129957.
42. Lee, C. L., Strong, R., & Dooley, K. E. (2021). Analyzing precision agriculture adoption across the globe: A systematic review of scholarship from 1999–2020. *Sustainability*, 13(18), 10295.
43. Lee, S., Bae, J. H., Lee, G., Shin, H., Yang, J. E., Kim, J., & Lim, K. J. (2024). Predicting Nutrient Runoff Load in Cultivated Lands using Best Management Practices with Machine Learning Models. *Journal of Agricultural, Life and Environmental Sciences*, 36(2), 152-163.
44. Lehmann, J., Bossio, D. A., Kögel-Knabner, I., & Rillig, M. C. (2020). The concept and future prospects of soil health. *Nature Reviews Earth & Environment*, 1(10), 544-553.
45. Liakos, K. G., Busato, P., Moshou, D., Pearson, S., & Bochtis, D. (2018). Machine learning in agriculture: A review. *Sensors*, 18(8), 2674.
46. Liang, S., Ma, A., Zhang, T., Fan, X., Qian, D., Shao, J., ... & Xun, W. (2026). Soil organic matter enhances rhizosphere nutrient cycling by shaping bacterial life-history strategy and carbon metabolic preference. *Applied Soil Ecology*, 223, 107018.
47. Liu, J., Wang, H., Penuelas, J., Mou, J., Delgado-Baquerizo, M., Sardans, J., ... & Fang, L. (2025). Global-scale prevalence of low nutrient use efficiency across major crops. *Nature Communications*, 16(1), 11036.
48. Lyu, J., Wang, X., Hou, S., Zeb, A., Zhu, H., & Xu, Y. (2023). Content variation and potential runoff loss risk of nutrients in surface water of saline-alkali paddy in response to the application of different nitrogen fertilizer types. *Sustainability*, 15(9), 7040.
49. Maaz, T. M., Dobermann, A., Lyons, S. E., & Thomson, A. M. (2025). Review of research and innovation on novel fertilizers for crop nutrition. *npj Sustainable Agriculture*, 3(1), 25.
50. Mahmud, K., Panday, D., Mergoum, A., & Missaoui, A. (2021). Nitrogen losses and potential mitigation strategies for a sustainable agroecosystem. *Sustainability*, 13(4), 2400.
51. Mamatha, B., Mudigiri, C., Ramesh, G., Saidulu, P., Meenakshi, N., & Prasanna, C. L. (2024). Enhancing soil health and fertility management for sustainable agriculture: A review. *Asian J. Soil Sci. Plant Nutr*, 10(3), 182-190.
52. Miao, Y. (2023). Precision nutrient management. In *Encyclopedia of Digital Agricultural Technologies* (pp. 1054-1061). Cham: Springer International Publishing.

53. Miller, T., Mikiciuk, G., Durlik, I., Mikiciuk, M., Łobodzińska, A., & Śnieg, M. (2025). The IoT and AI in agriculture: The time is now—A systematic review of smart sensing technologies. *Sensors*, 25(12), 3583.
54. Mishra, H., & Mishra, D. (2024). AI for data-driven decision-making in smart agriculture: from field to farm management. In *Artificial intelligence techniques in smart agriculture* (pp. 173-193). Singapore: Springer Nature Singapore.
55. Mohan, R. N. V., Rayanoothala, P. S., & Sree, R. P. (2025). Next-gen agriculture: integrating AI and XAI for precision crop yield predictions. *Frontiers in Plant Science*, 15, 1451607.
56. Ngo, V. M., Le-Khac, N. A., & Kechadi, M. T. (2019, June). Designing and implementing data warehouse for agricultural big data. In *International conference on big data* (pp. 1-17). Cham: Springer International Publishing.
57. Ngo, V. M., Le-Khac, N. A., & Kechadi, M. T. (2020). Data warehouse and decision support on integrated crop big data. *International Journal of Business Process Integration and Management*, 10(1), 17-28.
58. Noulas, C., Torabian, S., & Qin, R. (2023). Crop nutrient requirements and advanced fertilizer management strategies. *Agronomy*, 13(8), 2017.
59. Padarian, J., & McBratney, A. B. (2020). A new model for intra-and inter-institutional soil data sharing. *Soil*, 6(1), 89-94.
60. Padarian, J., Minasny, B., & McBratney, A. B. (2020). Machine learning and soil sciences: A review aided by machine learning tools. *Soil*, 6(1), 35-52.
61. Pan, S. Y., He, K. H., Lin, K. T., Fan, C., & Chang, C. T. (2022). Addressing nitrogenous gases from croplands toward low-emission agriculture. *Npj Climate and Atmospheric Science*, 5(1), 43.
62. Pandey, U., Vashisth, A., Mishra, A., & Kumari, M. (2025). AI and IoT-driven soil health restoration: A machine learning approach for sustainable agriculture.
63. Pandey, V., Das, R., & Biswas, D. (2025). AgroSense: An Integrated Deep Learning System for Crop Recommendation via Soil Image Analysis and Nutrient Profiling. *arXiv preprint arXiv:2509.01344*.
64. Penuelas, J., Coello, F., & Sardans, J. (2023). A better use of fertilizers is needed for global food security and environmental sustainability. *Agriculture & Food Security*, 12(1), 5.
65. Pierre, J. F., & Regmi, N. (2026). Fertilizer form influences agronomic efficiency and environmental sustainability in nutrient management. *Discover Soil*, 3(1), 50.
66. Pincheira, M., Antonelli, F., & Vecchio, M. (2025). AGRICLIMA: Towards a Federated Platform for Spatiotemporal Risk Analysis in Agriculture. *Agriculture*, 15(23), 2450.
67. Reginald, P. J. (2024). Integrating IoT and AI for Precision Irrigation and Intelligent Nutrient Management in Smart Agriculture. *National Journal of Plant Sciences and Smart Horticulture*, 34-39.
68. Reisinger, A. J., Bean, E. Z., Clark, M., Levine, A. J., & Wilson, P. C. (2025). Fertilizer management approaches influence nutrient leaching from residential landscapes (Vol. 54, No. 1, pp. 289-302).
69. Rodrigues, G. C. (2022). Precision agriculture: Strategies and technology adoption. *Agriculture*, 12(9), 1474.
70. Rouzky, R., Sarrafzadeh, A., Sowell-Boone, E., Pasandi, H. B., & Goins, G. (2025). Data governance in smart farming: A systematic review of challenges and gaps. *Information Processing in Agriculture*.
71. San Emeterio de la Parte, M., Martínez-Ortega, J. F., Hernández Díaz, V., & Martínez, N. L. (2023). Big Data and precision agriculture: a novel spatio-temporal semantic IoT data management framework for improved interoperability. *Journal of Big Data*, 10(1), 52.

72. Shao, F., Tao, W., Wang, Q., Wu, J., Su, L., Sun, Y., & Zeng, S. (2021). A modified model for predicting nutrient loss in runoff using a time-varying mixing layer. *Journal of Hydrology*, 603, 127091.
73. Shrivastav, P., Prasad, M., Singh, T. B., Yadav, A., Goyal, D., Ali, A., & Dantu, P. K. (2020). Role of nutrients in plant growth and development. In *Contaminants in agriculture: Sources, impacts and management* (pp. 43-59). Cham: Springer International Publishing.
74. Soussi, A., Zero, E., Sacile, R., Trincherio, D., & Fossa, M. (2024). Smart sensors and smart data for precision agriculture: a review. *Sensors*, 24(8), 2647.
75. Talaviya, T., Shah, D., Patel, N., Yagnik, H., & Shah, M. (2020). Implementation of artificial intelligence in agriculture for optimisation of irrigation and application of pesticides and herbicides. *Artificial intelligence in agriculture*, 4, 58-73.
76. Vullaganti, N., Ram, B. G., & Sun, X. (2025). Precision agriculture technologies for soil site-specific nutrient management: A comprehensive review. *Artificial Intelligence in Agriculture*, 15(2), 147-161.
77. Wang, Q., & Wang, T. (2026). Machine learning-driven sensor fusion for precision agriculture: current applications and future perspectives. *Sensor Review*, 1-26.
78. Wang, Z., Xiao, S., Wang, J., Parab, A., & Patel, S. (2024). Intelligent Agricultural Management Considering N₂O Emission and Climate Variability with Uncertainties. *arXiv preprint arXiv:2402.08832*.
79. Wolfert, S., Ge, L., Verdouw, C., & Bogaardt, M. J. (2017). Big data in smart farming—a review. *Agricultural systems*, 153, 69-80.
80. Yanardağ, A. B. (2024). *Damage to Soil and Environment Caused by Excessive Fertilizer Use. Agriculture, Forestry and Aquaculture Sciences*. Ankara: Serüven Publishing, 203.
81. Yang, Y., Ge, X., Feng, W., Xian, L., Shao, G., Liao, J., & Zeng, S. (2025). Risk and driving factors of nitrogen and phosphorus runoff losses under extreme rainfall conditions: sludge application strategies in forest soils. *Catena*, 259, 109318.
82. Zheng, M., Zhu, P., Zheng, J., Xue, L., Zhu, Q., Cai, X., ... & Zhang, J. (2021). Effects of soil texture and nitrogen fertilisation on soil bacterial community structure and nitrogen uptake in flue-cured tobacco. *Scientific Reports*, 11(1), 22643.