

1. (a)

$$g = \left\langle \frac{\partial \varphi^{-1}}{\partial x}, \frac{\partial \varphi^{-1}}{\partial x} \right\rangle = \langle (1, f'(x)), (1, f'(x)) \rangle = 1 + (f'(x))^2$$

Thus the formula for arc length in this context is

$$\int_a^b \sqrt{1 + (f'(x))^2} dx$$

(b)

$$\begin{aligned} g = (\varphi^{-1})^* g_{\text{EUC}} &= \begin{bmatrix} \frac{\partial(\varphi^{-1})^x}{\partial x_1} & \frac{\partial(\varphi^{-1})^y}{\partial x_1} & \frac{\partial(\varphi^{-1})^z}{\partial x_1} \\ \frac{\partial(\varphi^{-1})^x}{\partial y_1} & \frac{\partial(\varphi^{-1})^y}{\partial y_1} & \frac{\partial(\varphi^{-1})^z}{\partial y_1} \end{bmatrix} \begin{bmatrix} \frac{\partial(\varphi^{-1})^x}{\partial x} & \frac{\partial(\varphi^{-1})^x}{\partial y} \\ \frac{\partial(\varphi^{-1})^y}{\partial x} & \frac{\partial(\varphi^{-1})^y}{\partial y} \\ \frac{\partial(\varphi^{-1})^z}{\partial x} & \frac{\partial(\varphi^{-1})^z}{\partial y} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & \frac{\partial f}{\partial x} \\ 0 & 1 & \frac{\partial f}{\partial y} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{\partial f}{\partial x}\right)^2 + 1 & \frac{\partial f}{\partial x} \frac{\partial f}{\partial y} \\ \frac{\partial f}{\partial x} \frac{\partial f}{\partial y} & \left(\frac{\partial f}{\partial y}\right)^2 + 1 \end{bmatrix} \end{aligned}$$

So that $\det(g_{ij}) = 1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2$. It follows that the surface area formula in this context simplifies to

$$\int_U \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dx dy$$

(c) The solution to this exercise relies on the crucial observation, that on $S \cap V$,

$$F(x, y, z) = 0 = z - f(x, y)$$

With this observation, the exercise is routine, first note that

$$\begin{aligned} \frac{\partial F}{\partial x} &= \frac{\partial}{\partial x}(z - f(x, y)) = -\frac{\partial f}{\partial x} \\ \frac{\partial F}{\partial y} &= \frac{\partial}{\partial y}(z - f(x, y)) = -\frac{\partial f}{\partial y} \\ \frac{\partial F}{\partial z} &= \frac{\partial}{\partial z}(z - f(x, y)) = 1 \end{aligned}$$

Now to verify the formula:

$$\begin{aligned} \int_U \frac{|\nabla F|}{\left|\frac{\partial F}{\partial z}\right|} dx dy &= \int_U \frac{\sqrt{\left(\frac{\partial F}{\partial x}\right)^2 + \left(\frac{\partial F}{\partial y}\right)^2 + \left(\frac{\partial F}{\partial z}\right)^2}}{\left|\frac{\partial F}{\partial z}\right|} dx dy \\ &= \int_U \sqrt{1 + \left(\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}}\right)^2 + \left(\frac{\frac{\partial F}{\partial y}}{\frac{\partial F}{\partial z}}\right)^2} dx dy \\ &= \int_U \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dx dy \end{aligned}$$

As in part (b). Now checking the formula:

$$\begin{aligned}
 \int_U \frac{|\nabla F|}{|\frac{\partial}{\partial z} F|} dx dy &= \int_U 2 \frac{\sqrt{x^2 + y^2 + z^2}}{2z} dx dy = \int_U \sqrt{1 + \frac{x^2}{z^2} + \frac{y^2}{z^2}} dx dy \\
 &= \int_U \sqrt{1 + \frac{x^2}{1 - x^2 - y^2} + \frac{y^2}{1 - x^2 - y^2}} dx dy \\
 &= \int_U \sqrt{\frac{1}{1 - x^2 - y^2}} dx dy \\
 &= \int_{y=-1}^1 \int_{x=-\sqrt{1-y^2}}^{\sqrt{1-y^2}} \frac{1}{\sqrt{1 - x^2 - y^2}} dx dy \\
 &= \int_{y=-1}^1 \arctan \frac{x}{\sqrt{1 - x^2 - y^2}} \Big|_{x=-\sqrt{1-y^2}}^{x=\sqrt{1-y^2}} dy \\
 &= \int_{y=-1}^1 \lim_{x \rightarrow \infty} \arctan(x) - \lim_{x \rightarrow -\infty} \arctan(x) dy \\
 &= \int_{-1}^1 \pi dy = 2\pi
 \end{aligned}$$

As desired.

2. (a) We compute

$$A = \begin{bmatrix} \frac{\partial}{\partial u} f^x & \frac{\partial}{\partial u} f^y & \frac{\partial}{\partial u} f^z \\ \frac{\partial}{\partial v} f^x & \frac{\partial}{\partial v} f^y & \frac{\partial}{\partial v} f^z \end{bmatrix} = \begin{bmatrix} -b \sin u \cos v & -b \sin u \sin v & b \cos u \\ -(a + b \cos u) \sin v & (a + b \cos u) \cos v & 0 \end{bmatrix}$$

Then in (u, v) coordinates, we have $g = f^* g_{\text{EUC}}$, i.e.

$$g = AA^T = \begin{bmatrix} b^2 & 0 \\ 0 & (a + b \cos u)^2 \end{bmatrix}$$

(b) Write $c_1 = f \circ \gamma_1$, $c_2 = f \circ \gamma_2$. Then we have $\dot{\gamma}_1^1 = 0$, $\dot{\gamma}_1^2 = 1$, so that $|\dot{\gamma}_1|_g^2 = g_{2,2} = (a + b \cos \pi)^2 = (a - b)^2$. It follows that

$$L(c_1) = \int_0^{2\pi} a - b dt = 2\pi(a - b)$$

Similarly, $\dot{\gamma}_2^1 = 1$, $\dot{\gamma}_2^2 = 0$, so that $|\dot{\gamma}_2|_g^2 = g_{1,1} = b^2$. It follows that

$$L(c_2) = \int_0^{2\pi} b = 2b\pi$$

(c) $dA = \sqrt{\det g} du dv = b(a + b \cos u) du dv$, hence the surface area is given by

$$SA = \int_0^{2\pi} \int_0^{2\pi} b(a + b \cos u) du dv = \int_0^{2\pi} uab + b^2 \sin u \Big|_0^{2\pi} dv = \int_0^{2\pi} 2\pi ab dv = (2\pi)^2 ab$$