

1 Classification of Semi-Simple Lie Algebras

1.1 Basic Examples

1.1.1 $\mathbf{A}_\ell \sim \mathfrak{sl}_{\ell+1}$

Definition 1.

$$\mathfrak{sl}_{\ell+1} = \mathfrak{gl}_{\ell+1} \cap \{\text{tr} = 0\} \quad (1)$$

The special linear lie algebra has dimension $(\ell + 1)^2 - 1$, an easy way to see this is to apply the submersion theorem to the determinant map on the lie group. A basis is given by $\{e_{i,j} \mid i \neq j\} \cup \{e_{ii} - e_{jj}\}$.

Remark 1. A thought that comes to mind is that defining a lie algebra using structure constants (with some level of care about prime characteristic) the dimension is preserved, so in some sense the manifold structure for the real lie algebra can tell us about the dimensionality over an arbitrary field.

1.1.2 $\mathbf{B}_\ell \sim \mathfrak{so}(2\ell + 1)$

Definition 2. Define the symmetric matrix $s = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & \mathbf{1}_\ell \\ 0 & \mathbf{1}_\ell & 0 \end{pmatrix}$ which defines a symmetric bilinear form

B on $k^{2\ell+1}$ defined as $B(v, w) = vsw$.

Definition 3. The naïve definition of $\mathfrak{so}(2\ell + 1)$ is to take the tangent space to $O(2\ell + 1)$, which gives us $\mathfrak{gl} \cap \{A^t = -A\}$. This is equivalent to skew symmetric matrices with respect to the symmetric bilinear form defined by $\mathbf{1}_{2\ell+1}$. The drawback of this is it contains no diagonal elements, so that in particular, the Cartan subalgebra is not easily understood. Instead we define $\mathfrak{so}(2\ell + 1)$ as skew symmetric matrices with respect to the binary form B given in **Definition 2**. Explicitly $B(\xi(v), w) = -B(v, \xi(w))$.

Proposition 1.1.1.

$$\mathfrak{so}(2\ell + 1) = \mathfrak{gl} \cap \{\xi \mid s\xi = -\xi^t s\} = \left\{ \begin{pmatrix} 0 & \mathbf{b}_1 & \mathbf{b}_2 \\ -\mathbf{b}_1^t & \mathbf{m} & \mathbf{n} \\ -\mathbf{b}_2^t & \mathbf{p} & -\mathbf{m}^t \end{pmatrix} \mid \mathbf{b}_1, \mathbf{b}_2, \mathbf{m}, \mathbf{n}, \mathbf{p}, -\mathbf{n}^t = \mathbf{n}, -\mathbf{p}^t = \mathbf{p} \right\} \quad (2)$$

Proof. It suffices to show that $\xi \in \mathfrak{so}(2\ell + 1) \iff s\xi = -\xi^t s$, since we read the third condition off of this one. If $\xi \in \mathfrak{so}(2\ell + 1)$, then for all $v, w \in k^{2\ell+1}$

$$v^t \xi^t s w = \xi(v)^t s w = -v^t s \xi(w) = -v^t s \xi w \quad (3)$$

Since this holds for arbitrary v, w it is only possible if $\xi^t s = -s\xi$. The converse is trivial. $\square$

The third equivalent form in **Proposition 1.1.1** lets us count the dimension of $\mathfrak{so}(2\ell + 1)$ by elementary means of 2ℓ for the $\mathbf{b}_j$, ℓ^2 for the $\mathbf{m}$ and $2 \cdot \frac{\ell(\ell-1)}{2}$ for $\mathbf{p}, \mathbf{n}$ by counting the strictly upper triangular entries, in total the dimension is $2\ell^2 + \ell$. This derivation gives us a basis.

1.1.3 $\mathbf{C}_\ell \sim \mathfrak{spl}(2\ell)$

Definition 4. Let B be a skew symmetric non-degenerate bilinear form on the finite dimensional vector space V , existence of such a form implies V has even dimension (**Proposition 4.0.2**), at the risk of circular reasoning, the geometry of this is such a form defines an almost complex structure. Then we can take some isomorphism $V \cong k^{2\ell}$

$$\mathfrak{spl}(2\ell) = \mathfrak{gl}(V) \cap \{\xi \mid B(\xi(v), w) = -B(v, \xi(w))\} \quad (4)$$

In practice we would like to define B by the skew symmetric matrix $s = \begin{pmatrix} 0 & \mathbf{1}_\ell \\ -\mathbf{1}_\ell & 0 \end{pmatrix}$. Following the proof of **Proposition 1.1.1** this is matrices ξ satisfying $-\xi^t s = s\xi$.

We can observe that

$$\mathfrak{spl}(2\ell) = \left\{ \begin{pmatrix} \mathbf{m} & \mathbf{p} \\ \mathbf{q} & -\mathbf{m}^t \end{pmatrix} \mid \mathbf{p} = \mathbf{p}^t, \mathbf{q} = \mathbf{q}^t \right\} \quad (5)$$

And the dimension is $2\ell^2 + \ell$

1.1.4 $\mathbf{D}_\ell \sim \mathfrak{so}(2\ell)$

Definition 5. Once again, we don't take the naive skew symmetric approach, and define the lie algebra to be the maps satisfying $B(\xi(v), w) = -B(v, \xi(w))$ where here B is defined by the matrix $s = \begin{pmatrix} 0 & \mathbf{1}_\ell \\ \mathbf{1}_\ell & 0 \end{pmatrix}$. Thus we find that

$$\mathfrak{so}(2\ell) = \{ \xi \in \mathfrak{gl}(2\ell) \mid -\xi^t s = s \xi \} = \left\{ \begin{pmatrix} \mathbf{m} & \mathbf{p} \\ \mathbf{q} & -\mathbf{m}^t \end{pmatrix} \mid \mathbf{p} = -\mathbf{p}^t, \mathbf{q} = \mathbf{q}^t \right\} \quad (6)$$

Once again it is easy to see the dimension is $2\ell^2 + \ell$ by using the explicit form and counting skew symmetric matrices.

1.2 Definitions and Major Theorems

Definition 6. An ideal of a Lie algebra is **TODO**

Definition 7. The center $\mathfrak{z}$ of a Lie algebra is **TODO**

Definition 8. A subalgebra of a Lie algebra is **TODO**

Definition 9. A Lie algebra homomorphism is **TODO**

Definition 10. A representation of a Lie algebra is **TODO**

Definition 11. The derived subalgebra of a Lie algebra is **TODO**

Definition 12. A Lie algebra is called solveable when **TODO**

Definition 13. The radical of a lie algebra is **TODO**

Definition 14. The normalizer $N_{\mathfrak{g}}(\mathfrak{h})$ of a subalgebra $\mathfrak{h}$ of a Lie algebra $\mathfrak{g}$ is **TODO**

Definition 15. A derivation $D : \mathfrak{g} \rightarrow \mathfrak{g}$ is an endomorphism satisfying

$$D[\xi, \eta] = [D(\xi), \eta] + [\xi, D(\eta)] \quad (7)$$

Definition 16. In characteristic zero, the exponential of a nilpotent derivation (i.e. $D^j = 0$ for some j) is the formal power series $\exp(D) = \sum_0^\infty D^j/j!$. By the Liebniz rule $\exp(D)$ is a derivation.

Definition 17. The killing form κ is a bilinear form on $\mathfrak{g}$ defined as

$$\kappa : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g} \quad (8)$$

$$(\xi, \eta) \mapsto \text{tr}(\text{ad}_\xi \circ \text{ad}_\eta) \quad (9)$$

Bilinearity follows from linearity of the arguments in the adjoint representation (i.e. bilinearity of lie bracket).

Definition 18. A bilinear form $B(-, -)$ on $\mathfrak{g}$ is called ad invariant, or simply invariant, when

$$B([x, \xi], \eta) + B(\xi, [x, \eta]) = 0 \quad (10)$$

for all x, ξ, η .

Definition 19. The Lie Algebra of inner derivations of $\mathfrak{g}$, denoted $\text{Inn}(\mathfrak{g})$ is given by

$$\text{Inn}(\mathfrak{g}) = \{ \text{ad}_\xi \mid \xi \in \mathfrak{g} \} \quad (11)$$

The Lie bracket is inhereted from $\text{Der}(\mathfrak{g})$, which is inhereted from $\mathfrak{gl}(\mathfrak{g})$

1.3 Basic Results on Nilpotence and Solveability

Proposition 1.3.1. 1. If $I \subset \mathfrak{g}$ is a solveable ideal, then $\mathfrak{g}/I$ is solveable iff $\mathfrak{g}$ is.

2. The sum of solveable ideals is solveable

Proof. 1. One direction is trivial. In the other direction let $\pi : \mathfrak{g} \rightarrow \mathfrak{g}/I$ be the quotient map. Then $\pi(\mathfrak{g}^{(n)}) = \pi(\mathfrak{g})^{(n)} = 0$ so that $\mathfrak{g}^{(n)} \subset I$, this is obviously sufficient.

2. $I + J/J \cong I/I \cap J$ is solveable, apply part (i). □

Proposition 1.3.2. 1. If $\mathfrak{g}/\mathfrak{z}$ is nilpotent then so is $\mathfrak{g}$

2. A nonzero nilpotent Lie algebra has nontrivial center

Proof. 1. Let π be the quotient map, then $\pi([\mathfrak{g}^n, g]) = [\pi(\mathfrak{g}^n), \pi(g)] = \pi(\mathfrak{g})^{n+1}$. Thus $\mathfrak{g}^{n+1} \in \mathfrak{z}$. Thus $\mathfrak{g}^{n+2} = 0$

2. Suppose n is the least n such that $\mathfrak{g}^n = 0$, then $n > 0$ and $\mathfrak{g}^{n-1} \in \mathfrak{z}$. □

Proposition 1.3.3. A subalgebra $\mathfrak{g} \subset \mathfrak{gl}(V)$ of nilpotent endomorphisms has a common eigenvector.

Proof. Induct on dimension of the lie algebra. $\dim = 1$ is trivial since k contains all of the roots of the characteristic polynomial.

Now let $\mathfrak{h} \subset \mathfrak{g}$ be a proper subalgebra, $\mathfrak{h}$ is thus also a nilpotent subalgebra of $\mathfrak{gl}(V)$. Then we consider the vectorspace $U = \mathfrak{g}/\mathfrak{h}$, the adjoint representation of $\mathfrak{h}$ on U is nilpotent, thus by induction has a common eigenvector. This shows that $N_{\mathfrak{g}}(\mathfrak{h}) \supsetneq \mathfrak{h}$.

Every solveable lie algebra has a codimension 1 subalgebra, to see this take $\pi : \mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{g}'$, then $\mathfrak{g}/\mathfrak{g}'$ is abelian so we can take a codimension 1 subalgebra $\bar{\mathfrak{h}}$, then $\mathfrak{h} := \pi^{-1}(\bar{\mathfrak{h}})$ is a codimension 1 subalgebra of $\mathfrak{g}$.

Combining the preceding two arguments we have $\mathfrak{h} \subset \mathfrak{g}$ with $N_{\mathfrak{g}}(\mathfrak{h}) = \mathfrak{g}$ so that $\mathfrak{h}$ is a codimension 1 ideal. We apply the induction hypothesis, i.e. let $W \subset V$ be the nonzero subspace of eigenvectors for $\mathfrak{h}$, since $\mathfrak{h}$ is an ideal we have for $w \in W$, $\xi \in \mathfrak{g}$, $\eta \in \mathfrak{h}$

$$\eta\xi(w) = \xi\eta(w) + [\eta, \xi](w) = 0 \quad (12)$$

So that $\mathfrak{g} : W \rightarrow W$, now writing $\mathfrak{g} = \mathfrak{h} + k\xi$, we know the nilpotent endomorphism $\xi : W \rightarrow W$ has an eigenvector, this is already a common eigenvector for $\mathfrak{h}$. □

Theorem 1 (Engel). If every element of $\mathfrak{g}$ is ad-nilpotent, then $\mathfrak{g}$ is nilpotent.

Proof. Consider $\text{ad}(\mathfrak{g}) \subset \mathfrak{gl}(\mathfrak{g})$, by **Proposition 1.3.3** $\mathfrak{z}(\mathfrak{g}) \neq 0$, so that $\dim \mathfrak{g}^1 < \dim \mathfrak{g}$. Now proceeding by induction the ad action of any element of $\mathfrak{g}$ on $\mathfrak{g}^n$ is still nilpotent for any n , thus $\mathfrak{g}^n \cap \mathfrak{z} \neq 0$ and $\dim \mathfrak{g}^{n+1} < \dim \mathfrak{g}^n$. Since $\mathfrak{g}$ is finite dimensional it must be nilpotent. □

Proposition 1.3.4. Let V be a finite dimensional vectorspace and $\mathfrak{g} \subset \text{End}(V)$ a subalgebra of nilpotent transformations, then there is a basis for V in which every matrix in $\mathfrak{g}$ is upper triangular

Proof. Start with a common eigenvector $v \in V$, then take the second basis vector to be an eigenvector of $V/\text{span}(v)$, continuing this process gives a basis. With $\mathfrak{g}$ upper triangular. □

1.4 The Killing Form

See Definition 17.

Proposition 1.4.1. The killing form is a symmetric bilinear form.

Proof. tr is symmetric. □

Proposition 1.4.2. κ is automorphism invariant. I.e. Let $\phi \in \text{Aut}(\mathfrak{g})$, then

$$\kappa(\phi(\xi), \phi(\eta)) = \kappa(\xi, \eta) \quad (13)$$

Proof.

$$\text{ad}_{\phi(\xi)} = \phi \circ \text{ad}_\xi \circ \phi^{-1} \quad (14)$$

and trace is basis independent. □

Proposition 1.4.3. The killing form is an invariant binary form See Definition 18. More generally it is invariant (in the same sense) under all derivations.

Proof. Choose a path in $\text{Aut}(\mathfrak{g})$ which differentiates to the desired inner derivation. Then differentiate equation for automorphism invariance using proposition 2.2.1, apply bilinearity of the killing form to take the derivative. □

A second proof is much less elegant, but proves the result for general derivations.

Proof. By the definition of derivations,

$$\text{ad}_{D(x)} = D \circ \text{ad}_x - \text{ad}_x \circ D \quad (15)$$

Thus we find that

$$\kappa(D(\xi), \eta) = \text{tr}(\text{ad}_{D(\xi)} \circ \text{ad}_\eta) \quad (16)$$

$$\stackrel{(15)}{=} \text{tr}(D \circ \text{ad}_\xi \circ \text{ad}_\eta) - \text{tr}(\text{ad}_\xi \circ D \circ \text{ad}_\eta) \quad (17)$$

$$= -(\text{tr}(\text{ad}_\xi \circ D \circ \eta) - \text{tr}(\text{ad}_\xi \circ \text{ad}_\eta \circ D)) \quad (18)$$

$$= \text{tr}(\text{ad}_\xi \circ (D \circ \text{ad}_\eta - \text{ad}_\eta \circ D)) \quad (19)$$

$$\stackrel{(15)}{=} -\text{tr}(\text{ad}_\xi \circ \text{ad}_{D(\eta)}) = -\kappa(\xi, D(\eta)) \quad (20)$$

Where (18) follows from cyclicity of trace. □

Proposition 1.4.4. If the killing form κ of $\mathfrak{g}$ is nondegenerate, then every derivation of $\mathfrak{g}$ is inner. I.e.

$$\text{Der}(\mathfrak{g}) = \text{Inn}(\mathfrak{g}) \quad (21)$$

Proof. Since κ is nondegenerate, the map $\mathfrak{g} \rightarrow \mathfrak{g}^*$ via $\xi \mapsto \kappa(\xi, -)$ is an isomorphism. Now let D be a derivation, we define the functional

$$\eta \mapsto \text{tr}(D \circ \text{ad}_\eta) \quad (22)$$

which must be equal to $\eta \mapsto \kappa(\xi, \eta)$ for some ξ . Fixing $x \in \mathfrak{g}$ we find that

$$\eta \mapsto \kappa((D - \text{ad}_\xi)(x), \eta) = \kappa(D(x), \eta) - \kappa(\text{ad}_\xi(x), \eta) \quad (23)$$

$$= \text{tr}(\text{ad}_{D(x)} \circ \text{ad}_\eta) - \text{tr}(\text{ad}_{\text{ad}_\xi(x)} \circ \text{ad}_\eta) \quad (24)$$

$$\stackrel{(15)}{=} \text{tr}([D, \text{ad}_x] \circ \text{ad}_\eta) - \text{tr}([\text{ad}_\xi, \text{ad}_x] \circ \text{ad}_\eta) \quad (25)$$

$$= \text{tr}(D \circ [\text{ad}_x, \text{ad}_\eta]) - \text{tr}(\text{ad}_\xi \circ [\text{ad}_x, \text{ad}_\eta]) \quad (26)$$

$$\stackrel{(15)}{=} \text{tr}(D \circ \text{ad}_{\text{ad}_x(\eta)}) - \kappa(\xi, \text{ad}_x(\eta)) \quad (27)$$

$$= \text{tr}(0) = 0 \quad (28)$$

Where (26) follows from cyclicity of trace. □

2 Lie Groups

2.1 General Theory

Theorem 2. A closed subgroup of a Lie Group is a Lie Group

2.2 Lie Groups For studying Lie Algebras

Proposition 2.2.1. $\text{Aut}(\mathfrak{g})$ is a Lie group, with Lie algebra $\text{Inn}(\mathfrak{g})$.

Proof. $\text{GL}(\mathfrak{g})$ is certainly a Lie group, and $\text{Aut}(\mathfrak{g})$ is a subgroup which is closed by continuity of $[-, -]$ so **Theorem 2** applies. By bilinearity, differentiation of the equation if $\phi(t)$ is a path in $\text{Aut}(\mathfrak{g})$ with $\phi(0) = \text{id}$ then

$$\left. \frac{d}{dt} \right|_{t=0} \phi(t)([\xi, \eta]) = \left. \frac{d}{dt} \right|_{t=0} [\phi(t)(\xi), \phi(t)(\eta)] \quad (29)$$

$$= \left[\left. \frac{d}{dt} \right|_{t=0} \phi(t)(\xi), \eta \right] + \left[\xi, \left. \frac{d}{dt} \right|_{t=0} \phi(t)(\eta) \right] \quad (30)$$

□

The converse direction follows since the automorphism equation integrates the derivation equation.

3 Exercises

3.1 Humphreys Ch1.

3.1.1. Compute the structure constants of the cross product on $\mathbb{R}^3$

Proof. $a_{12}^3 = 1, a_{31}^2 = 1, a_{23}^1 = 1$, else zero. □

3.1.2. Verify that the bilinearity and "antisymmetry" axioms alongside

$$[x, y] = z, [x, z] = y, [y, z] = 0 \quad (31)$$

define a Lie algebra.

Proof. It suffices to check the Jacobi equations

$$[x, [y, z]] - [y, [x, z]] + [z, [x, y]] = 0 - [y, y] + [z, z] = 0 \quad (32)$$

□

3.1.3. Let $e, h, f = e_{12}, e_{11} - e_{22}, e_{21}$ be the standard basis of $\mathfrak{sl}(2)$, compute the matrices ad_-

Proof.

$$\text{ad}_e = -2e_{12} + e_{23} \quad (33)$$

$$\text{ad}_f = -e_{21} + 2e_{32} \quad (34)$$

$$\text{ad}_h = 2e_{11} - 2e_{33} \quad (35)$$

□

3.1.4. Find a linear lie algebra $\mathfrak{g}$ with $\mathfrak{g} = \text{span}_k \{x, y\}$ and $[x, y] = x$.

Proof. We exploit the symmetry of the adjoint representation $\text{ad}_{[x,y]} = [\text{ad}_x, \text{ad}_y]$, and take x, y to their adjoint representations. This works since $\text{Inn}(\mathfrak{g})$ is also a 2 dimensional lie algebra.

$$x = e_{12}, y = e_{11} \quad (36)$$

□

3.1.5. Compute the dimensions of the diagonal, upper triangular and strictly upper triangular lie algebras. Verify that $[\mathfrak{d}, \mathfrak{n}] = \mathfrak{n}$.

Proof. The dimensiona are $n, \frac{n(n+1)}{2}, \frac{n(n-1)}{2}$. The bracket condition follows from $[e_{jj}, e_{k,r}] = \delta_{jk}e_{jr} - \delta_{pr}e_{kj}$. But by assuming strictly upper triangular $j = k$ implies $r > j$ and $r = j$ implies $j > k$. □

3.1.6. Suppose $x \in \mathfrak{gl}(n, k)$ has n distinct eigenvalues a_j . Show that the eigenvalues of ad_x are $a_i - a_j$.

Proof. Work in the basis of eigenvectors for x , then

$$\text{ad}_x(e_{ij})(e_k) = \begin{cases} 0 & k \neq j \\ x(e_i) - e_{ij}(a_j e_j) = (a_i - a_j)e_i & k = j \end{cases} \quad (37)$$

Since the e_{ij} form a basis, each is an eigenvector with eigenvalue $a_i - a_j$. □

3.1.7. Show that $\mathfrak{gl}(n, k) = \mathfrak{sl}(n, k) + \text{span}_k \{\mathbf{1}_n\}$ where $\text{char } k$ is not a prime dividing n

Proof. Let $\xi \in \mathfrak{gl}(n)$, then $\xi - n^{-1} \text{tr}(\xi) \mathbf{1}_n \in \mathfrak{sl}(n)$ □

3.1.8. Verify the dimension of $\mathbf{D}_\ell$ is $\ell^2 + \ell$

Proof. See definition 5. □

3.1.9. Show that in characteristic zero, the classical lie algebras are equal to their derived subalgebras.

Proof. Notice that by defining the orthogonal algebras as we did, all of the classical lie algebras have diagonal elements. Thus $\text{tr} : \mathfrak{g} \times \mathfrak{g} \rightarrow k$ by $\xi, \eta \mapsto \text{tr}(\xi\eta)$ is a nondegenerate bilinear form. Moreover by cyclicity it is an invariant bilinear form. This tells us that

$$\text{tr}([\mathfrak{g}, \mathfrak{g}], -) = \text{tr}(\mathfrak{g}, [\mathfrak{g}, -]) \quad (38)$$

Thus $\xi \in [\mathfrak{g}, \mathfrak{g}]^\perp \implies \xi \in \mathfrak{z}(\mathfrak{g})$. Semisimple lie algebras have trivial center so we are done. □

3.1.10. Exhibit the exceptional isomorphisms $A_1 \cong B_1 \cong C_1$, show D_1 has dimension 1, show $B_2 \cong C_2$, $D_3 \cong A_3$. What about D_2 ?

Proof. $\mathbf{A}_1 \cong \mathbf{B}_1$. Consider $\text{ad} : \mathfrak{sl}(2) \rightarrow \mathfrak{gl}(3)$, on $\mathfrak{sl}(2)$, tr is a nondegenerate symmetric bilinear form, moreover, cyclicity gives us $\text{tr}(\text{ad}_\xi(x), y) = -\text{tr}(x, \text{ad}_\xi(y))$. Thus by equivalence of bilinear forms (we are using here the field is algebraically closed), we get $\text{ad}(\mathfrak{sl}(2)) \subset \mathfrak{so}(3)$. Note that this proof fails due to the killing form on $\mathfrak{sl}(2, \mathbb{R})$ not being positive definite.

$\mathbf{A}_1 \cong \mathbf{C}_1$. $-\xi^t s = s\xi$ if and only if $\text{tr}(\xi) = 0$

$\mathbf{D}_1 \cong \mathbf{U}(1)$ By direct computation $D_1 = \left\{ \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix} \right\}$

TODO: Prove the remaining exceptional isomorphisms □

3.1.11. Verify the commutator of two derivations is a derivation, show that the composition of derivations on an algebra need not be one.

Proof.

$$[D, E][\xi, \eta] = D([E\xi, \eta] + [\xi, E\eta]) - E([D\xi, \eta] + [\xi, D\eta]) \quad (39)$$

$$= [DE\xi, \eta] + [E\xi, D\eta] + [D\xi, E\eta] + [\xi, DE\eta] - ([E\xi, D\eta] + [D\xi, E\eta]) - [ED\xi, \eta] - [\xi, ED\eta] \quad (40)$$

$$= [[D, E]\xi, \eta] + [\xi, [D, E]\eta] \quad (41)$$

Take the example of the formal derivative on $k[X]$. □

3.1.12. Show that $\{\eta \in \mathfrak{g} \mid \exists \lambda \in k, \text{ad}_\xi(\eta) = \lambda\eta\} \subset \mathfrak{g}$ is a subalgebra.

Proof. Apply the Jacobi identity to get

$$\text{ad}_\xi[\eta, x] = [\eta, \text{ad}_\xi x] + [x, -\text{ad}_\xi \eta] \quad (42)$$

$$= [\eta, \lambda_x x] + [x, -\lambda_\eta \eta] \quad (43)$$

$$= (\lambda_x + \lambda_\eta)[\eta, x] \quad (44)$$

□

3.2 Humphreys Ch2.

3.2.1. Show that $\text{Inn}(\mathfrak{g}) \subset \text{Der}(\mathfrak{g})$ is an ideal

Proof. Just compute

$$[\text{ad}_\xi, D](\eta) = [x, D(\xi)] - D([x, \xi]) \quad (45)$$

$$= [x, D(\xi)] - [x, D(\xi)] - [D(x), \xi] \quad (46)$$

$$= \text{ad}_{-D(x)}(\xi) \quad (47)$$

□

3.2.2. Show that $\mathfrak{sl}(n) = [\mathfrak{gl}(n), \mathfrak{gl}(n)]$

Proof. We have shown in Humphreys 1.9 that $\mathfrak{sl}$ is its own derived subalgebra, thus $\mathfrak{sl} \subset [\mathfrak{gl}, \mathfrak{gl}]$. Now notice by symmetry and bilinearity of trace, $\text{tr}([\xi, \eta]) = 0$ for any ξ, η . Thus $[\mathfrak{gl}, \mathfrak{gl}] \subset \mathfrak{sl}$. □

3.2.3. Show the center of $\mathfrak{gl}(n)$ is scalar matrices, show the center of $\mathfrak{sl}(n)$ is trivial, but in characteristic $p \mid n$ is scalar matrices.

Proof. Suppose that $\xi \in \mathfrak{z}(\mathfrak{gl})$, then

$$0 = [\xi, e_{ij}] = \sum_k \xi_{ki} e_{kj} - \xi_{jk} e_{ik} \quad (48)$$

□

We read from this equation that ξ is diagonal. Then $\xi = \lambda^k e_{kk}$ so that

$$0 = [\lambda^k e_{kk}, e_{ij}] = \lambda_i e_{ij} - \lambda_j e_{ij} \implies \lambda_i = \lambda_j \quad (49)$$

so that ξ is a scalar.

3.2.4. Show there is a unique isomorphism class of lie algebra $\mathfrak{g}$ with dimension 3 and $\mathfrak{g}' = [\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{z}(\mathfrak{g})$, and $\dim \mathfrak{g}' = 1$.

Proof. Since the derived subalgebra is one dimensional, there must be two elements x, y with $[x, y] = \mathfrak{g}'$. Then $\{x, y, [x, y]\}$ is a basis and the structure constants are fixed by $[x, y] = \mathfrak{z}$. □

3.2.5. Show that $\dim \mathfrak{g} = 3$ and $\mathfrak{g}' = \mathfrak{g}$ implies $\mathfrak{g}$ is simple.

Proof. Assume that $\mathfrak{g}$ is not simple, then a nontrivial ideal implies a homomorphism $\phi : \mathfrak{g} \rightarrow L$ for $\dim L = 1, 2$. If $\dim L = 1$ then $\phi(\mathfrak{g})$ is abelian, thus $0 = \phi(\mathfrak{g}') = \phi(\mathfrak{g})$ a contradiction. If $\dim L = 2$, then $L \cong \text{span}\{x, y\}$ with $[x, y] = x$, thus $\text{span}\{x\} = \phi(\mathfrak{g}') = \phi(\mathfrak{g}) = \text{span}\{x, y\}$ a contradiction. $\square$

3.2.6. Show $\mathfrak{sl}(3)$ is simple unless characteristic $k = 3$.

Proof. We work with the basis

$$h_1, h_2, e_1, e_2, f_1, f_2, [e_1, e_2], [f_1, f_2] \quad (50)$$

Since $\mathfrak{sl}(2)$ has no ideals, any ideal must contain a copy of either $\mathfrak{sl}(2)$ and thus also the bracket terms. But then $[e_{21}, e_{13}] = e_{23}$ so any ideal containing the first copy of $\mathfrak{sl}$ must contain an element of the second and we are done, conclude by symmetry. $\square$

3.2.7. Show that strictly upper triangular and diagonal matrices are self normalizing in $\mathfrak{gl}$, show that upper triangular matrices are normalized by strictly upper triangular matrices.

Proof. **TODO** $\square$

3.2.8. If k is a field of characteristic zero then the diagonal matrices are self normalizing subalgebras of the classical lie algebras.

Proof. The fact they are subalgebras is obvious. To see that they are self normalizing we use that for every non diagonal entry, there is a matrix in each classical lie algebra having it nonzero. Now supposing that $\xi_{ii} \neq \xi_{jj}$ for ξ diagonal (we can always do this since $\text{tr}(\xi) \neq 0$), the equation (49) gives bracketing this matrix with ξ gives a matrix with nonzero off diagonal coordinates. $\square$

3.2.9. Prove the isomorphism theorems

Proof. 1. It is clear that $\bar{\phi}$ must satisfy $\bar{\phi}(x + \ker) = \phi(x)$, so it suffices to check that such a map is a lie algebra homomorphism.

$$\bar{\phi}[x + \ker, y + \ker] = \bar{\phi}([x, y] + [\ker, y] + [x, \ker] + [\ker, \ker]) = \bar{\phi}([x, y] + \ker) = \phi[x, y] \quad (51)$$

$$[\bar{\phi}(x + \ker), \bar{\phi}(y + \ker)] = [\phi(x), \phi(y)] = \phi([x, y]) \quad (52)$$

injectivity and surjectivity are immediate.

2. Take the map $\phi : \mathfrak{g}/\mathfrak{k} \rightarrow \mathfrak{g}/\mathfrak{h}$ via $x + \mathfrak{k} \mapsto x + \mathfrak{h}$ this is a well defined homomorphism with kernel $\mathfrak{h}/\mathfrak{k}$. Apply the first isomorphism theorem.

3. There's nothing new in the third theorem either... $\square$

3.2.10. for $g \in GL(n)$ consider $\mathfrak{sl}(n) \rightarrow \mathfrak{sl}(n)$ via $x \mapsto -gx^t g^{-1}$, show this is an automorphism of $\mathfrak{sl}(n)$. When $n = 2$ and $g = 1$ show this is an inner automorphism.

Proof. **TODO** $\square$

3.2.11. Prove that for $g \in O(n)$ the map $x \mapsto gxg^{-1}$ defines an automorphism of an orthogonal lie algebra.

Proof. **TODO** $\square$

3.3 Humphrey's Ch3.

3.3.1. Let $I \subset \mathfrak{g}$ be an ideal, prove that $I^{(n)}$ and I^n are ideals for all n .

4 Appendix

Proposition 4.0.1. All non-degenerate skew symmetric Bilinear forms are equivalent.

Over an algebraically closed field, all non-degenerate symmetric binary forms are equivalent. The obstruction is existence of nonsquares (i.e. over $\mathbb{R}$ we classify which signs we act positively or negatively, over $\mathbb{Q}$ there are many ways forms can be non equivalent).

Proposition 4.0.2. A nondegenerate skew symmetric bilinear form can only exist in even dimensions.

Proof. Assume that $\dim V = 2\ell + 1$ has a nondegenerate bilinear form B . Then by induction and graham schmidt, we get a basis for a 2ℓ dimensional subspace $e_1, \dots, e_\ell, d_1, \dots, d_\ell$ with $B(e_j, d_j) = 1$ and all other products zero. Then there is another linearly independent element f , define f' as follows

$$f' = f + \sum_j B(f, e_j)d_j - B(f, d_j)e_j \quad (53)$$

Then $B(f', -) = 0$ so that $f' = 0$ contradicting V being $2\ell + 1$ dimensional. Construction of the basis from induction follows a similar technique so I won't make it explicit. $\square$