

1. We first find an expression for $\zeta(s)\Gamma(s)$ on $\Re(s) > 1$, to do so we will do a substitution $u = x/n$, valid for any $n \in \mathbb{Z}_{>0}$

$$\Gamma(s) = \int_0^\infty e^{-nu} (nu)^{s-1} n du = n^s \int_0^\infty u^{s-1} (e^{-u})^n du$$

Now since I don't like u I will switch back to x ; Multiplying n^{-s} on both sides and summing over $n \in \mathbb{Z}_{>0}$ yields

$$\zeta(s)\Gamma(s) = \sum_1^\infty \int_0^\infty x^{s-1} (e^{-x})^n dx \stackrel{\text{DCT}}{=} \int_0^\infty x^{s-1} \sum_1^\infty (e^{-x})^n dx = \int_0^\infty x^{s-1} \frac{1}{e^x - 1} dx \quad (1)$$

Where DCT is taken with respect to $|x^{s-1}| \sum_1^\infty e^{-nx}$. We can use $\Gamma(s)\Gamma(1-s) = \frac{\pi}{\sin(\pi s)}$ to rewrite (1):

$$\zeta(s) = \frac{\Gamma(1-s)\sin(\pi s)}{\pi} \int_0^\infty \frac{x^{s-1}}{e^x - 1} dx \quad (2)$$

Now we deal with the contour integral, letting $C(\epsilon)$ denote the curve described in the problem for fixed $\epsilon \in \mathbb{R}_{>0}$. The expression $e^z - 1$ has no poles away from $\{z \mid e^z = 1\} = \{2\pi ki \mid k \in \mathbb{Z}\}$, whence if we take the branch cut of \log away from the non-negative reals the integral is not dependent on ϵ for $\epsilon < 2\pi$, since (for $\delta < \epsilon < 2\pi$) we have the area enclosed between the two curves is a quotient of holomorphic functions with the denominator non-vanishing in between the curves. This independence is a result of taking separating the curve $C(\epsilon) - C(\delta)$ into two curves (see picture), the first of which has integral zero by Cauchy's theorem, and the second being arbitrarily small depending on where we take the cut.

$$\int_{C(\epsilon)} f - \int_{C(\delta)} f = \int_{C(\epsilon, \delta, M)} f + \int_{\gamma_1(\epsilon, \delta, M)} f + \int_{\gamma_2(\epsilon, \delta, M)} f$$

INSERT ILLUSTRATION HERE

Then for f holomorphic away from the real line, the $C(\epsilon, \delta, M)$ term vanishes. Notice now that using the standard arclength inequality for large M we have

$$\begin{aligned} \left| \int_{\gamma_j(\epsilon, \delta, M)} \frac{(-z)^{s-1}}{e^z - 1} dz \right| &\leq (\epsilon - \delta) \left| \frac{M^{\Re(s)-1}}{e^M - 1} \right| + \left| \int_M^\infty \frac{(-x - i\epsilon)^{s-1}}{e^{x+i\epsilon} - 1} dx - \int_M^\infty \frac{(-x - i\delta)^{s-1}}{e^{x+i\delta} - 1} dx \right| \\ &\leq (\epsilon - \delta) \left| \frac{M^{\Re(s)-1}}{e^M - 1} \right| + 2 \int_M^\infty \frac{|x + i\epsilon|^{\Re(s)-1}}{|e^x| - 1} dx \end{aligned}$$

The right hand side clearly converges to zero. as $M \rightarrow \infty$ using basic limits of exponentials and DCT. This gives the desired invariance.

$$\left| \int_{C(\epsilon)} \frac{(-z)^{s-1}}{e^z - 1} dz - \int_{C(\delta)} \frac{(-z)^{s-1}}{e^z - 1} dz \right| \leq \left| \int_{\gamma_1(\epsilon, \delta, M)} \frac{(-z)^{s-1}}{e^z - 1} dz \right| + \left| \int_{\gamma_2(\epsilon, \delta, M)} \frac{(-z)^{s-1}}{e^z - 1} dz \right| = 0$$

Now, we can compute the value of the integral along this curve by letting $\epsilon \rightarrow 0$, to get $C(0)$, a ray from ∞ to 0 where $(-z)^{s-1} = x^{s-1}e^{-(s-1)\pi i}$, and a ray from 0 to ∞ where $(-z)^{s-1} = x^{s-1}e^{(s-1)\pi i}$, to see that we can indeed pass to this limit, once again decompose $C = C(\epsilon)$ into three curves, with C_1 the ray in the upper half plane, C_2 the ray in the lower half plane and C_3 the circular portion, then once again using the arc length inequality and the fact that $e^z - 1 = \mathcal{O}(z)$

$$\left| \int_{C_3} \frac{(-z)^{s-1}}{e^z - 1} dz \right| \leq 2\pi\epsilon \sup_{|z|=\epsilon} \frac{|z^{s-1}|}{|e^z - 1|} = 2\pi\epsilon \mathcal{O}(\epsilon^{\Re(s)-2}) = 2\pi\mathcal{O}(\epsilon^{\Re(s)-1}) \xrightarrow{\Re(s)>1} 0$$

We can check convergence on C_1 explicitly

$$\begin{aligned} & \left| \int_0^\infty \frac{x^{s-1} e^{-(s-1)\pi i}}{e^x - 1} dx - \int_\epsilon^\infty \frac{(-x - i\epsilon)^{s-1}}{e^{x+i\epsilon} - 1} dx \right| \\ & \leq \left| \int_0^\epsilon \frac{x^{s-1} e^{-(s-1)\pi i}}{e^x - 1} dx \right| + \left| \int_\epsilon^\infty \frac{x^{s-1} e^{-(s-1)\pi i}}{e^x - 1} dx - \int_\epsilon^\infty \frac{(-x - i\epsilon)^{s-1}}{e^{x+i\epsilon} - 1} dx \right| \\ & \leq \left| \int_0^\epsilon \frac{x^{s-1} e^{-(s-1)\pi i}}{e^x - 1} dx \right| + \left| \int_\epsilon^\infty \frac{(e^{x+i\epsilon} - 1)x^{s-1} e^{-(s-1)\pi i} - (e^x - 1)(\sqrt{x^2 + \epsilon^2})^{s-1} e^{i(s-1) \arctan \frac{\epsilon}{x} - (s-1)\pi i}}{(e^x - 1)(e^{x+i\epsilon} - 1)} dx \right| \end{aligned}$$

Convergence as $\epsilon \rightarrow 0$ of the big ugly term to zero is actually simple from convergence of each of the terms in the two expressions in the product. Convergence of the first term follows from $\Re(s) > 1$, so writing the bounds of integration as $\chi_{(0,\epsilon)}$ we can just apply DCT to the absolute value of the integrand. The proof of convergence for C_2 is similar to C_1 .

Now we finally established that (due to taking the limit in ϵ and invariance with respect to ϵ)

$$\int_C \frac{(-z)^{s-1}}{e^z - 1} dz = - \int_0^\infty \frac{x^{s-1} e^{-(s-1)\pi i}}{e^x - 1} dx + \int_0^\infty \frac{x^{s-1} e^{(s-1)\pi i}}{e^x - 1} dx \quad (3)$$

$$= \int_0^\infty \frac{x^{s-1}}{e^x - 1} 2i \sin((s-1)\pi) = - \int_0^\infty \frac{x^{s-1}}{e^x - 1} 2i \sin(s\pi) \quad (4)$$

Multiplying (2) by $1 = \frac{\int_C \frac{(-z)^{s-1}}{e^z - 1} dz}{-\int_0^\infty \frac{x^{s-1}}{e^x - 1} 2i \sin(s\pi)}$ yields the desired equality

$$\zeta(s) = - \frac{\Gamma(1-s)}{2\pi i} \int_C \frac{(-z)^{s-1}}{e^z - 1} dz \quad (5)$$

Now, using the right side of (5) as an analytic continuation, we first compute $(e^z - 1)^{-1} = \frac{1}{z} - \frac{1}{2} + \frac{z}{12} + \mathcal{O}(z^2)$, since we know it has a simple pole at zero, hence is meromorphic this expression just comes from evaluating the systems of equations given for the coefficients for $(a_{-1}z^{-1} + a_0 + a_1z + \dots)(\sum_0^\infty \frac{z^n}{n!}) = 1$. This is everything we need to evaluate $\zeta(0)$. First using a similar separation of curves into two parts as was used for invariance, we find that $\int_C \frac{1}{z(e^z - 1)} dz = \int_\gamma \frac{1}{z(e^z - 1)} dz$ where γ is a piecewise C^1 closed curve, this allows us to use the residue theorem

$$\zeta(0) = \frac{\Gamma(1)}{2\pi i} \int_\gamma \frac{1}{z(e^z - 1)} dz = \frac{1}{2\pi i} \int_\gamma \frac{1}{z} \left(\frac{1}{z} - \frac{1}{2} + \frac{z}{12} + \mathcal{O}(z^2) \right) dz \quad (6)$$

$$= \text{Res} \left(\frac{1}{z} \left(\frac{1}{z} - \frac{1}{2} + \frac{z}{12} + \mathcal{O}(z^2) \right) \right) = -\frac{1}{2} \quad (7)$$

□

2. (a)

(b) Let $z \in \mathbb{C}$, then by **Insert EQ number here ...**

$$f'(z) = \frac{1}{2\pi i} \int_{|w-z|=R} \frac{f(w)}{(w-z)^2} dz$$

Applying the standard arc-length inequality yields

$$|f'(z)| \leq \frac{1}{2\pi} 2\pi R \frac{\sup_{\mathbb{C}} |f|}{R^2} = \frac{\sup_{\mathbb{C}} |f|}{R} \xrightarrow{R \rightarrow \infty} 0 \quad (8)$$

Therefore $f' \equiv 0$, for any two points in \mathbb{C} , to see this implies f is constant we use the fundamental theorem of calculus, if $z_0, z_1 \in \mathbb{C}$ take γ to be the straight line starting at z_0 and ending at z_1 so that

$$|f(z_1) - f(z_0)| = \left| \int_\gamma f'(z) dz \right| \leq \ell(\gamma) \sup_{\mathbb{C}} |f'| = 0$$

(c) Suppose a polynomial P does not have a root. Then $\frac{1}{P}$ is entire, hence $1/P$ is constant by Liouville's theorem. We conclude any polynomial without a root is of degree zero. □

□