

THESIS

BY

TIGRAN SAIDNIA

Emission kernel of parton shower

Emission kernel of parton shower



Karlsruhe institute for Technology (KIT)

Institute of theoretical physics

Referents: PD Dr. Stefan Gieseke

Dr. Simon Plätzer

Supervisor: Emma Simpson

statement of originality

I hereby confirm that I have written the accompanying thesis by myself, without contributions from any sources other than those cited in the text and acknowledgements. This applies also to all graphics, drawings, maps and images included in the thesis.

Karlsruhe, 15. Januar 2019

Tigran Saidnia



Inhaltsverzeichnis


Literaturverzeichnis	2
0.1 parametrisation	1
1 Quark antiquark gluon emission kernel	2
1.1 $qg-\bar{q}$	3
1.2 $\bar{q}g-q$	7
1.3 $M_1M_2^\dagger$	9
1.4 $ M^2 $	11
2 Gluon quark gluon emission kernel	12
2.1 $gg-q$	13
2.1.1 One-loop corrections to the gluon self-energy diagram	14
2.2 $qg-g$	15
2.3 $M_1M_2^\dagger$	16
2.4 $ M^2 $	19


0.1 parametrisation

$$\left. \begin{aligned}
 q_i^\mu &= zp_i^\mu + y(1-z)p_j^\mu + \sqrt{zy(1-z)}m_\perp \\
 q^\mu &= (1-z)p_i^\mu + yzp_j^\mu - \sqrt{zy(1-z)}m_\perp \\
 q_j^\mu &= (1-y)p_j^\mu \\
 y &= \frac{q_i q}{p_i p_j} \\
 q_i + q &= p_i + yp_j \\
 q_j + q &= (1-z)p_i^\mu + (1+yz-y)p_j^\mu - \sqrt{zy(1-z)}m_\perp \\
 q_i \cdot q &= y(1-2z+2z^2)(p_i \cdot p_j) \\
 q_i \cdot q_j &= z(1-y)(p_i \cdot p_j) \\
 q_j \cdot q &= (1-z)(1-y)(p_i \cdot p_j)
 \end{aligned} \right\} \text{parametrisation} \quad (1)$$


Kapitel 1

Quark antiquark gluon emission kernel




1.1 $qg\text{-}\bar{q}$ 

$$M_1 = [\bar{u}_\sigma(q_i)(-ig_s\gamma^\mu \times [T^a]_o^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q)] [v_\tau(q_j)] \quad (1.1)$$



$$M_1^\dagger = [\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s\gamma^{\mu'} \times [T^b]_{o'}^k) u_{\sigma'}(q_i) \varepsilon^{\lambda_2}_{\mu'}(q)] [\bar{v}_{\tau'}(q_j)] \quad (1.2)$$



$$|M_1|^2 = M_1 M_1^\dagger = [\bar{u}_\sigma(q_i) (-ig_s \gamma^\mu \times [T^a]_{o'}^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q) [v_\tau(q_j)]$$

$$[\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s \gamma^{\mu'} \times [T^b]_{o'}^k) u_{\sigma'}(q_i) \varepsilon^{\lambda_2}_{\mu'}(q) [\bar{v}_{\tau'}(q_j)]] \quad (1.3)$$

$$|M_1|^2 = [\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s \gamma^{\mu'} \times [T^b]_{o'}^k) \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) \varepsilon^{\lambda_2*}_{\mu'}(q) \varepsilon^{\lambda_1}_\mu(q)$$

$$\times (-ig_s \gamma^\mu \times [T^a]_{o'}^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} [\bar{v}_{\tau'}(q_j) v_\tau(q_j)] \quad (1.4)$$

and after sum over the lorenz index (σ, σ') and (τ, τ') and unsing the spin addition relation:

$$\sum_{\sigma, \sigma'} \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) = \not{q}_i,$$

$$\sum_{\tau, \tau'} \bar{v}_\tau(q_j) v_{\tau'}(q_j) = \not{q}_j \quad (1.5)$$

and sum over polarization index (λ_1, λ_2) :

$$\sum_{\mu, \mu'} \varepsilon^{\lambda_2*}_{\mu'}(q) \varepsilon^{\lambda_1}_\mu(q) = -g_{\mu\mu'} \quad (1.6)$$

$$|M_1|^2 = \frac{-g_s^2 [T^b]_{o'}^k [T^a]_{o'}^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \gamma^{\mu'} \not{q}_i g_{\mu'\mu} \gamma^\mu (\not{q}_i + q)] [\not{q}_j] \quad (1.7)$$

from here and after contraction between all indices we can actually make statements about the last result.

$$|M_1|^2 = \frac{-g_s^2 [T^b]_{o'}^k [T^a]_{o'}^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \gamma^{\mu'} \not{q}_i \gamma_{\mu'} (\not{q}_i + q)] [\not{q}_j] \quad (1.8)$$

In other words we expect the tree level diagram from LO and a number: Which means:

$$|M^2| = \left| \text{diagram} \right|^2 \otimes \left| \text{diagram} \right|^2$$

contribution from LO
a complex number

$$|M_1|^2 = \frac{-g_s^2 [T^b]_{o'k} [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [P_i][P_j] \otimes (\text{a complex number}) \quad (1.9)$$

Let's calculate the contribution and compare the final result with this expectation:

$$\begin{aligned}
 N &=: \gamma^{\mu'} \not{q}_i \gamma_{\mu'} = q_{i\sigma} \gamma^{\mu'} \gamma^\sigma \gamma_{\mu'} \\
 &= q_{i\sigma} (\{\gamma^{\mu'}, \gamma^\sigma\} - \gamma^\sigma \gamma^{\mu'}) \gamma_{\mu'} \\
 &= q_{i\sigma} 2g^{\mu'\sigma} \gamma_{\mu'} - d \gamma^\sigma \\
 &= (2 - d) \not{q}_i
 \end{aligned} \quad (1.10)$$

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^b]_{o'k} [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \not{q}_i (\not{q}_i + q)] [\not{q}_j] \quad (1.11)$$

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^b]_{o'k} [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [\not{q}_i \not{q}_i \not{q}_i + \not{q}_i \not{q}_i \not{q} + \not{q} \not{q}_i \not{q}_i + \not{q} \not{q}_i \not{q}] [\not{q}_j] \quad (1.12)$$

For the momenta are on-shell which means:

$$\begin{aligned}
 \not{q}_i \not{q}_i &= q_i^2 = m_i^2 \\
 \not{q} \not{q} &= q^2 = m^2 \\
 \not{q}_j \not{q}_j &= q_j^2 = m_j^2
 \end{aligned} \quad (1.13)$$

we can first neglect the mass of patrons and ignore each term with $\not{q}_i \not{q}_i$ and $\not{q} \not{q}$ as well.

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^b]_{o'k} [T^a]_o^l}{(2q_i q)(2q_i q)} [\not{q} \not{q}_i \not{q}] [\not{q}_j] \quad (1.14)$$

$$\begin{aligned}
L &= \not{q}_i \not{q}_\mu (\{\gamma^\mu, \gamma^\sigma\} - \gamma^\sigma \gamma^\mu) \\
&= \not{q}_i [2q_i^\mu q_\mu - q_{i\sigma} q_\mu \gamma^\mu \gamma^\sigma] \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q_\mu [\gamma^\mu \gamma^\mu \gamma^\sigma] \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q_\mu \left[\frac{\gamma^\mu \gamma^\mu}{2} + \frac{\gamma^\mu \gamma^\mu}{2} \right] \gamma^\sigma \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q_\mu [g^{\mu\mu}] \gamma^\sigma \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q^\mu \gamma^\sigma \\
&= \not{q}_i (2q_i q) - q^2 \not{q}_i \\
&= \not{q}_i (2q_i q)
\end{aligned} \tag{1.15}$$

After inserting the last result of L and simplify the term $(2q_i q)$ from the denominator and nominator because , we get:

$$|M_1|^2 = -(2-d) \frac{g_s^2 [T^b]_{o'}^k [T^a]_o^l}{2y(1-2z+z^2)(p_i \cdot p_j)} [\not{q}_i][\not{q}_j] \tag{1.16}$$

Now we are going to use the parametrisation from equation (1) to reduce the 3-member matrix element to 2-member and take out the singularity term from the amplitude.


$$|M_1|^2 = (d-2) \frac{g_s^2 [T^b]_{o'}^k [T^a]_o^l}{2y(1-2z+z^2)(p_i \cdot p_j)} [(1-z) \not{p}_i + zy \not{p}_j - \sqrt{zy(1-z)} \not{m}_\perp][(1-y) \not{p}_j] \tag{1.17}$$

Multiplying the both sides


$$\begin{aligned}
|M_1|^2 &= (d-2) \frac{g_s^2 [T^b]_{o'}^k [T^a]_o^l}{2y(1-2z+z^2)(p_i \cdot p_j)} [(1-z)(1-y) \not{p}_i \not{p}_j \\
&\quad + zy(1-y) \not{p}_j \not{p}_j + (1-y)\sqrt{zy(1-z)} \not{m}_\perp \not{p}_j]
\end{aligned} \tag{1.18}$$

and under consideration of the fact that p_i and p_j are the on-shell momenta of the emitter and spectator partons, we can ignore the terms with $\not{p}_i \not{p}_i$ and $\not{p}_j \not{p}_j$. The $p_i \cdot m_\perp$ and $p_j \cdot m_\perp$ are always 0 because the p_i and p_j are lightlike, i.e. zero transverse component. So those terms can be neglected.


$$|M_1|^2 = (d-2)(1-z)(1-y) \frac{g_s^2 [T^b]_{o'}^k [T^a]_o^l}{2y(1-2z+z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \tag{1.19}$$

1.2 $\bar{q}g$ -q

$$M_2 = \left[\frac{i(\not{q}_j + \not{q})}{(q_j + q)^2} (-ig_s \gamma^\nu \times [T^c]_f^m) v_\tau(q_j) \varepsilon^{\lambda_3}_\nu(q) [u_\sigma(q_i)] \right] \quad (1.20)$$



$$M_2^\dagger = [\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q) [\bar{u}_{\sigma'}(q_i)] \quad (1.21)$$



$$|M_2|^2 = M_2 M_2^\dagger = \left[\frac{i(\not{q}_j + \not{q})}{(q_j + q)^2} (-ig_s \gamma^\nu \times [T^c]_f^m) v_\tau(q_j) \varepsilon^{\lambda_3}_\nu(q) [u_\sigma(q_i)] \right] \quad (1.22)$$

$$\left[\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q) [\bar{u}_{\sigma'}(q_i)] \right]$$

$$|M_2|^2 = \frac{g_s^2 [T^c]_f^m [T^d]_{f'}^n}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\nu v_\tau(q_j) \bar{v}_{\tau'}(q_j) \varepsilon^{\lambda_3}_\nu(q) \varepsilon^{\lambda_4}_{\nu'}(q) \gamma^{\nu'} (\not{q}_j + \not{q})] \quad (1.23)$$

$$[u_\sigma(q_i)] [\bar{u}_{\sigma'}(q_i)]$$

and after sum over the lorenz index (σ, σ') and (τ, τ') and unsing the spin addition relation:

$$\sum_{\sigma, \sigma'} \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) = \not{q}_i, \quad (1.24)$$

$$\sum_{\tau, \tau'} \bar{v}_\tau(q_j) v_{\tau'}(q_j) = \not{q}_j$$

and sum over polarization index (λ_3, λ_4) :

$$\sum_{\nu, \nu'} \varepsilon^{\lambda_4*}_{\nu'}(q) \varepsilon^{\lambda_3}_\nu(q) = -g_{\nu\nu'} \quad (1.25)$$


$$|M_2|^2 = \frac{g_s^2 [T^c]_f^m [T^d]_{f'}^n}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\nu \not{q}_j (-g_{\nu\nu'}) \gamma^{\nu'} (\not{q}_j + \not{q})] [\not{q}_i] \quad (1.26)$$

After the same calculation from the last part, we'll get:

$$|M_2|^2 = (d - 2) \frac{g_s^2 [T^c]_f^m [T^d]_{f'}^n}{(2qq_j)} [\not{q}] [\not{q}_i] \quad (1.27)$$

finally:

$$|M_2|^2 = -(d - 2) y z^2 \frac{g_s^2 [T^c]_f^m [T^d]_{f'}^n}{2(1 - z)(1 - y)(p_i \cdot p_j)} [\not{p}_i] [\not{p}_j] \quad (1.28)$$

1.3 $M_1 M_2^\dagger$ 

$$M_1 M_2^\dagger = [\bar{u}_\sigma(q_i) (-ig_s \gamma^\mu \times [T^a]_o^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q)] [v_\tau(q_j)]$$

$$[\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q)] [u_{\sigma'}(q_i)] \quad (1.29)$$

$$M_1 M_2^\dagger = \frac{g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} [\not{q}_i \gamma^\mu (\not{q}_i + \not{q})] \varepsilon^{\lambda_1}_\mu(q) \varepsilon^{\lambda_4}_{\nu'}(q)$$

$$[\not{q}_j \gamma^{\nu'} (\not{q}_j + \not{q})] \quad (1.30)$$

$$M_1 M_2^\dagger = \frac{g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} [\not{q}_i \gamma^\mu (\not{q}_i + \not{q})] - g_{\mu\nu'}$$

$$[\not{q}_j \gamma^{\nu'} (\not{q}_j + \not{q})] \quad (1.31)$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} [\not{q}_i \gamma^\mu (\not{q}_i + \not{q})] [\not{q}_j \gamma_\mu (\not{q}_j + \not{q})] \quad (1.32)$$

Expectation:

$$|M^2| = \left| \begin{array}{c} \text{Diagram with two shaded circles connected by two horizontal lines labeled } P_i \text{ and } P_j \\ \text{contribution from LO} \end{array} \right|^2 \otimes \left| \begin{array}{c} \text{Diagram with a shaded circle and a wavy line} \\ \text{a complex number} \end{array} \right|^2$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} [(\not{q}_i + \not{q}) \gamma^\mu \not{q}_i] [(\not{q}_j + \not{q}) \gamma_\mu \not{q}_j] \quad (1.33)$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} [-(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu + 2(\not{q}_i + \not{q}) q_i^\mu] \quad (1.34)$$

$$[-(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu + 2(\not{q}_j + \not{q}) q_{j\mu}]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} \quad (1.35)$$

$$[(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu] [(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu]$$

$$-2[(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu] [(\not{q}_j + \not{q}) q_{j\mu}]$$

$$-2[(\not{q}_i + \not{q}) q_i^\mu] [(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu]$$

$$+4[(\not{q}_i + \not{q}) q_i^\mu] [(\not{q}_j + \not{q}) q_{j\mu}]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} \quad (1.36)$$

$$[\not{q} \not{q}_i \gamma^\mu] [\not{q} \not{q}_j \gamma_\mu]$$

$$-2[\not{q} \not{q}_i \gamma^\mu] [(\not{q} + \not{q}_j) q_{j\mu}]$$

$$-2[(\not{q} + \not{q}_j) q_i^\mu] [\not{q} \not{q}_j \gamma_\mu]$$

$$+4[(\not{q} + \not{q}_j) q_i^\mu] [(\not{q} + \not{q}_j) q_{j\mu}]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)(p_i \cdot p_j)} \quad (1.37)$$

$$[y(1-2z+z^2) \not{p}_i \not{p}_j \gamma^\mu] [(1-z)(1-y) \not{p}_i \not{p}_j \gamma_\mu]$$

$$+4(q_i^\mu \cdot q_{j\mu}) [(\not{q}_i + \not{q})] [(\not{q}_j + \not{q})]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)(p_i \cdot p_j)} \quad (1.38)$$

$$[y(1-2z+z^2) \not{p}_i \not{p}_j \gamma^\mu] [(1-z)(1-y) \not{p}_i \not{p}_j \gamma_\mu]$$


$$+4(p_i \cdot p_j) [(\not{p}_i + y \not{p}_j)] [(1-z) \not{p}_i + (1+yz-y) \not{p}_j - \sqrt{zy(1-z)} \not{m}]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)} z(1-y) [\not{p}_i] [\not{p}_j] \quad (1.39)$$


$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(1-z)y(1-2z+z^2)(p_i \cdot p_j)} z [\not{p}_i] [\not{p}_j] \quad (1.40)$$

1.4 $|M^2|$

$$|M|^2 = |M_1|^2 + |M_2|^2 + M_1 M_2^\dagger + M_1^\dagger M_2 \quad (1.41)$$




$$|M|^2 = |M_1|^2 + |M_2|^2 + 2RE(M_1 M_2^\dagger) \quad (1.42)$$




$$\begin{aligned}
 |M|^2 = & (d-2)(1-z)(1-y) \frac{g_s^2 [T^b]_{o'}^k [T^a]_o^l}{2y(1-2z+z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\
 & - (d-2)yz^2 \frac{g_s^2 [T^c]_f^m [T^d]_{f'}^n}{2(1-z)(1-y)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\
 & + 2RE\left(\left(\frac{-2z}{z-1}\right) \frac{g_s^2 [T^a]_o^l [T^d]_{f'}^n}{2y(1-2z+z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j]\right)
 \end{aligned} \quad (1.43)$$

Kapitel 2

Gluon quark gluon emission kernel




2.1 gg-q




$$M_1 = \left[\frac{-i}{(q_i + q)^2} (-g_s f^{a o l} (g^{\mu \zeta} (-q_i + q)^\eta + g^{\zeta \eta} (-q - (q_i + q))^\mu + g^{\eta \mu} (q_i + q + q_i)^\zeta) \right. \\ \left. \varepsilon^{\lambda_1}_\mu(q) \varepsilon^{\lambda_6}_\zeta(q_i) \right] [\bar{u}_\tau(q_j)] \quad (2.1)$$

$$M_1 = \left[\frac{-i}{(q_i + q)^2} (-g_s f^{a o l} (g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q_i + q)^\mu + g^{\eta \mu} (2q_i + q)^\zeta) \right. \\ \left. \varepsilon^{\lambda_1}_\mu(q) \varepsilon^{\lambda_6}_\zeta(q_i) \right] [\bar{u}_\tau(q_j)] \quad (2.2)$$



$$M_1^\dagger = \left[\frac{i}{(q_i + q)^2} (-g_s f^{a' o' k} (g^{\mu' \zeta'} (q - q_i)^{\eta'} - g^{\zeta' \eta'} (2q_i + q)^{\mu'} + g^{\eta' \mu'} (2q_i + q)^{\zeta'}) \right. \\ \left. \varepsilon^{\lambda_2}_{\mu'}(q) \varepsilon^{\lambda_5}_{\zeta'}(q_i) \right] [u_{\tau'}(q_j)] \quad (2.3)$$




$$\begin{aligned}
 |M_1|^2 = & \left[\frac{-i}{(q_i + q)^2} (-g_s f^{a o l} (g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q_i + q)^\mu + g^{\eta \mu} (2q_i + q)^\zeta) \right. \\
 & \left. \varepsilon^{\lambda_1}_{\mu}(q) \varepsilon^{\lambda_2}_{\mu'}(q) \varepsilon^{\lambda_6}_{\zeta}(q_i) \varepsilon^{\lambda_5}_{\zeta'}(q_i) \right. \\
 & \left. (-g_s f^{a' o' k} (g^{\mu' \zeta'} (q - q_i)^{\eta'} - g^{\zeta' \eta'} (2q_i + q)^{\mu'} + g^{\eta' \mu'} (2q_i + q)^{\zeta'}) \frac{i}{(q_i + q)^2} [\bar{u}_\tau(q_j) u_{\tau'}(q_j)] \right] \quad (2.4)
 \end{aligned}$$

$$\begin{aligned}
 |M_1|^2 = & \frac{g_s^2 f^{a o l} f^{a' o' k}}{(q_i + q)^2 (q_i + q)^2} \\
 & [(g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q_i + q)^\mu + g^{\eta \mu} (2q_i + q)^\zeta) \\
 & g_{\mu \mu'} g^{\zeta \zeta'} (g_{\mu' \zeta'} (q - q_i)_{\eta'} - g_{\zeta' \eta'} (2q_i + q)_{\mu'} + g_{\eta' \mu'} (2q_i + q)_{\zeta'})] [\not{A}_j] \quad (2.5)
 \end{aligned}$$


$$\begin{aligned}
 |M_1|^2 = & \frac{g_s^2 f^{a o l} f^{a' o' k}}{(q_i + q)^2 (q_i + q)^2} \\
 & [(g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q_i + q)_{\mu'} + g^{\eta \mu'} (2q_i + q)^\zeta) \\
 & (g^{\eta \mu'} (q - q_i)_{\eta'} - g^{\zeta \eta'} (2q_i + q)_{\mu'} + g_{\eta' \mu'} (2q_i + q)^\zeta)] [\not{A}_j] \quad (2.6)
 \end{aligned}$$

$$\begin{aligned}
 |M_1|^2 = & \frac{g_s^2 f^{a o l} f^{a' o' k}}{(q_i + q)^2 (q_i + q)^2} \\
 & [g^{\mu' \zeta} (q - q_i)^\eta g^{\eta \mu'} (q - q_i)_{\eta'} - g^{\mu' \zeta} (q - q_i)^\eta g^{\zeta \eta'} (2q_i + q)_{\mu'} \\
 & + g^{\mu' \zeta} (q - q_i)^\eta g_{\eta' \mu'} (2q_i + q)^\zeta - g^{\zeta \eta} (2q_i + q)_{\mu'} g^{\eta \mu'} (q - q_i)_{\eta'} \\
 & + g^{\zeta \eta} (2q_i + q)_{\mu'} g^{\zeta \eta'} (2q_i + q)_{\mu'} - g^{\zeta \eta} (2q_i + q)_{\mu'} g_{\eta' \mu'} (2q_i + q)^\zeta \\
 & + g^{\eta \mu'} (2q_i + q)^\zeta g^{\eta \mu'} (q - q_i)_{\eta'} - g^{\eta \mu'} (2q_i + q)^\zeta g^{\zeta \eta'} (2q_i + q)_{\mu'} \\
 & + g^{\eta \mu'} (2q_i + q)^\zeta g_{\eta' \mu'} (2q_i + q)^\zeta] [\not{A}_j] \quad (2.7)
 \end{aligned}$$

2.1.1 One-loop corrections to the gluon self-energy diagram



2.2 qg-g



$$|M_2|^2 = M_2 M_2^\dagger = [\bar{u}_\sigma(q_j) (-ig_s \gamma^\Upsilon \times [T^c]_o^l) \frac{i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_7 \Upsilon}(q)] [\varepsilon^{\lambda_9 \nu}(q_i)]$$

$$[\frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} (ig_s \gamma^{\Upsilon'} \times [T^d]_{o'}^k) u_{\sigma'}(q_j) \varepsilon^{\lambda_8 \Upsilon'}(q)] [\varepsilon^{\lambda_{10} \nu'}(q_i)] \quad (2.8)$$

$$|M_2|^2 = [\frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} (-ig_s \gamma^\Upsilon \times [T^c]_o^l) \bar{u}_\sigma(q_j) u_{\sigma'}(q_j) \varepsilon^{\lambda_7 \Upsilon}(q) \varepsilon^{\lambda_8 \Upsilon'}(q)]$$

$$\times (ig_s \gamma^{\Upsilon'} \times [T^d]_{o'}^k) \frac{i(\not{q}_j + \not{q})}{(q_j + q)^2} [\varepsilon^{\lambda_{10} \nu'}(q_i) \varepsilon^{\lambda_9 \nu}(q_i)] \quad (2.9)$$

$$|M_2|^2 = \frac{g_s^2 [T^c]_o^l [T^d]_{o'}^k}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\Upsilon \not{q}_j (-g_{\Upsilon \Upsilon'})$$

$$\gamma^{\Upsilon'} (\not{q}_j + \not{q})] [-g_{\nu \nu'}] \quad (2.10)$$

Expect:

$$|M_2|^2 = \left| \text{diagram} \right|^2 \otimes \left| \text{diagram} \right|^2$$

contribution from LO a complex number

The diagram on the left shows two shaded circles representing hard subprocesses. A quark line with momentum \$p_i\$ connects the top of the circles, and a gluon line with momentum \$q_i\$ connects the bottom. The gluon line is labeled \$g_{\nu \nu'}\$. The diagram on the right shows a quark line with momentum \$q_i + q\$ entering a vertex, from which a gluon with momentum \$q\$ is emitted. The quark line continues with momentum \$q_i + q\$. A gluon with momentum \$q_i\$ is exchanged between this vertex and another vertex where a quark line with momentum \$q_i + q\$ enters.


$$|M_2|^2 = -\frac{g_s^2 [T^c]_o^l [T^d]_{o'}^k}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\Upsilon \not{q}_j \gamma^\Upsilon (\not{q}_j + \not{q})] [-g_{\nu'\nu}] \quad (2.11)$$

$$|M_2|^2 = (d-2) \frac{g_s^2 [T^c]_o^l [T^d]_{o'}^k}{(2qq_j)} [\not{q}] [-g_{\nu'\nu}] \quad (2.12)$$

$$|M_2|^2 = (d-2) \frac{g_s^2 [T^c]_o^l [T^d]_{o'}^k}{2(1-z)(1-y)(p_i \cdot p_j)} [(1-z) \not{p}_i + zy \not{p}_j - \sqrt{zy(1-z)} \not{m}_\perp] [-g_{\nu'\nu}] \quad (2.13)$$

$$|M_2|^2 = (d-2)zy \frac{g_s^2 [T^c]_o^l [T^d]_{o'}^k}{2(1-z)(1-y)(p_i \cdot p_j)} [\not{p}_j] [-g_{\nu'\nu}] \quad (2.14)$$

2.3 $M_1 M_2^\dagger$




$$M_1 M_2^\dagger = \left[\frac{-ig_\eta \zeta}{(q_i + q)^2} (-g_s f^{aol} (g^{\mu\zeta} (q - q_i)^\eta - g^{\zeta\eta} (2q_i + q)^\mu + g^{\eta\mu} (2q_i + q)^\zeta) \right. \\ \left. \varepsilon^{\lambda_1}_\mu(q_i) \varepsilon^{\lambda_6}_\zeta(q) [\bar{u}_\tau(q_j)] \left[\frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} (ig_s \gamma^{\mu'} \times [T^d]_{o'}^k) u_{\sigma'}(q_j) \varepsilon^{\lambda_8}_{\Upsilon'}(q) [\varepsilon^{\lambda_{10}}_{\nu'}(q_i)] \right] \right] \quad (2.15)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(qq_i)(qq_j)} [(g^{\mu\zeta} (q - q_i)^\eta - g^{\zeta\eta} (2q_i + q)^\mu + g^{\eta\mu} (2q_i + q)^\zeta)] \\ (-g_{\zeta\nu'}) (-g_{\mu\Upsilon'}) [(\not{q}_j + \not{q}) \gamma^{\Upsilon'} \not{q}_j] \quad (2.16)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(qq_i)(qq_j)} [(g^{\mu\nu'} (q - q_i)^\eta - g^{\eta\nu'} (2q_i + q)^\mu + g^{\eta\mu} (2q_i + q)_{\nu'})] \\ [(\not{q}_j + \not{q}) \gamma_\mu \not{q}_j] \quad (2.17)$$

Expect:





$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(qq_i)(qq_j)} [-g^{\eta}_{\nu'}(2q_i + q)^\mu] [-(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu + 2(\not{q}_j + \not{q}) q_{j\mu}] \quad (2.18)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(qq_i)(qq_j)} [-g^{\eta}_{\nu'}(2q_i + q)^\mu] [-(qq_j) \gamma_\mu + 2(\not{q}_j + \not{q}) q_{j\mu}] \quad (2.19)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(qq_i)(qq_j)} [(qq_j) g^{\eta}_{\nu'}(2 \not{q}_i + \not{q}) - 2((2q_i + q) \cdot q_j) g^{\eta}_{\nu'}(\not{q}_j + \not{q})] \quad (2.20)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)(p_i \cdot p_j)} [g^{\eta}_{\nu'}][(qq_j)(2 \not{q}_i + \not{q}) - 2((2q_i + q) \cdot q_j) (\not{q}_j + \not{q})] \quad (2.21)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)(p_i \cdot p_j)} [g^{\eta}_{\nu'}][(1-z)(1-y)(p_i \cdot p_j)(2 \not{q}_i + \not{q}) - (4q_i \cdot q_j + 2q \cdot q_j) (\not{q}_j + \not{q})] \quad (2.22)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)(p_i \cdot p_j)} [g^{\eta}_{\nu'}][(1-z)(1-y)(p_i \cdot p_j)(2 \not{q}_i + \not{q}) - (4(z(1-y)(p_i \cdot p_j)) + 2((1-z)(1-y)(p_i \cdot p_j))) (\not{q}_j + \not{q})] \quad (2.23)$$

$$M_1 M_2^\dagger = \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)} [g^{\eta}_{\nu'}][(1-z)(1-y)(2 \not{q}_i + \not{q}) - (4(z(1-y)) + 2((1-z)(1-y))) (\not{q}_j + \not{q})] \quad (2.24)$$

$$\begin{aligned}
M_1 M_2^\dagger = & \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)} \\
& [g^\eta_{\nu'}][(1-z)(1-y)(2y(1-z) \not{p}_j + yz \not{p}_j) \\
& -2(1+z)(1-y)((1+yz-z) \not{p}_j)]
\end{aligned} \tag{2.25}$$

$$\begin{aligned}
M_1 M_2^\dagger = & \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(1-z)(1-y)y(1-2z+z^2)(p_i \cdot p_j)} \\
& [g^\eta_{\nu'}][-2(1+z)(1-y)(1+yz-z) \not{p}_j]
\end{aligned} \tag{2.26}$$

$$M_1 M_2^\dagger = -2(1+z)(1+yz-z) \frac{ig_s^2 f^{aol} [T^d]_{o'}^k}{4(1-z)y(1-2z+z^2)(p_i \cdot p_j)} [g^\eta_{\nu'}][\not{p}_j] \tag{2.27}$$

2.4 $|M^2|$

$$|M^2| = \left| \begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \end{array} \right|^2 + 2 \operatorname{Re} E \left(\begin{array}{c} \text{Diagram 4} \end{array} \right)$$

The equation shows the squared magnitude of the amplitude M^2 as the sum of the squared magnitudes of three diagrams plus twice the real part of the interference term between the first and third diagrams. The diagrams involve two shaded circular regions representing nuclei, with quark and gluon lines connecting them. Momenta are labeled as q_i , $q_i + q$, q_j , and $q_j + q$. A vertical dashed line separates the two nuclei in each diagram.