

# THESIS

BY


TIGRAN SAIDNIA

---

## Emission kernel of parton shower

Emission kernel of parton shower

---



Karlsruhe institute for Technology (KIT)

Institute of theoretical physics

Referents: PD Dr. Stefan Gieseke

Dr. Simon Plätzer

Korreferent: Prof. Dr. Dieter Zeppenfeld

Supervisor: Emma Simpson Dore

statement of originality

I hereby confirm that I have written the accompanying thesis by myself, without contributions from any sources other than those cited in the text and acknowledgements. This applies also to all graphics, drawings, maps and images included in the thesis.

Karlsruhe, April 25, 2019

---

Tigran Saidnia



# Abstract

Emission kernel of parton shower



# Contents

<b>Table of contents</b>	<b>4</b>
<b>1 Theoretical Basics</b>	<b>2</b>
1.1 Brief history of particle physics . . . . .	2
1.2 Standard model . . . . .	4
1.3 Quantum chromo dynamics . . . . .	4
1.4 QCD Lagrangian . . . . .	9
1.5 Colour factor calculation . . . . .	10
<b>2 Catani Seymour</b>	<b>14</b>
2.1 IR and Collinear Divergences . . . . .	14
2.2 Subtraction method . . . . .	17
2.3 Hard scattering . . . . .	22
2.4 Mapping 3 partons to 2 . . . . .	24
2.5 Old mapping . . . . .	24
2.6 new kinematic . . . . .	25
2.7 Single emission part . . . . .	26
2.8 Common scalar products . . . . .	27
2.9 Recipe for the use of the new parametrisation . . . . .	27
2.9.1 Parametrization in terms of $(k_1 \cdot q_i)(k_1 \cdot q_k)$ . . . . .	27
2.9.2 Parametrization in terms of $(k_1 \cdot q_i)(k_1 \cdot q_i)$ . . . . .	28
<b>3 Quark anti-quark gluon emission kernel</b>	<b>31</b>
3.1 Matrix element of a quark with a gluon radiation $ M_1 ^2$ . . . . .	32
3.2 Matrix element of an anti-quark with a gluon radiation $ M_2 ^2$ . . . . .	36
3.3 Interference contribution . . . . .	37
3.4 Final result . . . . .	39
3.5 Double-check the results with the new kinematic . . . . .	41
<b>4 Gluon radiation from a parent quark</b>	<b>43</b>
4.1 Gluon-Emitter Bubble . . . . .	44
4.1.1 One-loop corrections to the gluon self-energy diagram(Gluon-Emitter Bubble) . . . . .	46
4.2 Gluon-Spectator Bubble . . . . .	52

4.2.1	One-loop corrections to the gluon self-energy diagram (Gluon-Spectator Bubble)	53
4.3	Interference term $M_1 M_2^\dagger$	53
4.4	Interference term of inverse $M_1 M_2^{\dagger'}$	59
4.5	Parametrization in terms of $(k_1 \cdot q_i)(q_i \cdot q_k)$	59
4.6	$ M^2 $	60
<b>5</b>	<b>Quark gluon quark emission kernel</b>	<b>62</b>
5.1	Quark loop	63
5.2	Spectator Quark loop	65
5.3	Interference term	66
5.4	$ M^2 $	66
<b>6</b>	<b>Gluon quark quark emission kernel</b>	<b>67</b>
6.1	$M_1$	68
6.2	$M_2$	69
6.3	$M_1 M_2^\dagger$	70

# Introduction

## Introduction



# Chapter 1

## Theoretical Basics

### 1.1 Brief history of particle physics

Knowledge is a human need. For thousands of years we have been trying to understand the secrets of the universe. Such riddles fascinated even Johann Wolfgang von Goethe, as he wrote in his book *Faust* [14]; eine Tragedie, "What holds the world together in its innermost." Almost 400 years before Christ, an ancient Greek philosopher, Democritus, and his teacher Leukipp claimed that matter cannot be divided at will. Rather, there must be an Atomos (Greek: indivisible) that could no longer be subdivided. Democritus was of the opinion that there were infinitely many atoms with different geometric forms that were in contact in a certain way. He pointed out that a thing has a color, taste or even soul, based on the apparent effect of the composition of these small grains. [3]

This statement of Democritus was first laughed at by the renowned philosopher Aristotiles. It took about 2000 years for a chemist named John Dalton to deal with the subject. Based on various test series, he summarized his conclusion in his book *A New System of Chemical Philosophy*, that all substances consist of spherical indivisible atoms. The atoms of different elements have different masses and volumes. This was exactly the most striking difference to Democritus's atomic world.[6]

The discovery of the periodic system by **D. Mendeleev** and **P. Meyer** enabled us to arrange the atoms according to their mass in such a way that their properties occur in a certain order.[17]

In 1897 Joseph Thompson was able to obtain a stream of particles by heating metals and deflecting them by a magnetic field. This electron beam was 200 times lighter than the lightest atom, hydrogen. His conclusion was that atoms cannot be indivisible. He suggested that each atom consists of an electrically positively charged sphere in which electrically negatively charged electrons are stored - like raisins in a cake.

furthermore, renowned scientists as well as Marie and Pierre Curie have contributed much to the development of atomic theory by discovering radioactivity, Boltzmann by kinetic gas theory and Plank, the founder of quantum physics. However, one of the most important steps in the atomic model was taken by the British physicist Rutherford. He bombarded a thin aluminium foil with a radioactive sample. If Thompson's cake model were correct, only a few alpha particles would be detected behind the aluminium foil.



Surprisingly, many particles were visible, which could only be explained by the assumption that the majority of atoms consisted of empty spaces. Another miracle was that some particles could be seen above or below the target sample. Since we knew that the alpha particles were positively charged, we could assume the electric repulsive force of two positive charges. From the ideas of Planck and Rutherford, the Danish physicist Bohr (1885-1962) developed a planetary atomic model. The electrons then move around the nucleus in certain orbits, like planets orbit the sun. The orbits are also called shells. The special thing about it was that the distances of the electron orbits follow strict mathematical laws.

At first, however, it remained unclear what this core should consist of. [7, 17] In 1912, the Austrian physicist Victor Hess discovered during his balloon flights that the ionization rate of the Earth's atmosphere increases with altitude. This result was not expected because until then the Earth's radioactivity was known as the only source of air ionization. Therefore, he postulated this new type of radiation as cosmic radiation, which must originate outside the Earth's atmosphere [10].

Further investigations two years later confirmed the thesis of a cosmic background of such radiation. After this new discovery, it was discovered that the radiation consists of charged particles. In 1932, the American physicist Carl David Anderson was able to prove the postulated particle of Dirac, the positron, as a component of an air shower through his cloud chamber. For a long time, cosmic rays were the only way to analyse such exotic particles.[1] This changed when particle accelerators were able to generate particles in collisions. But even today, cosmic rays are the only way to study particles of the highest energies, since these energies cannot be reached by today's particle accelerators, such as the LHC. The LHC, the world's largest accelerator at CERN, produces particles with centre-of-mass energy equivalent to a cosmic particle of nearly  $10^{17} \text{eV}$ , with the energy spectrum of cosmic particles reaching up to  $10^{20} \text{eV}$ . However, we can only analyse such exotic particles in detail by increasing the luminosity and procession of the particle accelerators at the nucleus. The discovery of the neutron by **Chadwick** (1932) showed that atomic nuclei are made up of protons and neutrons. It was also clear that, in addition to gravitation and the electromagnetic force, there should exist two short-range forces in nature; a strong force which binds the nucleons together and a weak force which is responsible for radioactive. In the meantime it was agreed that a new theory was needed for the classification and grouping of this particle zoo. This is how the current standard model came into being.

The SLAC experiments indicate that the electrons can be scattered as quasi-free point-like constituents within the proton structure, which actually meant that the protons or neutrons are not point-like and must consist of other constituents. Through the bubble chamber a huge number of previously invisible particles (Gell-Mann's eightfold path) could suddenly be made visible, which represented contradictions to the previous physics. To explain this, the physicist Gell-Mann found basic building blocks from which all previously known atomic particles should be built. The components are later identified with quarks.



## 1.2 Standard model

The Standard Model in particle physics encompasses all of the Elementary particles and their interactions. It is a gauge theory spontaneously broken by the Higgs mechanism with the gauge group  $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ .

From a theoretical point of view, the Standard Model is a quantum field theory that is based on local gauge invariance and consists of two rough parts. The electroweak sector  $SU(2)_L \otimes U(1)_Y$  is called **GWS (Glashow-Weinberg-Salam)** theory and describes the gauge bosons  $W^\pm, Z^0, \gamma$ , the Higgs sector and its interaction with the leptons and quarks. In contrast to the other gauge bosons, the exchange particles of the weak interaction carry mass, which also affects the properties of the interactions. The color-charged sector  $SU(3)_C$ , the chromodynamics, deals with quarks and contains the eight massless, electrically neutral gluons as gauge bosons. The gauge groups  $SU(3)_C$  and  $SU(2)_L$  are non-abel gauge theories, more precisely **Yang Mills theories**. The massive particles, fermions, will be divided into two groups, leptons and quarks. Each group is arranged in 3 generations. Within the leptons there are three electrically neutral neutrinos. The mass of the particles increases from generation to generation. Neutrinos only interact weakly, whereas the charged leptons interact both weakly and electromagnetically. Quarks are characterized by the fact that they can also interact strongly [8].


## 1.3 Quantum chromo dynamics

Nowadays, we know there are four types of interactions.

Interaction	Energy scale	Range [m]	Mediators
Strong	$\sim 1$	$10^{-15}$	$g$
Electromagnetic	$\sim 10^{-2}$	$\infty$	$\gamma$
Weak	$\sim 10^{-6}$	$10^{-18}$	$W^\pm, Z$
Gravity	$\sim 10^{-38}$	$\infty$	maybe graviton

Otherwise, it's clear that nucleons are made up of quark and gluons. Whereby, the gluons are the exchange bosons for this short ranged interaction. To explain the short range of the strong interaction Yukawa (1934) postulated mesons as a mediator for this force by the exchange of this massive field quanta. Three years later a candidate ( $\pi$  meson) was found in cosmic rays. Later on it was shown that massive gauge field quanta break the gauge symmetry so that the mediator must be massless. If it is based on the  $SU(3)$  gauge symmetry of the QCD<sup>1</sup> massless Lagrangian how can the strong sector be short range? Another question came from a series of experiments at SLAC. Through high-energy electron-proton scattering there was a search for evidence of the existence of quarks and their behaviour like free particles despite the energetically bound inner proton. The solution to these question was explained by Gross, Politzer and Wilczek through asymptotic freedom. This effect can be proved by the running coupling and anti screen-

<sup>1</sup>The quantum field theory which describes this area is called Quantum chromodynamics short QCD.



**Figure 1.1:** A schematic picture of neutron structure. at the left side of the resolution is too low to see. The 3 quarks picture allows us to interpretate the quantum numbers of the neutron in the valence band. We also obtain a high-resolution picture for a large  $Q^2$ . Here we have a lot of gluons (gluon sea) and quarks pairs. [2]

The interesting thing is, it doesn't matter in which energy scale we observe the quantum number of a neutron, because it is always the same.

ing in QCD. For the calculation of the propagator loop correction in QCD we have to consider both quark loops (negative contribution  $\rightarrow$  screening) and gluon loops (positive contribution  $\rightarrow$  anti screening).



**Figure 1.2:** Running coupling compared for QED, with a positive and QCD with a negative beta function. The quark loop vacuum polarization diagram gives a negative contribution to  $\beta_0 \sim n_f$  and the gluon loop gives a positive contribution to  $\beta_0 \sim N_c$ . The second contribution is bigger than the first, so that  $\beta_0 > 0$  in QCD. In contrast to this, the function in QED is negative because the second contribution does not exist  $N_C = 0$ .

The one loop running coupling in QCD is:


$$\alpha_s(Q^2) = \frac{\alpha_s(\mu^2)}{1 + \beta_0 \alpha_s(\mu^2) \ln(\frac{Q^2}{\mu^2})} \quad (1.1)$$

Where  $\beta_0 = \frac{11N_c - 2n_f}{12\pi}$ ,  $n_f$  comes from the first diagram and causes screening.  $n_f$  is the number of quarks and  $N_c$  the number of colours and comes from the second diagram (anti screening).

Obviously, with  $n_f = 6$  and  $N_c = 3$  in standard model we will get  $\beta_0 > 0$ . The Beta function is defined as:

$$\beta(\alpha) = -(\beta_0 \alpha^2 + \beta_1 \alpha^3 + \beta_2 \alpha^4 + \dots) = \frac{d\alpha(Q^2)}{d \ln(Q^2)} \quad (1.2)$$

e.g.  $-(\beta_0 \alpha^2) < 0$  will be negative, which is actually the opposite of QED with  $\beta_0 = -\frac{\pi}{3} \rightarrow -(\beta_0 \alpha^2) > 0$  ! That means the coupling constant in QCD will increase with decreasing  $Q^2$  (increasing distance), In QED vice versa.



Asymptotic freedom allows us to use perturbation theory <sup>2</sup>. Quarks have not yet been observed as free particles. With increasing separation it will be easier to produce a quark-antiquark pair than to isolate a quark because the coupling between them is too strong.

<sup>2</sup>Actually there is need of two more things, if we want to make the connection between theory and experiment: either infrared safety or factorisation. That becomes discussed in the next chapter

This mechanism is called confinement. Confinement It has been confirmed in Lattice QCD, but not yet mathematically. And it belongs to nonperturbative theory. Quarks prefer to bind into hadrons which can be classified into baryons with three quarks state and mesons with a quark-antiquark state. As we know, the wave function of fermions must be antisymmetric according to the Pauli exclusion principle under the exchange of two quarks. Interestingly, there are resonance states with spin  $\frac{3}{2}$  like  $\Delta^{++}$ . The spins of the three up quarks are parallel to each other, have the same flavour and orbital angular momentum  $L=0$ . This means that an exchange of flavour, spin and space (orbital angular momentum) does not lead to any change. This problem is solved with the additional degree of freedom, the so-called color charge. Each quark comes in one of three colours red, green or blue and also anticolour  $\bar{r}, \bar{b}, \bar{g}$  for antiquarks. The hadrons are colour singlets with regard to the hypothesis and so are invariant under rotations in colour space. The colour hypothesis describes the existence of mesons with  $q\bar{q}$  and baryons with  $qqq$ . because if the wave function is odd in color, we have solved the spin statistical problem. The total wave function for each particle can be expressed in terms of:

$$\Psi_{3q} = \psi_{space} \times \chi_{spin} \times \theta_{colour} \times \phi_{flavour} \quad (1.3)$$

$$O(3) \quad SU(2) \quad SU(3) \quad SU(6)$$

Now we can compute all possible colour states with Young Tableaux [15]. One uses group theory methods, for instance the Young Tableaux technique, to decompose products of irreducible representations into sums.

$$\begin{array}{c}
 \boxed{3} \otimes \boxed{3} \otimes \boxed{3} = \boxed{\begin{smallmatrix} 3/3 & 4/2 & 5/1 \end{smallmatrix}} \oplus \boxed{\begin{smallmatrix} 3/3 & 4/1 \\ 2/1 \end{smallmatrix}} \oplus \boxed{\begin{smallmatrix} 3/3 & 4/1 \\ 2/1 \end{smallmatrix}} \oplus \boxed{\begin{smallmatrix} 3/3 \\ 2/2 \\ 1/1 \end{smallmatrix}} \\
 \text{Totally} \quad \text{Mixed} \quad \text{Mixed} \quad \text{Totally} \\
 \text{symmetric} \quad \text{symmetric} \quad \text{symmetric} \quad \text{antisymmetric} \\
 \\
 = 10 \oplus 8 \oplus 8 \oplus 1
 \end{array}$$

After using The same procedure for  $SU(2)$  and  $SU(6)$  for spins and flavours of the three quarks we will get:

$$\begin{aligned}
 2 \otimes 2 \otimes 2 &= 4 \oplus 2 \oplus 2 \oplus 0 \\
 6 \otimes 6 \otimes 6 &= 56 \oplus 70 \oplus 70 \oplus 20
 \end{aligned} \quad (1.4)$$

As we can see, the total wave function is most complicated in the QCD region. That is the reason why the Lagrangian of QCD is always given in the shortened form. Before


QCD is formulated as a gauge theory, an experiment should be pointed out which makes it clear why there is an additional degree of freedom in the QCD and why there is no U(1)-symmetry. Looking at the electron-positron scattering again, it is important to realize that not only  $\mu^+\mu^-$ , but also  $e^+e^- \rightarrow \tau^+\tau^-$  and also  $q\bar{q}$  can arise, when the quark pairs fragment into hadrons. For the ratio:

$$R = \frac{\sigma(e^+e^- \rightarrow \text{Hadrons})}{\sigma(e^+e^- \rightarrow \mu^+\mu^-)} \quad (1.5)$$

one would expect, due to the fact that the coupling takes place between two charged particles, that only the sum over the square of the quark charges (because  $e_\mu^2 = 1$ ) contributes. However, there is an additional factor  $N_C$  that can be determined experimentally

$$R = N_C \sum_q e_q^2 \quad (1.6)$$

Without this factor one would expect for  $u, d, s$ ,  $u, d, s, c$  and  $u, d, s, c, b$  Respectively  $\frac{2}{3}, \frac{10}{9}, \frac{11}{10}$ . The experiment showed a third of the expected results (i.e.  $N_C = 3$ ):



[25]

## 1.4 QCD Lagrangian

QCD like QED and the weak interaction theory is described by representations of a symmetry group. From the condition that the Lagrangian must be invariant under arbitrary global and local symmetry transformations (Noether's theorem) follows the interaction terms. The Lagrangian of QCD is invariant under  $U(3) = U(1) \times SU(3)$  global transformation. Here only  $SU(3)$  is discussed. The three Pauli matrices from  $SU(2)$  can be replaced by the eight Gell-Mann  $\lambda^a$  with using the following relation:

$$\begin{aligned} T^a &= \frac{1}{2}\lambda^a \\ [T^a, T^b] &= if^{abc}T^c && \text{fundamental representation} \\ (T^a_{adj})_{bc} &= -if^{abc} && \text{adjoint representation} \end{aligned} \quad (1.7)$$

To quantize QCD theory, the Faddeev-Popov method [11] is usually used in the path integral to fix a gauge and define a gluon propagator. The Lagrangian is given:

$$\begin{aligned} \mathcal{L} &= \mathcal{L}_{free} + \mathcal{L}_{int} \\ \mathcal{L} &= \sum_f \bar{\psi}_{if}(i\gamma^\mu \partial_\mu - m_f)\psi^{if} - \frac{1}{4}F_a^{\mu\nu}F_{\mu\nu}^a - \frac{1}{2\xi}(\partial^\mu A_\mu^a)(\partial^\nu A_\nu^a) + (\partial^\mu \chi^{a*})(\partial_\mu \chi^a) \\ &\quad - g_s \bar{\psi}_i T^a_{ij} \psi_j \gamma^\mu A_\mu^a - \frac{g_s}{2} f^{abc}(\partial_\mu A_\nu^a - \partial_\nu A_\mu^a)A_b^\mu A_c^\nu - \frac{g_s^2}{4} f^{abc}(A_b^\mu A_c^\nu) f^{ade}(A_\mu^d A_\nu^e) \\ &\quad - g_s f^{abc}(\partial^\mu \chi^{a*})\chi^b A_\mu^c \end{aligned} \quad (1.8)$$

Here  $i, j$  are color indices in the fundamental representation,  $a$  color index in the adjoint representation of  $SU(3)$ .  $f$  labels the six flavours of the quarks.  $g_s$  describes the strong coupling constant and  $A_\mu^a$  is the gluon field and it corresponds to a non-abelian gauge theory with structure constants  $f^{abc}$ .  $\chi^a$  is a scalar field under Lorentz group, but anti-commuting with the field strength tensor of QCD by [21, 26]

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g_s f_{abc} A_\mu^b A_\nu^c \quad (1.9)$$

It can be shown that the above Lagrangian is invariant under the following  $SU(3)$  gauge transformations:

$$\begin{aligned} \psi'(x) &\rightarrow \exp(i\eta_a(x) T^a) \psi(x) \\ D' &\rightarrow \partial_\mu + ig_s T_a A_\mu'^a \\ A_\mu'^a &\rightarrow A_\mu^a - \frac{1}{g_s} \partial_\mu \eta^a(x) + f^{abc} \eta_b(x) A_{c\mu}(x) \end{aligned} \quad (1.10)$$

$$L_{q,free} = \sum_{i, \alpha} \bar{\psi}_{i f} (i \gamma^\mu \partial_\mu - m_f) \delta_{ij} \psi^j f$$

$$\begin{array}{c} \bullet \longrightarrow \bullet \\ k, m_f \end{array} = \left( \frac{i}{\not{k} - m_f} \right)_{\alpha\beta} \delta_{ij}$$

$$L_{g,free} = -\frac{1}{4} F_a^{\mu\nu} F^a_{\mu\nu} - \frac{1}{2\xi} (\partial^\mu A^a_\mu) (\partial^\nu A^a_\nu)$$

$$\begin{array}{c} \bullet \text{-----} \bullet \\ k \end{array} = \frac{-i}{k^2} (-g_{\mu\nu} - (1 - \zeta) \frac{k_\mu k_\nu}{k^2}) \delta^{ab}$$

$$L_{ghost,free} = (\partial^\mu \chi^{a*}) (\partial_\mu \chi^a)$$

$$\begin{array}{c} \bullet \text{-----} \bullet \\ k \end{array} = \frac{i}{k^2} \delta^{ab}$$

## 1.5 Colour factor calculation

In this section we will calculate the Casimir operators of the respective diagrams which will be used later. The fundamental. representation in  $SU(3)$  are given by [24, 26]

$$T^a = \vartheta^a \equiv \frac{\lambda^2}{2} \quad \text{with Gell - Mann matrices } \lambda^a \quad (1.11)$$

$$\lambda^1 = \begin{pmatrix} 0 & 1 & \\ 1 & 0 & \\ & & 0 \end{pmatrix}, \quad \lambda^2 = \begin{pmatrix} 0 & -i & \\ i & 0 & \\ & & 0 \end{pmatrix}, \quad \lambda^3 = \begin{pmatrix} 1 & & \\ & -1 & \\ & & 0 \end{pmatrix}, \quad \lambda^4 = \begin{pmatrix} & & 1 \\ & 0 & \\ 1 & & \end{pmatrix}$$

$$\lambda^5 = \begin{pmatrix} & -i & \\ & 0 & \\ i & & \end{pmatrix}, \quad \lambda^6 = \begin{pmatrix} & & \\ & 0 & 1 \\ & 1 & 0 \end{pmatrix}, \quad \lambda^7 = \begin{pmatrix} & & \\ & 0 & -i \\ & i & 0 \end{pmatrix}, \quad \lambda^8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & & \\ & 1 & \\ & & -2 \end{pmatrix} \quad (1.12)$$

As we can see,  $\lambda^3$  and  $\lambda^8$  are diagonal. These generators satisfy:



$$L_{qg\bar{q},int} = g_s \bar{\psi}_i T^a_{ij} \psi_j \gamma^\mu A^a_\mu$$

$$= i g_s \gamma^\mu \otimes T^a_{ij}$$

$$L_{ggg,int} = -\frac{g_s}{2} f^{abc} (\partial_\mu A^a_\nu - \partial_\nu A^a_\mu) A_b^\mu A_c^\nu$$

$$= -g_s f^{bca} [(k_1 - k_2)^\rho g^{\mu\nu} + (k_2 - k_3)^\mu g^{\nu\rho} + (k_3 - k_1)^\nu g^{\rho\mu}]$$

(all Momenta are incoming!)

$$L_{gggg,int} = -\frac{g_s^2}{4} f^{abc} (A_b^\mu A_c^\nu) f^{ade} (A^d_\mu A^e_\nu)$$

$$= -i g_s^2 [f^{abe} f^{cde} (g^{\mu\rho} g^{\nu\sigma} - g^{\mu\sigma} g^{\nu\rho}) + f^{ade} f^{bce} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\sigma} g^{\nu\rho}) + f^{ade} f^{bce} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\rho} g^{\nu\sigma})]$$

$$L_{\chi g \bar{\chi}} = -g_s f^{abc} (\partial^\mu \chi^{a*}) \chi^b A^c_\mu$$

$$= g_s f^{abc} k^\mu$$

Or in the adjoint representation:

$$[T^a, T^b] = i f^{abc} T^c \Rightarrow T^a T^b - T^b T^a = i f^{abc} T^c$$



$$[F^a, F^b] = if^{abc} F^c \Rightarrow \text{diagram 1} - \text{diagram 2} = \text{diagram 3}$$

The most common convention for the normalization of the generators in physics is:

$$\sum_{c,d} f^{acd} f^{bcd} = N \delta^{ab} \quad (1.13)$$

One of the most important equations for the colour factor calculation is the Jaccobi-Identity:

$$[T^a, [T^b, T^c]] + [T^c, [T^a, T^b]] + [T^b, [T^c, T^a]] = 0 \quad (1.14)$$

Writing this in terms of the structure constant, it turns out:

$$f^{axd} f^{bcx} + f^{cxd} f^{abx} + f^{bxd} f^{cax} = 0 \quad (1.15)$$

So we are able to compute:

$$f^{abc} = -2i \operatorname{tr}(T^a [T^b, T^c]) \quad (1.16)$$

Which generalises to:

$$f^{abc} f^{xcd} = 4i \operatorname{tr}(T^a [T^b, [T^c, T^d]]) \quad (1.17)$$

With these relations we can calculate all Casimir operators:

Fundamental representation 3:

Adjoint representation 8:


$$\begin{array}{c} \text{diagram: horizontal line with two vertices, each connected to a loop} \\ T^a_{ik} \quad T^a_{kj} \end{array} \quad \sum_a (T^a T^a)_{ij} = C_F \delta_{ij} \Rightarrow C_F \rightarrow \text{arrow} \\ C_F = \frac{N_c^2 - 1}{2N_c} = C_F \sim \frac{N_c}{2}$$

$$\begin{array}{c} \text{diagram: horizontal line with two vertices, each connected to a loop, with a vertical dashed line through the center} \\ f^{axy} \quad f^{bxy} \end{array} \quad \sum_{xy} f^{axy} f^{bxy} = C_A \delta^{ab} \Rightarrow C_A \text{ loop} \\ C_A = N_C$$


Which means the charge of the gluon is twice that of the quark because:

$$C_A = N_c = 2C_F \sim 2\left(\frac{N_C}{2}\right) \quad (1.18)$$

Trace identities:



$$T_f \delta^{ab}$$




$$\text{tr}(T^a) = 0$$

One of the most important relation in this case is the Fierz identity. It shows the difference between QED and QCD!

$$\sum_a T_{ij}^a T_{kl}^a = \frac{1}{2}(\delta_{il}\delta_{kj} - \frac{1}{N}\delta_{ij}\delta_{kl}) \quad (1.19)$$

Graphically it means: The charge transfer in QED takes place along the Fermion line



because photons cannot transport charges. On the other hand, the gluons can transfer color charges because they have color charges themselves.

The main relation we will use later for SU(N):

$$\text{tr}(T^a T^b) = T_{ij}^a T_{ji}^b = T_F \delta^{ab} \quad (1.20)$$

$$\sum_a (T^a T^a) = C_F \delta^{ij} \quad (1.21)$$

$$f^{acd} f^{bcd} = C_A \delta^{ab} \quad (1.22)$$


With  $T_F = \frac{1}{2}$ ,  $C_A = N$  and  $C_F = \frac{N^2-1}{2N}$ .

# Chapter 2


## Catani Seymour

### 2.1 IR and Collinear Divergences

Beyond the LO (Leading order) diagrams it happens singularities. To discuss about these consider first the process  $e^-e^+ \rightarrow q\bar{q}g$



In order to calculate the cross section of this diagram, we have to consider the gluon emission from the antiquark. Since the calculation is quite long, we concentrate on the final result:



**Figure 2.1:** Left diagram  $e^-e^+ \rightarrow qq\bar{q}$  and right  $e^-e^+ \rightarrow q\bar{q}g$

$$\begin{aligned}
 A &= \frac{\bar{u}(k_1)(-ig_s\gamma^\nu \times T^a)[-i(\not{k}_1 + \not{k}_3)](-iee_q\gamma^\mu)v(k_2)\epsilon_\mu^{\lambda_1}\epsilon_\nu^{\lambda_2*}}{(k_1 + k_3)^2} \\
 &- \frac{\bar{u}(k_1)(-iee_q\gamma^\mu)[i(\not{k}_2 + \not{k}_3)](-ig_s\gamma^\nu \times T^a)v(k_2)\epsilon_\mu^{\lambda_1}\epsilon_\nu^{\lambda_2*}}{(k_1 + k_3)^2} \\
 \Rightarrow A &= -g_s T^a \left[ \frac{\bar{u} \not{\epsilon} (\not{k}_1 + \not{k}_3) \Gamma v}{(k_1 + k_3)^2} - \frac{\bar{u} \Gamma (\not{k}_2 + \not{k}_3) \not{\epsilon} v}{(k_2 + k_3)^2} \right] \quad \text{with } \Gamma = (-iee_q\gamma^\mu)\epsilon_\mu^{\lambda_1}
 \end{aligned} \tag{2.1}$$

Under consideration that the partons are on-shell, we get:

$$A = -g_s T^a \left[ \frac{\bar{u} \not{\epsilon} (\not{k}_1 + \not{k}_3) \Gamma v}{2k_1 \cdot k_3} - \frac{\bar{u} \Gamma (\not{k}_2 + \not{k}_3) \not{\epsilon} v}{2k_2 \cdot k_3} \right] \tag{2.2}$$

In the soft limit with  $k_0 \rightarrow 0$  we can factorize  $A_{soft}$  the amplitude in two parts:

$$A = -g_s T^a \left[ \frac{k_1 \cdot \epsilon}{k_1 \cdot k_3} - \frac{k_2 \cdot \epsilon}{k_2 \cdot k_3} \right] A_{born} \quad \text{with } A_{born} = \bar{u} \Gamma v \tag{2.3}$$

Which one contains all information about colour and momenta and  $A_{born}$  with all spin information. If one calculates the cross section for it, one gets:

$$\begin{aligned}
 A &= C_F g_s^2 \sigma^{born} \int \frac{d^3k}{2k_0(2\pi)^3} 2 \left( \frac{k_1 \cdot k_2}{(k_1 \cdot k_3)(k_2 \cdot k_3)} \right) \\
 &C_F g_s^2 \sigma^{born} \int d\cos\theta \frac{dk_0}{k_0} \frac{4}{(1 - \cos\theta)(1 + \cos\theta)}
 \end{aligned} \tag{2.4}$$

We define the energy fraction by:

$$x_i = \frac{2E_i}{\sqrt{s}} = \frac{2q \cdot k_i}{s} \tag{2.5}$$


One can show that  $\sum x_i = 2$  and thus, that only two of the  $x_i$  are independent. The final result is:

$$\frac{d^2\sigma}{dx_1 dx_2} = \left( \frac{4\pi\alpha}{s} \right) \sum e_i^2 \frac{2\alpha_s}{3\pi} \frac{x_1^2 + x_2^2}{(1 - x_1)(1 - x_2)} \tag{2.6}$$


There are three singularities in regard with the final result. If the emitted photon is collinear to the outgoing quark or anti-quark ( $x_1 \rightarrow 1$  or  $x_2 \rightarrow 1$ ) and When the emitted gluon is very soft ( $x_1 \rightarrow 1$  and  $x_2 \rightarrow 1$ ). The singularities come from the quark propagator in each diagram. The denominators contain according Feynmann rules terms with  $\sim \frac{1}{(k_i+k_j)^2}$ . We can eliminate the quark mass under on-shell condition so that:

$$\frac{1}{(k_i+k_j)^2} = \frac{1}{2k_i \cdot k_j} = \frac{1}{2E_i E_j (1 - \cos\theta_{ij})} = \frac{1}{s(1-x_k)} \quad (2.7)$$

One can show all possibilities for three partons through a triangle:



fortunately, According to KLN-Theorem, IR singularities must cancel when summing the transition rate over all degenerate (initial and final) states. The sum of the integrals  $int_R$  and  $int_V$  above is finite. However, this is not true for the individual contributions. The real part contains divergences due to soft and collinear radiation of massless particles. While  $M_{real}$  itself is a tree level amplitude and thus finite, the divergences show up upon integration over the phase space  $\int d\phi_3$ . In R V, the phase space is the same as for the Born amplitude, but the loop integrals contained in  $M_{virt}$  contain IR singularities. We will use deep inelastic scattering (DIS) to show how



the infra-red singularities are absorbed in the parton distributions [2].

## 2.2 Subtraction method

$$|A|^2 = |A^{(0)}_m|^2 + |A^{(0)}_{m+1}|^2 + 2Re(A^{(0)*}_m A^{(1)}_m) \quad (2.8)$$

Whereby,  $|A^{(0)}_m|^2$  is the tree level contribution (Born sector) from Lo and has no divergences,  $|A^{(0)}_{m+1}|^2 + 2Re(A^{(0)*}_m A^{(1)}_m)$  comes from the NLO and they are separately divergent. The problem in this case is that Integrations cannot be combined due to different phase space dimensions:

$$\sigma^{NLO} = \int_{m+1} \partial\sigma^R + \int_m \partial\sigma^V \quad (2.9)$$

The real and virtual contributions are both IR divergent and need to regularize in  $d = 4 - 2\epsilon$  dim. To tackle this problem one can use the subtraction method in that way one add and subtract a local counter term  $\partial\sigma^A$  with same singularity structure as terms  $\partial\sigma^R$  to the integral.  $\partial\sigma^A$  approximates soft and collinear singularities of  $\partial\sigma^R$ .

$$\sigma^{NLO} = \int_{m+1} [\partial\sigma^R - \partial\sigma^A] + \int_m [\partial\sigma^V + \int_1 \partial\sigma^A] \quad (2.10)$$

In this case, we can safely set  $\epsilon \rightarrow 0$  for  $\partial\sigma^R|_{\epsilon \rightarrow 0} - \partial\sigma^A|_{\epsilon \rightarrow 0}$  and run the integral numerically in 4-dim. On the other side, integrate over one-parton PS analytically explicitly cancel poles and then set  $\epsilon \rightarrow 0$ .

$$\sigma^{NLO} = \int_{m+1} [\partial\sigma^R|_{\epsilon \rightarrow 0} - \partial\sigma^A|_{\epsilon \rightarrow 0}] + \int_m [\partial\sigma^V + \int_1 \partial\sigma^A]_{\epsilon \rightarrow 0} \quad (2.11)$$

The virtual contribution must be UV-finite:

$$\int_m \partial\sigma^V = \int_m [\int_{loop} \partial\sigma^V_{bare} + \sigma^V_{Counter term}] \quad (2.12)$$

The addition of  $\int_1 \partial\sigma^A$  to the  $\int_m \partial\sigma^V$  ensures that IR poles are cancelled. The bare and counter contribution are separately divergent and have also different integral dimensions. One can use the same idea with the subtraction method to solve this problem [4, 5]

$$\int_m \partial\sigma^V + \int_{loop} \partial\sigma^L - \int_{loop} \partial\sigma^L = \int_m \int_{loop} [\partial\sigma^V_{bare} - \partial\sigma^L] + \int_m [\sigma^V_{Counter term} + \int_{loop} \partial\sigma^L] \quad (2.13)$$

$$\sigma^{NLO} = \int_{m+1} [\partial\sigma^R - \partial\sigma^A] + \int_m \int_{loop} [\partial\sigma^V_{bare} - \partial\sigma^L] + \int_m [\sigma^V_{Counter term} + \int_{loop} \partial\sigma^L + \partial\sigma^A] \quad (2.14)$$

## Determination of emission kernels

Now we want to introduce the properties of the counter term  $\partial\sigma_A$ . This term need to have the same behaviour like  $\partial\sigma_R$  in d dimension. This process and specific observable independent term has to be obtained in a way that is independent of the particular jet observable considered. It must be exactly integrable analytically over one-parton phase space in d dim and  $\partial\sigma_R - \partial\sigma_A$  has to be integrable via Monte Carlo methods.  $\partial\sigma_A$  acts as a local counter-term for  $\partial\sigma_B$ . At this point, one should derive improved factorization formulae which called dipole formulae. Note that the notation below is symbolic:

$$\partial\sigma_A = \sum_{dipoles} \partial\sigma_B \otimes \partial V_{dipoles} \quad (2.15)$$

Whereby, the sum is over dipoles for all  $m+1$  configurations with consideration to given m-parton state.  $\partial\sigma_B$  describes the color/spin projection of Born-level exclusive cross section. The symbol  $\otimes$  is for phase space convolutions and sums over colour and spin indices.  $\partial V_{dipoles}$  will compute just once for all and match the singular property of the real part. They are universal in the sense that they do not depend on the hard scattering. The good news is that we are now allowed to use a factorizable mapping from the  $m+1$ -parton phase space to an m-parton subspace. That will be more clearer when we will use the parametrisation in the next chapter. If we integrate over all  $m+1$ , we can write:

$$\int_{m+1} \partial\sigma_A = \int_m \partial\sigma_B \otimes \sum_{dipoles} \int_1 \partial V_{dipoles} \quad (2.16)$$

That's exactly the most important thing we got because so is distinguished between the known m-sector from LO and the second the universal factor which contains all  $\epsilon$ -poles.

## Singularity Structure

before we begin with the collinear limit or soft limit respectively we are going to pull up the matrix element from Lo which has this below general form:

$$\mathcal{M}_m^{c_1, \dots, c_m; s_1, \dots, s_m}(p_1, \dots, p_m) \quad (2.17)$$

$c_i$ ,  $s_i$  and  $p_i$  denotes respectively the colour, spin indices and the momenta for each m-parton in the tree level matrix element in the final state. A common way at the site is to define a basis in colour+helicity space.

$$\mathcal{M}_m^{c_1, \dots, c_m; s_1, \dots, s_m}(p_1, \dots, p_m) \equiv (\langle c_i, \dots, c_m | \otimes \langle s_1, \dots, s_m |) | 1, \dots, m \rangle_m \quad (2.18)$$

With  $\langle c_i, \dots, c_m | \otimes \langle s_1, \dots, s_m |$  as the basis and  $| 1, \dots, m \rangle_m$  as a vector in this space. Thus, for the matrix element squared:

$$\begin{aligned} |\mathcal{M}_m|^2 &= (\langle c_i, \dots, c_m | \otimes \langle s_1, \dots, s_m |) (| c_i, \dots, c_m \rangle \otimes | s_1, \dots, s_m \rangle) \langle 1, \dots, m | 1, \dots, m \rangle \\ &= \delta_{c_1 c_1} \dots \delta_{c_m c_m} \otimes \delta_{s_1 s_1} \dots \delta_{s_m s_m} \langle 1, \dots, m | 1, \dots, m \rangle_m \end{aligned} \quad (2.19)$$

Define a colour-charge operator  $T_i$  with the emission of a gluon from each parton i:

$$T_i = T_i^c |c\rangle \quad (2.20)$$

Its action onto the colour space is defined by:

$$\langle c_1, \dots, c_i, \dots, c_m, c | T_i | b_1, \dots, b_i, \dots, b_m \rangle = \delta_{c_1 b_1} \dots T_{c_i b_i}^c \dots \delta_{c_m b_m} \quad (2.21)$$

Where  $T_{c_i b_i}^c$  the colour-charge matrix in the adjoint representation in the case of gluon emission or colour-charge matrix in the fundamental representation for quark/antiquark emission case. The following properties must be taken into account for that:

$$\begin{aligned} T_i \cdot T_j &= T_j \cdot T_i && \text{if } i \neq j, \text{ commutative property} \\ T_i^2 &= C_i && C_i = C_A \text{ for gluon and } C_i = C_F \text{ for (anti)quark} \\ \sum_{i=1}^m T_i |1, \dots, m\rangle_m &= 0 && \text{for single state} \end{aligned} \quad (2.22)$$

Thus, the square of colour-correlated tree-amplitudes for the indices I, J refer either to final-state or initial-state partons will be [4, 5]

$$|\mathcal{M}_{m,a\dots}^{I,J}|^2 = \langle 1, \dots, m; a, \dots | T_I \cdot T_J | 1, \dots, m; a, \dots \rangle_{m,a\dots} \quad (2.23)$$

## Dipole factorisation

Consider (m+1)-partons with the general matrix element [5, 27]

$$|\mathcal{M}_{m+1}(Q; p_1, \dots, p_i, \dots, p_j, \dots, p_{m+1})|^2 \quad (2.24)$$

We need to take collinear and soft limits which allow factorisation. In the soft region it is used the kinematic with  $p_j \rightarrow \lambda q$ ,  $\lambda \rightarrow 0$ , where  $q$  is a arbitrary four vector and  $\lambda$  a scale parameter. The matrix element is characterise with  $|\mathcal{M}|^2 \sim \frac{1}{\lambda^2}$ . and if  $p_i$  and  $p_j$  become collinear, we parametrise  $p_j = \frac{z}{1-z} p_i$ . So the matrix element will be  $|\mathcal{M}|^2 \sim \frac{1}{p_i \cdot p_j}$ . I am going to go into more details in the next chapter. That's here should be a summery of the behaviour of the matrix elements in different regions. Based on the Catani-Seymour method for (m + 1) partons matrix element, it's possible to factorise out parton k to give  $|\mathcal{M}_m|^2$ :

$$|\mathcal{M}_{m+1}|^2 \rightarrow \sum |\mathcal{M}_m|^2 \otimes V_{ij,k} \Rightarrow |\mathcal{M}_{m+1}|^2 \rightarrow \sum |\mathcal{M}_m|^2 \otimes V_{ij,k} \quad (2.25)$$







$V_{ij,k}$  singular factor including parton k and it's interaction with partons i and j from the m parton amplitude. This situation can be represented by the bottom diagram.

Here i and k are the emitters and k plays the role of a spectator. The blobs denote the tree-level matrix elements and their complex conjugate. The dots on the right-hand side stand for non-singular terms both in the soft and collinear limits. When the partons i and j become soft and/or collinear, the singularities are factorized into the term  $V_{ij,k}$  (the dashed box on the right-hand side) which embodies correlations with a single additional parton k.



The diagram shows an equation between two multi-parton states. On the left, a state with  $m+1$  partons is shown as a circle with  $m+1$  external lines. A vertical dashed line separates the first parton (line 1) from the rest. The first parton is labeled 'non QCD partons' with a lightning bolt symbol. On the right, the same state is expressed as a sum over  $\{ij, k\}$  of a state with  $m$  partons. The  $m$ -parton state is a circle with  $m$  external lines, where the first parton is labeled  $i$ , the last is  $j$ , and the middle one is  $k$ . The sum is followed by a plus sign and an ellipsis. The reference [4, 5] is shown to the right.

The diagram shows an equation between two multi-parton states. On the left, a state with  $m+1$  partons is shown as a circle with  $m+1$  external lines. On the right, the same state is expressed as a sum over  $\{ij, k\}$  of a state with  $m$  partons. The  $m$ -parton state is a circle with  $m$  external lines, where the first parton is labeled  $i$ , the last is  $j$ , and the middle one is  $k$ . The sum is followed by a plus sign and an ellipsis.



In this context the different dipole factorisation for both initial states and final states shall be presented. All these different possibilities can be seen in the diagram below.

The circle in the center of each sub diagram presents the  $m$ -partons matrix element and the tilde labels the collinear splitting process for the initial or final states. For this work, the first upper diagram with final-state singularities without initial-state partons, is completely sufficient and is discussed here in detail with its formula. The matrix element

for this is written:

$$|\mathcal{M}_{m+1}|^2 = \langle 1, \dots, m+1 | 1, \dots, m+1 \rangle = \sum_{k \neq i, j} \mathcal{D}_{ij,k}(p_1, \dots, p_{m+1}) + \text{finite terms} \quad (2.26)$$

The first term with the sum over dipoles is divergent as  $p_i \cdot p_j \rightarrow 0$ . It needs to drop all finite terms. These dipole terms are explicitly given by:

$$\mathcal{D}_{ij,k}(p_1, \dots, p_{m+1}) = \frac{-1}{2p_i \cdot p_j} {}_m \langle 1, \dots, \tilde{i}j, \dots, k, \dots, m+1 | \frac{T_k \cdot T_{ij}}{T_{ij}^2} V_{ij,k} | 1, \dots, \tilde{i}j, \dots, k, \dots, m+1 \rangle_m \quad (2.27)$$

Where  $T_k \cdot T_{ij}$  are the color charges of spectator and emitter

$V_{ij,k}$  splitting kernel in helicity space of emitter explicit form depends on parton type become proportional to Altarelli-Parisi splitting functions and Eikonal factors in collinear and soft limits.

Using the kinematic:

$$\begin{aligned} \tilde{p}_{ij}^\mu &= p_i^\mu + p_j^\mu - \frac{y_{ij,k}}{1 - y_{ij,k}} p_k^\mu \\ \tilde{p}_k^\mu &= \frac{1}{1 - y_{ij,k}} p_k^\mu \\ \text{with } y_{ij,k} &= \frac{p_i \cdot p_j}{p_i \cdot p_j + p_j \cdot p_k + p_k \cdot p_i} \end{aligned} \quad (2.28)$$


Note, that momenta are on-shell. Due to momenta conservation  $p_i^\mu + p_j^\mu + p_k^\mu = \tilde{p}_k^\mu + \tilde{p}_{ij}^\mu$ .

## 2.3 Hard scattering

The hadron hadron scattering can be written as:

$$\sigma = \sum_{ij} \int dx_1 dx_2 f_i(x_1, \mu^2) f_j(x_2, \mu^2) \sigma_{ij}(x_1, x_2, Q^2/\mu^2 \dots) \quad (2.29)$$

Here the (arbitrary) factorisation scale  $\mu$  can be thought of as the scale which separates



the long and short-distance physics. Roughly speaking, a parton with a transverse momentum less than  $\mu$  is then considered to be part of the hadron structure and is absorbed in the parton distribution. Partons with larger transverse momenta participate in the hard scattering process with a short-distance partonic cross-section. The factorisation theorem also applies to deep inelastic scattering. It can be shown how the IR singularities are absorbed in the PDFs. The DIS cross section can be written as [9]

$$\frac{d^2\sigma}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} [(1-y)F_2(x, Q^2) + xy^2 F_1(x, Q^2)] \quad (2.30)$$

In this case we need to introduce the structure function, is defined as the charge weighted sum of the parton momentum densities, the probability that the parton carries a momentum fraction between  $x$  and  $\partial x$  of the proton momentum. The index  $i$  denotes the quark flavour. Parton distributions are nonperturbative and have not been obtained from experiment.

$$F_2^{exp}(x) = \sum_i e_i^2 x f_i(x) \quad (2.31)$$

The evolution of a parton distribution due to splitting is called the DGLAP evolution equation [28].

$$\frac{\partial f(x, \mu^2)}{\partial \ln \mu^2} = \frac{\alpha_s}{2\pi} \int_x^1 \frac{dy}{y} f_j(y, \mu^2) P_{ij}\left(\frac{x}{y}\right) + O(\alpha_s^2) \quad (2.32)$$


This is a system of coupled integral or differential equations.  $P_{ij}(\frac{x}{y})$  represents the probability that a daughter parton  $i$  with momentum fraction  $\frac{x}{y}$  splits from a parent parton  $j$ . The above convolution in compact notation (Mellin Convolution) in the general case:

$$\frac{\partial f_i(x, \mu^2)}{\partial \ln \mu^2} = \sum_{j=-n_f}^{n_f} P_{ij} \otimes f_j(\mu^2) \quad (2.33)$$

There four splitting possibilities according to below diagrams and the transitions lead to set of  $2n_f + 1$  coupled evolution equations.

$$\frac{\partial}{\partial \ln \mu^2} \begin{pmatrix} q_s \\ g \end{pmatrix} = \frac{\alpha_s}{2\pi} \begin{bmatrix} P_{qq} & 2n_f P_{qg} \\ P_{gq} & P_{gg} \end{bmatrix} \otimes \begin{pmatrix} q_s \\ g \end{pmatrix} \quad (2.34)$$

A parton distribution changes when a different parton splits or the parton itself splits. we can not compute parton densities, but we can predict how they evolve from one scale to another. we can measure PDFs in one process and use them as an input for another process.



$$\left. \begin{aligned} \langle \hat{P}_{qq} \rangle &= C_F \left[ \frac{1+z^2}{1-z} - \varepsilon(1-z) \right] \\ \langle \hat{P}_{gq} \rangle &= T_R \left[ 1 - \frac{2z(1-z)}{1-\varepsilon} \right] \\ \langle \hat{P}_{qg} \rangle &= C_F \left[ \frac{1+(1-z)^2}{z} - \varepsilon z \right] \\ \langle \hat{P}_{gg} \rangle &= 2C_A \left[ \frac{z}{1-z} + \frac{1-z}{z} + z(1-z) \right] \end{aligned} \right\} \text{splitting functions} \quad (2.35)$$

## 2.4 Mapping 3 partons to 2

As we have already seen in the last section, the dipole factorization is obtained from the square matrix element. Now we want to take a closer look at the four possible parton showers from the previous section. For this goal we first calculate the respective matrix elements and the complex-conjugated one of them in the known 3 parton evaluation. This results in the dipole-term, which contains the soft- and collinear singularities. Then this is parametrized with a certain kinematics in order to separate the finite terms from infinities. First of all, the Emmitter and the spectator are defined. After the substitution of the new momenta we get splitting kernel in helicity space and color charges of spectator and emitter. Then we ignore the finite terms because we are looking for the singular terms. It should be noted that at the beginning of this thesis a parametrisation was used, which unfortunately only works for LO. This was recognized later and therefore a new kinematics was used. This theoretical description will become clearer as we begin to implement the mappings. First of all the kinematics have to be introduced [13, 22, 23, 24].

## 2.5 Old mapping

First we start with the old mapping, as promised.

$$\left. \begin{aligned} q_i^\mu &= zp_i^\mu + y(1-z)p_j^\mu + \sqrt{zy(1-z)}m^\mu_\perp \\ q^\mu &= (1-z)p_i^\mu + yzp_j^\mu - \sqrt{zy(1-z)}m^\mu_\perp \\ q_j^\mu &= (1-y)p_j^\mu \end{aligned} \right\} \text{parametrisation} \quad (2.36)$$

Where  $q$  is the radiated soft momentum,  $q_i$  the momenta of the emitter and  $q_j$  the momentum of the spectator is. the dimensionless variable is given by  $y = \frac{q_i \cdot q}{p_i \cdot p_j}$ . Note that both the emitter and the spectator are on-shell. momentum conservation is implemented exactly:

$$q_i^\mu + q^\mu + q_j^\mu = p_i^\mu + p_j^\mu + m^\mu_\perp \quad (2.37)$$

For this mapping it is useful to calculate some common relation:

$$\begin{aligned} q_i + q &= p_i + yp_j \\ q_j + q &= (1-z)p_i^\mu + (1+yz-y)p_j^\mu - \sqrt{zy(1-z)}m^\mu_\perp \\ q_i \cdot q &= y(1-2z+2z^2)(p_i \cdot p_j) \\ q_i \cdot q_j &= z(1-y)(p_i \cdot p_j) \\ q_j \cdot q &= (1-z)(1-y)(p_i \cdot p_j) \end{aligned} \quad (2.38)$$

## 2.6 new kinematic

For the general  $m$  emission case it must be defined a new mapping. The parametrisation of the splitting momenta is formalized as:

$$\begin{aligned} k_l^\mu &= \alpha_l \alpha \Lambda^\mu{}_\nu p_i^\nu + y \beta n^\mu + \sqrt{y \alpha_l \beta_l} n_{\perp,l}^\mu & l = 1, \dots, m \\ q_i^\mu &= (1 - \sum_{l=1}^m \alpha_l) \alpha \Lambda^\mu{}_\nu p_i^\nu + y (1 - \sum_{l=1}^m \beta_l) n^\mu - \sqrt{y \alpha_l \beta_l} n_{\perp,l}^\mu \\ q_k^\mu &= \alpha \Lambda^\mu{}_\nu p_k^\nu & k = 1, \dots, n \quad k \neq i \end{aligned} \quad (2.39)$$

$k = 1, \dots, n$  labels the emission momenta and is taken to be massless  $k_l^2 = 0$ . Where the label  $l$  denotes the count of emissions. In this work we just want to considerate the one-emission kernels. The other important issue here is that all hard momenta are on-shell,  $p_k^2 = q_k^2 = 0$ .

$n^\mu$  is an auxiliary light-like vector which is necessary to specify the transverse component of  $n_{\perp,l}^\mu$ . To absorb the recoil we define  $n^\mu$  as:

$$n^\mu = Q^\mu - \frac{Q^2}{2p_i \cdot Q} p_i^\mu \quad (2.40)$$

Whereby  $Q$  is the total momentum with:

$$Q^\mu = q_i^\mu + \sum_{l=1}^m k_l^\mu + \sum_{k=1}^m q_k^\mu = p_i^\mu + \sum_{k=1}^m p_k^\mu \quad (2.41)$$

To fulfil the condition that the emission momenta are massless, we need the following condition:

$$\begin{aligned} n_{\perp,l}^\mu \Lambda^\mu{}_\nu p_i^\nu &= n_{\perp,l} \cdot n = n_{\perp,l} \cdot Q = 0 \\ n_{\perp,l}^\mu \cdot p_k &\neq 0 \end{aligned} \quad (2.42)$$

$n_{\perp,l}^2 = -2\alpha \Lambda^\mu{}_\nu p_i^\nu n_\mu$  is not on-shell and in terms of single emission case we get  $n_{\perp,1}^2 = -2p_i \cdot Q$ . The parameter  $y$  is related to the virtuality of the splitting parton:

$$q_i^\mu + \sum_{l=1}^m k_l^\mu = \alpha \Lambda^\mu{}_\nu p_i^\nu + y n^\mu \quad (2.43)$$

With  $\alpha = \sqrt{1-y}$ .

### Lorentz transformation of momenta

In order to be able to work with the parametrisation, we have to do the Lorentz transformation of the Emitters, Spectator and total momentum first.

$$\begin{aligned} \alpha \Lambda^\mu{}_\nu &= p_i^\mu p_{i\nu} \frac{-y^2 Q^2}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + p_i^\mu Q_\nu \frac{y(1 + \sqrt{1-y})}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \\ &+ Q^\mu p_{i\nu} \frac{(y^2 - y - y\sqrt{1-y})}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} \eta^\mu{}_\nu \end{aligned} \quad (2.44)$$

In the collinear limit of  $y \rightarrow 0, \alpha \rightarrow 1$  this transformation reduces to trivial  $\eta^\mu{}_\nu$ . Finally we are going to compute the Lorentz transformation of the Momenta. The detailed calculation of them can be found in Appendix A.

$$\boxed{\hat{p}_i^\mu = \alpha \Lambda^\mu{}_\nu p_i^\nu = p_i^\mu} \quad (2.45)$$

$$\boxed{\hat{Q}^\mu = \frac{Q^2}{2p_i \cdot Q} y p_i^\mu + (1-y) Q^\mu} \quad (2.46)$$

$$\boxed{\hat{p}_k^\mu = A_1 p_i^\mu + A_2 Q^\mu + \sqrt{1-y} p_k^\mu} \quad (2.47)$$

with

$$\begin{aligned} A_1 &\equiv \frac{-y^2 Q^2 (p_i \cdot p_k)}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + \frac{y(1 + \sqrt{1-y})(Q \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \\ A_2 &\equiv \frac{(y^2 - y - y\sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \end{aligned}$$

## 2.7 Single emission part

In terms of one emission where  $l = 1$  the mapping will be simplified as:

$$\begin{aligned} k_1^\mu &= (\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) p_i^\mu + y\beta_1 Q^\mu + \sqrt{y\alpha_1\beta_1} n_{\perp,1}^\mu \\ q_i^\mu &= (\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) p_i^\mu + y\alpha_1 Q^\mu - \sqrt{y\alpha_1\beta_1} n_{\perp,l}^\mu \\ q_k^\mu &= \alpha \Lambda^\mu{}_\nu p_k^\nu \quad k = 1, \dots, n \quad k \neq i \end{aligned} \quad (2.48)$$

$$\begin{aligned} k_1^\mu &= \zeta_1 p_i^\mu + \lambda_1 Q^\mu + \sqrt{y\alpha_1\beta_1} n_{\perp,1}^\mu \\ q_i^\mu &= \zeta_q p_i^\mu + \lambda_q Q^\mu - \sqrt{y\alpha_1\beta_1} n_{\perp,l}^\mu \\ q_k^\mu &= A_1 p_i^\mu + A_2 Q^\mu + \sqrt{1-y} p_k^\mu \end{aligned}$$

## 2.8 Common scalar products

To investigate the mapping it is useful to determine the dot products between these four vectors. To understand the often occurring pre-factor products one should look them up in the appendix A.

$$\boxed{k_1 \cdot q_i = y(\alpha_1 + \beta_1)^2 p_i \cdot Q = y p_i \cdot Q} \quad (2.49)$$

$$\boxed{k_1 \cdot q_k = [\alpha_1(1 - y) + y\beta_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\beta_1 Q \cdot p_k + \sqrt{\alpha_1\beta_1 y(1 - y)} p_k \cdot n_{\perp,1}} \quad (2.50)$$

$$\boxed{q_i \cdot q_k = [\beta_1(1 - y) + y\alpha_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\alpha_1 Q \cdot p_k - \sqrt{\alpha_1\beta_1 y(1 - y)} p_k \cdot n_{\perp,1}} \quad (2.51)$$

## 2.9 Recipe for the use of the new parametrisation

In the previous chapter we have discussed that the singularities come from the propagators in each diagram since the denominators contain according Feynmann rules terms with  $\sim \frac{1}{2q_a \cdot q_b}$ . Whereby a and b here place holder the respective momenta. Since the calculations are sometimes very complicated and confusing, the procedure for eliminating the finite terms is as follows:

In the calculating of the square matrix elements always appear products in the form of  $p_a \cdot p_b$  both in the numerator and denominator. The denominator shows which pre-factor causes the singularity. As we know, if , we get zero in the denominator. These terms from the numerator with the same prefix can be omitted from the beginning because they appear in both the denominator and the numerator and are therefore finite. This is explicitly shown below for two common denominators.

### 2.9.1 Parametrization in terms of $(k_1 \cdot q_i)(k_1 \cdot q_k)$

$$\boxed{(k_1 \cdot q_i)(k_1 \cdot q_k) \approx y(1 - \beta_1)(1 - y) (p_i \cdot p_k)(p_i \cdot Q)} \quad (2.52)$$

Here you can quickly see that this term converges for  $y \rightarrow 0$  and  $\beta_1 \rightarrow 1$  towards zero. That means, you could ignore all terms with  $y(1 - \beta_1)$ . However, since the equation becomes rather large quickly if we first use all the momenta products and then drop the terms with the pre-factor out of the denominator, this is already done for the scalar products. And this is exactly the biggest simplification in the calculation. The result



looks like this:

$$\begin{aligned}
k_1^\eta k_1^{\eta'} &= [(1 - \beta_1)^2 - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})^2] p_i^\eta p_i^{\eta'} - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q}) p_i^\eta Q^{\eta'} - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q}) Q^\eta p_i^{\eta'} \\
k_1^\eta q_i^{\eta'} &= [\beta_1(1 - \beta_1) - y \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})] p_i^\eta p_i^{\eta'} + y \beta_1^2 Q^\eta p_i^{\eta'} \\
q_i^\eta k_1^{\eta'} &= [\beta_1(1 - \beta_1) - y \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})] p_i^\eta p_i^{\eta'} + y \beta_1^2 p_i^\eta Q^{\eta'} \\
q_i^\eta q_i^{\eta'} &= \beta_1^2 p_i^\eta p_i^{\eta'} \\
k_1^\eta q_k^{\eta'} &= [(1 - \beta_1) - y \beta_1 (\frac{Q^2}{2p_i \cdot Q})] \sqrt{1 - y} p_i^\eta p_k^{\eta'} - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_1 p_i^\eta p_i^{\eta'} \\
&\quad - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_2 p_i^\eta Q^{\eta'} + y \beta_1 A_1 Q^\eta p_i^{\eta'} + y \beta_1 A_2 Q^\eta Q^{\eta'} + y \beta_1 \sqrt{1 - y} Q^\eta p_k^{\eta'} \\
q_i^\eta q_k^{\eta'} &= A_1 \beta_1 p_i^\eta p_i^{\eta'} + A_2 \beta_1 p_i^\eta Q^{\eta'} + \beta_1 \sqrt{1 - y} p_i^\eta p_k^{\eta'} \\
q_k^\eta k_1^{\eta'} &= [(1 - \beta_1) - y \beta_1 (\frac{Q^2}{2p_i \cdot Q})] \sqrt{1 - y} p_k^\eta p_i^{\eta'} - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_1 p_i^\eta p_i^{\eta'} \\
&\quad - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_2 Q^\eta p_i^{\eta'} + y \beta_1 A_1 p_i^\eta Q^{\eta'} + y \beta_1 A_2 Q^\eta Q^{\eta'} + y \beta_1 \sqrt{1 - y} p_k^\eta Q^{\eta'} \\
q_k^\eta q_i^{\eta'} &= A_1 \beta_1 p_i^\eta p_i^{\eta'} + A_2 \beta_1 Q^\eta p_i^{\eta'} + \beta_1 \sqrt{1 - y} p_k^\eta p_i^{\eta'}
\end{aligned} \tag{2.53}$$

### 2.9.2 Parametrization in terms of $(k_1 \cdot q_i)(k_1 \cdot q_i)$

$$\boxed{(k_1 \cdot q_i)(k_1 \cdot q_i) = y^2 (p_i \cdot Q)(p_i \cdot Q)} \tag{2.54}$$

With the same interpretation from above one could say that this term converges just for

$y \rightarrow 0$  towards zero. That's why we will remove all product terms with  $y^2$ .


$$\begin{aligned}
k_1^\eta k_1^{\eta'} &= [(1 - \beta_1)^2 - 2y\beta_1(\frac{Q^2}{2p_i \cdot Q})]p_i^\eta p_i^{\eta'} + y\beta_1(1 - \beta_1)(\frac{Q^2}{2p_i \cdot Q})p_i^\eta Q^{\eta'} \\
&\quad + y\beta_1(1 - \beta_1)(\frac{Q^2}{2p_i \cdot Q})Q^\eta p_i^{\eta'} \\
k_1^\eta q_i^{\eta'} &= [\beta_1(1 - \beta_1) - y(1 - \beta_1)^2(\frac{Q^2}{2p_i \cdot Q}) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})]p_i^\eta p_i^{\eta'} + y(1 - \beta_1)^2 Q^\eta p_i^{\eta'} \\
q_i^\eta k_1^{\eta'} &= [\beta_1(1 - \beta_1) - y(1 - \beta_1)^2(\frac{Q^2}{2p_i \cdot Q}) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})]p_i^\eta p_i^{\eta'} + y(1 - \beta_1)^2 p_i^\eta Q^{\eta'} \\
q_i^\eta q_i^{\eta'} &= [\beta_1^2 - 2y\beta_1(1 - \beta_1)(\frac{Q^2}{2p_i \cdot Q})]p_i^\eta p_i^{\eta'} + y\beta_1(1 - \beta_1)(\frac{Q^2}{2p_i \cdot Q})p_i^\eta Q^{\eta'} \\
&\quad + y\beta_1(1 - \beta_1)(\frac{Q^2}{2p_i \cdot Q})Q^\eta p_i^{\eta'} \\
k_1^\eta q_k^{\eta'} &= (1 - \beta_1)A_1 p_i^\eta p_i^{\eta'} + (1 - \beta_1)A_2 p_i^\eta Q^{\eta'} + (1 - \beta_1)\sqrt{1 - y}p_i^\eta p_k^{\eta'} \\
q_i^\eta q_k^{\eta'} &= A_1 \beta_1 p_i^\eta p_i^{\eta'} + A_2 \beta_1 p_i^\eta Q^{\eta'} + \beta_1 \sqrt{1 - y}p_i^\eta p_k^{\eta'} \\
q_k^\eta k_1^{\eta'} &= (1 - \beta_1)A_1 p_i^\eta p_i^{\eta'} + (1 - \beta_1)A_2 Q^\eta p_i^{\eta'} + (1 - \beta_1)\sqrt{1 - y}p_k^\eta p_i^{\eta'} \\
q_k^\eta q_i^{\eta'} &= A_1 \beta_1 p_i^\eta p_i^{\eta'} + A_2 \beta_1 Q^\eta p_i^{\eta'} + \beta_1 \sqrt{1 - y}p_k^\eta p_i^{\eta'}
\end{aligned} \tag{2.55}$$

## This is how we go about it

- First of all, let's take a look at a possible splitting. For this we have to make sure that all possible meaningful diagrams have been considered. All  $M_1$ ,  $M_2$ ,  $M_1^\dagger$  and  $M_2^\dagger$  diagrams are indexed independently of each other. When setting up the interference terms, however, we will cut out the respective results from the above-mentioned diagrams. To determine the matrix elements, we will use the Feynmann Rules from the appendix.
- In the next step you have to decide for an observer with  $q$ . It is necessary to take care that the observer must be selected so that when the diagrams of  $M_1$  and  $M_2$  or  $M_2$  and  $M_1$  are next to each other, you get a full but cut diagram so that you can use the cutting rules. Whereby we don't use cutting rules here, but separate diagrams. This point will be clarified in the next chapter.
- The square matrix element is a complex number and therefore:
- we have to keep all indices as they are defined at  $M_1$  and  $M_2$ . This is very important for the calculation of the interference term in order to get a meaningful result.
- In the case of the gluon loop, you have to add the corresponding ghost diagram.
- for all parts of the matrix element, use the parametrisation and the recipe for sending the kinematics from the last section. After the addition of all terms we get all singularities (soft and collinear).
- To determine whether everything was calculated correctly, one goes to the collinear limits, so to speak for  $y$ , because here one has to get the Alterali-Parisi splitting function known in the parton shower

## Chapter 3

### Quark anti-quark gluon emission kernel




First we are going to consider a daughter quark from the splitting of a parent quark into a quark and a gluon with an arbitrary spectator like an anti-quark, see the left picture above. where  $q_j + q$  is the momentum of the quark before splitting,  $q$  the momentum of the gluon and  $q_j$  of daughter quark respectively. The momentum of the spectator is  $q_i$ . The distinction between daughter and parent vanishes, when the gluon becomes soft, and a singularity develops. The other possibility to get a singularity is surely if the gluon will be collinear to quark. The splitting functions are flavour independent since the strong interaction is flavour independent. Furthermore, leading order splitting cannot change the flavour of a quark, thus we can write for the splitting functions In leading order QCD:

$$P_{\bar{q}_i \bar{q}_j} = P_{q_i q_j} = P_{qq} \delta_{ij} \quad (3.1)$$

For this aim we have to take any diagram to the account which can have the same splitting. Since there is no distinction between quark and anti-quark, one can imagine exact the same splitting variation for anti-quark with a quark as a spectator, see the right picture above.


### 3.1 Matrix element of a quark with a gluon radiation $|M_1|^2$



If one simply calculates the amplitude of this diagram, one gets:


$$M_1 = [\bar{u}_\sigma(q_i)(-ig_s\gamma^\mu \times [T^a]_o^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q)] [v_\tau(q_j)] \quad (3.2)$$

For the quadratic matrix element we need the dagger of  $M_1$  as well.



$$M_1^\dagger = [\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s\gamma^{\mu'} \times [T^b]_{o'}^k) u_{\sigma'}(q_i) \varepsilon^{\lambda_2}_{\mu'}(q)] [\bar{v}_{\tau'}(q_j)] \quad (3.3)$$

After multiplying  $M_1^\dagger$  and  $M_1$  we get the desired result.



$$|M_1|^2 = M_1 M_1^\dagger = [\bar{u}_\sigma(q_i) (-ig_s \gamma^\mu \times [T^a]_{o'}^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q)] [v_\tau(q_j)]$$

$$[\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s \gamma^{\mu'} \times [T^b]_{o'}^k) u_{\sigma'}(q_i) \varepsilon^{\lambda_2}_{\mu'}^*(q)] [\bar{v}_{\tau'}(q_j)] \quad (3.4)$$

Now it's time to connect those terms which are related to each other.

$$|M_1|^2 = [\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s \gamma^{\mu'} \times [T^b]_{o'}^k) \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) \varepsilon^{\lambda_2*}_{\mu'}(q) \varepsilon^{\lambda_1}_\mu(q)$$

$$\times (-ig_s \gamma^\mu \times [T^a]_{o'}^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2}] [\bar{v}_{\tau'}(q_j) v_\tau(q_j)] \quad (3.5)$$

Sum over the lorenz index  $(\sigma, \sigma')$  and  $(\tau, \tau')$  and spin addition relation leads to:

$$\sum_{\sigma, \sigma'} \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) = \not{q}_i \delta^{\sigma\sigma'},$$

$$\sum_{\tau, \tau'} \bar{v}_\tau(q_j) v_{\tau'}(q_j) = \not{q}_j \delta^{\tau\tau'} \quad (3.6)$$

Sum over polarization index  $(\lambda_1, \lambda_2)$  :

$$\sum_{\mu, \mu'} \varepsilon^{\lambda_2*}_{\mu'}(q) \varepsilon^{\lambda_1}_\mu(q) = -g_{\mu\mu'} \delta^{ab} \quad (3.7)$$

The matrix element will be simplified with:

$$|M_1|^2 = \frac{-g_s^2 [T^a]_{o'}^k [T^a]_{o'}^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \gamma^{\mu'} \not{q}_i g_{\mu'\mu} \gamma^\mu (\not{q}_i + q)] [\not{q}_j] \quad (3.8)$$

When we contract all related indices we will be actually able to make a statements about the last result.

$$|M_1|^2 = \frac{-g_s^2 [T^a]_{o'}^k [T^a]_{o'}^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \gamma^{\mu'} \not{q}_i \gamma_{\mu'} (\not{q}_i + q)] [\not{q}_j] \quad (3.9)$$

In other words we expect the tree level diagram from LO and a number:

Which graphically means:

$$|M^2| = \left| \begin{array}{c} \text{diagram with two shaded circles and momenta } P_i, P_j \\ \text{contribution from LO} \end{array} \right|^2 \otimes \left| \begin{array}{c} \text{diagram with a loop and momenta } q_i, q, q_i+q \\ \text{a complex number} \end{array} \right|^2$$

$$|M_1|^2 = \frac{-g_s^2 [T^a]_o^k [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [P_i][P_j] \otimes (\text{a complex number}) \quad (3.10)$$

Let's calculate the contribution and compare the final result with this expectation:

$$\begin{aligned} N &= : \gamma^{\mu'} \not{q}_i \gamma_{\mu'} = q_{i\sigma} \gamma^{\mu'} \gamma^\sigma \gamma_{\mu'} \\ &= q_{i\sigma} (\{\gamma^{\mu'}, \gamma^\sigma\} - \gamma^\sigma \gamma^{\mu'}) \gamma_{\mu'} \\ &= q_{i\sigma} 2g^{\mu'\sigma} \gamma_{\mu'} - d \gamma^\sigma \\ &= (2 - d) \not{q}_i \end{aligned} \quad (3.11)$$

Simplification of the bracket:

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \not{q}_i (\not{q}_i + q)] [\not{q}_j] \quad (3.12)$$

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [\not{q}_i \not{q}_i \not{q}_i + \not{q}_i \not{q}_i \not{q} + \not{q} \not{q}_i \not{q}_i + \not{q} \not{q}_i \not{q}] [\not{q}_j] \quad (3.13)$$

Momenta are on-shell, so:

$$\begin{aligned} \not{q}_i \not{q}_i &= q_i^2 = m_i^2 \\ \not{q} \not{q} &= q^2 = m^2 \\ \not{q}_j \not{q}_j &= q_j^2 = m_j^2 \end{aligned} \quad (3.14)$$

we can first neglect the mass of patrons:

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{(2q_i q)(2q_i q)} [\not{q} \not{q}_i \not{q}] [\not{q}_j] \quad (3.15)$$

Here we need to make the terms in the brackets simpler and:

$$\begin{aligned}
L &= \not{q}_i \not{q} \not{q} = \not{q}[q_{i\sigma}q_\mu (\{\gamma^\mu, \gamma^\sigma\} - \gamma^\sigma \gamma^\mu)] \\
&= \not{q}[2q_i^\mu q_\mu - q_{i\sigma}q_\mu \gamma^\mu \gamma^\sigma] \\
&= \not{q}(2q_i q) - q_\mu q_{i\sigma}q_\mu [\gamma^\mu \gamma^\mu \gamma^\sigma] \\
&= \not{q}(2q_i q) - q_\mu q_{i\sigma}q_\mu \left[ \frac{\gamma^\mu \gamma^\mu}{2} + \frac{\gamma^\mu \gamma^\mu}{2} \right] \gamma^\sigma \\
&= \not{q}(2q_i q) - q_\mu q_{i\sigma}q_\mu [g^{\mu\mu}] \gamma^\sigma \\
&= \not{q}(2q_i q) - q_\mu q_{i\sigma}q^\mu \gamma^\sigma = \not{q}(2q_i q) - q^2 \not{q}_i \\
&= \not{q}(2q_i q)
\end{aligned} \tag{3.16}$$

After inserting the last result of  $L$  and simplify the term  $(2q_i q)$  from the denominator and nominator, we get:

$$|M_1|^2 = -(2-d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{q}][\not{q}_j] \tag{3.17}$$

Now we are going to use the parametrisation from equation (1) to reduce the 3-member matrix element to 2-member and take out the singularity term from the amplitude.

$$|M_1|^2 = (d-2) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [(1-z) \not{p}_i + zy \not{p}_j - \sqrt{zy(1-z)} \not{m}_\perp][(1-y) \not{p}_j] \tag{3.18}$$

Multiplying the both sides

$$\begin{aligned}
|M_1|^2 &= (d-2) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [(1-z)(1-y) \not{p}_i \not{p}_j \\
&\quad + zy(1-y) \not{p}_j \not{p}_j + (1-y)\sqrt{zy(1-z)} \not{m}_\perp \not{p}_j]
\end{aligned} \tag{3.19}$$

Under consideration of the fact that  $p_i$  and  $p_j$  are the on-shell momenta of the emitter and spectator partons, we can ignore the terms with  $\not{p}_i \not{p}_i$  and  $\not{p}_j \not{p}_j$ . The  $p_i \cdot m_\perp$  and  $p_j \cdot m_\perp$  are always 0 because the  $p_i$  and  $p_j$  are lightlike, i.e. zero transverse component. So those terms can be neglected.


$$|M_1|^2 = \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \otimes \frac{(d-2)(1-z)(1-y)}{2y(1-2z+2z^2)} \tag{3.20}$$

As discussed, we get a contribution from the LO a complex number. As you can see, the number is just for  $y \rightarrow 0$  singular and not for  $z \rightarrow 1$ .



### 3.2 Matrix element of an anti-quark with a gluon radiation $|M_2|^2$

the same procedure is used to obtain the matrix element for an anti-quark with a single gluon emission.



$$|M_2|^2 = M_2 M_2^\dagger = \left[ \frac{i(\not{q}_j + \not{q})}{(q_j + q)^2} (-ig_s \gamma^\nu \times [T^c]_f^m) v_\tau(q_j) \varepsilon^{\lambda_3}_\nu(q) [u_\sigma(q_i)] \right. \\ \left. [\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q) [\bar{u}_{\sigma'}(q_i)] \right] \quad (3.21)$$

$$|M_2|^2 = \frac{g_s^2 [T^c]_f^m [T^d]_{f'}^n}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\nu v_\tau(q_j) \bar{v}_{\tau'}(q_j) \varepsilon^{\lambda_3}_\nu(q) \varepsilon^{\lambda_4}_{\nu'}(q) \gamma^{\nu'} (\not{q}_j + \not{q})] \\ [u_\sigma(q_i)] [\bar{u}_{\sigma'}(q_i)] \quad (3.22)$$

and after sum over the lorenz and polarization indexes like  $(\sigma, \sigma')$ ,  $(\tau, \tau')$  and  $(\lambda_3, \lambda_4)$  as well and using the spin addition relation:

$$|M_2|^2 = \frac{g_s^2 [T^c]_f^m [T^c]_f^n}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\nu \not{q}_j (-g_{\nu\nu'}) \gamma^{\nu'} (\not{q}_j + \not{q})] [\not{q}_i] \quad (3.23)$$

Analogous to the last calculation from the previous section:

$$|M_2|^2 = (d - 2) \frac{g_s^2 [T^c]_f^m [T^c]_f^n}{(2qq_j)} [\not{q}] [\not{q}_i] \quad (3.24)$$


finally, we achieve:

$$|M_2|^2 = \frac{g_s^2 [T^c]_f^m [T^c]_f^n}{(p_i \cdot p_j)} [\not{p}_i] [\not{p}_j] \otimes \frac{-(d - 2)yz^2}{2(1 - z)(1 - y)} \quad (3.25)$$

Interestingly, here is a term with  $y$  concerning the gluon radiation from an anti-quark. This means that this result cannot contribute to the collinear limit for soft gluon  $y \rightarrow 0$ .

### 3.3 Interference contribution

So far most of the work is done and we just have to put the results of  $M_1$  and  $M_2^\dagger$  next to each other, as we can see in the diagram. So we still get the interference contribution. Using the results from the previous section and we received:



$$M_1 M_2^\dagger = [\bar{u}_\sigma(q_i) (-ig_s \gamma^\mu \times [T^a]_o^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q)] [v_\tau(q_j)] \quad (3.26)$$

$$[\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q)] [u_{\sigma'}(q_i)]$$

Expectation:

$$|M^2| = \left| \begin{array}{c} \text{Diagram with two shaded circles connected by two horizontal lines labeled } P_i \text{ and } P_j \\ \text{contribution from LO} \end{array} \right|^2 \otimes \left| \begin{array}{c} \text{Diagram with a shaded circle and a gluon line} \\ \text{a complex number} \end{array} \right|^2$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} [-(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu + 2(\not{q}_i + \not{q}) q_i^\mu]$$

$$[-(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu + 2(\not{q}_j + \not{q}) q_{j\mu}]$$

$$\Rightarrow M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} [(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu] [(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu]$$

$$- 2[(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu] [(\not{q}_j + \not{q}) q_{j\mu}] - 2[(\not{q}_i + \not{q}) q_i^\mu] [(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu]$$

$$+ 4[(\not{q}_i + \not{q}) q_i^\mu] [(\not{q}_j + \not{q}) q_{j\mu}] \quad (3.27)$$

The middle terms disappear due to the influence of the gamma matrices if the sequence of the matrices is reversed, so that the two terms become the same, because  $AB = -BA$ :

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} [(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu] [(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu] + 4[(\not{q}_i + \not{q}) q_i^\mu][(\not{q}_j + \not{q}) q_{j\mu}] \quad (3.28)$$

Now we use the old parametrization to collect the singularities.

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{4(1-z)(1-y)y(1-2z+2z^2)(p_i \cdot p_j)(p_i \cdot p_j)} [y(1-2z+2z^2) \not{p}_i \not{p}_j \gamma^\mu] [(1-z)(1-y) \not{p}_i \not{p}_j \gamma_\mu] + 4(p_i \cdot p_j)[(\not{p}_i + y \not{p}_j)][(1-z) \not{p}_i + (1+yz-y) \not{p}_j - \sqrt{zy(1-z)} \not{n}] \quad (3.29)$$

Here we can use the singular term in the denominator  $y(1-z)$  to drop the term with the same pre-factor and thus obtain:

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \otimes \frac{z}{1-z} \quad (3.30)$$

### 3.4 Final result

One could assume that for a complete result the contribution  $M_1^\dagger M_2$  is still missing.

$$|M|^2 = |M_1|^2 + |M_2|^2 + M_1 M_2^\dagger + M_1^\dagger M_2 \quad (3.31)$$

It should be noted that it is completely sufficient to calculate  $M_1 M_2^\dagger$ , because we know

it from the quadratic amount of the complex numbers, we can calculate double of real part of  $2RE(M_1 M_2^\dagger)$  instead of  $M_1 M_2^\dagger + M_1^\dagger M_2$  and that is exactly what is preferred here.

$$|M|^2 = |M_1|^2 + |M_2|^2 + 2RE(M_1 M_2^\dagger) \quad (3.32)$$

Let's just add up the results from the previous sections and get:

$$\begin{aligned} |M|^2 &= (d-2)(1-z)(1-y) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ &\quad - (d-2)yz^2 \frac{g_s^2 [T^c]_f^m [T^c]_f^n}{2(1-z)(1-y)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ &\quad + 2RE\left(\left(\frac{-2z}{z-1}\right) \frac{g_s^2 [T^a]_o^l [T^a]_f^n}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j]\right) \end{aligned} \quad (3.33)$$

Now we use the knowledge from the introduction about the calculation of the Colour factor. With Fritz equation:

$$T^a_{ok} T^a_{lo} = \frac{1}{2}(\delta_{oo}\delta_{lk} - \frac{1}{N}\delta_{ok}\delta_{lo}) = \frac{1}{2}(N\delta_{lk} - \frac{1}{N}\delta_{lk}) = C_F\delta_{lk} \quad (3.34)$$

After summation over the final colour states and averaging over initial colour states we get:

$$T^a_{ok} T^a_{lo} = C_F\delta_{lk} = \frac{1}{N} \sum_{l=1}^N \delta_{lk} C_F = C_F \quad (3.35)$$

The same calculation for  $T^c_{mf} T^c_{fn}$  and  $T^a_{ol} T^a_{fn}$  turns  $C_F$  out as the colour factor. Now we are going to compute the splitting function in the case of the colinearity, wich means, if:

$$y \longrightarrow 0 \quad (3.36)$$

$$|M|^2 = \frac{g_s^2 C_F}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \otimes ((d-2)(1-z) - \frac{4z}{z-1}) \quad (3.37)$$

for  $d = 4 - 2\epsilon$

$$\begin{aligned} |M|^2 &= \frac{g_s^2}{y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \otimes C_F \left( \frac{(1+z^2)}{1-z} - \epsilon(1-z) \right) \\ &= \frac{g_s^2}{q_i \cdot q} [\not{p}_i][\not{p}_j] \otimes \langle \hat{P}_{qq} \rangle \end{aligned} \quad (3.38)$$

With Alterali-Parisi splitting function  $\langle \hat{P}_{qq} \rangle$  in the collinear limes, which was mentioned in the previous chapter. This is exactly the confirmation of our calculation that our calculation was actually performed correctly, otherwise we would not have received the same splitting function for soft gluons.

### 3.5 Double-check the results with the new kinematic

One could do exactly the same calculation for the new kinematics to see if you get the same result in the collinear limit. From the next chapter we will explicitly work with the new parametrisation, because we found that the old kinematics only work in NLO and one-single emission.

$$|M_1|^2$$

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{(2k_1 \cdot q_i)} [k_1] [\not{q}_k] \quad (3.39)$$

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{2y p_i \cdot Q} [(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\beta_1 \not{Q} + \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1}] [A_1 \not{p}_i + A_2 \not{Q} + \sqrt{1-y} \not{p}_k] \quad (3.40)$$

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{2y p_i \cdot Q} [(A_2(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) + A_1 y\beta_1) p_i \cdot Q + (\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \sqrt{1-y} p_i \cdot p_k + A_2 y\beta_1 Q^2 + \sqrt{1-y} \sqrt{y\alpha_1\beta_1} n_{\perp,1} \cdot p_k] \quad (3.41)$$

For the collinearity  $y \rightarrow 0$  we'll get:

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{2y p_i \cdot Q} [(A_2(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) + A_1 y\beta_1) \not{p}_i \not{Q} + (\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \sqrt{1-y} \not{p}_i \not{p}_k + A_2 y\beta_1 Q^2 + \sqrt{1-y} \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1} \not{p}_k] \quad (3.42)$$

$$|M_1|^2 = (d-2)(1-\beta_1) \sqrt{1-y} \frac{g_s^2 C_F}{2y p_i \cdot Q} [\not{p}_i \not{p}_k] \quad (3.43)$$

$$|M_2|^2$$

$$|M_2|^2 = (d-2) \frac{g_s^2 C_F}{2k_1 \cdot q_k} [k_1] [\not{q}_i] \quad (3.44)$$

$$|M_2|^2 = (d-2) \frac{g_s^2 C_F}{2k_1 \cdot q_k} [(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\beta_1 \not{Q} + \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1}] [(\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\alpha_1 \not{Q} - \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1}] \quad (3.45)$$

Which means:

$$|M_2|^2 \sim (d-2) \frac{g_s^2 C_F}{2k_1 \cdot q_k} y [\dots] \quad (3.46)$$

$$|M_2|^2 \rightarrow 0 \quad \text{for } y \rightarrow 0$$

$M_1 M_2^\dagger$ 

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{4y(1-\beta_1)(1-y) (p_i \cdot p_k)(p_i \cdot Q)} \quad (3.47)$$


$$4(\beta_1\sqrt{1-y}p_i \cdot p_k)[\beta_1\sqrt{1-y} \not{p}_i \not{p}_k + (1-\beta_1)\sqrt{1-y} \not{p}_i \not{p}_k]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{y(1-\beta_1) (p_i \cdot p_k)(p_i \cdot Q)} \beta_1(p_i \cdot p_k)[\beta_1 \not{p}_i \not{p}_k + (1-\beta_1) \not{p}_i \not{p}_k] \quad (3.48)$$

$$M_1 M_2^\dagger = \frac{\beta_1}{(1-\beta_1)} \frac{-g_s^2 C_F}{y (p_i \cdot Q)} [\not{p}_i \not{p}_k] \quad (3.49)$$

## Chapter 4


### Gluon radiation from a parent quark



Now consider a daughter gluon from the splitting of a parent gluon with radiation an another gluon. When the gluon becomes soft, the distinction between daughter and parent vanishes, and a singularity develops. In this chapter we are going to keep the same procedure with a difference that we won't use the old parametrisation since it only works in LO. One of the mainly challenges about this emission kernel is that the calculations are long and complicated. Before we get started, it must be mentioned that we will work here with the recipe from section 3.6, in which the respective mathematical tools were calculated in detail.




## 4.1 Gluon-Emitter Bubble



If we start with the first diagram and use the necessary Feynmann rules, we get:

$$M_1 = \left[ \frac{-i}{(q_i + q)^2} (-g_s f^{a o l} (g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q + q_i)^\mu + g^{\eta \mu} (2q_i + q)^\zeta) \right. \\ \left. \varepsilon^{\lambda_1}_\mu(q_i) \varepsilon^{\lambda_6}_\zeta(q) \right] [\varepsilon^{\theta}_{\tau'}(q_j)] \quad (4.1)$$


It has to be emphasized that here only the impulse from the parent gluon is incoming and the two further impulses from the vertex are pointing out. This is why the minus signs appear in the equation. If the upper equation is daggered, you get the following diagram and the corresponding following amplitude:



$$M_1^\dagger = \left[ \frac{i}{(q_i + q)^2} (-g_s f^{a' \kappa o'} (-g^{\mu' \eta'} (2q_i + q)^{\zeta'} + g^{\eta' \zeta'} (2q + q_i)^{\mu'} + g^{\zeta' \mu'} (q_i - q)^{\eta'}) \right. \\ \left. \varepsilon^{\lambda_2}_{\mu'}(q_i) \varepsilon^{\lambda_5}_{\zeta'}(q) \right] [\varepsilon^{\theta'}_{\tau'}(q_j)] \quad (4.2)$$

Let us now calculate the matrix element, in which we place the results from above next

to each other, it turns out:



$$\begin{aligned}
 |M_1|^2 = & \left[ \frac{-i}{(q_i + q)^2} (-g_s f^{a o l} (g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q + q_i)^\mu + g^{\eta \mu} (2q_i + q)^\zeta) \right. \\
 & \varepsilon^{\lambda_1}_\mu(q_i) \varepsilon^{\lambda_2}_{\mu'}(q_i) \varepsilon^{\lambda_6}_\zeta(q) \varepsilon^{\lambda_5}_{\zeta'}(q) (-g_s f^{a' k o'} (-g^{\mu' \eta'} (2q_i + q)^{\zeta'} + g^{\eta' \zeta'} (2q + q_i)^{\mu'}) \\
 & \left. + g^{\zeta' \mu'} (q_i - q)^{\eta'}) \frac{i}{(q_i + q)^2} \right] [g^{\gamma \delta}] \quad (4.3)
 \end{aligned}$$

After the summation over the spin- just as well polarization indices can be obtained:

$$|M_1|^2 = \frac{g_s^2 f^{a o l} f^{a k o}}{(q_i + q)^2 (q_i + q)^2} [N^{\eta \eta'}] [g^{\gamma \delta}] \quad (4.4)$$

The tensor  $N^{\eta \eta'}$  appears exactly when you contract across all possible indices.

The goal of the next step is to evaluate this tensor, which is avoided here and instead of that, it only presents the final result. The more detailed calculation can be found in Appendix A.


$$\begin{aligned}
 N^{\eta \eta'} \equiv & [(6 - d) q^\eta q^{\eta'} + (d + 3) q^\eta q_i^\eta q_i^{\eta'} + (d + 3) q_i^\eta q^{\eta'} + (6 - d) q_i^\eta q_i^{\eta'} \\
 & - g^{\eta \eta'} (5q^2 + 5q_i^2 + 8q q_i)] [g^{\gamma \delta}] \quad (4.5)
 \end{aligned}$$

Replace this result in the equation

$$\begin{aligned}
 |M_1|^2 = & \frac{g_s^2 f^{a o l} f^{a k o}}{(q_i + q)^2 (q_i + q)^2} [(6 - d) q^\eta q^{\eta'} + (d + 3) q^\eta q_i^\eta q_i^{\eta'} + (d + 3) q_i^\eta q^{\eta'} + (6 - d) q_i^\eta q_i^{\eta'} \\
 & - g^{\eta \eta'} (5q^2 + 5q_i^2 + 8q q_i)] [g^{\gamma \delta}] \quad (4.6)
 \end{aligned}$$

This Gluon self-energy diagram has to be corrected by Ghost Loop to get the complete result. That is exactly what we are going to do in the next section.

### 4.1.1 One-loop corrections to the gluon self-energy diagram(Gluon-Emitter Bubble)



In order to get a meaningful result and correct the gluon loop, the same indices must be used. For this the same diagram with a fine difference is used, in which you replace the cut off gluon propagators with ghost propagators and the rest remain as in the previous diagram.

$$|M_1|_{Ghost\ loop}^2 = \frac{g_s^2 f^{aol} f^{ako}}{(q_i + q)^2 (q_i + q)^2} [-q_i^\eta q^{\eta'} - q^\eta q_i^{\eta'}] [g^{\gamma\delta}] \quad (4.7)$$

$$|M_1'|^2 = |M_1|^2 + |M_1|_{Ghost\ loop}^2 = \frac{g_s^2 f^{aol} f^{ako}}{(q_i + q)^2 (q_i + q)^2} [(6 - d)q^\eta q^{\eta'} + (d + 3)q^\eta q_i^{\eta'} + (d + 3)q_i^\eta q^{\eta'} + (6 - d)q_i^\eta q_i^{\eta'} - g^{\eta\eta'} (5q^2 + 5q_i^2 + 8qq_i) - q_i^\eta q^{\eta'} - q^\eta q_i^{\eta'}] [g^{\gamma\delta}] \quad (4.8)$$

After simplification and addition over the same terms:

$$|M_1'|^2 = \frac{g_s^2 f^{aol} f^{ako}}{(q_i + q)^2 (q_i + q)^2} [(6 - d)q^\eta q^{\eta'} + (d + 2)q^\eta q_i^{\eta'} + (d + 2)q_i^\eta q^{\eta'} + (6 - d)q_i^\eta q_i^{\eta'} - g^{\eta\eta'} (8qq_i)] [g^{\gamma\delta}] \quad (4.9)$$

Implementation the new parametrisation:

$$|M_1'|^2 = \frac{g_s^2 f^{aol} f^{ako}}{4y^2(\alpha_1 + \beta_1)^2 (p_i \cdot Q) (p_i \cdot Q)} [(6 - d)(\zeta_1 p_i^\eta + \lambda_1 Q^\eta + \sqrt{y\alpha_1\beta_1} n_{\perp,1}^\eta)(\zeta_1 p_i^{\eta'} + \lambda_1 Q^{\eta'} + \sqrt{y\alpha_1\beta_1} n_{\perp,1}^{\eta'}) + (d + 2)(\zeta_1 p_i^\eta + \lambda_1 Q^\eta + \sqrt{y\alpha_1\beta_1} n_{\perp,1}^\eta)(\zeta_q p_i^{\eta'} + \lambda_q Q^{\eta'} - \sqrt{y\alpha_1\beta_1} n_{\perp,1}^{\eta'}) + (d + 2)(\zeta_q p_i^\eta + \lambda_q Q^\eta - \sqrt{y\alpha_1\beta_1} n_{\perp,1}^\eta)(\zeta_1 p_i^{\eta'} + \lambda_1 Q^{\eta'} + \sqrt{y\alpha_1\beta_1} n_{\perp,1}^{\eta'}) + (6 - d)(\zeta_q p_i^\eta + \lambda_q Q^\eta - \sqrt{y\alpha_1\beta_1} n_{\perp,1}^\eta)(\zeta_q p_i^{\eta'} + \lambda_q Q^{\eta'} - \sqrt{y\alpha_1\beta_1} n_{\perp,1}^{\eta'}) - 8g^{\eta\eta'} [(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))n_{\perp,1} \cdot n_{\perp,1}]] [g^{\gamma\delta}] \quad (4.10)$$

Note that here the short version of the kinematics was used to increase the overview. Now you have to multiply the terms in brackets and simplify the matrix element.

$$\begin{aligned}
 |M'_1|^2 = & \frac{g_s^2 f^{aol} f^{ako}}{y^2 (p_i \cdot Q) (p_i \cdot Q)} [(6-d)[\zeta_1 \zeta_1 p_i^\eta p_i^{\eta'} + \zeta_1 \lambda_1 p_i^\eta Q^{\eta'} + \zeta_1 \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} \\
 & + \lambda_1 \zeta_1 Q^\eta p_i^{\eta'} + \lambda_1 \lambda_1 Q^\eta Q^{\eta'} + \lambda_1 \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} + \zeta_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} + \lambda_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} \\
 & + \sqrt{y\alpha_1 \beta_1} \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} n^{\eta'}_{\perp,1}] [(d+2)[\zeta_1 \zeta_q p_i^\eta p_i^{\eta'} + \zeta_1 \lambda_q p_i^\eta Q^{\eta'} - \zeta_1 \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} \\
 & + \lambda_1 \zeta_q Q^\eta p_i^{\eta'} + \lambda_1 \lambda_q Q^\eta Q^{\eta'} - \lambda_1 \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} + \zeta_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} + \lambda_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} \\
 & - \sqrt{y\alpha_1 \beta_1} \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} n^{\eta'}_{\perp,1}] [(d+2)[\zeta_q \zeta_1 p_i^\eta p_i^{\eta'} + \zeta_q \lambda_1 p_i^\eta Q^{\eta'} + \zeta_q \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} \\
 & + \lambda_q \zeta_1 Q^\eta p_i^{\eta'} + \lambda_q \lambda_1 Q^\eta Q^{\eta'} + \lambda_q \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} - \zeta_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} - \lambda_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} \\
 & - \sqrt{y\alpha_1 \beta_1} \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} n^{\eta'}_{\perp,1}] [(6-d)[\zeta_q \zeta_q p_i^\eta p_i^{\eta'} + \zeta_q \lambda_q p_i^\eta Q^{\eta'} - \zeta_q \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} \\
 & + \lambda_q \zeta_q Q^\eta p_i^{\eta'} + \lambda_q \lambda_q Q^\eta Q^{\eta'} - \lambda_q \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} - \zeta_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} - \lambda_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} \\
 & + \sqrt{y\alpha_1 \beta_1} \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} n^{\eta'}_{\perp,1} - 8g^{\eta\eta'} [(\alpha_1^2 + \beta_1^2) p_i \cdot Q - (\beta_1(1 - \beta_1)) n_{\perp,1} \cdot n_{\perp,1}]] [g^{\gamma\delta}]
 \end{aligned} \tag{4.11}$$

$$\begin{aligned}
 |M'_1|^2 = & \frac{g_s^2 f^{aol} f^{ako}}{4y^2 (p_i \cdot Q) (p_i \cdot Q)} [(6-d)\{(\alpha_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q})) p_i^\eta p_i^{\eta'} + y\alpha_1\beta_1 p_i^\eta Q^{\eta'} \\
 & + \zeta_1 \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} + y\beta_1\alpha_1 Q^\eta p_i^{\eta'} + \lambda_1 \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} + \zeta_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} \\
 & + \lambda_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} + y\alpha_1\beta_1 n^\eta_{\perp,1} n^{\eta'}_{\perp,1}\} + (d+2)\{(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q})) p_i^\eta p_i^{\eta'} \\
 & + y\alpha_1^2 p_i^\eta Q^{\eta'} - \zeta_1 \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} + y\beta_1^2 Q^\eta p_i^{\eta'} - \lambda_1 \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} + \zeta_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} \\
 & + \lambda_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} - y\alpha_1\beta_1 n^\eta_{\perp,1} n^{\eta'}_{\perp,1}\} \\
 & + (d+2)\{(\beta_1\alpha_1 - y(\beta_1^2 + \alpha_1^2)(\frac{Q^2}{2p_i \cdot Q})) p_i^\eta p_i^{\eta'} + y\beta_1^2 p_i^\eta Q^{\eta'} + \zeta_q \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} \\
 & + y\alpha_1^2 Q^\eta p_i^{\eta'} + \lambda_q \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} - \zeta_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} - \lambda_1 \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} \\
 & - y\alpha_1\beta_1 n^\eta_{\perp,1} n^{\eta'}_{\perp,1}\} + (6-d)\{(\beta_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q})) p_i^\eta p_i^{\eta'} + y\beta_1\alpha_1 p_i^\eta Q^{\eta'} \\
 & - \zeta_q \sqrt{y\alpha_1 \beta_1} p_i^\eta n^{\eta'}_{\perp,1} + y\alpha_1\beta_1 Q^\eta p_i^{\eta'} - \lambda_q \sqrt{y\alpha_1 \beta_1} Q^\eta n^{\eta'}_{\perp,1} - \zeta_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} p_i^{\eta'} \\
 & - \lambda_q \sqrt{y\alpha_1 \beta_1} n^\eta_{\perp,1} Q^{\eta'} + y\alpha_1\beta_1 n^\eta_{\perp,1} n^{\eta'}_{\perp,1}\} - 8g^{\eta\eta'} [(\alpha_1^2 + \beta_1^2) p_i \cdot Q - (\beta_1(1 - \beta_1)) n_{\perp,1} \cdot n_{\perp,1}]] [g^{\gamma\delta}]
 \end{aligned} \tag{4.12}$$

$$\begin{aligned}
|M'_1|^2 &= \frac{g_s^2 f^{aol} f^{ak}}{4y^2 (p_i \cdot Q) (p_i \cdot Q)} \\
&[(6-d)\{(\alpha_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1\beta_1 p_i^\eta Q^{\eta'} + y\beta_1\alpha_1 Q^\eta p_i^{\eta'} \\
&\quad + y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
&\quad + (d+2)\{(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1^2 p_i^\eta Q^{\eta'} + y\beta_1^2 Q^\eta p_i^{\eta'} \\
&\quad - y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'}\} + (d+2)\{(\beta_1\alpha_1 - y(\beta_1^2 + \alpha_1^2)(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\beta_1^2 p_i^\eta Q^{\eta'} \\
&\quad + y\alpha_1^2 Q^\eta p_i^{\eta'} - y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'}\} + (6-d)\{(\beta_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} \\
&\quad + y\beta_1\alpha_1 p_i^\eta Q^{\eta'} + y\alpha_1\beta_1 Q^\eta p_i^{\eta'} + y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'}\} - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))n_{\perp,1} \cdot n_{\perp,1}]] \\
&\quad (4.13)
\end{aligned}$$

$$\begin{aligned}
|M'_1|^2 &= \frac{g_s^2 f^{aol} f^{ako}}{4y^2 (p_i \cdot Q) (p_i \cdot Q)} \\
&[(6-d)(\alpha_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q})) + 2(d+2)(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q})) \\
&\quad + (6-d)(\beta_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q}))]p_i^\eta p_i^{\eta'} \\
&\quad + [2(6-d)y\alpha_1\beta_1 + (d+2)y(\alpha_1^2 + \beta_1^2)]p_i^\eta Q^{\eta'} \\
&\quad + [2(6-d)y\beta_1\alpha_1 + (d+2)y(\alpha_1^2 + \beta_1^2)]Q^\eta p_i^{\eta'} \\
&\quad + [2(6-d) - 2(d+2)]y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))n_{\perp,1} \cdot n_{\perp,1}][g^{\gamma\delta}] \\
&\quad (4.14)
\end{aligned}$$

$$\begin{aligned}
|M'_1|^2 &= \frac{g_s^2 f^{aol} f^{ako}}{4y^2 (p_i \cdot Q) (p_i \cdot Q)} \\
&[(6-d)(\alpha_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q})) + 2(d+2)(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q})) \\
&\quad + (6-d)(\beta_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q}))]p_i^\eta p_i^{\eta'} \\
&\quad + y[(4d-8)\alpha_1^2 + (8-4d)\alpha_1 + (d+2)]p_i^\eta Q^{\eta'} \\
&\quad + y[(4d-8)\alpha_1^2 + (8-4d)\alpha_1 + (d+2)]Q^\eta p_i^{\eta'} \\
&\quad + y[8-4d](\alpha_1 - \alpha_1^2)n_{\perp,1}^\eta n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))n_{\perp,1} \cdot n_{\perp,1}][g^{\gamma\delta}] \\
&\quad (4.15)
\end{aligned}$$

$$\begin{aligned}
|M'_1|^2 &= \frac{g_s^2 f^{aol} f^{ako}}{4y (p_i \cdot Q) (p_i \cdot Q)} \\
&[[8-4d]\beta_1(1 - \beta_1)n_{\perp,1}^\eta n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))n_{\perp,1} \cdot n_{\perp,1}][g^{\gamma\delta}]] \\
&\quad (4.16)
\end{aligned}$$

$$|M'_1|^2 = \frac{g_s^2 f^{aol} f^{ako}}{4y (p_i \cdot Q) (p_i \cdot Q)} [8[\epsilon - 1]\beta_1(1 - \beta_1)n_{\perp,1}^{\eta}n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))(-2p_i \cdot Q)][g^{\gamma\delta}]] \quad (4.17)$$

$$|M'_1|^2 = \frac{g_s^2 f^{aol} f^{ako}}{4y (p_i \cdot Q) (p_i \cdot Q)} [8[\epsilon - 1]\beta_1(1 - \beta_1)n_{\perp,1}^{\eta}n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q + 2\alpha_1\beta_1p_i \cdot Q]][g^{\gamma\delta}] \quad (4.18)$$

$$|M'_1|^2 = \frac{g_s^2 f^{aol} f^{ako}}{4y (p_i \cdot Q) (p_i \cdot Q)} [8[\epsilon - 1]\beta_1(1 - \beta_1)n_{\perp,1}^{\eta}n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}[(\alpha_1 + \beta_1)^2p_i \cdot Q]][g^{\gamma\delta}] \quad (4.19)$$

$$|M'_1|^2 = \frac{g_s^2 f^{aol} f^{ako}}{y (p_i \cdot Q)} [2[\epsilon - 1]\beta_1(1 - \beta_1)n_{\perp,1}^{\eta}n_{\perp,1}^{\eta'} - 2g^{\eta\eta'}][g^{\gamma\delta}] \quad (4.20)$$

Another way:

$$\begin{aligned}
k_1^\eta k_1^{\eta'} &= (\alpha_1^2 - 2\alpha_1\beta_1 y(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1\beta_1 p_i^\eta Q^{\eta'} + y\alpha_1\beta_1 Q^\eta p_i^{\eta'} + y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
k_1^\eta q_i^{\eta'} &= (\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1^2 p_i^\eta Q^{\eta'} + y\beta_1^2 Q^\eta p_i^{\eta'} - y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
q_i^\eta k_1^{\eta'} &= (\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\beta_1^2 p_i^\eta Q^{\eta'} + y\alpha_1^2 Q^\eta p_i^{\eta'} - y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
q_i^\eta q_i^{\eta'} &= (\beta_1^2 - 2\alpha_1\beta_1 y(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1\beta_1 p_i^\eta Q^{\eta'} + y\alpha_1\beta_1 Q^\eta p_i^{\eta'} + y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'}
\end{aligned} \tag{4.21}$$

$$\begin{aligned}
N &\equiv (6-d)(\alpha_1^2 - 2\alpha_1\beta_1 y(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1\beta_1 p_i^\eta Q^{\eta'} + y\alpha_1\beta_1 Q^\eta p_i^{\eta'} + y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
&\quad + (d+2)(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1^2 p_i^\eta Q^{\eta'} + y\beta_1^2 Q^\eta p_i^{\eta'} - y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
&\quad + (d+2)(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\beta_1^2 p_i^\eta Q^{\eta'} + y\alpha_1^2 Q^\eta p_i^{\eta'} - y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
&\quad + (6-d)(\beta_1^2 - 2\alpha_1\beta_1 y(\frac{Q^2}{2p_i \cdot Q}))p_i^\eta p_i^{\eta'} + y\alpha_1\beta_1 p_i^\eta Q^{\eta'} + y\alpha_1\beta_1 Q^\eta p_i^{\eta'} + y\alpha_1\beta_1 n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
&\quad - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))n_{\perp,1} \cdot n_{\perp,1}]
\end{aligned} \tag{4.22}$$


$$\begin{aligned}
N &\equiv [(6-d)(\alpha_1^2 - 2\alpha_1\beta_1 y(\frac{Q^2}{2p_i \cdot Q})) + (d+2)(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q})) \\
&\quad + (d+2)(\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q})) + (6-d)(\beta_1^2 - 2\alpha_1\beta_1 y(\frac{Q^2}{2p_i \cdot Q}))]p_i^\eta p_i^{\eta'} \\
&\quad + [(6-d)y\alpha_1\beta_1 + (d+2)y\alpha_1^2 + (d+2)y\beta_1^2 + (6-d)y\alpha_1\beta_1]p_i^\eta Q^{\eta'} \\
&\quad + [(6-d)y\alpha_1\beta_1 + (d+2)y\beta_1^2 + (d+2)y\alpha_1^2 + (6-d)y\alpha_1\beta_1]Q^\eta p_i^{\eta'} \\
&\quad + [(6-d)y\alpha_1\beta_1 - (d+2)y\alpha_1\beta_1 - (d+2)y\alpha_1\beta_1 + (6-d)y\alpha_1\beta_1]n_{\perp,1}^\eta n_{\perp,1}^{\eta'} \\
&\quad - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - (\beta_1(1 - \beta_1))n_{\perp,1} \cdot n_{\perp,1}]
\end{aligned} \tag{4.23}$$

$$\begin{aligned}
|M_1'|^2 &= \frac{g_s^2 f^{aol} f^{ako}}{4y(p_i \cdot Q)^2} [(12-2d)y\alpha_1\beta_1 - 2(d+2)y\alpha_1\beta_1]n_{\perp,1}^\eta n_{\perp,1}^{\eta'} - 8yg^{\eta\eta'} p_i \cdot Q [g_{\gamma\delta}] \\
\Rightarrow |M_1'|^2 &= \frac{g_s^2 f^{aol} f^{ako}}{4y(p_i \cdot Q)^2} [(12-2d)\alpha_1\beta_1 - 2(d+2)\alpha_1\beta_1]n_{\perp,1}^\eta n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}(\alpha_1^2 + \beta_1^2)p_i \cdot Q [g_{\gamma\delta}] \\
|M_1'|^2 &= \frac{g_s^2 f^{aol} f^{ako}}{4y(p_i \cdot Q)(p_i \cdot Q)} \\
&[8[\epsilon-1]\beta_1(1-\beta_1)n_{\perp,1}^\eta n_{\perp,1}^{\eta'} - 8g^{\eta\eta'}[(\alpha_1^2 + \beta_1^2)p_i \cdot Q - \beta_1\alpha_1(-2p_i \cdot Q)]] [g_{\gamma\delta}]
\end{aligned} \tag{4.24}$$

$$|M'_1|^2 = \frac{g_s^2 f^{a o l} f^{a k o}}{y(p_i \cdot Q)} [2[\epsilon - 1]\beta_1(1 - \beta_1)n_{\perp,1}^{\eta} n_{\perp,1}^{\eta'} - 2g^{\eta\eta'}][g^{\gamma\delta}] \quad (4.25)$$



## 4.2 Gluon-Spectator Bubble




$$|M_2|^2 = \left[ \frac{-i}{(q_j + q)^2} (-g_s f^{b f m} (g^{\tau\gamma} (-2q_j - q)^\rho + g^{\gamma\rho} (2q + q_j)^\tau + g^{\rho\tau} (q_j - q)^\gamma) \right. \\ \left. g_{\tau\tau'} g_{\rho\rho'} (-g_s f^{b' n f'} (g^{\rho'\delta} (-2q - q_j)^{\tau'} + g^{\delta\tau'} (2q_j + q)^{\rho'} + g^{\tau'\rho'} (q - q_j)^\delta) \frac{i}{(q_j + q)^2} [g^{\eta\eta'}] \right] \quad (4.26)$$

$$|M_2|^2 = \frac{g_s^2 f^{b f m} f^{b' n f'} \delta^{aa'} \delta^{ff'} \delta^{bb'}}{(q_j + q)^2 (q_j + q)^2} [g_{\tau\tau'} g_{\rho\rho'} (g^{\tau\gamma} (2q_j + q)^\rho g^{\rho'\delta} (2q + q_j)^{\tau'} \\ - g^{\tau\gamma} (2q_j + q)^\rho g^{\delta\tau'} (2q_j + q)^{\rho'} - g^{\tau\gamma} (2q_j + q)^\rho g^{\tau'\rho'} (q - q_j)^\delta - g^{\gamma\rho} (2q + q_j)^\tau g^{\rho'\delta} (2q + q_j)^{\tau'} \\ + g^{\gamma\rho} (2q + q_j)^\tau g^{\delta\tau'} (2q_j + q)^{\rho'} + g^{\gamma\rho} (2q + q_j)^\tau g^{\tau'\rho'} (q - q_j)^\delta - g^{\rho\tau} (q_j - q)^\gamma g^{\rho'\delta} (2q + q_j)^{\tau'} \\ + g^{\rho\tau} (q_j - q)^\gamma g^{\delta\tau'} (2q_j + q)^{\rho'} + g^{\rho\tau} (q_j - q)^\gamma g^{\tau'\rho'} (q - q_j)^\delta) [g^{\eta\eta'}] \quad (4.27)$$

$$|M_2|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [(2q + q_j)^\gamma (2q_j + q)^\delta \\ - g^{\delta\gamma} (2q_j + q)^\rho (2q_j + q)_\rho - (2q_j + q)^\gamma (q - q_j)^\delta - g^{\delta\gamma} (2q + q_j)^\tau (2q + q_j)_\tau \\ + (2q_j + q)^\gamma (2q + q_j)^\delta + (2q + q_j)^\gamma (q - q_j)^\delta - (q_j - q)^\gamma (2q + q_j)^\delta \\ + (q_j - q)^\gamma (2q_j + q)^\delta + d(q_j - q)^\gamma (q - q_j)^\delta] [g^{\eta\eta'}] \quad (4.28)$$

$$|M_2|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [(3 + d) q^\gamma q_j^\delta + (6 - d) q^\gamma q^\delta \\ + (6 - d) q_j^\gamma q_j^\delta + (3 + d) q_j^\gamma q^\delta - g^{\delta\gamma} (5q_j^2 + 5q^2 + 8qq_j) \\ ] [g^{\eta\eta'}] \quad (4.29)$$

### 4.2.1 One-loop corrections to the gluon self-energy diagram (Gluon-Spectator Bubble)



$$|M_2|_{Ghost\ loop}^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [-q_j^\gamma q^\delta - q^\delta q_j^\gamma] [g^{\eta\eta'}] \quad (4.30)$$

$$|M_2'|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [(2 + d)q^\gamma q_j^\delta + (6 - d)q^\gamma q^\delta + (6 - d)q_j^\gamma q_j^\delta + (2 + d)q_j^\gamma q^\delta - g^{\delta\gamma}(8qq_j)] [g^{\eta\eta'}] \quad (4.31)$$


$$|M_2'|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{4(q_j \cdot q)(q_j \cdot q)} [-8g^{\delta\gamma}(q \cdot q_j)] [g^{\eta\eta'}] \quad (4.32)$$

$$|M_2'|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j \cdot q)} [-2g^{\delta\gamma}] [g^{\eta\eta'}] \quad (4.33)$$

$$|M_2'|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(1 - \beta_1)(1 - y)(p_i \cdot p_k)} [-2g^{\delta\gamma}] [g^{\eta\eta'}] \quad (4.34)$$

### 4.3 Interference term $M_1 M_2^\dagger$

$$\begin{aligned} M_1 M_2^\dagger = & \left[ \frac{-i}{(q_i + q)^2} (-g_s f^{l a o} (g^{\eta\mu} (2q_i + q)^\zeta + g^{\mu\zeta} (q - q_i)^\eta - g^{\zeta\eta} (2q + q_i)^\mu) \varepsilon^{\lambda_1}_\mu(q_i) \varepsilon^{\lambda_6}_\zeta(q)) \right. \\ & \left. [\varepsilon^{\theta^*}_{\tau'}(q_j)] \right. \\ & \left[ \frac{i}{(q + q_j)^2} (-g_s f^{f' b' n} (g^{\tau'\rho'} (q_j - q)^\delta + g^{\rho'\delta} (2q + q_j)^{\tau'} - g^{\delta\tau'} (2q_j + q)^{\rho'}) \varepsilon^{\theta'}_{\tau'}(q_j) \varepsilon^{\lambda_8}_{\rho'}(q)) \right. \\ & \left. [\varepsilon^{\lambda_2}_{\mu'}(q_i)] \right] \end{aligned} \quad (4.35)$$



$$M_1 M_2^\dagger = \frac{g_s^2 f^{l a o} f^{f' b' n} \delta^{a a'} \delta^{o b'} \delta^{f f'}}{(q_i + q)^2 (q_j + q)^2} [g_{\mu}^{\eta'} g_{\tau \tau'} (g^{\eta \mu} (2q_i + q)^\zeta + g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q + q_i)^\mu) \\ g_{\zeta \rho'} (g^{\tau' \rho'} (q_j - q)^\delta + g^{\rho' \delta} (2q + q_j)^{\tau'} - g^{\delta \tau'} (2q_j + q)^{\rho'})] \quad (4.36)$$

$$M_1 M_2^\dagger = \frac{g_s^2 f^{l a o} f^{f' b' n} \delta^{a a'} \delta^{o b'} \delta^{f f'}}{(q_i + q)^2 (q_j + q)^2} [g^{\eta \mu} (2q_i + q)^\gamma (q_j - q)^\delta + g^{\eta \mu} (2q + q_j)^\gamma (2q_i + q)^\delta - g^{\eta \mu} g^{\gamma \delta} (2q_i + q) \cdot (2q_j + q) \\ + g^{\eta \mu} (q - q_i)^\eta (q_j - q)^\delta + g^{\eta \mu} (q - q_i)^\eta (2q + q_j)^\gamma - g^{\gamma \delta} (q - q_i)^\eta (2q_j + q)^{\eta'} \\ - g^{\gamma \eta} (2q + q_i)^{\eta'} (q_j - q)^\delta - g^{\eta \delta} (2q + q_i)^{\eta'} (2q + q_j)^\gamma + g^{\gamma \delta} (2q_j + q)^\eta (2q + q_i)^{\eta'}] \quad (4.37)$$

$$M_1 M_2^\dagger = \frac{g_s^2 f^{l a o} f^{f o n}}{4(q \cdot q_i)(q \cdot q_j)} \{g^{\eta \mu} [2q_i^\gamma q_j^\delta + 2q_i^\gamma q^\delta + q^\gamma q_j^\delta + q^\gamma q^\delta + 4q^\gamma q_i^\delta + 2q^\gamma q^\delta + 2q_j^\gamma q_i^\delta + q_j^\gamma q^\delta] \\ - g^{\eta \mu} g^{\gamma \delta} (2q \cdot q_j + q \cdot q + 4q_i \cdot q_j + 2q_i \cdot q) + g^{\eta \mu} [q^\eta q_j^\delta - q^\eta q^\delta - q_i^\eta q_j^\delta + q_i^\eta q^\delta] \\ + g^{\eta \mu} [2q^\eta q^\gamma + q^\eta q_j^\gamma + q_i^\eta q^\gamma + q_i^\eta q_j^\gamma] - g^{\gamma \delta} [2q^\eta q_j^{\eta'} + q^\eta q^{\eta'} - 2q_i^\eta q_j^{\eta'} - q_i^\eta q^{\eta'}] \\ - g^{\gamma \eta} [2q^{\eta'} q_j^\delta - 2q^{\eta'} q^\delta + q_i^{\eta'} q_j^\delta - q_i^{\eta'} q^\delta] - g^{\eta \delta} [4q^{\eta'} q^\gamma + 2q^{\eta'} q_j^\gamma + 2q_i^{\eta'} q^\gamma + q_i^{\eta'} q_j^\gamma] \\ + g^{\gamma \delta} [4q_j^\eta q^{\eta'} + 2q_j^\eta q_i^{\eta'} + q^\eta q^{\eta'} + q^\eta q_i^{\eta'}]\} \quad (4.38)$$

$$\begin{aligned}
k_1^\eta k_1^{\eta'} &= [(1 - \beta_1)^2 - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})^2] p_i^\eta p_i^{\eta'} - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q}) p_i^\eta Q^{\eta'} - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q}) Q^\eta p_i^{\eta'} \\
k_1^\eta q_i^{\eta'} &= [\beta_1(1 - \beta_1) - y \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})] p_i^\eta p_i^{\eta'} + y \beta_1^2 Q^\eta p_i^{\eta'} \\
q_i^\eta k_1^{\eta'} &= [\beta_1(1 - \beta_1) - y \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})] p_i^\eta p_i^{\eta'} + y \beta_1^2 p_i^\eta Q^{\eta'} \\
q_i^\eta q_i^{\eta'} &= \beta_1^2 p_i^\eta p_i^{\eta'} \\
k_1^\eta q_k^{\eta'} &= [(1 - \beta_1) - y \beta_1 (\frac{Q^2}{2p_i \cdot Q})] \sqrt{1 - y} p_i^\eta p_k^{\eta'} - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_1 p_i^\eta p_i^{\eta'} - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_2 p_i^\eta Q^{\eta'} \\
&\quad + y \beta_1 A_1 Q^\eta p_i^{\eta'} + y \beta_1 A_2 Q^\eta Q^{\eta'} + y \beta_1 \sqrt{1 - y} Q^\eta p_k^{\eta'} \\
q_i^\eta q_k^{\eta'} &= A_1 \beta_1 p_i^\eta p_i^{\eta'} + A_2 \beta_1 p_i^\eta Q^{\eta'} + \beta_1 \sqrt{1 - y} p_i^\eta p_k^{\eta'} \\
q_k^\eta k_1^{\eta'} &= [(1 - \beta_1) - y \beta_1 (\frac{Q^2}{2p_i \cdot Q})] \sqrt{1 - y} p_k^\eta p_i^{\eta'} - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_1 p_i^\eta p_i^{\eta'} - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_2 Q^\eta p_i^{\eta'} \\
&\quad + y \beta_1 A_1 p_i^\eta Q^{\eta'} + y \beta_1 A_2 Q^\eta Q^{\eta'} + y \beta_1 \sqrt{1 - y} p_k^\eta Q^{\eta'} \\
q_k^\eta q_i^{\eta'} &= A_1 \beta_1 p_i^\eta p_i^{\eta'} + A_2 \beta_1 Q^\eta p_i^{\eta'} + \beta_1 \sqrt{1 - y} p_k^\eta p_i^{\eta'}
\end{aligned} \tag{4.39}$$

### Calculation of the first Term

$$\begin{aligned}
g^{\eta\eta'} [2\{A_1 \beta_1 p_i^\gamma p_i^\delta + A_2 \beta_1 p_i^\gamma Q^\delta + \beta_1 \sqrt{1 - y} p_i^\gamma p_k^\delta\} \\
+ 2\{[\beta_1(1 - \beta_1) - y \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})] p_i^\gamma p_i^\delta + y \beta_1^2 p_i^\gamma Q^\delta\} \\
+ \{[(1 - \beta_1) - y \beta_1 (\frac{Q^2}{2p_i \cdot Q})] \sqrt{1 - y} p_i^\gamma p_k^\delta - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_1 p_i^\gamma p_i^\delta - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_2 p_i^\gamma Q^\delta \\
+ y \beta_1 A_1 Q^\gamma p_i^\delta + y \beta_1 A_2 Q^\gamma Q^\delta + y \beta_1 \sqrt{1 - y} Q^\gamma p_k^\delta\} \\
+ 3\{[(1 - \beta_1)^2 - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})^2] p_i^\gamma p_i^\delta - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q}) p_i^\gamma Q^\delta - y^2 \beta_1^2 (\frac{Q^2}{2p_i \cdot Q}) Q^\gamma p_i^\delta\} \\
+ 4\{[\beta_1(1 - \beta_1) - y \beta_1^2 (\frac{Q^2}{2p_i \cdot Q})] p_i^\gamma p_i^\delta + y \beta_1^2 Q^\gamma p_i^\delta\} \\
+ 2\{A_1 \beta_1 p_i^\gamma p_i^\delta + A_2 \beta_1 Q^\gamma p_i^\delta + \beta_1 \sqrt{1 - y} p_k^\gamma p_i^\delta\} \\
+ \{[(1 - \beta_1) - y \beta_1 (\frac{Q^2}{2p_i \cdot Q})] \sqrt{1 - y} p_k^\gamma p_i^\delta - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_1 p_i^\gamma p_i^\delta - y \beta_1 (\frac{Q^2}{2p_i \cdot Q}) A_2 Q^\gamma p_i^\delta \\
+ y \beta_1 A_1 p_i^\gamma Q^\delta + y \beta_1 A_2 Q^\gamma Q^\delta + y \beta_1 \sqrt{1 - y} p_k^\gamma Q^\delta\}
\end{aligned} \tag{4.40}$$

$$\begin{aligned}
& g^{\eta\eta'} \{ [2A_1\beta_1 + 2[\beta_1(1 - \beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] \\
& + 4[\beta_1(1 - \beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] + 3[(1 - \beta_1)^2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2] \\
& + 2A_1\beta_1 - y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_1 - y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_1 ] p_i^\gamma p_i^\delta \\
& + [2A_2\beta_1 + 2y\beta_1^2 - y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_2 - 3y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) + y\beta_1 A_1] p_i^\gamma Q^\delta \\
& + [2\beta_1 + [(1 - \beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})]] \sqrt{1 - y} p_i^\gamma p_k^\delta \\
& + [y\beta_1 A_1 + 4y\beta_1^2 + 2A_2\beta_1 - 3y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_2 ] Q^\gamma p_i^\delta \\
& + [y\beta_1 A_2 + y\beta_1 A_2] Q^\gamma Q^\delta + y\beta_1 \sqrt{1 - y} Q^\gamma p_k^\delta \\
& + [2\beta_1 + [(1 - \beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})]] \sqrt{1 - y} p_k^\gamma p_i^\delta + y\beta_1 \sqrt{1 - y} p_k^\gamma Q^\delta \}
\end{aligned} \tag{4.41}$$

Calculation of the second term

$$-g^{\eta\eta'} g^{\gamma\delta} (2q \cdot q_j + q \cdot q + 4q_i \cdot q_j + 2q_i \cdot q) \tag{4.42}$$

$$\begin{aligned}
& -g^{\eta\eta'} g^{\gamma\delta} [2([\alpha_1(1 - y) + y\beta_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\beta_1 Q \cdot p_k + \sqrt{\alpha_1\beta_1 y(1 - y)} p_k \cdot n_{\perp,1}) \\
& 4([\beta_1(1 - y) + y\alpha_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\alpha_1 Q \cdot p_k - \sqrt{\alpha_1\beta_1 y(1 - y)} p_k \cdot n_{\perp,1}) \\
& + 2(y p_i \cdot Q)]
\end{aligned} \tag{4.43}$$

Calculation of the third term

$$\begin{aligned}
& + g^{\eta\eta'} \{ [(1 - \beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})] \sqrt{1 - y} p_i^\eta p_k^\delta - y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_1 p_i^\eta p_i^\delta - y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_2 p_i^\eta Q^\delta \\
& + y\beta_1 A_1 Q^\eta p_i^\delta + y\beta_1 A_2 Q^\eta Q^\delta + y\beta_1 \sqrt{1 - y} Q^\eta p_k^\delta \\
& - [[(1 - \beta_1)^2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2] p_i^\eta p_i^\delta - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) p_i^\eta Q^\delta - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) Q^\eta p_i^\delta] \\
& - [A_1\beta_1 p_i^\eta p_i^\delta + A_2\beta_1 p_i^\eta Q^\delta + \beta_1 \sqrt{1 - y} p_i^\eta p_k^\delta] \\
& + [\beta_1(1 - \beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] p_i^\eta p_i^{\eta'} + y\beta_1^2 p_i^\eta Q^{\eta'} \}
\end{aligned} \tag{4.44}$$

## Calculation of the fourth term

$$\begin{aligned}
& + g^{\eta'\delta} \{ [(1 - \beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q}) - \beta_1] \sqrt{1 - y} p_i^\eta p_k^\gamma \\
& + [2[(1 - \beta_1)^2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2] - y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_1 + A_1\beta_1 + \\
& [\beta_1(1 - \beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] p_i^\eta p_i^\gamma \} \quad (4.45) \\
& + [-2y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) - y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_2 + A_2\beta_1 + y\beta_1^2] p_i^\eta Q^\gamma \\
& + [y\beta_1 A_1 + 2y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] Q^\eta p_i^\gamma + y\beta_1 A_2 Q^\eta Q^\gamma + y\beta_1 \sqrt{1 - y} Q^\eta p_k^\gamma \}
\end{aligned}$$

## Calculation of the fifth term

$$\begin{aligned}
& - g^{\gamma\delta} \{ [2[(1 - \beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})] - 2\beta_1] \sqrt{1 - y} p_i^\eta p_k^{\eta'} \\
& [-2y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_1 + [(1 - \beta_1)^2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2] - 2A_1\beta_1 \\
& - [\beta_1(1 - \beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] p_i^\eta p_i^{\eta'} \} \quad (4.46) \\
& [-2y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) - y\beta_1^2 - 2A_2\beta_1] p_i^\eta Q^{\eta'} \\
& + [2y\beta_1 A_1 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] Q^\eta p_i^{\eta'} + 2y\beta_1 A_2 Q^\eta Q^{\eta'} + 2y\beta_1 \sqrt{1 - y} Q^\eta p_k^{\eta'} \}
\end{aligned}$$

## Calculation of the sixth term

$$\begin{aligned}
& - g^{\gamma\eta} \{ [2[(1 - \beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})] + \beta_1] \sqrt{1 - y} p_i^{\eta'} p_k^\delta \\
& [-2y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_1 - 2[(1 - \beta_1)^2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2] \\
& - [\beta_1(1 - \beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] + A_1\beta_1] p_i^{\eta'} p_i^\delta \} \quad (4.47) \\
& [-2y\beta_1(\frac{Q^2}{2p_i \cdot Q}) A_2 + 2y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) + A_2\beta_1 - y\beta_1^2] p_i^{\eta'} Q^\delta \\
& + [2y\beta_1 A_1 + 2y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] Q^{\eta'} p_i^\delta + 2y\beta_1 A_2 Q^{\eta'} Q^\delta + 2y\beta_1 \sqrt{1 - y} Q^{\eta'} p_k^\delta \}
\end{aligned}$$

## Calculation of the seventh term

$$\begin{aligned}
& -g^{\eta\delta} \{ [2[(1-\beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})] + \beta_1] \sqrt{1-y} p_i^{\eta'} p_k^\gamma \\
& [4[(1-\beta_1)^2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2] - 2y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_1 + A_1\beta_1 \\
& + 2[\beta_1(1-\beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] p_i^{\eta'} p_i^\gamma \\
& + [-4y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) - 2y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_2 + 2y\beta_1^2 + A_2\beta_1] p_i^{\eta'} Q^\gamma \\
& + [-4y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) + 2y\beta_1 A_1] Q^\eta p_i^{\eta'} + 2y\beta_1 A_2 Q^\eta Q^{\eta'} + 2y\beta_1 \sqrt{1-y} Q^{\eta'} p_k^\gamma \}
\end{aligned} \tag{4.48}$$

## Calculation of the eighth term


$$\begin{aligned}
& +g^{\gamma\delta} \{ [4[(1-\beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})] + 2\beta_1] \sqrt{1-y} p_k^\eta p_i^{\eta'} \\
& + [-4y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_1 + 2A_1\beta_1 + [\beta_1(1-\beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] \\
& + [(1-\beta_1)^2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2] p_i^\eta p_i^{\eta'} \\
& + [4y\beta_1 A_1 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] p_i^\eta Q^{\eta'} + 4y\beta_1 A_2 Q^\eta Q^{\eta'} + 4y\beta_1 \sqrt{1-y} p_k^\eta Q^{\eta'} \\
& + [2A_2\beta_1 - 4y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_2 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q}) + y\beta_1^2] Q^\eta p_i^{\eta'} \}
\end{aligned} \tag{4.49}$$

## Final result

$$\begin{aligned}
M_1 M_2^\dagger &= \frac{g_s^2 C_A}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} g^{\eta\eta'} g^{\gamma\delta} \\
& [2([\alpha_1(1-y) + y\beta_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\beta_1 Q \cdot p_k + \sqrt{\alpha_1\beta_1 y(1-y)} p_k \cdot n_{\perp,1}) \\
& 4([\beta_1(1-y) + y\alpha_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\alpha_1 Q \cdot p_k - \sqrt{\alpha_1\beta_1 y(1-y)} p_k \cdot n_{\perp,1}) \\
& + 2(y p_i \cdot Q)]
\end{aligned} \tag{4.50}$$

$$\begin{aligned}
M_1 M_2^\dagger &= g_s^2 C_A g^{\eta\eta'} g^{\gamma\delta} \left[ \frac{1}{2y(p_i \cdot Q)} + \frac{\beta_1(\frac{Q^2}{2p_i \cdot Q})}{2y(1-\beta_1)(1-y)(p_i \cdot Q)} \right. \\
& \left. + \frac{\beta_1 Q \cdot p_k}{2y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} + \frac{\beta_1}{y(1-\beta_1)(p_i \cdot Q)} + \frac{1}{2(1-\beta_1)(1-y)(p_i \cdot p_k)} \right]
\end{aligned} \tag{4.51}$$

#### 4.4 Interference term of inverse $M_1 M_2^{\dagger'}$



$$M_1 M_2^{\dagger} = \frac{g_s^2 f^{l o a} f^{f' a' n} \delta^{a a'} \delta^{o b'} \delta^{f f'}}{(q_i + q)^2 (q_j + q_i)^2} [g_{\zeta}^{\eta'} g_{\tau'}^{\gamma} (g^{\eta \zeta} (2q + q_i)^{\mu} + g^{\zeta \mu} (q_i - q)^{\eta} - g^{\mu \eta} (2q_i + q)^{\zeta}) \\ g_{\mu \mu'} (g^{\tau' \mu'} (q_j - q_i)^{\delta} + g^{\mu' \delta} (2q_i + q_j)^{\tau'} - g^{\delta \tau'} (2q_j + q_i)^{\mu'})] \quad (4.52)$$

$$M_1 M_2^{\dagger} = \frac{g_s^2 f^{l o a} f^{f a n}}{4(q \cdot q_i)(q_i \cdot q_j)} [g^{\eta \eta'} (2q + q_i)^{\gamma} (q_j - q_i)^{\delta} + g^{\eta \eta'} (2q_i + q_j)^{\gamma} (2q + q_i)^{\delta} - g^{\eta \eta'} g^{\gamma \delta} (2q + q_i) \cdot (2q_j + q_i) \\ + g^{\gamma \eta'} (q_i - q)^{\eta} (q_j + q_i)^{\delta} + g^{\eta' \delta} (q_i - q)^{\eta} (2q_i + q_j)^{\gamma} - g^{\gamma \delta} (q_i - q)^{\eta} (2q_j + q_i)^{\eta'} \\ - g^{\gamma \eta} (2q_i + q)^{\eta'} (q_j - q_i)^{\delta} - g^{\eta \delta} (2q_i + q)^{\eta'} (2q_i + q_j)^{\gamma} + g^{\gamma \delta} (2q_j + q_i)^{\eta} (2q_i + q)^{\eta'}] \quad (4.53)$$

#### 4.5 Parametrization in terms of $(k_1 \cdot q_i)(q_i \cdot q_k)$

$$(k_1 \cdot q_i)(k_1 \cdot q_k) \approx y \beta_1 (1 - y) (p_i \cdot Q)(p_i \cdot Q) \quad (4.54)$$

Calculation of the third term

$$-g^{\eta \eta'} g^{\gamma \delta} \{4k_1 \cdot q_j + 2k_1 \cdot q_i + 2q_i \cdot q_k\} \quad (4.55)$$

$$M_1 M_2^{\dagger} = \frac{g_s^2 C_A}{4y \beta_1 (1 - y) (p_i \cdot p_k)(p_i \cdot Q)} g^{\eta \eta'} g^{\gamma \delta} [4([\alpha_1 (1 - y) + y \beta_1 (\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y \beta_1 Q \cdot p_k + \sqrt{\alpha_1 \beta_1 y (1 - y)} p_k \cdot n_{\perp, 1}) \\ 2([\beta_1 (1 - y) + y \alpha_1 (\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y \alpha_1 Q \cdot p_k - \sqrt{\alpha_1 \beta_1 y (1 - y)} p_k \cdot n_{\perp, 1}) \\ + 2(y p_i \cdot Q)] \quad (4.56)$$



$$\begin{aligned}
& -g^{\eta\eta'} g^{\gamma\delta} [4([\alpha_1(1-y) + y\beta_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\beta_1 Q \cdot p_k + \sqrt{\alpha_1\beta_1 y(1-y)} p_k \cdot n_{\perp,1}) \\
& 2([\beta_1(1-y) + y\alpha_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\alpha_1 Q \cdot p_k - \sqrt{\alpha_1\beta_1 y(1-y)} p_k \cdot n_{\perp,1}) \\
& + 2(y p_i \cdot Q)]
\end{aligned} \tag{4.57}$$

$$\begin{aligned}
M_1 M_2^\dagger = & g_s^2 C_A g^{\eta\eta'} g^{\gamma\delta} [\frac{1-\beta_1}{y\beta_1(p_i \cdot Q)} + \frac{1}{2y(p_i \cdot Q)} + \frac{(1-\beta_1)(\frac{Q^2}{2p_i \cdot Q})}{2y\beta_1(1-y)(p_i \cdot Q)} \\
& + \frac{(1-\beta_1) Q \cdot p_k}{2y\beta_1(1-y)(p_i \cdot p_k)(p_i \cdot Q)} + \frac{1}{2(1-\beta_1)(1-y)(p_i \cdot p_k)}]
\end{aligned} \tag{4.58}$$

## 4.6 $|M^2|$

$$\begin{aligned}
|M|^2 = & |M'_2|^2 + |M'_1|^2 + 2RE(M_1 M_2^\dagger + M_1 M_2^{\dagger'}) \\
|M|^2 = & \frac{g_s^2 C_A}{y(p_i \cdot Q)} [2[\epsilon - 1]\beta_1(1-\beta_1)n_{\perp,1}^\eta n_{\perp,1}^{\eta'} - 2g^{\eta\eta'}][g^{\gamma\delta}] \\
& + \frac{g_s^2 C_A}{(1-\beta_1)(1-y)(p_i \cdot p_k)} [-2g^{\delta\gamma}][g^{\eta\eta'}] \\
& + 2Re(g_s^2 C_A g^{\eta\eta'} g^{\gamma\delta} [\frac{1}{2y(p_i \cdot Q)} + \frac{\beta_1(\frac{Q^2}{2p_i \cdot Q})}{2y(1-\beta_1)(1-y)(p_i \cdot Q)} \\
& + \frac{\beta_1 Q \cdot p_k}{2y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} + \frac{\beta_1}{y(1-\beta_1)(p_i \cdot Q)} + \frac{1}{2(1-\beta_1)(1-y)(p_i \cdot p_k)}] \\
& + g_s^2 C_A g^{\eta\eta'} g^{\gamma\delta} [\frac{1-\beta_1}{y\beta_1(p_i \cdot Q)} + \frac{1}{2y(p_i \cdot Q)} + \frac{(1-\beta_1)(\frac{Q^2}{2p_i \cdot Q})}{2y\beta_1(1-y)(p_i \cdot Q)} \\
& + \frac{(1-\beta_1) Q \cdot p_k}{2y\beta_1(1-y)(p_i \cdot p_k)(p_i \cdot Q)} + \frac{1}{2(1-\beta_1)(1-y)(p_i \cdot p_k)}])
\end{aligned} \tag{4.59}$$


$$\begin{aligned}
|M|^2 = & |M'_2|^2 + |M'_1|^2 + 2RE(M_1 M_2^\dagger + M_1 M_2^{\dagger'}) \\
|M|^2 = & g_s^2 C_A g^{\eta\eta'} g^{\gamma\delta} [2[\epsilon - 1]\beta_1(1-\beta_1)n_{\perp,1}^\eta n_{\perp,1}^{\eta'} + \frac{\beta_1(\frac{Q^2}{2p_i \cdot Q})}{y(1-\beta_1)(1-y)(p_i \cdot Q)} \\
& + \frac{\beta_1 Q \cdot p_k}{y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} + \frac{2\beta_1}{y(1-\beta_1)(p_i \cdot Q)} \\
& + \frac{2(1-\beta_1)}{y\beta_1(p_i \cdot Q)} + \frac{(1-\beta_1)(\frac{Q^2}{2p_i \cdot Q})}{y\beta_1(1-y)(p_i \cdot Q)} + \frac{(1-\beta_1) Q \cdot p_k}{y\beta_1(1-y)(p_i \cdot p_k)(p_i \cdot Q)}]
\end{aligned} \tag{4.60}$$

$$\begin{aligned}
|M|^2 = & g_s^2 C_A g^{\eta\prime} g^{\gamma\delta} \left[ 2\beta_1(1 - \beta_1) + \frac{2\beta_1}{y(1 - \beta_1)(p_i \cdot Q)} + \frac{2(1 - \beta_1)}{y\beta_1(p_i \cdot Q)} + \frac{\left(\frac{Q^2}{2p_i \cdot Q}\right)}{y\beta_1(1 - y)(p_i \cdot Q)} \right. \\
& \left. + \frac{Q \cdot p_k}{y\beta_1(1 - y)(p_i \cdot p_k)(p_i \cdot Q)} \right]
\end{aligned} \tag{4.61}$$


$$|M|^2 = 2 \frac{g_s^2 C_A}{y(p_i \cdot Q)} g^{\eta\prime} g^{\gamma\delta} \left[ \beta_1(1 - \beta_1) + \frac{\beta_1}{1 - \beta_1} + \frac{1 - \beta_1}{\beta_1} \right] \tag{4.62}$$

## Chapter 5

### Quark gluon quark emission kernel



## 5.1 Quark loop



$$|M_1|^2 = \left[ \frac{-i}{(q_i + q)^2} \not{q}_i (-ig_s \gamma^\eta \times [T^l]_a^o) \not{q} (ig_s \gamma^{\eta'} \times [T^k]_{o'}^{a'}) \frac{i}{(q_i + q)^2} \right] [\not{q}_j] \quad (5.1)$$

$$|M_1|^2 = \frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4(k_1 \cdot q_i)(k_1 \cdot q_i)} [\not{q}_i \gamma^\eta \not{k}_1 \gamma^{\eta'}] [\not{q}_k] \quad (5.2)$$

$$|M_1|^2 = -\frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4y^2(p_i \cdot Q)(p_i \cdot Q)} [\not{q}_i \not{k}_1 \gamma^\eta \gamma^{\eta'}] [\not{q}_k] \quad (5.3)$$

$$\begin{aligned} |M_1|^2 = & -\frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4y^2(p_i \cdot Q)(p_i \cdot Q)} \\ & [((\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\alpha_1 \not{Q} - \sqrt{y\alpha_1\beta_1} \not{p}_{\perp,l}) \\ & ((\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\beta_1 \not{Q} + \sqrt{y\alpha_1\beta_1} \not{p}_{\perp,1}) \gamma^\eta \gamma^{\eta'}] \\ & [A_1 \not{p}_i + A_2 \not{Q} + \sqrt{1-y} \not{p}_k] \end{aligned} \quad (5.4)$$


$$\begin{aligned} |M_1|^2 = & -\frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4y^2(p_i \cdot Q)(p_i \cdot Q)} \\ & [(y\beta_1(\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i \not{Q} + y\alpha_1(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i \not{Q} + y^2\alpha_1\beta_1 \not{Q} \not{Q}) g^{\eta\eta'}] \\ & [A_1 \not{p}_i + A_2 \not{Q} + \sqrt{1-y} \not{p}_k] \end{aligned} \quad (5.5)$$

$$|M_1|^2 = -\frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4y^2(p_i \cdot Q)(p_i \cdot Q)} [(y\beta_1^2 \not{p}_i \not{Q} + y\alpha_1^2 \not{Q} \not{p}_i) g^{m'}] [A_1 \not{p}_i + A_2 \not{Q} + \sqrt{1-y} \not{p}_k] \quad (5.6)$$

$$|M_1|^2 = -\frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4y^2(p_i \cdot Q)(p_i \cdot Q)} [y(\beta_1^2 - \alpha_1^2) \not{p}_i \not{Q} g^{m'}] [A_1 \not{p}_i + A_2 \not{Q} + \sqrt{1-y} \not{p}_k] \quad (5.7)$$


$$|M_1|^2 = -\frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4y(p_i \cdot Q)(p_i \cdot Q)} [g^{m'}] [\sqrt{1-y}(\beta_1^2 - \alpha_1^2) \not{p}_i \not{Q} \not{p}_k] \quad (5.8)$$

## 5.2 Spectator Quark loop



$$|M_2|^2 = \frac{g_s^2 [T^m]_f^b [T^n]_f^b}{4(k_1 \cdot q_k)(k_1 \cdot q_k)} [\not{A}_k \gamma^\gamma \not{k}_1 \gamma^\delta] [\not{A}_i] \quad (5.9)$$

### 5.3 Interference term



$$M_1 M_2^\dagger = \frac{g_s^2 [T^l]_a^o [T^n]_f^o}{4(qq_i)(qq_j)} [\not{q}_i \gamma^\eta \not{q} \gamma^\delta \not{q}_j] \quad (5.10)$$

$$M_1 M_2^\dagger = -\frac{g_s^2 [T^l]_a^o [T^n]_f^o}{4(k_1 \cdot q_i)(k_1 \cdot q_k)} [\not{q}_i \not{k}_1 \not{q}_k] [g^{\eta\delta}] \quad (5.11)$$

$$\begin{aligned} M_1 M_2^\dagger &= -\frac{g_s^2 [T^l]_a^o [T^n]_f^o}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} [g^{\eta\delta}] \\ &[ ((\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\alpha_1 \not{Q} - \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,l}) \\ &((\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\beta_1 \not{Q} + \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,l}) \\ &(A_1 \not{p}_i + A_2 \not{Q} + \sqrt{1-y} \not{p}_k) ] \end{aligned} \quad (5.12)$$


$$M_1 M_2^\dagger = -\frac{g_s^2 [T^l]_a^o [T^n]_f^o}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} [g^{\eta\delta}] [\beta_1 \sqrt{1-y} \not{p}_i \not{Q} \not{p}_k] \quad (5.13)$$

### 5.4 $|M^2|$


$$\begin{aligned} |M|^2 &= |M_2|^2 + |M_1|^2 + 2RE(M_1 M_2^\dagger) \\ &- \frac{g_s^2 [T^l]_a^o [T^k]_{o'}^{a'}}{4y(p_i \cdot Q)(p_i \cdot Q)} [g^{\eta\eta'}] [\sqrt{1-y}(\beta_1^2 - \alpha_1^2) \not{p}_i \not{Q} \not{p}_k] \\ &+ 2RE(-\frac{g_s^2 [T^l]_a^o [T^n]_f^o}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} [g^{\eta\delta}] [\beta_1 \sqrt{1-y} \not{p}_i \not{Q} \not{p}_k]) \end{aligned} \quad (5.14)$$


## Chapter 6


### Gluon quark quark emission kernel





6.1  $M_1$ 

6.2  $M_2$ 

6.3  $M_1 M_2^\dagger$ 

# Appendix A

## MATHEMATICAL TOOLS

Lorentz transformation of momenta  $\hat{p}_i^\mu, \hat{p}_k^\mu$  and  $\hat{Q}^\mu$

$$\begin{aligned}
 \hat{p}_i^\mu &= \alpha \Lambda^\mu{}_\nu p_i^\nu = p_i^\mu p_{i\nu} p_i^\nu \frac{-y^2 Q^2}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + p_i^\mu Q_\nu p_i^\nu \frac{y(1 + \sqrt{1-y})}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \\
 &+ Q^\mu p_{i\nu} p_i^\nu \frac{(y^2 - y - y\sqrt{1-y})}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} \eta^\mu{}_\nu p_i^\nu \\
 \hat{p}_i^\mu &= p_i^\mu (Q \cdot p_i) \frac{y(1 + \sqrt{1-y})}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} p_i^\mu \\
 &= p_i^\mu \left[ \frac{y(1 + \sqrt{1-y})}{(2 + 2\sqrt{1-y} - y)} + \sqrt{1-y} \right] = p_i^\mu \\
 \boxed{\hat{p}_i^\mu &= \alpha \Lambda^\mu{}_\nu p_i^\nu = p_i^\mu} \tag{6.1}
 \end{aligned}$$

$$\begin{aligned}
 \hat{p}_k^\mu &= \alpha \Lambda^\mu{}_\nu p_k^\nu = p_i^\mu \left[ \frac{-y^2 Q^2 (p_i \cdot p_k)}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + \frac{y(1 + \sqrt{1-y})(Q \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] \\
 &+ Q^\mu \left[ \frac{(y^2 - y - y\sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] + \sqrt{1-y} p_k^\mu \\
 \hat{p}_k^\mu &= \alpha \Lambda^\mu{}_\nu p_k^\nu = p_i^\mu \left[ \frac{-y^2 Q^2 (p_i \cdot p_k)}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + \frac{y(1 + \sqrt{1-y})(Q \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] \\
 &+ Q^\mu \left[ \frac{(y^2 - y - y\sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] + \sqrt{1-y} p_k^\mu
 \end{aligned}$$

with

$$\begin{aligned}
 A_1 &\equiv \frac{-y^2 Q^2 (p_i \cdot p_k)}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + \frac{y(1 + \sqrt{1-y})(Q \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \\
 A_2 &\equiv \frac{(y^2 - y - y\sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})}
 \end{aligned}$$

$$\boxed{\hat{p}_k^\mu = A_1 p_i^\mu + A_2 Q^\mu + \sqrt{1-y} p_k^\mu} \quad (6.2)$$

$$\begin{aligned} \hat{Q}^\mu &= \alpha \Lambda^\mu_\nu Q^\nu = p_i^\mu \left[ \frac{-y^2 Q^2 (p_i \cdot Q)}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + \frac{y(1 + \sqrt{1-y}) Q^2}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] \\ &+ Q^\mu \left[ \frac{(y^2 - y - y\sqrt{1-y})(p_i \cdot Q)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] + \sqrt{1-y} Q^\mu \end{aligned}$$

with

$$\begin{aligned} S_1 &\equiv \frac{Q^2}{2p_i \cdot Q} \left[ \frac{-y^2}{2(1 + \sqrt{1-y} - \frac{y}{2})} + \frac{y(1 + \sqrt{1-y})}{(1 + \sqrt{1-y} - \frac{y}{2})} \right] = \frac{Q^2}{2p_i \cdot Q} y \\ S_2 &\equiv \frac{(y^2 - y - y\sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} = 1 - y \end{aligned}$$

$$\boxed{\hat{Q}^\mu = \frac{Q^2}{2p_i \cdot Q} y p_i^\mu + (1 - y) Q^\mu} \quad (6.3)$$

The often occurring pre-factor products

$$\begin{aligned} \zeta_1 \zeta_1 &= (\alpha_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q}) + y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})^2) \\ \zeta_1 \lambda_1 &= (y\alpha_1\beta_1 - y^2\beta_1^2(\frac{Q^2}{2p_i \cdot Q})) \\ \zeta_1 \zeta_q &= (\alpha_1\beta_1 - y(\alpha_1^2 + \beta_1^2)(\frac{Q^2}{2p_i \cdot Q}) + y^2\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q})^2) \\ \zeta_1 \lambda_q &= (y\alpha_1^2 - y^2\beta_1\alpha_1(\frac{Q^2}{2p_i \cdot Q})) \\ \zeta_q \zeta_q &= (\beta_1^2 - 2y\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q}) + y^2\alpha_1^2(\frac{Q^2}{2p_i \cdot Q})^2) \\ \zeta_q \lambda_1 &= (y\beta_1^2 - y^2\alpha_1\beta_1(\frac{Q^2}{2p_i \cdot Q})) \\ \zeta_q \lambda_q &= (y\beta_1\alpha_1 - y^2\alpha_1^2(\frac{Q^2}{2p_i \cdot Q})) \\ \lambda_1 \lambda_1 &= y^2\beta_1^2 \quad \lambda_1 \lambda_q = y^2\beta_1\alpha_1 \quad \lambda_q \lambda_q = y^2\alpha_1^2 \end{aligned} \quad (6.4)$$

## Common scalar products

$$\begin{aligned}
k_1 \cdot q_i &= (\zeta_1 \lambda_q + \lambda_1 \zeta_q) p_i \cdot Q + \lambda_1 \lambda_q Q^2 - y \alpha_1 \beta_1 n_{\perp,1}^2 \\
&= [(\alpha_1 - y \beta_1 (\frac{Q^2}{2 p_i \cdot Q})) y \alpha_1 + y \beta_1 (\beta_1 - \alpha_1 y (\frac{Q^2}{2 p_i \cdot Q}))] p_i \cdot Q \\
&\quad y^2 \beta_1 \alpha_1 Q^2 + 2 y \alpha_1 \beta_1 p_i Q \\
\Rightarrow k_1 \cdot q_i &= [y \alpha_1^2 - y^2 \alpha_1 \beta_1 (\frac{Q^2}{2 p_i \cdot Q}) + y \beta_1^2 - y^2 \alpha_1 \beta_1 (\frac{Q^2}{2 p_i \cdot Q})] p_i \cdot Q \\
&\quad y^2 \beta_1 \alpha_1 Q^2 + 2 y \alpha_1 \beta_1 p_i Q
\end{aligned} \tag{6.5}$$

$$\boxed{k_1 \cdot q_i = y(\alpha_1 + \beta_1)^2 p_i \cdot Q = y p_i \cdot Q} \tag{6.6}$$

$$\begin{aligned}
k_1 \cdot q_k &= (\zeta_1 A_2 + \lambda_1 A_1) p_i \cdot Q + \zeta_1 \sqrt{1-y} p_i \cdot p_k + \lambda_1 A_2 Q^2 + \lambda_1 \sqrt{1-y} Q \cdot p_k \\
&\quad + \sqrt{\alpha_1 \beta_1 y (1-y)} p_k \cdot n_{\perp,1} \\
&= \{[(\alpha_1 - y \beta_1 (\frac{Q^2}{2 p_i \cdot Q})) \frac{(y^2 - y - y \sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})}] \\
&\quad - y^2 Q^2 (p_i \cdot p_k) \frac{y(1 + \sqrt{1-y})(Q \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})}\} p_i \cdot Q \\
&\quad + (\alpha_1 - y \beta_1 (\frac{Q^2}{2 p_i \cdot Q})) \sqrt{1-y} p_i \cdot p_k + y \beta_1 \frac{(y^2 - y - y \sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} Q^2 \\
&\quad + y \beta_1 \sqrt{1-y} Q \cdot p_k + \sqrt{\alpha_1 \beta_1 y (1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.7}$$

$$\begin{aligned}
k_1 \cdot q_k &= \alpha_1 \frac{(y^2 - y - y \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) - y \beta_1 (\frac{Q^2}{2 p_i \cdot Q}) \frac{(y^2 - y - y \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) \\
&\quad + y \beta_1 \frac{-y^2 Q^2}{4(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) + y \beta_1 \frac{y(1 + \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} Q \cdot p_k \\
&\quad + \alpha_1 \sqrt{1-y} p_i \cdot p_k - y \beta_1 (\frac{Q^2}{2 p_i \cdot Q}) \sqrt{1-y} p_i \cdot p_k \\
&\quad + y \beta_1 (\frac{Q^2}{2 p_i \cdot Q}) \frac{(y^2 - y - y \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) + y \beta_1 \sqrt{1-y} (Q \cdot p_k) \\
&\quad + \sqrt{\alpha_1 \beta_1 y (1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.8}$$

$$\begin{aligned}
k_1 \cdot q_k &= [\alpha_1 \frac{(y^2 - y - y \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + y \beta_1 \frac{-y^2 Q^2}{4(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} + \alpha_1 \sqrt{1-y} \\
&\quad - y \beta_1 (\frac{Q^2}{2 p_i \cdot Q}) \sqrt{1-y}] p_i \cdot p_k + [y \beta_1 \frac{y(1 + \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + y \beta_1 \sqrt{1-y}] (Q \cdot p_k) \\
&\quad + \sqrt{\alpha_1 \beta_1 y (1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.9}$$

$$\begin{aligned}
k_1 \cdot q_k &= \left\{ \alpha_1 \left[ \frac{(y^2 - y - y\sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} \right] \right. \\
&\quad + y\beta_1 \left( \frac{Q^2}{p_i \cdot Q} \right) \left[ \frac{-y^2}{4(1 + \sqrt{1-y} - \frac{y}{2})} - \sqrt{1-y} \right] \left. \right\} p_i \cdot p_k \\
&\quad + y\beta_1 \left[ \frac{y(1 + \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} \right] (Q \cdot p_k) \\
&\quad + \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.10}$$

$$\boxed{k_1 \cdot q_k = [\alpha_1(1-y) + y\beta_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\beta_1 Q \cdot p_k + \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1}} \tag{6.11}$$

$$\begin{aligned}
q_i \cdot q_k &= (\zeta_q A_2 + \lambda_q A_1) p_i \cdot Q + \zeta_q \sqrt{1-y} p_i \cdot p_k + \lambda_q A_2 Q^2 + \lambda_q \sqrt{1-y} Q \cdot p_k \\
&\quad - \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1} \\
&= \left\{ \left[ (\beta_1 - y\alpha_1(\frac{Q^2}{2p_i \cdot Q})) \frac{(y^2 - y - y\sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] \right. \\
&\quad + y\alpha_1 \left[ \frac{-y^2 Q^2 (p_i \cdot p_k)}{4(p_i \cdot Q)^2 (1 + \sqrt{1-y} - \frac{y}{2})} + \frac{y(1 + \sqrt{1-y})(Q \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} \right] \left. \right\} p_i \cdot Q \\
&\quad + (\beta_1 - y\alpha_1(\frac{Q^2}{2p_i \cdot Q})) \sqrt{1-y} p_i \cdot p_k + y\alpha_1 \frac{(y^2 - y - y\sqrt{1-y})(p_i \cdot p_k)}{2(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} Q^2 \\
&\quad + y\alpha_1 \sqrt{1-y} Q \cdot p_k - \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.12}$$

$$\begin{aligned}
q_i \cdot q_k &= \beta_1 \frac{(y^2 - y - y\sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) - y\alpha_1 \left( \frac{Q^2}{2p_i \cdot Q} \right) \frac{(y^2 - y - y\sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) \\
&\quad + y\alpha_1 \frac{-y^2 Q^2}{4(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) + y\alpha_1 \frac{y(1 + \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} Q \cdot p_k \\
&\quad + \beta_1 \sqrt{1-y} p_i \cdot p_k - y\alpha_1 \left( \frac{Q^2}{2p_i \cdot Q} \right) \sqrt{1-y} p_i \cdot p_k \\
&\quad + y\alpha_1 \left( \frac{Q^2}{2p_i \cdot Q} \right) \frac{(y^2 - y - y\sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} (p_i \cdot p_k) + y\alpha_1 \sqrt{1-y} (Q \cdot p_k) \\
&\quad - \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.13}$$

$$\begin{aligned}
q_i \cdot q_k &= \left[ \beta_1 \frac{(y^2 - y - y\sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + y\alpha_1 \frac{-y^2 Q^2}{4(p_i \cdot Q)(1 + \sqrt{1-y} - \frac{y}{2})} + \beta_1 \sqrt{1-y} \right. \\
&\quad - y\alpha_1 \left( \frac{Q^2}{2p_i \cdot Q} \right) \sqrt{1-y} \left. \right] p_i \cdot p_k + \left[ y\alpha_1 \frac{y(1 + \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + y\alpha_1 \sqrt{1-y} \right] (Q \cdot p_k) \\
&\quad - \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.14}$$

$$\begin{aligned}
k_1 \cdot q_k = & \left\{ \beta_1 \left[ \frac{(y^2 - y - y\sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} \right] \right. \\
& + y\alpha_1 \left( \frac{Q^2}{p_i \cdot Q} \right) \left[ \frac{-y^2}{4(1 + \sqrt{1-y} - \frac{y}{2})} - \sqrt{1-y} \right] \left. \right\} p_i \cdot p_k \\
& + y\alpha_1 \left[ \frac{y(1 + \sqrt{1-y})}{2(1 + \sqrt{1-y} - \frac{y}{2})} + \sqrt{1-y} \right] (Q \cdot p_k) \\
& - \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1}
\end{aligned} \tag{6.15}$$

$$\boxed{q_i \cdot q_k = [\beta_1(1-y) + y\alpha_1(\frac{Q^2}{2p_i \cdot Q})] p_i \cdot p_k + y\alpha_1 Q \cdot p_k - \sqrt{\alpha_1 \beta_1 y(1-y)} p_k \cdot n_{\perp,1}} \tag{6.16}$$



## Detailed calculation of the gluon radiation of a quark

$$|M_1|^2$$

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{(2k_1 \cdot q_i)} [k_1] [\not{q}_k] \quad (6.17)$$

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{2y p_i \cdot Q} [(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\beta_1 \not{Q} + \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1}] [A_1 \not{p}_i + A_2 \not{Q} + \sqrt{1-y} \not{p}_k] \quad (6.18)$$

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{2y p_i \cdot Q} [(A_2(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) + A_1 y\beta_1) p_i \cdot Q + (\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \sqrt{1-y} p_i \cdot p_k + A_2 y\beta_1 Q^2 + \sqrt{1-y} \sqrt{y\alpha_1\beta_1} n_{\perp,1} \cdot p_k] \quad (6.19)$$

For the collinearity  $y \rightarrow 0$  we'll get:

$$|M_1|^2 = (d-2) \frac{g_s^2 C_F}{2y p_i \cdot Q} [(A_2(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) + A_1 y\beta_1) \not{p}_i \not{Q} + (\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \sqrt{1-y} \not{p}_i \not{p}_k + A_2 y\beta_1 Q^2 + \sqrt{1-y} \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1} \not{p}_k] \quad (6.20)$$

$$|M_1|^2 = (d-2)(1-\beta_1) \sqrt{1-y} \frac{g_s^2 C_F}{2y p_i \cdot Q} [\not{p}_i \not{p}_k] \quad (6.21)$$

$$|M_2|^2$$

$$|M_2|^2 = (d-2) \frac{g_s^2 [T^c]_f^m [T^c]_f^n}{2k_1 \cdot q_k} [k_1] [\not{q}_i] \quad (6.22)$$

$$|M_2|^2 = (d-2) \frac{g_s^2 C_F}{2k_1 \cdot q_k} [(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\beta_1 \not{Q} + \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1}] [(\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i + y\alpha_1 \not{Q} - \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1}] \quad (6.23)$$

$$|M_2|^2 = (d-2) \frac{g_s^2 C_F}{2k_1 \cdot q_k} [y\alpha_1(\alpha_1 - y\beta_1 (\frac{Q^2}{2p_i \cdot Q})) \not{p}_i \not{Q} + y\beta_1(\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \not{Q} \not{p}_i + y^2\alpha_1\beta_1 Q^2 - y\beta_1 \sqrt{y\alpha_1\beta_1} \not{Q} \not{n}_{\perp,1} + y\beta_1 \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1} \not{Q} - y\alpha_1\beta_1 n_{\perp,1}^2 + (\beta_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \sqrt{y\alpha_1\beta_1} \not{n}_{\perp,1} \not{p}_i - (\alpha_1 - \alpha_1 y (\frac{Q^2}{2p_i \cdot Q})) \sqrt{y\alpha_1\beta_1} \not{p}_i \not{n}_{\perp,1}] \quad (6.24)$$

Which means:

$$|M_2|^2 \sim (d-2) \frac{g_s^2 C_F}{2k_1 \cdot q_k} y[\dots] \quad (6.25)$$

$$|M_2|^2 \rightarrow 0 \quad \text{for} \quad y \rightarrow 0$$

$M_1 M_2^\dagger$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i k_1)(2q_k k_1)} [(\not{q}_i + \not{k}_1) \not{q}_i \gamma^\mu] [(\not{q}_k + \not{k}_1) \not{q}_k \gamma_\mu] \quad (6.26)$$

$$+ 4[(\not{q}_i + \not{k}_1) q_i^\mu][(\not{q}_k + \not{k}_1) q_{k\mu}]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} \quad (6.27)$$

$$[(\not{q}_i \not{q}_i + \not{k}_1 \not{q}_i) \gamma^\mu][(\not{q}_k \not{q}_k + \not{k}_1 \not{q}_k) \gamma_\mu] + 4(q_i^\mu q_{k\mu})[\not{q}_i + \not{k}_1][\not{q}_k + \not{k}_1]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} \quad (6.28)$$

$$[\not{k}_1 \not{q}_i \gamma^\mu][\not{k}_1 \not{q}_k \gamma_\mu] + 4(q_i \cdot q_k)[\not{q}_i \not{q}_k + \not{k}_1 \not{q}_k + \not{q}_i \not{k}_1]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} \quad (6.29)$$

$$4(A_1 \beta_1 p_i \cdot p_i + A_2 \beta_1 p_i \cdot Q + \beta_1 \sqrt{1-y} p_i \cdot p_k)$$

$$[A_1 \beta_1 \not{p}_i \not{p}_i + A_2 \beta_1 \not{p}_i \not{Q} + \beta_1 \sqrt{1-y} \not{p}_i \not{p}_k$$

$$+ [(1-\beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})]\sqrt{1-y} \not{p}_i \not{p}_k - y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_1 \not{p}_i \not{p}_i$$

$$- y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_2 \not{p}_i \not{Q} + y\beta_1 A_1 \not{Q} \not{p}_i + y\beta_1 A_2 \not{Q} \not{Q} + y\beta_1 \sqrt{1-y} \not{Q} \not{p}_k$$

$$+ [\beta_1(1-\beta_1) - y\beta_1^2(\frac{Q^2}{2p_i \cdot Q})] \not{p}_i \not{p}_i + y\beta_1^2 \not{p}_i \not{Q}]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} \quad (6.30)$$

$$4(A_2 \beta_1 p_i \cdot Q + \beta_1 \sqrt{1-y} p_i \cdot p_k)[A_2 \beta_1 \not{p}_i \not{Q} + \beta_1 \sqrt{1-y} \not{p}_i \not{p}_k$$

$$+ [(1-\beta_1) - y\beta_1(\frac{Q^2}{2p_i \cdot Q})]\sqrt{1-y} \not{p}_i \not{p}_k - y\beta_1(\frac{Q^2}{2p_i \cdot Q})A_2 \not{p}_i \not{Q}$$

$$+ y\beta_1 A_1 \not{Q} \not{p}_i + y\beta_1 A_2 \not{Q} \not{Q} + y\beta_1 \sqrt{1-y} \not{Q} \not{p}_k + y\beta_1^2 \not{p}_i \not{Q}]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{4y(1-\beta_1)(1-y)(p_i \cdot p_k)(p_i \cdot Q)} \quad (6.31)$$

$$4(\beta_1 \sqrt{1-y} p_i \cdot p_k)[\beta_1 \sqrt{1-y} \not{p}_i \not{p}_k + (1-\beta_1) \sqrt{1-y} \not{p}_i \not{p}_k]$$

$$M_1 M_2^\dagger = \frac{-g_s^2 C_F}{y(1-\beta_1)(p_i \cdot p_k)(p_i \cdot Q)} \beta_1(p_i \cdot p_k) [\beta_1 \not{p}_i \not{p}_k + (1-\beta_1) \not{p}_i \not{p}_k] \quad (6.32)$$

$$M_1 M_2^\dagger = \frac{\beta_1}{(1-\beta_1)} \frac{-g_s^2 C_F}{y(p_i \cdot Q)} [\not{p}_i \not{p}_k] \quad (6.33)$$

Evaluation of the tensor  $N^{\eta\eta'}$

$$\begin{aligned} N^{\eta\eta'} \equiv & g_{\mu\mu'} g_{\zeta\zeta'} [-g^{\mu\zeta} g^{\mu'\eta'} (q - q_i)^\eta (2q_i + q)^{\zeta'} + g^{\mu\zeta} g^{\eta'\zeta'} (q - q_i)^\eta (2q + q_i)^{\mu'} \\ & + g^{\mu\zeta} g^{\zeta'\mu'} (q - q_i)^\eta (q_i - q)^{\eta'} + g^{\zeta\eta} g^{\mu'\zeta'} (2q + q_i)^\mu (2q_i + q)^{\zeta'} \\ & - g^{\zeta\eta} g^{\eta'\zeta'} (2q + q_i)^\mu (2q + q_i)^{\mu'} - g^{\zeta\eta} g^{\zeta'\mu'} (2q + q_i)^\mu (q_i - q)^{\eta'} \\ & - g^{\eta\mu} g^{\mu'\eta'} (2q_i + q)^\zeta (2q_i + q)^{\zeta'} + g^{\eta\mu} g^{\eta'\zeta'} (2q_i + q)^\zeta (2q + q_i)^{\mu'} \\ & + g^{\eta\mu} g^{\zeta'\mu'} (2q_i + q)^\zeta (q_i - q)^{\eta'}] [g^{\gamma\delta}] \end{aligned} \quad (6.34)$$

$$\begin{aligned} N^{\eta\eta'} \equiv & [-(q - q_i)^\eta (2q_i + q)^{\eta'} + (q - q_i)^\eta (2q + q_i)^{\eta'} + d(q - q_i)^\eta (q_i - q)^{\eta'} \\ & + (2q + q_i)^{\eta'} (2q_i + q)^\eta - g^{\eta\eta'} (2q + q_i)^\mu (2q + q_i)_\mu - (2q + q_i)^\eta (q_i - q)^{\eta'} \\ & - g^{\eta\eta'} (2q_i + q)^\zeta (2q_i + q)_\zeta + (2q_i + q)^{\eta'} (2q + q_i)^\eta + (2q_i + q)^\eta (q_i - q)^{\eta'}] [g^{\gamma\delta}] \end{aligned} \quad (6.35)$$

$$\begin{aligned} N^{\eta\eta'} \equiv & [-(q^\eta q^{\eta'} + 2q^\eta q_i^{\eta'} - q_i^\eta q^{\eta'} - 2q_i^\eta q_i^{\eta'}) + (2q^\eta q^{\eta'} + q^\eta q_i^{\eta'} - 2q_i^\eta q^{\eta'} - q_i^\eta q_i^{\eta'}) \\ & + (dq^\eta q_i^{\eta'} - dq^\eta q^{\eta'} - dq_i^\eta q_i^{\eta'} + dq_i^\eta q^{\eta'}) + (4q^{\eta'} q_i^\eta + 2q^{\eta'} q^\eta + 2q_i^{\eta'} q_i^\eta + q_i^{\eta'} q^\eta) \\ & - (-2q^\eta q^{\eta'} + 2q^\eta q_i^{\eta'} - q_i^\eta q^{\eta'} + q_i^\eta q_i^{\eta'}) + (2q^{\eta'} q^\eta + q^{\eta'} q_i^\eta + 4q_i^{\eta'} q^\eta + 2q_i^{\eta'} q_i^\eta) \\ & + (-q^\eta q^{\eta'} + q^\eta q_i^{\eta'} - 2q_i^\eta q^{\eta'} + 2q_i^\eta q_i^{\eta'}) - g^{\eta\eta'} (5q^2 + 5q_i^2 + 8qq_i)] [g^{\gamma\delta}] \end{aligned} \quad (6.36)$$

# Bibliography

- [1] Johannes Blumer, Ralph Engel, and Jorg R. Horandel. Cosmic Rays from the Knee to the Highest Energies. *Prog. Part. Nucl. Phys.*, 63:293–338, 2009.
- [2] Michiel Botje. Lecture notes particle physics ii, quantum chromo dynamics, November 2013.
- [3] Wilhelm Capelle. *Die Vorsokratiker: die Fragmente und Quellenberichte*, volume 119. Kröner, 1968.
- [4] S. Catani and M. H. Seymour. A General algorithm for calculating jet cross-sections in NLO QCD. *Nucl. Phys.*, B485:291–419, 1997.
- [5] Stefano Catani, Stefan Dittmaier, Michael H. Seymour, and Zoltan Trocsanyi. The Dipole formalism for next-to-leading order QCD calculations with massive partons. *Nucl. Phys.*, B627:189–265, 2002.
- [6] John Dalton. *A new system of chemical philosophy*, volume 1. Cambridge University Press, 2010.
- [7] Wolfgang Demtröder. *Experimentalphysik*, volume 2. Springer, 2005.
- [8] L. Edelhäuser and A. Knochel. *Tutorium Quantenfeldtheorie: Was Sie schon immer über QFT wissen wollten, aber bisher nicht zu fragen wagten*. Springer Berlin Heidelberg, 2016.
- [9] R. Keith Ellis, W. James Stirling, and B. R. Webber. QCD and collider physics. *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.*, 8:1–435, 1996.
- [10] M. Ender. Radiodetektion von luftschauern unter dem einfluss starker elektronischer felder in der atmosphäre, 2009. Diplom Thesis.
- [11] L. D. Faddeev and V. N. Popov. Feynman Diagrams for the Yang-Mills Field. *Phys. Lett.*, B25:29–30, 1967. [,325(1967)].
- [12] Nadine Fischer, Stefan Gieseke, Simon Plätzer, and Peter Skands. Revisiting radiation patterns in  $e^+e^-$  collisions. *Eur. Phys. J.*, C74(4):2831, 2014.
- [13] Stefan Gieseke, P. Stephens, and Bryan Webber. New formalism for QCD parton showers. *JHEP*, 12:045, 2003.

- [14] Johann Wolfgang Goethe. *Faust*, volume 1. Ripol Classic, 1921.
- [15] Walter Greiner and Berndt Müller. *Representations of the Permutation Group and Young Tableaux*. Springer Berlin Heidelberg, Berlin, Heidelberg, 1989.
- [16] David Griffiths. *Introduction to elementary particles*. John Wiley & Sons, 2008.
- [17] Hermann Haken and Hans Christoph Wolf. *Atom-und Quantenphysik: Einführung in die experimentellen und theoretischen Grundlagen*. Springer-Verlag, 2013.
- [18] Francis Halzen and Alan D Martin. Quarks & leptons john wiley & sons. *New York*, 1984.
- [19] Francis Halzen, Alan D Martin, and Leptons Quarks. An introductory course in modern particle physics. *John and Wiley*, 1984.
- [20] Gudrun Heinrich. Introduction to quantum chromodynamics and loop calculations, SS 2018.
- [21] Michael E Peskin. *An introduction to quantum field theory*. CRC Press, 2018.
- [22] Simon Platzer and Stefan Gieseke. Coherent Parton Showers with Local Recoils. *JHEP*, 01:024, 2011.
- [23] Simon Platzer and Malin Sjödal. The Sudakov Veto Algorithm Reloaded. *Eur. Phys. J. Plus*, 127:26, 2012.
- [24] Simon Plätzer, Malin Sjödal, and Johan Thorén. Color matrix element corrections for parton showers. *JHEP*, 11:009, 2018.
- [25] Eva Popena. Hadron-kollider-experimente bei sehr hohen energien, WS 2016/2017.
- [26] Matthew D. Schwartz. *Quantum Field Theory and the Standard Model*. Cambridge University Press, 2014.
- [27] Michael H. Seymour. A Simple prescription for first order corrections to quark scattering and annihilation processes. *Nucl. Phys.*, B436:443–460, 1995.
- [28] B. F. L. Ward and S. Jadach. Dokshitser-gribov-lipatov-Altarelli-parisi evolution and the renormalization group improved yennie-frautschi-suura theory in QCD. In *High-energy physics. Proceedings, 29th International Conference, ICHEP'98, Vancouver, Canada, July 23-29, 1998. Vol. 1, 2*, pages 1628–1633, 1995. [Submitted to: Phys. Lett. B(1995)].

# Acknowledgement

First of all I would like to thank all those who supported me during the preparation of this work and who contributed a lot to the success of this work, in particular:

PD Dr. Stefan Gieseke for his excellent care and patience.

Prof. Dr. Dieter Zeppenfeld for the takeover of the second assessor.

Dr. Simon Plätzer who gave me a helpful feedback and took the time to discuss this work.

My great thanks also go to Emma Simpson Dore, who proofread my work in numerous hours. She pointed out to me the weaknesses of my letter and showed me the right paths to reach my goal at work.

Finally, I would like to thank my girlfriend Canan Kaman, who supported me in all things in this not always easy time.

