

THESIS

BY

TIGRAN SAIDNIA

Emission kernel of parton shower

Emission kernel of parton shower



Karlsruhe institute for Technology (KIT)

Institute of theoretical physics

Referents: PD Dr. Stefan Gieseke

Dr. Simon Plätzer

Supervisor: Emma Simpson

statement of originality

I hereby confirm that I have written the accompanying thesis by myself, without contributions from any sources other than those cited in the text and acknowledgements. This applies also to all graphics, drawings, maps and images included in the thesis.

Karlsruhe, February 4, 2019

Tigran Saidnia



Contents

| | |
|--|-----------|
| Table of contents | 2 |
| 0.1 Old parametrisation | 1 |
| 0.2 new parametrisation | 1 |
| 0.3 Altarelli-Parisi splitting functions | 2 |
| 0.4 Colour factor calculation | 3 |
| 1 Quark antiquark gluon emission kernel | 5 |
| 1.1 $qg\text{-}\bar{q}$ | 6 |
| 1.2 $\bar{q}g\text{-}q$ | 10 |
| 1.3 $M_1 M_2^\dagger$ | 12 |
| 1.4 $ M^2 $ | 14 |
| 2 Gluon gluon gluon emission kernel | 16 |
| 2.1 Gluon-Emitter Bubble | 17 |
| 2.1.1 One-loop corrections to the gluon self-energy diagram(Gluon-Emitter Bubble) | 19 |
| 2.2 Gluon-Spectator Bubble | 20 |
| 2.2.1 One-loop corrections to the gluon self-energy diagram (Gluon-Spectator Bubble) | 21 |
| 2.3 Interference term $M_1 M_2^\dagger$ | 21 |
| 2.4 $ M^2 $ | 23 |
| 3 Quark gluon quark emission kernel | 24 |
| 3.1 Gluon-Emitter Quark loop | 25 |
| 3.2 Gluon-Spectator Quark loop | 26 |
| 3.3 Gluon-Emitter Quark loop | 27 |
| 4 Gluon quark quark emission kernel | 28 |
| 4.1 M_1 | 29 |
| 4.2 M_2 | 30 |
| 4.3 $M_1 M_2^\dagger$ | 31 |

0.1 Old parametrisation

$$\left. \begin{aligned}
 q_i^\mu &= zp_i^\mu + y(1-z)p_j^\mu + \sqrt{zy(1-z)}m_\perp \\
 q^\mu &= (1-z)p_i^\mu + yzp_j^\mu - \sqrt{zy(1-z)}m_\perp \\
 q_j^\mu &= (1-y)p_j^\mu \\
 y &= \frac{q_i q}{p_i p_j} \\
 q_i + q &= p_i + yp_j \\
 q_j + q &= (1-z)p_i^\mu + (1+yz-y)p_j^\mu - \sqrt{zy(1-z)}m_\perp \\
 q_i \cdot q &= y(1-2z+2z^2)(p_i \cdot p_j) \\
 q_i \cdot q_j &= z(1-y)(p_i \cdot p_j) \\
 q_j \cdot q &= (1-z)(1-y)(p_i \cdot p_j)
 \end{aligned} \right\} \text{parametrisation} \quad (1)$$

0.2 new parametrisation

0.3 Altarelli-Parisi splitting functions

$$\left. \begin{aligned} \langle \hat{P}_{qq} \rangle &= C_F \left[\frac{1+z^2}{1-z} - \varepsilon(1-z) \right] \\ \langle \hat{P}_{gq} \rangle &= T_R \left[1 - \frac{2z(1-z)}{1-\varepsilon} \right] \\ \langle \hat{P}_{qg} \rangle &= C_F \left[\frac{1+(1-z)^2}{z} - \varepsilon z \right] \\ \langle \hat{P}_{gg} \rangle &= 2C_A \left[\frac{z}{1-z} + \frac{1-z}{z} + z(1-z) \right] \end{aligned} \right\} \text{splitting functions} \quad (2)$$

0.4 Colour factor calculation

fundamental representation in $SU(2)$ and $SU(3)$

$$\begin{aligned} T^a &= \tau^a \equiv \frac{\sigma^2}{2} \quad \text{with Pauli matrices } \sigma^a \\ T^a &= \vartheta^a \equiv \frac{\lambda^2}{2} \quad \text{with Gell - Mann matrices } \lambda^a \end{aligned} \quad (3)$$

$$\begin{aligned} \lambda^1 &= \begin{pmatrix} 0 & 1 & \\ 1 & 0 & \\ & & 0 \end{pmatrix}, \quad \lambda^2 = \begin{pmatrix} 0 & -i & \\ i & 0 & \\ & & 0 \end{pmatrix}, \quad \lambda^3 = \begin{pmatrix} 1 & & \\ & -1 & \\ & & 0 \end{pmatrix}, \quad \lambda^4 = \begin{pmatrix} & & 1 \\ & 0 & \\ 1 & & \end{pmatrix} \\ \lambda^5 &= \begin{pmatrix} & -i & \\ & 0 & \\ i & & \end{pmatrix}, \quad \lambda^6 = \begin{pmatrix} 0 & & \\ & 0 & 1 \\ & 1 & 0 \end{pmatrix}, \quad \lambda^7 = \begin{pmatrix} 0 & & \\ & 0 & -i \\ & i & 0 \end{pmatrix}, \quad \lambda^8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & & \\ & 1 & \\ & & -2 \end{pmatrix} \end{aligned} \quad (4)$$

As we can see, λ^3 and λ^8 are diagonal. These generators satisfy:

$$[T^a, T^b] = i\epsilon^{abc}T^c \quad (5)$$

The most common convention for the normalization of the generators in physics is:

$$\sum_{c,d} f^{acd} f^{bcd} = N\delta^{ab} \quad (6)$$

The main relation we will use later for $SU(N)$:

$$\text{tr}(T^a T^b) = T_{ij}^a T_{ji}^b = T_F \delta^{ab} \quad (7)$$

$$\sum_a (T^a T^a) = C_F \delta^{ij} \quad (8)$$

$$f^{acd} f^{bcd} = C_A \delta^{ab} \quad (9)$$

With $T_F = \frac{1}{2}$, $C_A = N$ and $C_F = \frac{N^2-1}{2N}$.

$$f^{abc} = -2i \text{tr}(T^a [T^b, T^c]) \quad (10)$$

$$d^{abc} = 2 \text{tr}(T^a T^b, T^c) \quad (11)$$

$$T^a T^b = \frac{1}{2} \left(\frac{1}{N} \delta_{ab} + (d^{abc} + i f^{abc}) T^c \right) \quad (12)$$

$$\text{tr}(T^a T^b T^c) = \frac{1}{4} (d^{abc} + i f^{abc}) \quad (13)$$

$$\text{tr}(T^a T^b T^a T^c) = \frac{-1}{4N} \delta_{bc} \quad (14)$$



$$f^{acd} f^{bcd} = N \delta^{ab} \quad (15)$$

$$f^{acd} d^{bcd} = 0 \quad (16)$$


$$f^{ade} f^{bef} f^{cfd} = \frac{N}{2} f^{abc} \quad (17)$$


Fierz identity:

$$\sum_a T_{ij}^a T_{kl}^a = \frac{1}{2} (\delta_{il} \delta_{kj} - \frac{1}{N} \delta_{ij} \delta_{kl}) \quad (18)$$


Chapter 1

Quark antiquark gluon emission kernel




1.1 $qg\text{-}\bar{q}$ 

$$M_1 = [\bar{u}_\sigma(q_i)(-ig_s\gamma^\mu \times [T^a]_{o^l}^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q)] [v_\tau(q_j)] \quad (1.1)$$



$$M_1^\dagger = [\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s\gamma^{\mu'} \times [T^b]_{o'^k}^{o'}) u_{\sigma'}(q_i) \varepsilon^{\lambda_2}_{\mu'}(q)] [\bar{v}_{\tau'}(q_j)] \quad (1.2)$$



$$|M_1|^2 = M_1 M_1^\dagger = [\bar{u}_\sigma(q_i) (-ig_s \gamma^\mu \times [T^a]_{o'}^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q) [v_\tau(q_j)]$$

$$[\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s \gamma^{\mu'} \times [T^b]_{o'}^k) u_{\sigma'}(q_i) \varepsilon^{\lambda_2*}_{\mu'}(q) [\bar{v}_{\tau'}(q_j)]] \quad (1.3)$$

$$|M_1|^2 = [\frac{-i(\not{q}_i + \not{q})}{(q_i + q)^2} (ig_s \gamma^{\mu'} \times [T^b]_{o'}^k) \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) \varepsilon^{\lambda_2*}_{\mu'}(q) \varepsilon^{\lambda_1}_\mu(q)$$

$$\times (-ig_s \gamma^\mu \times [T^a]_{o'}^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} [\bar{v}_{\tau'}(q_j) v_\tau(q_j)] \quad (1.4)$$

and after sum over the lorenz index (σ, σ') and (τ, τ') and unsing the spin addition relation:

$$\sum_{\sigma, \sigma'} \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) = \not{q}_i \delta^{\sigma\sigma'},$$

$$\sum_{\tau, \tau'} \bar{v}_\tau(q_j) v_{\tau'}(q_j) = \not{q}_j \delta^{\tau\tau'} \quad (1.5)$$

and sum over polarization index (λ_1, λ_2) :

$$\sum_{\mu, \mu'} \varepsilon^{\lambda_2*}_{\mu'}(q) \varepsilon^{\lambda_1}_\mu(q) = -g_{\mu\mu'} \delta^{ab} \quad (1.6)$$

$$|M_1|^2 = \frac{-g_s^2 [T^a]_{o'}^k [T^a]_{o'}^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \gamma^{\mu'} \not{q}_i g_{\mu'\mu} \gamma^\mu (\not{q}_i + \not{q})] [\not{q}_j] \quad (1.7)$$

from here and after contraction between all indices we can actually make statements about the last result.

$$|M_1|^2 = \frac{-g_s^2 [T^a]_{o'}^k [T^a]_{o'}^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \gamma^{\mu'} \not{q}_i \gamma_{\mu'} (\not{q}_i + \not{q})] [\not{q}_j] \quad (1.8)$$

In other words we expect the tree level diagram from LO and a number: Which means:

$$|M^2| = \left| \begin{array}{c} \text{diagram 1} \end{array} \right|^2 \otimes \left| \begin{array}{c} \text{diagram 2} \end{array} \right|^2$$

contribution from LO
a complex number

$$|M_1|^2 = \frac{-g_s^2 [T^a]_o^k [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [P_i][P_j] \otimes (\text{a complex number}) \quad (1.9)$$

Let's calculate the contribution and compare the final result with this expectation:

$$\begin{aligned}
 N &=: \gamma^{\mu'} \not{q}_i \gamma_{\mu'} = q_{i\sigma} \gamma^{\mu'} \gamma^\sigma \gamma_{\mu'} \\
 &= q_{i\sigma} (\{\gamma^{\mu'}, \gamma^\sigma\} - \gamma^\sigma \gamma^{\mu'}) \gamma_{\mu'} \\
 &= q_{i\sigma} 2g^{\mu'\sigma} \gamma_{\mu'} - d \gamma^\sigma \\
 &= (2 - d) \not{q}_i
 \end{aligned} \quad (1.10)$$

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [(\not{q}_i + \not{q}) \not{q}_i (\not{q}_i + q)] [\not{q}_j] \quad (1.11)$$

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{(q_i + q)^2 (q_i + q)^2} [\not{q}_i \not{q}_i \not{q}_i + \not{q}_i \not{q}_i \not{q} + \not{q} \not{q}_i \not{q}_i + \not{q} \not{q}_i \not{q}] [\not{q}_j] \quad (1.12)$$

For the momenta are on-shell which means:

$$\begin{aligned}
 \not{q}_i \not{q}_i &= q_i^2 = m_i^2 \\
 \not{q} \not{q} &= q^2 = m^2 \\
 \not{q}_j \not{q}_j &= q_j^2 = m_j^2
 \end{aligned} \quad (1.13)$$

we can first neglect the mass of patrons and ignore each term with $\not{q}_i \not{q}_i$ and $\not{q} \not{q}$ as well.

$$|M_1|^2 = -(2 - d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{(2q_i q)(2q_i q)} [\not{q} \not{q}_i \not{q}] [\not{q}_j] \quad (1.14)$$

$$\begin{aligned}
L &= \not{q}_i \not{q}_\mu (\{\gamma^\mu, \gamma^\sigma\} - \gamma^\sigma \gamma^\mu) \\
&= \not{q}_i [2q_i^\mu q_\mu - q_{i\sigma} q_\mu \gamma^\mu \gamma^\sigma] \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q_\mu [\gamma^\mu \gamma^\mu \gamma^\sigma] \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q_\mu \left[\frac{\gamma^\mu \gamma^\mu}{2} + \frac{\gamma^\mu \gamma^\mu}{2} \right] \gamma^\sigma \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q_\mu [g^{\mu\mu}] \gamma^\sigma \\
&= \not{q}_i (2q_i q) - q_\mu q_{i\sigma} q^\mu \gamma^\sigma \\
&= \not{q}_i (2q_i q) - q^2 \not{q}_i \\
&= \not{q}_i (2q_i q)
\end{aligned} \tag{1.15}$$

After inserting the last result of L and simplify the term $(2q_i q)$ from the denominator and nominator because , we get:

$$|M_1|^2 = -(2-d) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{q}_i] [\not{q}_j] \tag{1.16}$$

Now we are going to use the parametrisation from equation (1) to reduce the 3-member matrix element to 2-member and take out the singularity term from the amplitude.


$$|M_1|^2 = (d-2) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [(1-z) \not{p}_i + zy \not{p}_j - \sqrt{zy(1-z)} \not{m}_\perp] [(1-y) \not{p}_j] \tag{1.17}$$

Multiplying the both sides


$$\begin{aligned}
|M_1|^2 &= (d-2) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [(1-z)(1-y) \not{p}_i \not{p}_j \\
&\quad + zy(1-y) \not{p}_j \not{p}_j + (1-y) \sqrt{zy(1-z)} \not{m}_\perp \not{p}_j]
\end{aligned} \tag{1.18}$$

and under consideration of the fact that p_i and p_j are the on-shell momenta of the emitter and spectator partons, we can ignore the terms with $\not{p}_i \not{p}_i$ and $\not{p}_j \not{p}_j$. The $p_i \cdot m_\perp$ and $p_j \cdot m_\perp$ are always 0 because the p_i and p_j are lightlike, i.e. zero transverse component. So those terms can be neglected.


$$|M_1|^2 = (d-2)(1-z)(1-y) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i] [\not{p}_j] \tag{1.19}$$

1.2 $\bar{q}g$ -q

$$M_2 = \left[\frac{i(\not{q}_j + \not{q})}{(q_j + q)^2} (-ig_s \gamma^\nu \times [T^c]_f^m) v_\tau(q_j) \varepsilon^{\lambda_3}_\nu(q) [u_\sigma(q_i)] \right] \quad (1.20)$$



$$M_2^\dagger = [\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q) [\bar{u}_{\sigma'}(q_i)] \quad (1.21)$$



$$|M_2|^2 = M_2 M_2^\dagger = \left[\frac{i(\not{q}_j + \not{q})}{(q_j + q)^2} (-ig_s \gamma^\nu \times [T^c]_f^m) v_\tau(q_j) \varepsilon^{\lambda_3}_\nu(q) [u_\sigma(q_i)] \right] \quad (1.22)$$

$$\left[\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q) [\bar{u}_{\sigma'}(q_i)] \right]$$

$$|M_2|^2 = \frac{g_s^2 [T^c]_f^m [T^d]_{f'}^n}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\nu v_\tau(q_j) \bar{v}_{\tau'}(q_j) \varepsilon^{\lambda_3}_\nu(q) \varepsilon^{\lambda_4}_{\nu'}(q) \gamma^{\nu'} (\not{q}_j + \not{q})] \quad (1.23)$$

$$[u_\sigma(q_i)] [\bar{u}_{\sigma'}(q_i)]$$

and after sum over the lorenz index (σ, σ') and (τ, τ') and unsing the spin addition relation:

$$\sum_{\sigma, \sigma'} \bar{u}_\sigma(q_i) u_{\sigma'}(q_i) = \not{q}_i \delta^{\sigma\sigma'}, \quad (1.24)$$

$$\sum_{\tau, \tau'} \bar{v}_\tau(q_j) v_{\tau'}(q_j) = \not{q}_j \delta^{\tau\tau'}$$

and sum over polarization index (λ_3, λ_4) :

$$\sum_{\nu, \nu'} \varepsilon^{\lambda_4*}_{\nu'}(q) \varepsilon^{\lambda_3}_\nu(q) = -g_{\nu\nu'} \delta^{cd} \quad (1.25)$$


$$|M_2|^2 = \frac{g_s^2 [T^c]_f^m [T^c]_{f'}^n}{(q_j + q)^2 (q_j + q)^2} [(\not{q}_j + \not{q}) \gamma^\nu \not{q}_j (-g_{\nu\nu'}) \gamma^{\nu'} (\not{q}_j + \not{q})] [\not{q}_i] \quad (1.26)$$

After the same calculation from the last part, we'll get:

$$|M_2|^2 = (d - 2) \frac{g_s^2 [T^c]_f^m [T^c]_{f'}^n}{(2qq_j)} [\not{q}] [\not{q}_i] \quad (1.27)$$

finally:

$$|M_2|^2 = -(d - 2) y z^2 \frac{g_s^2 [T^c]_f^m [T^c]_{f'}^n}{2(1 - z)(1 - y)(p_i \cdot p_j)} [\not{p}_i] [\not{p}_j] \quad (1.28)$$

1.3 $M_1 M_2^\dagger$ 

$$M_1 M_2^\dagger = [\bar{u}_\sigma(q_i) (-ig_s \gamma^\mu \times [T^a]_o^l) \frac{i(\not{q}_i + \not{q})}{(q_i + q)^2} \varepsilon^{\lambda_1}_\mu(q)] [v_\tau(q_j)]$$

$$[\bar{v}_{\tau'}(q_j) (ig_s \gamma^{\nu'} \times [T^d]_{f'}^n) \frac{-i(\not{q}_j + \not{q})}{(q_j + q)^2} \varepsilon^{\lambda_4}_{\nu'}(q)] [u_{\sigma'}(q_i)] \quad (1.29)$$

$$M_1 M_2^\dagger = \frac{g_s^2 [T^a]_o^l [T^d]_{f'}^n}{(2q_i q)(2q_j q)} [\not{q}_i \gamma^\mu (\not{q}_i + \not{q})] \varepsilon^{\lambda_1}_\mu(q) \varepsilon^{\lambda_4}_{\nu'}(q)$$

$$[\not{q}_j \gamma^{\nu'} (\not{q}_j + \not{q})] \quad (1.30)$$

$$M_1 M_2^\dagger = \frac{g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} [\not{q}_i \gamma^\mu (\not{q}_i + \not{q})] - g_{\mu\nu'}$$

$$[\not{q}_j \gamma^{\nu'} (\not{q}_j + \not{q})] \quad (1.31)$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} [\not{q}_i \gamma^\mu (\not{q}_i + \not{q})] [\not{q}_j \gamma_\mu (\not{q}_j + \not{q})] \quad (1.32)$$

Expectation:

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} [(\not{q}_i + \not{q}) \gamma^\mu \not{q}_i] [(\not{q}_j + \not{q}) \gamma_\mu \not{q}_j] \quad (1.33)$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} [-(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu + 2(\not{q}_i + \not{q}) q_i^\mu]$$

$$[-(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu + 2(\not{q}_j + \not{q}) q_{j\mu}] \quad (1.34)$$

$$|M^2| = \left| \begin{array}{c} \text{diagram with two shaded circles connected by two horizontal lines labeled } P_i \text{ and } P_j \\ \text{contribution from LO} \end{array} \right|^2 \otimes \left| \begin{array}{c} \text{diagram with a shaded circle and a wavy line} \\ \text{a complex number} \end{array} \right|^2$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} \begin{aligned} & [(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu][(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu] \\ & - 2[(\not{q}_i + \not{q}) \not{q}_i \gamma^\mu][(\not{q}_j + \not{q}) q_{j\mu}] \\ & - 2[(\not{q}_i + \not{q}) q_i^\mu][(\not{q}_j + \not{q}) \not{q}_j \gamma_\mu] \\ & + 4[(\not{q}_i + \not{q}) q_i^\mu][(\not{q}_j + \not{q}) q_{j\mu}] \end{aligned} \quad (1.35)$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(2q_i q)(2q_j q)} \begin{aligned} & [\not{q} \not{q}_i \gamma^\mu][\not{q} \not{q}_j \gamma_\mu] \\ & - 2[\not{q} \not{q}_i \gamma^\mu][(\not{q} + \not{q}_j) q_{j\mu}] \\ & - 2[(\not{q}_i + \not{q}) q_i^\mu][\not{q} \not{q}_j \gamma_\mu] \\ & + 4[(\not{q}_i + \not{q}) q_i^\mu][(\not{q}_j + \not{q}) q_{j\mu}] \end{aligned} \quad (1.36)$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{4(1-z)(1-y)y(1-2z+2z^2)(p_i \cdot p_j)(p_i \cdot p_j)} \begin{aligned} & [y(1-2z+2z^2) \not{p}_i \not{p}_j \gamma^\mu][(1-z)(1-y) \not{p}_i \not{p}_j \gamma_\mu] \\ & + 4(q_i^\mu \cdot q_{j\mu})[(\not{q}_i + \not{q})][(\not{q}_j + \not{q})] \end{aligned} \quad (1.37)$$


$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{4(1-z)(1-y)y(1-2z+2z^2)(p_i \cdot p_j)(p_i \cdot p_j)} \begin{aligned} & [y(1-2z+2z^2) \not{p}_i \not{p}_j \gamma^\mu][(1-z)(1-y) \not{p}_i \not{p}_j \gamma_\mu] \\ & + 4(p_i \cdot p_j)[(\not{p}_i + y \not{p}_j)][(1-z) \not{p}_i + (1+yz-y) \not{p}_j - \sqrt{zy(1-z)} \not{m}] \end{aligned} \quad (1.38)$$

$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(1-z)(1-y)y(1-2z+2z^2)(p_i \cdot p_j)} z(1-y)[\not{p}_i][\not{p}_j] \quad (1.39)$$


$$M_1 M_2^\dagger = \frac{-g_s^2 [T^a]_o^l [T^a]_{f'}^n}{(1-z)y(1-2z+2z^2)(p_i \cdot p_j)} z[\not{p}_i][\not{p}_j] \quad (1.40)$$

1.4 $|M^2|$

$$|M|^2 = |M_1|^2 + |M_2|^2 + M_1 M_2^\dagger + M_1^\dagger M_2 \quad (1.41)$$



$$|M|^2 = |M_1|^2 + |M_2|^2 + 2RE(M_1 M_2^\dagger) \quad (1.42)$$



$$\begin{aligned}
 |M|^2 = & (d-2)(1-z)(1-y) \frac{g_s^2 [T^a]_o^k [T^a]_o^l}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\
 & - (d-2)yz^2 \frac{g_s^2 [T^c]_f^m [T^c]_f^n}{2(1-z)(1-y)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\
 & + 2RE\left(\left(\frac{-2z}{z-1}\right) \frac{g_s^2 [T^a]_o^l [T^a]_f^n}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j]\right)
 \end{aligned} \quad (1.43)$$

$$T^a_{ok} T^a_{lo} = \frac{1}{2}(\delta_{oo}\delta_{lk} - \frac{1}{N}\delta_{ok}\delta_{lo}) = \frac{1}{2}(N\delta_{lk} - \frac{1}{N}\delta_{lk}) = C_F\delta_{lk} \quad (1.44)$$

After summation over the final colour states and averaging over initial colour states we get:

$$T^a_{ok} T^a_{lo} = C_F \delta_{lk} = \frac{1}{N} \sum_{l=1}^N \delta_{lk} C_F = C_F \quad (1.45)$$

The same calculation for $T^c_{mf} T^c_{fn}$ and $T^a_{ol} T^a_{fn}$ turns C_F out as the colour factor. Now we are going to compute the splitting function in the case of the colinearity, wich means, if:

$$y \longrightarrow 0 \quad (1.46)$$

$$\begin{aligned} |M|^2 = & (d-2)(1-z)(1-y) \frac{g_s^2 C_F}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ & -(d-2)yz^2 \frac{g_s^2 C_F}{2(1-z)(1-y)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ & + 2RE\left(\left(\frac{-2z}{z-1}\right) \frac{g_s^2 C_F}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j]\right) \end{aligned} \quad (1.47)$$

$$|M|^2 = C_F \left((d-2)(1-z) - \frac{4z}{z-1} \right) \frac{g_s^2}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \quad (1.48)$$


for

$$d = 4 - 2\epsilon \quad (1.49)$$


$$\begin{aligned} |M|^2 = & C_F \left((4-2\epsilon-2)(1-z) + \frac{4z}{1-z} \right) \frac{g_s^2}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ = & C_F \left(\frac{2(1-\epsilon)(1-z)^2 + 4z}{1-z} \right) \frac{g_s^2}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ = & C_F \left(\frac{2-4z+2z^2-\epsilon(1-z)^2+4z}{1-z} \right) \frac{g_s^2}{2y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ = & C_F \left(\frac{(1+z^2)}{1-z} - \epsilon(1-z) \right) \frac{g_s^2}{y(1-2z+2z^2)(p_i \cdot p_j)} [\not{p}_i][\not{p}_j] \\ = & \langle \hat{P}_{qq} \rangle \frac{g_s^2}{q_i \cdot q} [\not{p}_i][\not{p}_j] \end{aligned} \quad (1.50)$$

Chapter 2

Gluon gluon gluon emission kernel




2.1 Gluon-Emitter Bubble




$$M_1 = \left[\frac{-i}{(q + q_i)^2} (-g_s f^{a o l} (g^{\mu \zeta} (q - q_i)^\eta + g^{\zeta \eta} (-q - (q + q_i))^\mu + g^{\eta \mu} (q_i + q_i + q)^\zeta) \right. \\ \left. \varepsilon^{\lambda_1}_\mu(q) \varepsilon^{\lambda_6}_\zeta(q) [\varepsilon^\theta_{\tau'}(q_j)] \right] \quad (2.1)$$

$$M_1 = \left[\frac{-i}{(q_i + q)^2} (-g_s f^{a o l} (g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q + q_i)^\mu + g^{\eta \mu} (2q_i + q)^\zeta) \right. \\ \left. \varepsilon^{\lambda_1}_\mu(q_i) \varepsilon^{\lambda_6}_\zeta(q) [\varepsilon^\theta_{\tau'}(q_j)] \right] \quad (2.2)$$



$$M_1^\dagger = \left[\frac{i}{(q_i + q)^2} (-g_s f^{a' k o'} (-g^{\mu' \eta'} (2q_i + q)^{\zeta'} + g^{\eta' \zeta'} (2q + q_i)^{\mu'} + g^{\zeta' \mu'} (q_i - q)^{\eta'}) \right. \\ \left. \varepsilon^{\lambda_2}_{\mu'}(q_i) \varepsilon^{\lambda_5}_{\zeta'}(q) [\varepsilon^{\theta'}_{\tau'}(q_j)] \right] \quad (2.3)$$

$$|M_1|^2 = \left[\frac{-i}{(q_i + q)^2} (-g_s f^{a o l} (g^{\mu \zeta} (q - q_i)^\eta - g^{\zeta \eta} (2q + q_i)^\mu + g^{\eta \mu} (2q_i + q)^\zeta) \right. \\ \left. \varepsilon^{\lambda_1}_\mu(q_i) \varepsilon^{\lambda_2}_{\mu'}(q_i) \varepsilon^{\lambda_6}_\zeta(q) \varepsilon^{\lambda_5}_{\zeta'}(q) \right. \\ \left. (-g_s f^{a' k o'} (-g^{\mu' \eta'} (2q_i + q)^{\zeta'} + g^{\eta' \zeta'} (2q + q_i)^{\mu'} + g^{\zeta' \mu'} (q_i - q)^{\eta'}) \frac{i}{(q_i + q)^2} [g_{\gamma \delta}] \right] \quad (2.4)$$



$$\begin{aligned}
N \equiv & g_{\mu\mu'} g_{\zeta\zeta'} [-g^{\mu\zeta} g^{\mu'\eta'} (q - q_i)^\eta (2q_i + q)^{\zeta'} + g^{\mu\zeta} g^{\eta'\zeta'} (q - q_i)^\eta (2q + q_i)^{\mu'} \\
& + g^{\mu\zeta} g^{\zeta'\mu'} (q - q_i)^\eta (q_i - q)^{\eta'} + g^{\zeta\eta} g^{\mu'\zeta'} (2q + q_i)^\mu (2q_i + q)^{\zeta'} \\
& - g^{\zeta\eta} g^{\eta'\zeta'} (2q + q_i)^\mu (2q + q_i)^{\mu'} - g^{\zeta\eta} g^{\zeta'\mu'} (2q + q_i)^\mu (q_i - q)^{\eta'} \\
& - g^{\eta\mu} g^{\mu'\eta'} (2q_i + q)^\zeta (2q_i + q)^{\zeta'} + g^{\eta\mu} g^{\eta'\zeta'} (2q_i + q)^\zeta (2q + q_i)^{\mu'} \\
& + g^{\eta\mu} g^{\zeta'\mu'} (2q_i + q)^\zeta (q_i - q)^{\eta'}] [g_{\gamma\delta}]
\end{aligned} \quad (2.5)$$


$$\begin{aligned}
N \equiv & [-(q - q_i)^\eta (2q_i + q)^{\eta'} + (q - q_i)^\eta (2q + q_i)^{\eta'} + d(q - q_i)^\eta (q_i - q)^{\eta'} \\
& + (2q + q_i)^{\eta'} (2q_i + q)^\eta - g^{\eta\eta'} (2q + q_i)^\mu (2q + q_i)_\mu - (2q + q_i)^\eta (q_i - q)^{\eta'} \\
& - g^{\eta\eta'} (2q_i + q)^\zeta (2q_i + q)_\zeta + (2q_i + q)^{\eta'} (2q + q_i)^\eta + (2q_i + q)^\eta (q_i - q)^{\eta'}] [g_{\gamma\delta}]
\end{aligned} \quad (2.6)$$

$$\begin{aligned}
N \equiv & [-(q^\eta q^{\eta'} + 2q^\eta q_i^{\eta'} - q_i^\eta q^{\eta'} - 2q_i^\eta q_i^{\eta'}) + (2q^\eta q^{\eta'} + q^\eta q_i^{\eta'} - 2q_i^\eta q^{\eta'} - q_i^\eta q_i^{\eta'}) \\
& + (dq^\eta q_i^{\eta'} - dq^\eta q^{\eta'} - dq_i^\eta q_i^{\eta'} + dq_i^\eta q^{\eta'}) + (4q^{\eta'} q_i^\eta + 2q^{\eta'} q^\eta + 2q_i^{\eta'} q_i^\eta + q_i^{\eta'} q^\eta) \\
& - (-2q^\eta q^{\eta'} + 2q^\eta q_i^{\eta'} - q_i^\eta q^{\eta'} + q_i^\eta q_i^{\eta'}) + (2q^{\eta'} q^\eta + q^{\eta'} q_i^\eta + 4q_i^{\eta'} q^\eta + 2q_i^{\eta'} q_i^\eta) \\
& + (-q^\eta q^{\eta'} + q^\eta q_i^{\eta'} - 2q_i^\eta q^{\eta'} + 2q_i^\eta q_i^{\eta'}) - g^{\eta\eta'} (5q^2 + 5q_i^2 + 8qq_i)] [g_{\gamma\delta}]
\end{aligned} \quad (2.7)$$

$$\begin{aligned}
N \equiv & [(6 - d)q^\eta q^{\eta'} + (d + 3)q^\eta q_i^{\eta'} + (d + 3)q_i^\eta q^{\eta'} + (6 - d)q_i^\eta q_i^{\eta'} \\
& - g^{\eta\eta'} (5q^2 + 5q_i^2 + 8qq_i)] [g_{\gamma\delta}]
\end{aligned} \quad (2.8)$$

$$\begin{aligned}
|M_1|^2 = & \frac{g_s^2 f^{aol} f^{ako}}{(q_i + q)^2 (q_i + q)^2} [(6 - d)q^\eta q^{\eta'} + (d + 3)q^\eta q_i^{\eta'} + (d + 3)q_i^\eta q^{\eta'} + (6 - d)q_i^\eta q_i^{\eta'} \\
& - g^{\eta\eta'} (5q^2 + 5q_i^2 + 8qq_i)] [g_{\gamma\delta}]
\end{aligned} \quad (2.9)$$

2.1.1 One-loop corrections to the gluon self-energy diagram(Gluon-Emitter Bubble)




$$|M_1|_{Ghost\ loop}^2 = \frac{g_s^2 f^{aol} f^{ako}}{(q_i + q)^2 (q_i + q)^2} [-q_i^\eta q^{\eta'} - q^\eta q_i^{\eta'}] [g_{\gamma\delta}] \quad (2.10)$$

$$\begin{aligned} |M_1'|^2 &= |M_1|^2 + |M_1|_{Ghost\ loop}^2 \\ &= \frac{g_s^2 f^{aol} f^{ako}}{(q_i + q)^2 (q_i + q)^2} [(6 - d)q^\eta q^{\eta'} + (d + 3)q^\eta q_i^{\eta'} \\ &\quad + (d + 3)q_i^\eta q^{\eta'} + (6 - d)q_i^\eta q_i^{\eta'} - g^{\eta\eta'} (5q^2 + 5q_i^2 + 8qq_i) - q_i^\eta q^{\eta'} - q^\eta q_i^{\eta'}] [g_{\gamma\delta}] \end{aligned} \quad (2.11)$$

$$\begin{aligned} |M_1'|^2 &= \frac{g_s^2 f^{aol} f^{ako}}{(q_i + q)^2 (q_i + q)^2} [(6 - d)q^\eta q^{\eta'} + (d + 2)q^\eta q_i^{\eta'} \\ &\quad + (d + 2)q_i^\eta q^{\eta'} + (6 - d)q_i^\eta q_i^{\eta'} - g^{\eta\eta'} (5q^2 + 5q_i^2 + 8qq_i)] [g_{\gamma\delta}] \end{aligned} \quad (2.12)$$

2.2 Gluon-Spectator Bubble




$$|M_2|^2 = \left[\frac{-i}{(q_j + q)^2} (-g_s f^{b f m} (g^{\tau\gamma} (-2q_j - q)^\rho + g^{\gamma\rho} (2q + q_j)^\tau + g^{\rho\tau} (q_j - q)^\gamma) \right. \\ \left. g_{\tau\tau'} g_{\rho\rho'} (-g_s f^{b' n f'} (g^{\rho'\delta} (-2q - q_j)^{\tau'} + g^{\delta\tau'} (2q_j + q)^{\rho'} + g^{\tau'\rho'} (q - q_j)^\delta) \frac{i}{(q_j + q)^2} \right] [g_{\eta\eta'}] \quad (2.13)$$

$$|M_2|^2 = \frac{g_s^2 f^{b f m} f^{b' n f'} \delta^{aa'} \delta^{ff'} \delta^{bb'}}{(q_j + q)^2 (q_j + q)^2} [g_{\tau\tau'} g_{\rho\rho'} (g^{\tau\gamma} (2q_j + q)^\rho g^{\rho'\delta} (2q + q_j)^{\tau'} \\ - g^{\tau\gamma} (2q_j + q)^\rho g^{\delta\tau'} (2q_j + q)^{\rho'} - g^{\tau\gamma} (2q_j + q)^\rho g^{\tau'\rho'} (q - q_j)^\delta - g^{\gamma\rho} (2q + q_j)^\tau g^{\rho'\delta} (2q + q_j)^{\tau'} \\ + g^{\gamma\rho} (2q + q_j)^\tau g^{\delta\tau'} (2q_j + q)^{\rho'} + g^{\gamma\rho} (2q + q_j)^\tau g^{\tau'\rho'} (q - q_j)^\delta - g^{\rho\tau} (q_j - q)^\gamma g^{\rho'\delta} (2q + q_j)^{\tau'} \\ + g^{\rho\tau} (q_j - q)^\gamma g^{\delta\tau'} (2q_j + q)^{\rho'} + g^{\rho\tau} (q_j - q)^\gamma g^{\tau'\rho'} (q - q_j)^\delta) [g_{\eta\eta'}] \quad (2.14)$$

$$|M_2|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [(2q + q_j)^\gamma (2q_j + q)^\delta \\ - g^{\delta\gamma} (2q_j + q)^\rho (2q_j + q)_\rho - (2q_j + q)^\gamma (q - q_j)^\delta - g^{\delta\gamma} (2q + q_j)^\tau (2q + q_j)_\tau \\ + (2q_j + q)^\gamma (2q + q_j)^\delta + (2q + q_j)^\gamma (q - q_j)^\delta - (q_j - q)^\gamma (2q + q_j)^\delta \\ + (q_j - q)^\gamma (2q_j + q)^\delta + d(q_j - q)^\gamma (q - q_j)^\delta] [g_{\eta\eta'}] \quad (2.15)$$

$$|M_2|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [(3 + d) q^\gamma q_j^\delta + (6 - d) q^\gamma q^\delta \\ + (6 - d) q_j^\gamma q_j^\delta + (3 + d) q_j^\gamma q^\delta - g^{\delta\gamma} (5q_j^2 + 5q^2 + 8qq_j) \\] [g_{\eta\eta'}] \quad (2.16)$$


2.2.1 One-loop corrections to the gluon self-energy diagram (Gluon-Spectator Bubble)



$$|M_1|_{Ghost\ loop}^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [-q_j^\gamma q^\delta - q^\delta q_j^\gamma] [g_{\gamma\delta}] \quad (2.17)$$

$$|M_2'|^2 = \frac{g_s^2 f^{b f m} f^{b n f}}{(q_j + q)^2 (q_j + q)^2} [(2 + d)q^\gamma q_j^\delta + (6 - d)q^\gamma q^\delta + (6 - d)q_j^\gamma q_j^\delta + (2 + d)q_j^\gamma q^\delta - g^{\delta\gamma} (5q_j^2 + 5q^2 + 8qq_j)] [g_{\eta\eta'}] \quad (2.18)$$

2.3 Interference term $M_1 M_2^\dagger$



$$\begin{aligned}
M_1 M_2^\dagger = & \left[\frac{-i}{(q_i + q)^2} (-g_s f^{a l o} (-g^{\mu\eta} (2q_i + q)^\zeta + g^{\eta\zeta} (2q + q_i)^\mu + g^{\zeta\mu} (q_i - q)^\eta) \right. \\
& \left. \varepsilon^{\lambda_1}{}_\mu(q_i) \varepsilon^{\lambda_6}{}_\zeta(q) [\varepsilon^{\theta}{}_\tau{}^*(q_j)] \right] \\
& \left[\frac{i}{(q + q_j)^2} (-g_s f^{f' n b'} (-g^{\tau'\delta} (2q_j + q)^{\rho'} + g^{\delta\rho'} (2q + q_j)^{\tau'} + g^{\rho'\tau'} (q_j - q)^\delta) \right. \\
& \left. \varepsilon^{\theta'}{}_{\tau'}{}^*(q_j) \varepsilon^{\lambda_8}{}_{\rho'}{}^*(q) [\varepsilon^{\lambda_2}{}_{\mu'}{}^*(q_i)] \right]
\end{aligned} \tag{2.19}$$

$$\begin{aligned}
M_1 M_2^\dagger = & \frac{g_s^2 f^{a l o} f^{f' n b'} \delta^{aa'} \delta^{ob'} \delta f f'}{(q_i + q)^2 (q_j + q)^2} [g_{\mu\mu'} g_{\zeta\rho'} g_{\tau\tau'} \\
& (-g^{\mu\eta} (2q_i + q)^\zeta + g^{\eta\zeta} (2q + q_i)^\mu + g^{\zeta\mu} (q_i - q)^\eta) \\
& (-g^{\tau'\delta} (2q_j + q)^{\rho'} + g^{\delta\rho'} (2q + q_j)^{\tau'} + g^{\rho'\tau'} (q_j - q)^\delta)]
\end{aligned} \tag{2.20}$$


$$\begin{aligned}
M_1 M_2^\dagger = & \frac{g_s^2 f^{a l o} f^{f n o}}{(q_i + q)^2 (q_j + q)^2} [g_{\mu\mu'} g_{\zeta\rho'} g_{\tau\tau'} \\
& (g^{\mu\eta} (2q_i + q)^\zeta g^{\tau'\delta} (2q_j + q)^{\rho'} - g^{\mu\eta} (2q_i + q)^\zeta g^{\delta\rho'} (2q + q_j)^{\tau'} - g^{\mu\eta} (2q_i + q)^\zeta g^{\rho'\tau'} (q_j - q)^\delta \\
& - g^{\eta\zeta} (2q + q_i)^\mu g^{\tau'\delta} (2q_j + q)^{\rho'} + g^{\eta\zeta} (2q + q_i)^\mu g^{\delta\rho'} (2q + q_j)^{\tau'} + g^{\eta\zeta} (2q + q_i)^\mu g^{\rho'\tau'} (q_j - q)^\delta \\
& - g^{\zeta\mu} (q_i - q)^\eta g^{\tau'\delta} (2q_j + q)^{\rho'} + g^{\zeta\mu} (q_i - q)^\eta g^{\delta\rho'} (2q + q_j)^{\tau'} + g^{\zeta\mu} (q_i - q)^\eta g^{\rho'\tau'} (q_j - q)^\delta)]
\end{aligned} \tag{2.21}$$

$$\begin{aligned}
M_1 M_2^\dagger = & \frac{g_s^2 f^{a l o} f^{f n o}}{(q_i + q)^2 (q_j + q)^2} [g_{\eta'}{}^\eta g_\gamma{}^\delta (2q_i + q)_{\rho'} (2q_j + q)^{\rho'} \\
& - g_{\eta'}{}^\eta (2q_i + q)^\eta (2q + q_j)_\gamma - g_{\eta'}{}^\eta (2q_i + q)^\delta (q_j - q)^\delta \\
& - g_\gamma{}^\delta (2q + q_i)_{\eta'} (2q_j + q)^\eta + g^{\eta\delta} (2q + q_i)_{\eta'} (2q + q_j)_\gamma \\
& + g_\gamma{}^\eta (2q + q_i)_{\eta'} (q_j - q)^\eta - g_\gamma{}^\delta (q_i - q)^\eta (2q_j + q)_{\eta'} \\
& + g_{\eta'}{}^\delta (q_i - q)^\eta (2q + q_j)_\gamma + g_{\eta'}{}^{\tau'} g_{\gamma\tau'} (q_i - q)^\eta (q_j - q)^\delta]
\end{aligned} \tag{2.22}$$


2.4 $|M^2|$

Chapter 3


Quark gluon quark emission kernel




3.1 Gluon-Emitter Quark loop



3.2 Gluon-Spectator Quark loop





3.3 Gluon-Emitter Quark loop




Chapter 4

Gluon quark quark emission kernel



4.1 M_1 

4.2 M_2 

4.3 $M_1 M_2^\dagger$ 