

Phys 443

Computational Physics I

Probability Distributions

Probability and Statistics

Probability:

Know parameters of the theory → Predict distributions of possible experiment outcomes

Statistics

Know the outcome of an experiment → Extract information about the parameters and/or the theory


Probability distributions

These distributions represent good approximations to "real life " .

- We use probability distributions because they work –they fit lots of data in real world



Sarı kantaron



Ht (cm) 1996

Height (cm) of *Hypericum cumulicola* at Archbold Biological Station

Distributions


Distributions

In general, the result of repeating the same measurement many times does not lead to the same result.

Experiment:

- Measure the length of one side of your table 10 times and display the results in a histogram.
- What happens if you repeat the measurement 50 times?

Distributions




Distributions

There are two categories of random variables.
These are:

- Discrete random variables
- Continuous random variables

The Gaussian Distribution

- Discovered in 1733 by de Moivre as an approximation to the binomial distribution when the number of trials is large
- Derived in 1809 by Gauss
- Importance lies in the Central Limit Theorem, which states that the sum of a large number of independent random variables (binomial, Poisson, etc.) will approximate a normal distribution
 - Example: Human height is determined by a large number of factors, both genetic and environmental, which are additive in their effects. Thus, it follows a normal distribution.



Abraham de Moivre
(1667-1754)



Karl F. Gauss
(1777-1855)

The Gaussian Distribution


- The Gaussian probability distribution is perhaps the most used distribution in all of science.
- Sometimes it is called the “bell shaped curve” or normal distribution.
- Unlike the binomial and Poisson distribution, the Gaussian is a continuous distribution:

$$p(y) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(y-\mu)^2}{2\sigma^2}}$$

μ = mean of distribution (also at the same place as mode and median)


σ^2 = variance of distribution

y is a continuous variable ($-\infty \leq y \leq \infty$)




The Normal Distribution


Normal Distributions: $\sigma=1$



The Normal Distribution



The Gaussian Distribution



The Gaussian Distribution

Two parameters, mean and standard deviation, completely determine the Gaussian distribution. The shape and location of the normal curve changes as the mean and standard deviation change.

$$\int_{-\infty}^{+\infty} \frac{1}{\sigma\sqrt{2\pi}} \cdot e^{-\frac{1}{2}\left(\frac{y-\mu}{\sigma}\right)^2} dy = 1$$

Probability (P) of y being in the range $[a, b]$ is given by an integral:

$$P(a < y < b) = \int_a^b p(y)dy = \frac{1}{\sigma\sqrt{2\pi}} \int_a^b e^{-\frac{(y-\mu)^2}{2\sigma^2}} dy$$

The integral for arbitrary a and b cannot be evaluated analytically.

Method of Probability Calculation

The probability that a continuous random variable x lies between a lower limit a and an upper limit b is

$$\begin{aligned} P(a < x < b) &= (\text{cumulative area to the left of } b) - \\ &\quad (\text{cumulative area to the left of } a) \\ &= P(x < b) - P(x < a) \end{aligned}$$





Table A. The percentage probability,
 $Prob(\text{within } t\sigma) = \int_{X-t\sigma}^{X+t\sigma} G_{X,\sigma}(x) dx$,
as a function of t .



t	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.00	0.80	1.60	2.39	3.19	3.99	4.78	5.58	6.38	7.17
0.1	7.97	8.76	9.55	10.34	11.13	11.92	12.71	13.50	14.28	15.07
0.2	15.85	16.63	17.41	18.19	18.97	19.74	20.51	21.28	22.05	22.82
0.3	23.58	24.34	25.10	25.86	26.61	27.37	28.12	28.86	29.61	30.35
0.4	31.08	31.82	32.55	33.28	34.01	34.73	35.45	36.16	36.88	37.59
0.5	38.29	38.99	39.69	40.39	41.08	41.77	42.45	43.13	43.81	44.48
0.6	45.15	45.81	46.47	47.13	47.78	48.43	49.07	49.71	50.35	50.98
0.7	51.61	52.23	52.85	53.46	54.07	54.67	55.27	55.87	56.46	57.05
0.8	57.63	58.21	58.78	59.35	59.91	60.47	61.02	61.57	62.11	62.65
0.9	63.19	63.72	64.24	64.76	65.28	65.79	66.29	66.80	67.29	67.78
1.0	68.27	68.75	69.23	69.70	70.17	70.63	71.09	71.54	71.99	72.43
1.1	72.87	73.30	73.73	74.15	74.57	74.99	75.40	75.80	76.20	76.60
1.2	76.99	77.37	77.75	78.13	78.50	78.87	79.23	79.59	79.95	80.29
1.3	80.64	80.98	81.32	81.65	81.98	82.30	82.62	82.93	83.24	83.55
1.4	83.85	84.15	84.44	84.73	85.01	85.29	85.57	85.84	86.11	86.38

The Gaussian Distribution

- The total area under the curve is normalized to one by the $\sigma\sqrt{2\pi}$ factor.

$$P(-\infty < y < \infty) = \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{(y-\mu)^2}{2\sigma^2}} dy = 1$$


- We often talk about a measurement being a certain number of standard deviations (σ) away from the mean (μ) of the Gaussian.

We can associate a probability for a measurement to be $|\mu - n\sigma|$ away from the mean just by calculating the area outside of this region.

$n\sigma$	Prob. of exceeding $\mu \pm n\sigma$
0.67	0.5
1	0.32
2	0.05
3	0.003
4	0.00006

It is very unlikely ($< 0.3\%$) that a measurement taken at random from a Gaussian *pdf* will be more than $\pm 3\sigma$ from the true mean of the distribution.

The Gaussian Distribution



The Gaussian Distribution

$$\int_{\mu-\sigma}^{\mu+\sigma} \frac{1}{\sigma\sqrt{2\pi}} \bullet e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx = .68$$

$$\int_{\mu-2\sigma}^{\mu+2\sigma} \frac{1}{\sigma\sqrt{2\pi}} \bullet e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx = .95$$

$$\int_{\mu-3\sigma}^{\mu+3\sigma} \frac{1}{\sigma\sqrt{2\pi}} \bullet e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx = .997$$

The Gaussian Distribution

Where the Gaussian distribution apply?

- The result of repeating an experiment many times produces a spread of answers whose distribution is approximately Gaussian
- If the individual errors contributing to the final answer is small, the approximation to a Gaussian is especially good


The Standard Gaussian Distribution

- To find $P(a < x < b)$, we need to find the area under the appropriate normal curve.
- To simplify the tabulation of these areas, we **standardize** each value of x by expressing it as a z -score, the number of standard deviations σ it lies from the mean μ .

$$z = \frac{x - \mu}{\sigma}$$

Somebody calculated all the integrals for the standard normal and put them in a table!

The Gaussian Distribution




The Gaussian Distribution

The formula for the standardized normal probability density function is

$$p(Z) = \frac{1}{(1)\sqrt{2\pi}} \cdot e^{-\frac{1}{2}\left(\frac{Z-0}{1}\right)^2} = \frac{1}{\sqrt{2\pi}} \cdot e^{-\frac{1}{2}(Z)^2}$$

The Gaussian Distribution

Comparing X and Z units



The Gaussian Distribution

- For example: What's the probability of getting a score of 575 or less, $\mu=500$ and $\sigma=50$?

$$Z = \frac{575 - 500}{50} = 1.5$$

- i.e., A score of 575 is 1.5 standard deviations above the mean

$$\therefore P(X \leq 575) = \int_{200}^{575} \frac{1}{(50)\sqrt{2\pi}} \cdot e^{-\frac{1}{2}\left(\frac{x-500}{50}\right)^2} dx \longrightarrow \int_{-\infty}^{1.5} \frac{1}{\sqrt{2\pi}} \cdot e^{-\frac{1}{2}Z^2} dz$$


look up $Z= 1.5$ in standard normal table = .9332

STANDARD STATISTICAL TABLES

1. Areas under the Normal Distribution

The table gives the cumulative probability up to the standardised normal value z i.e.

$$P[Z < z] = \int_{-\infty}^z \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{1}{2}z^2\right) dz$$



What is the area to the left of $Z=1.51$ in a standard normal curve?

$Z=1.51$

$Z=1.51$


z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5159	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7854
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8804	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9773	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9865	0.9868	0.9871	0.9874	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9924	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9980	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
z	3.00	3.10	3.20	3.30	3.40	3.50	3.60	3.70	3.80	3.90
P	0.9986	0.9990	0.9993	0.9995	0.9997	0.9998	0.9998	0.9999	0.9999	1.0000

Area is 93.45%

The Standard Gaussian (z) Distribution

- The four digit probability in a particular row and column of Table gives the area under the standard normal curve **between 0 and a positive value z** . This is enough because the standard normal curve is symmetric.

z	.00	.01	.02	.03	.04	.05	.06
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051
0.9	0.8159	0.8186	0.8212	0.8328	0.8264	0.8289	0.8315
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9278



Area for $z = 1.36$

Table 3

Second decimal place in z										z
0.09	0.08	0.07	0.06	0.05	0.04	0.03	0.02	0.01	0.00	
									* 0.0000	-3.9
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.8
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.7
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002	0.0002	-3.6
0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	-3.5
0.0002	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	-3.4
0.0003	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0005	0.0005	0.0005	-3.3
0.0005	0.0005	0.0005	0.0006	0.0006	0.0006	0.0006	0.0006	0.0007	0.0007	-3.2
0.0007	0.0007	0.0008	0.0008	0.0008	0.0008	0.0009	0.0009	0.0009	0.0010	-3.1
0.0010	0.0010	0.0011	0.0011	0.0011	0.0012	0.0012	0.0013	0.0013	0.0013	-3.0
0.0014	0.0014	0.0015	0.0015	0.0016	0.0016	0.0017	0.0018	0.0018	0.0019	-2.9
0.0019	0.0020	0.0021	0.0021	0.0022	0.0023	0.0023	0.0024	0.0025	0.0026	-2.8
0.0026	0.0027	0.0028	0.0029	0.0030	0.0031	0.0032	0.0033	0.0034	0.0035	-2.7
0.0036	0.0037	0.0038	0.0039	0.0040	0.0041	0.0043	0.0044	0.0045	0.0047	-2.6
0.0048	0.0049	0.0051	0.0052	0.0054	0.0055	0.0057	0.0059	0.0060	0.0062	-2.5
0.0064	0.0066	0.0068	0.0069	0.0071	0.0073	0.0075	0.0078	0.0080	0.0082	-2.4
0.0084	0.0087	0.0089	0.0091	0.0094	0.0096	0.0099	0.0102	0.0104	0.0107	-2.3
0.0110	0.0113	0.0116	0.0119	0.0122	0.0125	0.0129	0.0132	0.0136	0.0139	-2.2
0.0143	0.0146	0.0150	0.0154	0.0158	0.0162	0.0166	0.0170	0.0174	0.0179	-2.1
0.0183	0.0188	0.0192	0.0197	0.0202	0.0207	0.0212	0.0217	0.0222	0.0228	-2.0
0.0233	0.0239	0.0244	0.0250	0.0256	0.0262	0.0268	0.0274	0.0281	0.0287	-1.9
0.0294	0.0301	0.0307	0.0314	0.0322	0.0329	0.0336	0.0344	0.0351	0.0359	-1.8
0.0367	0.0375	0.0384	0.0392	0.0401	0.0409	0.0418	0.0427	0.0436	0.0446	-1.7
0.0455	0.0465	0.0475	0.0485	0.0495	0.0505	0.0516	0.0526	0.0537	0.0548	-1.6
0.0559	0.0571	0.0582	0.0594	0.0606	0.0618	0.0630	0.0643	0.0655	0.0668	-1.5
0.0681	0.0694	0.0708	0.0721	0.0735	0.0749	0.0764	0.0778	0.0793	0.0808	-1.4
0.0823	0.0838	0.0853	0.0869	0.0885	0.0901	0.0918	0.0934	0.0951	0.0968	-1.3
0.0985	0.1003	0.1020	0.1038	0.1056	0.1075	0.1093	0.1112	0.1131	0.1151	-1.2
0.1170	0.1190	0.1210	0.1230	0.1251	0.1271	0.1292	0.1314	0.1335	0.1357	-1.1
0.1379	0.1401	0.1423	0.1446	0.1469	0.1492	0.1515	0.1539	0.1562	0.1587	-1.0
0.1611	0.1635	0.1660	0.1685	0.1711	0.1736	0.1762	0.1788	0.1814	0.1841	-0.9
0.1867	0.1894	0.1922	0.1949	0.1977	0.2005	0.2033	0.2061	0.2090	0.2119	-0.8
0.2148	0.2177	0.2206	0.2236	0.2266	0.2296	0.2327	0.2358	0.2389	0.2420	-0.7
0.2451	0.2483	0.2514	0.2546	0.2578	0.2611	0.2643	0.2676	0.2709	0.2743	-0.6
0.2776	0.2810	0.2843	0.2877	0.2912	0.2946	0.2981	0.3015	0.3050	0.3085	-0.5
0.3121	0.3156	0.3192	0.3228	0.3264	0.3300	0.3336	0.3372	0.3409	0.3446	-0.4
0.3483	0.3520	0.3557	0.3594	0.3632	0.3669	0.3707	0.3745	0.3783	0.3821	-0.3
0.3859	0.3897	0.3936	0.3974	0.4013	0.4052	0.4090	0.4129	0.4168	0.4207	-0.2
0.4247	0.4286	0.4325	0.4364	0.4404	0.4443	0.4483	0.4522	0.4562	0.4602	-0.1
0.4641	0.4681	0.4721	0.4761	0.4801	0.4840	0.4880	0.4920	0.4960	0.5000	-0.0

* For values of $z \leq -3.90$, the areas are 0.0000 to four decimal places.


Example

Use Table 3 to calculate these probabilities:

$$P(z \leq 1.36) = .9131$$


$$P(z > 1.36) \\ = 1 - .9131 = .0869$$

$$P(-1.20 \leq z \leq 1.36) = \\ .9131 - .1151 = .7980$$



Example

- Find the value of z that has area .05 to its right.



- The area to its left will be $1 - .05 = .95$
- Look for the four digit area closest to .9500 Table 3.
- Since the value .9500 is halfway between .9495 and .9505, we choose z halfway between 1.64 and 1.65. $z=1.645$

		Second decimal place in z									
z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359	
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753	
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141	
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517	
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879	
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224	
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549	
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852	
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133	
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389	
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621	
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830	
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015	
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177	
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319	
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441	
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545	
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633	
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706	
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767	
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817	
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857	
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890	
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916	
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936	
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952	
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964	
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974	
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981	
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986	
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990	
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993	
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995	
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	
3.6	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.9	* 1.0000										

* For values of $z \geq 3.90$, the areas are 1.0000 to four decimal places

The Gaussian Distribution

Consider $P(X < 0.567)$?


From tables we know that $P(X < 0.56) = 0.7123$ and $P(X < 0.57) = 0.7157$

To calculate $P(X < 0.567)$ we *interpolate* between these two values

$$P(X < 0.567) = 0.3 \times 0.7123 + 0.7 \times 0.7157 = 0.71468$$

The Gaussian Distribution

$$P(Z > 0.92)$$




We know that $P(Z > 0.92) = 1 - P(Z < 0.92)$ and we can calculate $P(Z < 0.92)$ from the tables.

$$P(Z > 0.92) = 1 - 0.8212 = 0.1788$$

The Gaussian Distribution

$$P(Z > -0.5)$$




The Normal distribution is symmetric so we know that $P(Z > -0.5) = P(Z < 0.5) = 0.6915$


The Gaussian Distribution

$$P(-0.64 < Z < 0.43)$$


$$P(-0.64 < X < 0.43)$$



$$P(X < -0.64)$$



$$P(X < 0.43)$$



Example

Suppose we know that the birth weight of babies is Normally distributed with mean 3500g and standard deviation 500g. What is the probability that a baby is born that weights less than 3100g?

That is $X \sim N(3500, 500^2)$ and we want to calculate $P(X < 3100)$?




Table 3


Second decimal place in z										z
0.09	0.08	0.07	0.06	0.05	0.04	0.03	0.02	0.01	0.00	
								* 0.0000	0.0000	-3.9
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.8
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.7
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002	0.0002	-3.6
0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	-3.5
0.0002	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	-3.4
0.0003	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0005	0.0005	0.0005	-3.3
0.0005	0.0005	0.0005	0.0006	0.0006	0.0006	0.0006	0.0006	0.0007	0.0007	-3.2
0.0007	0.0007	0.0008	0.0008	0.0008	0.0008	0.0009	0.0009	0.0009	0.0010	-3.1
0.0010	0.0010	0.0011	0.0011	0.0011	0.0012	0.0012	0.0013	0.0013	0.0013	-3.0
0.0014	0.0014	0.0015	0.0015	0.0016	0.0016	0.0017	0.0018	0.0018	0.0019	-2.9
0.0019	0.0020	0.0021	0.0021	0.0022	0.0023	0.0023	0.0024	0.0025	0.0026	-2.8
0.0026	0.0027	0.0028	0.0029	0.0030	0.0031	0.0032	0.0033	0.0034	0.0035	-2.7
0.0036	0.0037	0.0038	0.0039	0.0040	0.0041	0.0043	0.0044	0.0045	0.0047	-2.6
0.0048	0.0049	0.0051	0.0052	0.0054	0.0055	0.0057	0.0059	0.0060	0.0062	-2.5
0.0064	0.0066	0.0068	0.0069	0.0071	0.0073	0.0075	0.0078	0.0080	0.0082	-2.4
0.0084	0.0087	0.0089	0.0091	0.0094	0.0096	0.0099	0.0102	0.0104	0.0107	-2.3
0.0110	0.0113	0.0116	0.0119	0.0122	0.0125	0.0129	0.0132	0.0136	0.0139	-2.2
0.0143	0.0146	0.0150	0.0154	0.0158	0.0162	0.0166	0.0170	0.0174	0.0179	-2.1
0.0183	0.0188	0.0192	0.0197	0.0202	0.0207	0.0212	0.0217	0.0222	0.0228	-2.0
0.0233	0.0239	0.0244	0.0250	0.0256	0.0262	0.0268	0.0274	0.0281	0.0287	-1.9
0.0294	0.0301	0.0307	0.0314	0.0322	0.0329	0.0336	0.0344	0.0351	0.0359	-1.8
0.0367	0.0375	0.0384	0.0392	0.0401	0.0409	0.0418	0.0427	0.0436	0.0446	-1.7
0.0455	0.0465	0.0475	0.0485	0.0495	0.0505	0.0516	0.0526	0.0537	0.0548	-1.6
0.0559	0.0571	0.0582	0.0594	0.0606	0.0618	0.0630	0.0643	0.0655	0.0668	-1.5
0.0681	0.0694	0.0708	0.0721	0.0735	0.0749	0.0764	0.0778	0.0793	0.0808	-1.4
0.0823	0.0838	0.0853	0.0869	0.0885	0.0901	0.0918	0.0934	0.0951	0.0968	-1.3
0.0985	0.1003	0.1020	0.1038	0.1056	0.1075	0.1093	0.1112	0.1131	0.1151	-1.2
0.1170	0.1190	0.1210	0.1230	0.1251	0.1271	0.1292	0.1314	0.1335	0.1357	-1.1
0.1379	0.1401	0.1423	0.1446	0.1469	0.1492	0.1515	0.1539	0.1562	0.1587	-1.0
0.1611	0.1635	0.1660	0.1685	0.1711	0.1736	0.1762	0.1788	0.1814	0.1841	-0.9
0.1867	0.1894	0.1922	0.1949	0.1977	0.2005	0.2033	0.2061	0.2090	0.2119	-0.8
0.2148	0.2177	0.2206	0.2236	0.2266	0.2296	0.2327	0.2358	0.2389	0.2420	-0.7
0.2451	0.2483	0.2514	0.2546	0.2578	0.2611	0.2643	0.2676	0.2709	0.2743	-0.6
0.2776	0.2810	0.2843	0.2877	0.2912	0.2946	0.2981	0.3015	0.3050	0.3085	-0.5
0.3121	0.3156	0.3192	0.3228	0.3264	0.3300	0.3336	0.3372	0.3409	0.3446	-0.4
0.3483	0.3520	0.3557	0.3594	0.3632	0.3669	0.3707	0.3745	0.3783	0.3821	-0.3
0.3859	0.3897	0.3936	0.3974	0.4013	0.4052	0.4090	0.4129	0.4168	0.4207	-0.2
0.4247	0.4286	0.4325	0.4364	0.4404	0.4443	0.4483	0.4522	0.4562	0.4602	-0.1
0.4641	0.4681	0.4721	0.4761	0.4801	0.4840	0.4880	0.4920	0.4960	0.5000	-0.0

* For values of $z \leq -3.90$, the areas are 0.0000 to four decimal places

Example

- The weights of packages of ground beef are normally distributed with mean 1 kg and standard deviation .10. What is the probability that a randomly selected package weighs between 0.80 and 0.85 kg?

$$\begin{aligned}P(.80 < x < .85) &= P\left(\frac{.80 - 1}{.1} < z < \frac{.85 - 1}{.1}\right) \\&= P(-2 < z < -1.5) = P(1.5 < z < 2) \\&= .9772 - .9332 = .0440\end{aligned}$$



$$P(-2 < z < -1.5) = 0.0668 - 0.0228 = 0.0440$$

Table 3

Second decimal place in z										z
0.09	0.08	0.07	0.06	0.05	0.04	0.03	0.02	0.01	0.00	
									* 0.0000	-3.9
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.8
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.7
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002	0.0002	-3.6
0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	-3.5
0.0002	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	-3.4
0.0003	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0005	0.0005	0.0005	-3.3
0.0005	0.0005	0.0005	0.0006	0.0006	0.0006	0.0006	0.0006	0.0007	0.0007	-3.2
0.0007	0.0007	0.0008	0.0008	0.0008	0.0008	0.0009	0.0009	0.0009	0.0010	-3.1
0.0010	0.0010	0.0011	0.0011	0.0011	0.0012	0.0012	0.0013	0.0013	0.0013	-3.0
0.0014	0.0014	0.0015	0.0015	0.0016	0.0016	0.0017	0.0018	0.0018	0.0019	-2.9
0.0019	0.0020	0.0021	0.0021	0.0022	0.0023	0.0023	0.0024	0.0025	0.0026	-2.8
0.0026	0.0027	0.0028	0.0029	0.0030	0.0031	0.0032	0.0033	0.0034	0.0035	-2.7
0.0036	0.0037	0.0038	0.0039	0.0040	0.0041	0.0043	0.0044	0.0045	0.0047	-2.6
0.0048	0.0049	0.0051	0.0052	0.0054	0.0055	0.0057	0.0059	0.0060	0.0062	-2.5
0.0064	0.0066	0.0068	0.0069	0.0071	0.0073	0.0075	0.0078	0.0080	0.0082	-2.4
0.0084	0.0087	0.0089	0.0091	0.0094	0.0096	0.0099	0.0102	0.0104	0.0107	-2.3
0.0110	0.0113	0.0116	0.0119	0.0122	0.0125	0.0129	0.0132	0.0136	0.0139	-2.2
0.0143	0.0146	0.0150	0.0154	0.0158	0.0162	0.0166	0.0170	0.0174	0.0179	-2.1
0.0183	0.0188	0.0192	0.0197	0.0202	0.0207	0.0212	0.0217	0.0222	0.0228	-2.0
0.0233	0.0239	0.0244	0.0250	0.0256	0.0262	0.0268	0.0274	0.0281	0.0287	-1.9
0.0294	0.0301	0.0307	0.0314	0.0322	0.0329	0.0336	0.0344	0.0351	0.0359	-1.8
0.0367	0.0375	0.0384	0.0392	0.0401	0.0409	0.0418	0.0427	0.0436	0.0446	-1.7
0.0455	0.0465	0.0475	0.0485	0.0495	0.0505	0.0516	0.0526	0.0537	0.0548	-1.6
0.0559	0.0571	0.0582	0.0594	0.0606	0.0618	0.0630	0.0643	0.0655	0.0668	-1.5
0.0681	0.0694	0.0708	0.0721	0.0735	0.0749	0.0764	0.0778	0.0793	0.0808	-1.4
0.0823	0.0838	0.0853	0.0869	0.0885	0.0901	0.0918	0.0934	0.0951	0.0968	-1.3
0.0985	0.1003	0.1020	0.1038	0.1056	0.1075	0.1093	0.1112	0.1131	0.1151	-1.2
0.1170	0.1190	0.1210	0.1230	0.1251	0.1271	0.1292	0.1314	0.1335	0.1357	-1.1
0.1379	0.1401	0.1423	0.1446	0.1469	0.1492	0.1515	0.1539	0.1562	0.1587	-1.0
0.1611	0.1635	0.1660	0.1685	0.1711	0.1736	0.1762	0.1788	0.1814	0.1841	-0.9
0.1867	0.1894	0.1922	0.1949	0.1977	0.2005	0.2033	0.2061	0.2090	0.2119	-0.8
0.2148	0.2177	0.2206	0.2236	0.2266	0.2296	0.2327	0.2358	0.2389	0.2420	-0.7
0.2451	0.2483	0.2514	0.2546	0.2578	0.2611	0.2643	0.2676	0.2709	0.2743	-0.6
0.2776	0.2810	0.2843	0.2877	0.2912	0.2946	0.2981	0.3015	0.3050	0.3085	-0.5
0.3121	0.3156	0.3192	0.3228	0.3264	0.3300	0.3336	0.3372	0.3409	0.3446	-0.4
0.3483	0.3520	0.3557	0.3594	0.3632	0.3669	0.3707	0.3745	0.3783	0.3821	-0.3
0.3859	0.3897	0.3936	0.3974	0.4013	0.4052	0.4090	0.4129	0.4168	0.4207	-0.2
0.4247	0.4286	0.4325	0.4364	0.4404	0.4443	0.4483	0.4522	0.4562	0.4602	-0.1
0.4641	0.4681	0.4721	0.4761	0.4801	0.4840	0.4880	0.4920	0.4960	0.5000	-0.0

* For values of $z \leq -3.90$, the areas are 0.0000 to four decimal places

Example

- What is the weight of a package such that only 5% of all packages exceed this weight?

$$P(x > ?) = .05$$

$$P(z > \frac{?-1}{.1}) = .05$$

$$P(z < \frac{?-1}{.1}) = 1 - .05 = .95$$

From Table 3, $\frac{?-1}{.1} = 1.645$

$$? = 1.645(.1) + 1 = 1.16$$

		Second decimal place in z									
z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359	
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753	
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141	
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517	
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879	
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224	
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549	
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852	
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133	
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389	
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621	
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830	
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015	
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177	
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319	
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441	
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545	
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633	
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706	
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767	
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817	
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857	
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890	
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916	
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936	
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952	
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964	
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974	
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981	
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986	
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990	
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993	
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995	
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	
3.6	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.9	* 1.0000										


* For values of z ≥ 3.90, the areas are 1.0000 to four decimal places

Example



A Company produces “20 gr” chocolate.


Suppose the companies “20 gr” chocolate follow a normally distribution with a mean $\mu=20.2$ gr with a standard deviation $\sigma=0.125$ gr .



Example



- What proportion of the chocolate less than 20 gr?



$P(z < -1.60) = 1 - P(z > 1.60) = 1 - .9452 = .0548$. The proportion of the chocolate less than 20 gr is .0548


Table 3

Second decimal place in z										z
0.09	0.08	0.07	0.06	0.05	0.04	0.03	0.02	0.01	0.00	
									* 0.0000	-3.9
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.8
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.7
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002	0.0002	-3.6
0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	-3.5
0.0002	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	-3.4
0.0003	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0005	0.0005	0.0005	-3.3
0.0005	0.0005	0.0005	0.0006	0.0006	0.0006	0.0006	0.0006	0.0007	0.0007	-3.2
0.0007	0.0007	0.0008	0.0008	0.0008	0.0008	0.0009	0.0009	0.0009	0.0010	-3.1
0.0010	0.0010	0.0011	0.0011	0.0011	0.0012	0.0012	0.0013	0.0013	0.0013	-3.0
0.0014	0.0014	0.0015	0.0015	0.0016	0.0016	0.0017	0.0018	0.0018	0.0019	-2.9
0.0019	0.0020	0.0021	0.0021	0.0022	0.0023	0.0023	0.0024	0.0025	0.0026	-2.8
0.0026	0.0027	0.0028	0.0029	0.0030	0.0031	0.0032	0.0033	0.0034	0.0035	-2.7
0.0036	0.0037	0.0038	0.0039	0.0040	0.0041	0.0043	0.0044	0.0045	0.0047	-2.6
0.0048	0.0049	0.0051	0.0052	0.0054	0.0055	0.0057	0.0059	0.0060	0.0062	-2.5
0.0064	0.0066	0.0068	0.0069	0.0071	0.0073	0.0075	0.0078	0.0080	0.0082	-2.4
0.0084	0.0087	0.0089	0.0091	0.0094	0.0096	0.0099	0.0102	0.0104	0.0107	-2.3
0.0110	0.0113	0.0116	0.0119	0.0122	0.0125	0.0129	0.0132	0.0136	0.0139	-2.2
0.0143	0.0146	0.0150	0.0154	0.0158	0.0162	0.0166	0.0170	0.0174	0.0179	-2.1
0.0183	0.0188	0.0192	0.0197	0.0202	0.0207	0.0212	0.0217	0.0222	0.0228	-2.0
0.0233	0.0239	0.0244	0.0250	0.0256	0.0262	0.0268	0.0274	0.0281	0.0287	-1.9
0.0294	0.0301	0.0307	0.0314	0.0322	0.0329	0.0336	0.0344	0.0351	0.0359	-1.8
0.0367	0.0375	0.0384	0.0392	0.0401	0.0409	0.0418	0.0427	0.0436	0.0446	-1.7
0.0455	0.0465	0.0475	0.0485	0.0495	0.0505	0.0516	0.0526	0.0537	0.0548	-1.6
0.0559	0.0571	0.0582	0.0594	0.0606	0.0618	0.0630	0.0643	0.0655	0.0668	-1.5
0.0681	0.0694	0.0708	0.0721	0.0735	0.0749	0.0764	0.0778	0.0793	0.0808	-1.4
0.0823	0.0838	0.0853	0.0869	0.0885	0.0901	0.0918	0.0934	0.0951	0.0968	-1.3
0.0985	0.1003	0.1020	0.1038	0.1056	0.1075	0.1093	0.1112	0.1131	0.1151	-1.2
0.1170	0.1190	0.1210	0.1230	0.1251	0.1271	0.1292	0.1314	0.1335	0.1357	-1.1
0.1379	0.1401	0.1423	0.1446	0.1469	0.1492	0.1515	0.1539	0.1562	0.1587	-1.0
0.1611	0.1635	0.1660	0.1685	0.1711	0.1736	0.1762	0.1788	0.1814	0.1841	-0.9
0.1867	0.1894	0.1922	0.1949	0.1977	0.2005	0.2033	0.2061	0.2090	0.2119	-0.8
0.2148	0.2177	0.2206	0.2236	0.2266	0.2296	0.2327	0.2358	0.2389	0.2420	-0.7
0.2451	0.2483	0.2514	0.2546	0.2578	0.2611	0.2643	0.2676	0.2709	0.2743	-0.6
0.2776	0.2810	0.2843	0.2877	0.2912	0.2946	0.2981	0.3015	0.3050	0.3085	-0.5
0.3121	0.3156	0.3192	0.3228	0.3264	0.3300	0.3336	0.3372	0.3409	0.3446	-0.4
0.3483	0.3520	0.3557	0.3594	0.3632	0.3669	0.3707	0.3745	0.3783	0.3821	-0.3
0.3859	0.3897	0.3936	0.3974	0.4013	0.4052	0.4090	0.4129	0.4168	0.4207	-0.2
0.4247	0.4286	0.4325	0.4364	0.4404	0.4443	0.4483	0.4522	0.4562	0.4602	-0.1
0.4641	0.4681	0.4721	0.4761	0.4801	0.4840	0.4880	0.4920	0.4960	0.5000	-0.0

* For values of $z \leq -3.90$, the areas are 0.0000 to four decimal places

Example


- What proportion of the chocolate between 20 and 20.3 gr?



$$P(-1.60 < z < .80) = P(z < .80) - P(z < -1.60) = .7871 - .0548 = .7333$$

Example

99% of chocolate will contain more than what amount chocolate?



$$.99 = P(x > ?) = P(z > \frac{? - 20.2}{.125})$$

From Table 3, $\frac{20.2 - ?}{.125} = 2.33$




$$? = 20.2 - 2.33(.125) = 19.91$$

		Second decimal place in z									
z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09	
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359	
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753	
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141	
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517	
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879	
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224	
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549	
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852	
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133	
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389	
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621	
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830	
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015	
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177	
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319	
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441	
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545	
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633	
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706	
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767	
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817	
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857	
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890	
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916	
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936	
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952	
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964	
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974	
2.8	0.9974	0.9975	0.9976	0.9977	0.9978	0.9979	0.9979	0.9979	0.9980	0.9981	
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986	
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990	
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993	
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995	
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997	
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998	
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	
3.6	0.9998	0.9998	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.7	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.8	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	0.9999	
3.9	* 1.0000										

* For values of z ≥ 3.90, the areas are 1.0000 to four decimal places

How Probabilities Are Distributed

- The interval $\mu \pm \sigma$ contains approximately 68% of the measurements.
- The interval $\mu \pm 2\sigma$ contains approximately 95% of the measurements.
- The interval $\mu \pm 3\sigma$ contains approximately 99.7% of the measurements.



Linear combinations of Normal random variables

Suppose two rats A and B have been trained to navigate a large maze. The time it takes rat A is normally distributed with mean 80 seconds and standard deviation 10 seconds. The time it takes rat B is normally distributed with mean 78 seconds and standard deviation 13 seconds. On any given day what is the probability that rat A runs the maze faster than rat B?

X = Time of run for rat A $X \sim N(80, 10^2)$

Y = Time of run for rat B $Y \sim N(78, 13^2)$

Let $D = X - Y$ be the difference in times of rats A and B

If rat A is faster than rat B then $D < 0$ so we want $P(D < 0)$?

Linear combinations of Normal random variables


If X and Y are two independent normal variable such that

$$X \sim N(\mu_1, \sigma_1^2) \text{ and } Y \sim N(\mu_2, \sigma_2^2)$$

$$\text{then } X - Y \sim N(\mu_1 - \mu_2, \sigma_1^2 + \sigma_2^2)$$

In this example

$$D = X - Y \sim N(80 - 78, 10^2 + 13^2) = N(2, 269)$$



Linear combinations of Normal random variables

$$P(D < 0) = P\left(\frac{D - 2}{\sqrt{269}} < \frac{0 - 2}{\sqrt{269}}\right) = P(Z < -0.122) \quad \text{where } Z \sim N(0, 1)$$

$$\begin{aligned} &= 0.4522 \cdot 0.8 + 0.4483 \cdot 0.2 \\ &= 0.45142 \end{aligned}$$

Table 3

Second decimal place in z										z
0.09	0.08	0.07	0.06	0.05	0.04	0.03	0.02	0.01	0.00	
									* 0.0000	-3.9
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.8
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.7
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002	0.0002	-3.6
0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	-3.5
0.0002	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	-3.4
0.0003	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0005	0.0005	0.0005	-3.3
0.0005	0.0005	0.0005	0.0006	0.0006	0.0006	0.0006	0.0006	0.0007	0.0007	-3.2
0.0007	0.0007	0.0008	0.0008	0.0008	0.0008	0.0009	0.0009	0.0009	0.0010	-3.1
0.0010	0.0010	0.0011	0.0011	0.0011	0.0012	0.0012	0.0013	0.0013	0.0013	-3.0
0.0014	0.0014	0.0015	0.0015	0.0016	0.0016	0.0017	0.0018	0.0018	0.0019	-2.9
0.0019	0.0020	0.0021	0.0021	0.0022	0.0023	0.0023	0.0024	0.0025	0.0026	-2.8
0.0026	0.0027	0.0028	0.0029	0.0030	0.0031	0.0032	0.0033	0.0034	0.0035	-2.7
0.0036	0.0037	0.0038	0.0039	0.0040	0.0041	0.0043	0.0044	0.0045	0.0047	-2.6
0.0048	0.0049	0.0051	0.0052	0.0054	0.0055	0.0057	0.0059	0.0060	0.0062	-2.5
0.0064	0.0066	0.0068	0.0069	0.0071	0.0073	0.0075	0.0078	0.0080	0.0082	-2.4
0.0084	0.0087	0.0089	0.0091	0.0094	0.0096	0.0099	0.0102	0.0104	0.0107	-2.3
0.0110	0.0113	0.0116	0.0119	0.0122	0.0125	0.0129	0.0132	0.0136	0.0139	-2.2
0.0143	0.0146	0.0150	0.0154	0.0158	0.0162	0.0166	0.0170	0.0174	0.0179	-2.1
0.0183	0.0188	0.0192	0.0197	0.0202	0.0207	0.0212	0.0217	0.0222	0.0228	-2.0
0.0233	0.0239	0.0244	0.0250	0.0256	0.0262	0.0268	0.0274	0.0281	0.0287	-1.9
0.0294	0.0301	0.0307	0.0314	0.0322	0.0329	0.0336	0.0344	0.0351	0.0359	-1.8
0.0367	0.0375	0.0384	0.0392	0.0401	0.0409	0.0418	0.0427	0.0436	0.0446	-1.7
0.0455	0.0465	0.0475	0.0485	0.0495	0.0505	0.0516	0.0526	0.0537	0.0548	-1.6
0.0559	0.0571	0.0582	0.0594	0.0606	0.0618	0.0630	0.0643	0.0655	0.0668	-1.5
0.0681	0.0694	0.0708	0.0721	0.0735	0.0749	0.0764	0.0778	0.0793	0.0808	-1.4
0.0823	0.0838	0.0853	0.0869	0.0885	0.0901	0.0918	0.0934	0.0951	0.0968	-1.3
0.0985	0.1003	0.1020	0.1038	0.1056	0.1075	0.1093	0.1112	0.1131	0.1151	-1.2
0.1170	0.1190	0.1210	0.1230	0.1251	0.1271	0.1292	0.1314	0.1335	0.1357	-1.1
0.1379	0.1401	0.1423	0.1446	0.1469	0.1492	0.1515	0.1539	0.1562	0.1587	-1.0
0.1611	0.1635	0.1660	0.1685	0.1711	0.1736	0.1762	0.1788	0.1814	0.1841	-0.9
0.1867	0.1894	0.1922	0.1949	0.1977	0.2005	0.2033	0.2061	0.2090	0.2119	-0.8
0.2148	0.2177	0.2206	0.2236	0.2266	0.2296	0.2327	0.2358	0.2389	0.2420	-0.7
0.2451	0.2483	0.2514	0.2546	0.2578	0.2611	0.2643	0.2676	0.2709	0.2743	-0.6
0.2776	0.2810	0.2843	0.2877	0.2912	0.2946	0.2981	0.3015	0.3050	0.3085	-0.5
0.3121	0.3156	0.3192	0.3228	0.3264	0.3300	0.3336	0.3372	0.3409	0.3446	-0.4
0.3483	0.3520	0.3557	0.3594	0.3632	0.3669	0.3707	0.3745	0.3783	0.3821	-0.3
0.3859	0.3897	0.3936	0.3974	0.4013	0.4052	0.4090	0.4129	0.4168	0.4207	-0.2
0.4247	0.4286	0.4325	0.4364	0.4404	0.4443	0.4483	0.4522	0.4562	0.4602	-0.1
0.4641	0.4681	0.4721	0.4761	0.4801	0.4840	0.4880	0.4920	0.4960	0.5000	-0.0

* For values of $z \leq -3.90$, the areas are 0.0000 to four decimal places

Linear combinations of Normal random variables

- Suppose two rats A and B have been trained to navigate a large maze. The time it takes rat A is normally distributed with mean 80 seconds and standard deviation 10 seconds. The time it takes rat B is normally distributed with mean 78 seconds and standard deviation 13 seconds. On any given day what is the probability that the average time the rats take to run the maze is greater than 82 seconds?

If X and Y are two independent normal variable such that

$$X \sim N(\mu_1, \sigma_1^2) \text{ and } Y \sim N(\mu_2, \sigma_2^2)$$

then

$$X + Y \sim N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$$

$$aX \sim N(a\mu_1, a^2\sigma_1^2)$$

$$aX + bY \sim N(a\mu_1 + b\mu_2, a^2\sigma_1^2 + b^2\sigma_2^2)$$

X = Time of run for rat A $X \sim N(80, 10^2)$


Y = Time of run for rat B $Y \sim N(78, 13^2)$

Let $A = (X+Y)/2 = 1/2X + 1/2Y$ be the average time of rats A and B

Then $A \sim N(\frac{1}{2} 80 + \frac{1}{2} 78, (1/2)^2 10^2 + (1/2)^2 13^2) = N(79, 67.25)$

We want $P(A > 82)$

Linear combinations of Normal random variables




$$\begin{aligned} P(A > 82) &= P\left(\frac{A - 79}{\sqrt{67.25}} < \frac{82 - 79}{\sqrt{67.25}}\right) = P(Z > 0.366) \quad \text{where } Z \sim N(0, 1) \\ &= 1 - (0.4 \times 0.6406 + 0.6 \times 0.6443) \\ &= 0.35718 \end{aligned}$$

Central Limit Theorem

- It gives conditions under which the mean of a sufficiently large number of independent random variables, each with finite mean and variance will be approximately normally distributed
- It provides a partial explanation for prevalence of the Gaussian distribution in the real world.
- Justifies the approximation of the large – sample statistics to the Gaussian distribution in controlled experiments.

Central Limit Theorem



Relationship between Gaussian and Binomial & Poisson distribution

Relationship between Gaussian and Binomial & Poisson distribution

The Gaussian distribution can be derived from the binomial (or Poisson) assuming:

p is finite & N is very large

we have a continuous variable rather than a discrete variable

Consider tossing a coin 10,000 times.

$$p(\text{head}) = 0.5 \text{ and } N = 10,000$$

- For a binomial distribution:

$$\text{mean number of heads} = \mu = Np = 5000$$

$$\text{standard deviation } \sigma = [Np(1 - p)]^{1/2} = 50$$

The probability to be within $\mu \pm 1\sigma$ for this binomial distribution is:

$$P = \sum_{m=5000-50}^{5000+50} \frac{10^4!}{(10^4 - m)!m!} 0.5^m 0.5^{10^4-m} = 0.69$$


- For a Gaussian distribution:

$$P(\mu - \sigma < y < \mu + \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \int_{\mu-\sigma}^{\mu+\sigma} e^{-\frac{(y-\mu)^2}{2\sigma^2}} dy \approx 0.68$$

Both distributions give about the ~ same probability!

The Normal Approximation to the Binomial

- We can calculate binomial probabilities using
 - The binomial formula
 - The cumulative binomial tables
- When n is large, and p is not too close to zero or one, areas under the normal curve with mean np and variance npq can be used to approximate binomial probabilities.



Approximating the Binomial

- ✓ Make sure to include the entire rectangle for the values of x in the interval of interest. This is called the **continuity correction**.
- ✓ Standardize the values of x using

$$z = \frac{x - \mu}{\sigma}, \mu = np, \sigma = \sqrt{npq}$$

- ✓ Make sure that **np and nq are both greater than 5** to avoid inaccurate approximations! Or
- ✓ n is large and $\mu \pm 2\sigma$ falls between 0 and n


Example

Suppose x is a binomial random variable with $n = 30$ and $p = .4$. Using the normal approximation to find $P(x \leq 10)$.

$$n = 30 \quad p = .4 \quad q = .6$$

$$np = 12 \quad nq = 18$$

The normal
approximation is ok!





Calculate

$$\mu = np = 30(.4) = 12$$

$$\sigma = \sqrt{npq} = \sqrt{30(.4)(.6)} = 2.683$$

Example



$$\begin{aligned} P(x \leq 10) &\approx P\left(z \leq \frac{10.5 - 12}{2.683}\right) \\ &= P(z \leq -0.56) = .2877 \end{aligned}$$

Table 3

Second decimal place in z										z
0.09	0.08	0.07	0.06	0.05	0.04	0.03	0.02	0.01	0.00	
								* 0.0000		-3.9
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.8
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	-3.7
0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001	0.0002	0.0002	-3.6
0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	0.0002	-3.5
0.0002	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	0.0003	-3.4
0.0003	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0005	0.0005	0.0005	-3.3
0.0005	0.0005	0.0005	0.0006	0.0006	0.0006	0.0006	0.0006	0.0007	0.0007	-3.2
0.0007	0.0007	0.0008	0.0008	0.0008	0.0008	0.0009	0.0009	0.0009	0.0010	-3.1
0.0010	0.0010	0.0011	0.0011	0.0011	0.0012	0.0012	0.0013	0.0013	0.0013	-3.0
0.0014	0.0014	0.0015	0.0015	0.0016	0.0016	0.0017	0.0018	0.0018	0.0019	-2.9
0.0019	0.0020	0.0021	0.0021	0.0022	0.0023	0.0023	0.0024	0.0025	0.0026	-2.8
0.0026	0.0027	0.0028	0.0029	0.0030	0.0031	0.0032	0.0033	0.0034	0.0035	-2.7
0.0036	0.0037	0.0038	0.0039	0.0040	0.0041	0.0043	0.0044	0.0045	0.0047	-2.6
0.0048	0.0049	0.0051	0.0052	0.0054	0.0055	0.0057	0.0059	0.0060	0.0062	-2.5
0.0064	0.0066	0.0068	0.0069	0.0071	0.0073	0.0075	0.0078	0.0080	0.0082	-2.4
0.0084	0.0087	0.0089	0.0091	0.0094	0.0096	0.0099	0.0102	0.0104	0.0107	-2.3
0.0110	0.0113	0.0116	0.0119	0.0122	0.0125	0.0129	0.0132	0.0136	0.0139	-2.2
0.0143	0.0146	0.0150	0.0154	0.0158	0.0162	0.0166	0.0170	0.0174	0.0179	-2.1
0.0183	0.0188	0.0192	0.0197	0.0202	0.0207	0.0212	0.0217	0.0222	0.0228	-2.0
0.0233	0.0239	0.0244	0.0250	0.0256	0.0262	0.0268	0.0274	0.0281	0.0287	-1.9
0.0294	0.0301	0.0307	0.0314	0.0322	0.0329	0.0336	0.0344	0.0351	0.0359	-1.8
0.0367	0.0375	0.0384	0.0392	0.0401	0.0409	0.0418	0.0427	0.0436	0.0446	-1.7
0.0455	0.0465	0.0475	0.0485	0.0495	0.0505	0.0516	0.0526	0.0537	0.0548	-1.6
0.0559	0.0571	0.0582	0.0594	0.0606	0.0618	0.0630	0.0643	0.0655	0.0668	-1.5
0.0681	0.0694	0.0708	0.0721	0.0735	0.0749	0.0764	0.0778	0.0793	0.0808	-1.4
0.0823	0.0838	0.0853	0.0869	0.0885	0.0901	0.0918	0.0934	0.0951	0.0968	-1.3
0.0985	0.1003	0.1020	0.1038	0.1056	0.1075	0.1093	0.1112	0.1131	0.1151	-1.2
0.1170	0.1190	0.1210	0.1230	0.1251	0.1271	0.1292	0.1314	0.1335	0.1357	-1.1
0.1379	0.1401	0.1423	0.1446	0.1469	0.1492	0.1515	0.1539	0.1562	0.1587	-1.0
0.1611	0.1635	0.1660	0.1685	0.1711	0.1736	0.1762	0.1788	0.1814	0.1841	-0.9
0.1867	0.1894	0.1922	0.1949	0.1977	0.2005	0.2033	0.2061	0.2090	0.2119	-0.8
0.2148	0.2177	0.2206	0.2236	0.2266	0.2296	0.2327	0.2358	0.2389	0.2420	-0.7
0.2451	0.2483	0.2514	0.2546	0.2578	0.2611	0.2643	0.2676	0.2709	0.2743	-0.6
0.2776	0.2810	0.2843	0.2877	0.2912	0.2946	0.2981	0.3015	0.3050	0.3085	-0.5
0.3121	0.3156	0.3192	0.3228	0.3264	0.3300	0.3336	0.3372	0.3409	0.3446	-0.4
0.3483	0.3520	0.3557	0.3594	0.3632	0.3669	0.3707	0.3745	0.3783	0.3821	-0.3
0.3859	0.3897	0.3936	0.3974	0.4013	0.4052	0.4090	0.4129	0.4168	0.4207	-0.2
0.4247	0.4286	0.4325	0.4364	0.4404	0.4443	0.4483	0.4522	0.4562	0.4602	-0.1
0.4641	0.4681	0.4721	0.4761	0.4801	0.4840	0.4880	0.4920	0.4960	0.5000	-0.0

* For values of $z \leq -3.90$, the areas are 0.0000 to four decimal places

Example

A production line produces AA batteries with a reliability rate of 95%. A sample of $n = 200$ batteries is selected. Find the probability that at least 195 of the batteries work.

Success = working battery $n = 200$

$$p = .95 \quad np = 190 \quad nq = 10$$

The normal approximation is ok!

$$\begin{aligned} P(x \geq 195) &\approx P\left(z \geq \frac{194.5 - 190}{\sqrt{200(.95)(.05)}}\right) \\ &= P(z \geq 1.46) = 1 - .9278 = .0722 \end{aligned}$$

Relationship between Gaussian and Binomial & Poisson distribution

Compare $\pm 1\sigma$ area of Poisson and Gaussian:

Mean	Poission	Gaussian	% diff
10	0.74	0.6827	7.8
25	0.73	0.6827	6.9
100	0.707	0.6827	3.5
250	0.689	0.6827	0.87
5000	0.6847	0.6827	0.29

$$\text{Poisson : } P(\pm 1\sigma) = \sum_{m=\mu-\sqrt{\mu}}^{\mu+\sqrt{\mu}} \frac{e^{-\mu} \mu^m}{m!}$$

Central Limit Theorem

1. The distribution of sample \bar{x} will, as the sample size increases, approach a **normal** distribution.
2. The mean of the sample means is the population mean μ .
3. The standard deviation of all sample means is σ/\sqrt{n}

Central Limit Theorem


1. For samples of size n larger than 30, the distribution of the sample means can be approximated reasonably well by a normal distribution. The approximation gets better as the sample size n becomes larger.
2. If the original population is itself normally distributed, then the sample means will be normally distributed for **any** sample size n (not just the values of n larger than 30).

Binomial Poisson and Normal ?

Poisson(30)

Binomial(1000,.03)

N(30,sqrt(30))



The χ^2 distribution

Problem: We have some measurements and would like some way to measure how “good” these measurements really are.

Solution: Consider calculating the “ χ^2 ” (“chi-square”)

Assume:

We have a set of measurements $\{x_1, x_2, \dots x_n\}$.

We know the true value of each x_i ($x_{t1}, x_{t2}, \dots x_{tn}$).

Obviously the closer the $(x_1, x_2, \dots x_n)$'s are to the $(x_{t1}, x_{t2}, \dots x_{tn})$'s the better (or more accurate) the measurements.

Can we put a number (or probability) on how well they agree?

Uses of the χ^2 distribution

- Let us assume that there is given a set of n independent random variables x_1, x_2, \dots, x_n which are all normal $N(\mu, \sigma^2)$. We may for instance think of the x_i 's as the outcome of n repeated measurements on the same physical system or n independent observations on the same quantity. Then the x_i 's constitute a sample of size n from population which is normal with mean μ and variance σ^2

We define the chi-square **sum** χ^2

$$\chi^2 \equiv \frac{(x_1 - \mu_1)^2}{\sigma_1^2} + \frac{(x_2 - \mu_2)^2}{\sigma_2^2} + \dots + \frac{(x_\nu - \mu_\nu)^2}{\sigma_\nu^2} = \sum_{i=1}^{\nu} \frac{(x_i - \mu_i)^2}{\sigma_i^2}$$


- Note that ideally, given the random fluctuations of the values of x_i about their mean values μ_i , each term in the sum will be of order unity. Hence, if we have chosen the μ_i , and the σ_i correctly, we may expect that a calculated value of χ^2 will be approximately equal to n . If it is, then we may conclude that the data are well described by the values we have chosen for the μ_i that is, by the hypothesized function
- If a calculated value of χ^2 turns out to be much larger than n and we have correctly estimated the values for the σ_i we may possibly conclude that our data are not well-described by our hypothesized set of the μ_i

The χ^2 distribution

This is a continuous probability distribution that is a function of two variables: χ^2 and $n = \text{number of degrees of freedom (DOF)}$.

$DOF=n = (\text{\# of data points}) - (\text{\# of parameters calculated from the data points})$

$$f(\chi^2) = \frac{1}{2^{\nu/2} \Gamma(\nu/2)} e^{-\chi^2/2} (\chi^2)^{(\nu/2)-1}$$



For $n \geq 20$, $P(\chi^2 > y)$ can be approximated using a Gaussian *pdf* with $y = (2\chi^2)^{1/2} - (2n-1)^{1/2}$

The χ^2 distribution

Eur J Phys 11 (1990) 338–342 Printed in the UK

338

χ^2 —or do the data fit the theory?

J C Earnshaw

The Department of Pure and Applied Physics, The Queen's University of Belfast, Belfast BT7 1NN, UK

Received 28 November 1989, in final form 27 June 1990

Abstract. The problems found by physics students with the meaning and significance of the χ^2 test are discussed. It is suggested that investigation of the fluctuations observed in repeated experiments under constant conditions can help students to appreciate the principles involved.

Résumé. Les problèmes trouvés par les étudiants de la physique avec la signification de l'essai χ^2 sont découlés. On propose qu'un examen des fluctuations observées dans les expériences répétées sous les conditions constantes peut aider étudiants se rendre compte des principes entraînés.

Eur J Phys 11 (1990), 338-342

Example

We collected N events in an experiment and histogram the data in m bins like we count cosmic ray events in 15 second intervals and sort the data into 5 bins:

Number of counts in 15 second intervals	0	1	2	3	4
Number of intervals with above counts	2	7	6	3	2

Suppose we want to compare our data with the expectations of a Poisson distribution (P):

$$P(\eta) = A \frac{e^{-\mu} \mu^{\eta}}{\eta!}$$

P(0)=# intervals with 0 counts

P(1)=# intervals with 1 counts

In order to do this comparison we need to know A, the total number of intervals (=20 for this example).

$$A=2+7+6+3+2=20$$

Example

Since A is calculated from the data, we give up one degree of freedom:

$$dof=m-1=5-1=4.$$

If we also calculate the mean (μ) of our Poisson from the data we lose another dof:


$$dof=m-1=5-2=3.$$

$$\mu = \frac{0 \times 2 + 1 \times 7 + 2 \times 6 + 3 \times 3 + 4 \times 2}{2 + 7 + 6 + 3 + 2} = 1.8$$

Number of counts in 15 second intervals	0	1	2	3	4
Predicated Number of intervals with above counts	3.3	6.0	5.3	3.2	1.4

chi-square statistic

$$\chi^2 \approx \sum_{i=1}^{\text{datapoints}} \frac{(\text{measured}_i - \text{predicted}_i)^2}{\text{predicted}_i}$$



will be discussed later

Example


Example: Back to our cosmic ray example We can calculate the probability that our data agrees with the prediction from the Poisson using:

$$\chi^2 \approx \sum_{i=1}^{\text{data points}} \frac{(\text{measured}_i - \text{predicted}_i)^2}{\text{predicted}_i} = 1.03$$

$$P(\chi^2 \geq 1.03, n = 3) \approx 0.79.$$

Thus we say that the probability of getting a $\chi^2 \geq 1.03$ with 3 degrees of freedom by chance is 0.79 (79%).

Thus the data “looks” Poisson!




Example

The field of particle physics provides numerous situations where the χ^2 test can be applied. A particularly simple example involves measurements of the mass M_Z of the Z^0 boson by experimental groups at CERN. The results of measurements of M_Z made by four different detectors (L3, OPAL, Aleph and Delphi) are as follows:

Detector	Mass in GeV/c^2
L3	91.161 ± 0.013
OPAL	91.174 ± 0.011
Aleph	91.186 ± 0.013
Delphi	91.188 ± 0.013

Example

The listed uncertainties are estimates of the ie, the standard deviations for each of the measurements. The figure below shows these measurements plotted on a horizontal mass scale (vertically displaced for clarity).



Measurements of the Z^0 boson.

Example

The question arises: Can these data be well described by a single number, namely an estimate of M_Z made by determining the weighted mean of the four measurements?

We find the weighted mean \overline{M}_Z and its standard deviation $\sigma_{\overline{M}_Z}$

$$\overline{M}_Z = \frac{\sum M_i / \sigma_i^2}{\sum 1 / \sigma_i^2} \quad \text{and} \quad \sigma_{\overline{M}_Z}^2 = \frac{1}{\sum 1 / \sigma_i^2}$$


$$\overline{M}_Z \pm \sigma_{\overline{M}_Z} = 91.177 \pm 0.006$$

Example

$$\chi^2 = \sum_{i=1}^4 \frac{(M_i - \bar{M}_Z)^2}{\sigma_i^2} \approx 2.78$$

We expect this value of 2 to be drawn from a chi-square distribution with 3 degrees of freedom. The number is 3 (not 4) because we have used the mean of the four measurements to estimate the value of , the true mass of the Z^0 boson, and this uses up one degree of freedom. Hence the $\chi^2/n = 2.78/3 = 0.93$. Now from the graph we find that for 3 degrees of freedom, is about 0.42, meaning that if we were to repeat the experiments we would have about a 42 percent chance of finding a χ^2 for the new measurement set larger than 2.78, assuming our hypothesis is correct.

We have therefore no good reason to reject the hypothesis, and conclude that the four measurements of the Z^0 boson mass are consistent with each other.



Uses of the χ^2 distribution

The dominant use of the χ^2 statistics is for least squares fitting.

$$\chi^2 = \sum_{i=1}^N \left| \frac{y_i - f(x_i | \vec{\alpha})}{\sigma_i} \right|^2$$

The “best fit” values of the parameters α are those that minimize the χ^2 .

If there are m free parameters, and the deviation of the measured points from the model follows Gaussian distributions, then this statistic should be a χ^2 with $N-m$ degrees of freedom. More on this later.

χ^2 is also used to test the goodness of the fit

An exponential distribution

The random variable X that equals the distance between successive events of a Poisson process with mean $\lambda > 0$ is an **exponential random variable** with parameter λ . The probability density function of X is

$$f(x) = \lambda e^{-\lambda x} \quad \text{for } 0 \leq x < \infty$$

Consider for example the distribution of measured lifetimes for a decaying particle:

If the random variable X has an exponential distribution with parameter λ ,

$$\mu = E(X) = \frac{1}{\lambda} \quad \text{and} \quad \sigma^2 = V(X) = \frac{1}{\lambda^2}$$

An exponential distribution

The random variable X that equals the distance between successive events of a Poisson process with mean $\lambda > 0$ is an **exponential random variable** with parameter λ . The probability density function of X is


$$f(x) = \lambda e^{-\lambda x} \quad \text{for } 0 \leq x < \infty$$

Consider for example the distribution of measured lifetimes for a decaying particle:

If the random variable X has an exponential distribution with parameter λ ,

$$\mu = E(X) = \frac{1}{\lambda} \quad \text{and} \quad \sigma^2 = V(X) = \frac{1}{\lambda^2}$$

An exponential distribution



A Uniform distribution

A continuous random variable X with probability density function

$$f(x) = 1/(b - a), \quad a \leq x \leq b$$

is a **continuous uniform random variable**.



A Uniform distribution

Mean and Variance

If X is a continuous uniform random variable over $a \leq x \leq b$,

$$\mu = E(X) = \frac{(a + b)}{2} \quad \text{and} \quad \sigma^2 = V(X) = \frac{(b - a)^2}{12}$$



A Uniform distribution

Let the continuous random variable X denote the current measured in a thin copper wire in milliamperes. Assume that the range of X is $[0, 20 \text{ mA}]$, and assume that the probability density function of X is $f(x) = 0.05, 0 \leq x \leq 20$.

What is the probability that a measurement of current is between 5 and 10 milliamperes?

$$\begin{aligned} P(5 < X < 10) &= \int_5^{10} f(x) dx \\ &= 5(0.05) = 0.25 \end{aligned}$$

The mean and variance formulas can be applied with $a = 0$ and $b = 20$. Therefore,

$$E(X) = 10 \text{ mA} \quad \text{and} \quad V(X) = 20^2/12 = 33.33 \text{ mA}^2$$

Consequently, the standard deviation of X is 5.77 mA.

A Uniform distribution

The cumulative distribution function of a continuous uniform random variable is obtained by integration. If $a < x < b$,

$$F(x) = \int_a^x 1/(b - a) du = x/(b - a) - a/(b - a)$$

Therefore, the complete description of the cumulative distribution function of a continuous uniform random variable is

$$F(x) = \begin{cases} 0 & x < a \\ (x - a)/(b - a) & a \leq x < b \\ 1 & b \leq x \end{cases}$$

Useful NumPy Methods for Statistics

Let x be a numpy array of values.....

$x.mean()$ returns the mean of the values contained in array x .

$x.var()$ returns the variance of the values about their mean.

$x.std()$ returns the standard deviation of the values about their mean.

Useful NumPy Methods for Statistics

```
import numpy as np
import math
import matplotlib.pyplot as pl
from numpy.random import RandomState


r = RandomState()
Nexp = 10000 # number of experiments
Nsam = 100 # number of samples per experiment

# initialize array to hold experiment results
# experiment_results = np.zeros(nexp)
# now conduct nexp experiments with nsam samples
for experiment in range(nexp):
    x = r.randn(nsam)
    experiment_results[experiment] = x.mean()


fr = experiment_results.mean()
fe = experiment_results.std()
ee = 1./math.sqrt(nsam) # expected error on the mean

print ('Final results')
print ('mean ={0:8.5f} Expected = 0'.format(fr))
print ('error={0:8.5f} Expected ={1:8.5f}'.format(fe, ee))
```

For nexp experiments we take nsam samples from a normal distribution and compute the mean.



Find mean value and standard deviation of the experiment results.



In Class Exercise

Write a python code that

- reads data in page 67,
- calculate prediction assuming poisson distribution
- fit data with poisson distribution
- Calculate χ^2
- Calculate $\chi^2/\text{n.d.f}$
- Is the process poisson ? Give the probability and interpret your result