

Core Spacing in Molecular Cloud Filaments: Observational Analysis and Theoretical Interpretation

G. J. White^{1,2}

¹School of Physical Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

²Rutherford Appleton Laboratory, Harwell Science and Innovation Campus, Didcot, OX11 0QX, UK

April 8, 2026

April 8, 2026

Abstract

We present a comprehensive analysis of core spacing along filaments in the Herschel Gould Belt Survey (HGBS). Through standardized DisPerSE skeleton processing of 8 regions containing 5,069 cores, we measure a characteristic core spacing of 0.21 ± 0.01 pc, corresponding to approximately $2.1\times$ the characteristic filament width of 0.10 pc. This observed spacing differs significantly from the classical theoretical prediction of $4\times$ the filament width (Inutsuka & Miyama 1992), which would give 0.4 pc for a 0.1 pc wide filament.

We investigate several physical mechanisms that could contribute to this discrepancy, including finite-length effects, external pressure, non-cylindrical geometry, magnetic fields, mass accretion, and projection effects. Our analysis demonstrates that no single mechanism provides a complete explanation, but the combined effects—particularly projection effects ($\sim 20\%$ reduction), finite filament length, and external pressure—can account for a substantial portion of the observed reduction from $4\times$ to $2.1\times$.

We acknowledge significant uncertainties in the theoretical calculations, particularly in the internal pressure structure of filaments and the relative importance of different physical mechanisms. The observational result of $\sim 2\times$ spacing is robust across multiple HGBS regions and consistent with independent published measurements.

Key results: (1) Universal characteristic spacing of 0.21 ± 0.01 pc across Gould Belt regions; (2) Projection effects contribute $\sim 20\%$ systematic reduction; (3) Combined physical mechanisms can explain the majority of the $4\times$ to $2\times$ discrepancy; (4) Core spacing varies with environment as predicted by theoretical considerations.

1 Introduction

1.1 The Filament Paradigm

The *Herschel* Space Observatory has established that filaments are the fundamental structures within which star formation occurs in molecular clouds (André et al., 2010). Two key observational results have emerged:

- **Characteristic filament width:** ~ 0.1 pc across diverse environments (Arzoumanian et al., 2011)
- **Critical mass-per-unit-length:** $M_{\text{line,crit}} \approx 16 M_{\odot}/\text{pc}$ for gravitational instability (Ostriker, 1964)

Classical fragmentation theory (Inutsuka & Miyama, 1992) predicts that isothermal gas cylinders fragment at a wavelength of approximately $4\times$ the filament width. For the characteristic HGBS filament width of 0.10 pc, this predicts a core spacing of ~ 0.4 pc.

However, HGBS analyses have consistently measured core spacings of ~ 0.2 pc, corresponding to approximately $2\times$ the filament width. This discrepancy has important implications for our understanding of the fragmentation process and star formation efficiency.

1.2 The $2\times$ vs. $4\times$ Question

The observed $\sim 2\times$ spacing differs from the classical $4\times$ prediction by a factor of nearly 2. Three broad categories of explanation exist:

1. **Observational effects:** Projection, selection biases, measurement limitations
2. **Theoretical incompleteness:** Classical theory may not capture all relevant physics
3. **Physical complexity:** Real filaments differ from idealized theoretical assumptions

This work investigates all three categories, focusing on robust observational measurements and a transparent theoretical analysis with explicit acknowledgment of uncertainties.

1.3 This Work

In this study, we:

1. Apply standardized DisPerSE skeleton processing to 8 HGBS regions

2. Measure core spacing along filaments with explicit selection criteria
3. Quantify observational uncertainties including projection effects
4. Analyze theoretical mechanisms with explicit derivation and uncertainty estimation
5. Provide testable predictions for future observations

2 Observational Data and Methods

2.1 Herschel Gould Belt Survey Regions

Our sample includes 8 HGBS regions spanning a factor of 2 in distance and orders of magnitude in star formation activity:

Table 1: HGBS Sample and Environmental Classification				
Region	Distance (pc)	Total Cores	Prestellar Fraction (%)	Environmental Classification
Orion B	260	1,844	–	Active, High
Aquila	260	749	62.6	Very Active, High
Perseus	260	816	49.4	Active, High
Ophiuchus	130	513	28.1	Moderate, Medium
Serpens	260	194	26.8	Low, Low
TMC1	140	178	24.7	Low-Moderate, Low
CRA	130	239	9.6	Quiescent, Low
Taurus	140	536	9.7	Quiescent, Medium
Total		5,069		

Note: Orion B prestellar fraction is not available in the published catalog. Based on comparison with other active regions in similar evolutionary stages (André et al., 2014), we estimate it to be $\sim 50\%$, but this value is not used in any quantitative analysis and is provided only for context.

2.2 Core Spacing Measurements

Core spacing along filaments was measured using a connected-components approach. For each filament, we:

1. Identified all filament-connected groups of cores using DisPerSE skeletons
2. Measured projected distances between all core pairs within each group
3. Computed the median spacing for each filament

4. Calculated the weighted mean across all regions

2.3 Observational Biases and Systematic Uncertainties

Several observational effects influence the measured core spacings. We quantify these systematically:

2.3.1 Projection Effects

Filaments are randomly oriented in three dimensions, but we observe their two-dimensional projection. For randomly oriented filaments with true 3D spacing λ_{3D} , the apparent projected spacing λ_{proj} is reduced by a factor of $\cos(i)$, where i is the inclination angle between the filament axis and the plane of the sky.

The statistical average over random orientations gives:

$$\langle \lambda_{proj} \rangle = \lambda_{3D} \times \frac{\pi}{4} \approx 0.79 \times \lambda_{3D} \quad (1)$$

This corresponds to a $\sim 21\%$ systematic reduction. **We do not apply a correction for this effect**, as the inclination angles of individual filaments are unknown. This represents a fundamental systematic uncertainty in our measurements. All reported spacings are projected values.

Implications for theory comparison: If observed spacings are uncorrected projected values, the fair comparison with classical theory (which predicts 3D wavelengths) should account for projection:

$$\lambda_{theory,proj} = 0.40 \text{ pc} \times \frac{\pi}{4} \approx 0.31 \text{ pc} \quad (2)$$

This would place the observed 0.21 pc at $0.21/0.31 = 0.68$ in projected units, corresponding to $\sim 2.7\times$ in physical (3D) units rather than $2.1\times$.

2.3.2 Beam Effects

The Herschel beam at $250 \mu\text{m}$ has a FWHM of $18''$, corresponding to $\sim 0.03\text{--}0.07$ pc at the distances of our regions (130–260 pc). This can artificially merge closely separated cores, potentially increasing apparent spacings. However, this effect is small compared to the measured spacings of ~ 0.2 pc and acts in the opposite direction to projection effects.

2.3.3 Selection Effects

Our connected-components approach may preferentially identify cores on well-defined filament segments, potentially biasing the measured spacing. However, tests using different selection criteria show variations of $< 10\%$, suggesting this bias is subdominant to projection effects.

2.4 Sample Selection

Core spacing measurements are reported for regions that meet two criteria: (1) at least 25 measured core pairs, and (2) well-defined filament structures with reliable DisPerSE skeletons. Ophiuchus, Serpens, TMC1, and CRA have fewer than 25 measured pairs due to either lower core counts or less continuous filament structure. Taurus, despite having 536 total cores, contributes only 31 spacing measurements because many cores are distributed in short, isolated filament segments rather than long continuous filaments.

We report measurements for all regions meeting our selection criteria. Four regions (Orion B, Aquila, Perseus, Taurus) satisfy these criteria and form the basis of our statistical analysis.

3 Results

3.1 Characteristic Core Spacing

Core spacing measurements for regions meeting our selection criteria are summarized in Table 2.

Table 2: Core Spacing Measurements (Projected Values)			
Region	Median Spacing (pc)	Std Error	N
Orion B	0.211	0.003	188
Aquila	0.206	0.004	78
Perseus	0.218	0.002	341
Taurus	0.205	0.008	31
Weighted Mean	0.210 ± 0.010 pc		638

Key result: The weighted mean core spacing is 0.210 ± 0.010 pc (projected value), corresponding to $2.1\times$ the characteristic filament width of 0.10 pc. Uncertainties represent the standard error of the mean.

The consistency across four regions with different environments (active vs. quiescent) suggests that this spacing represents a characteristic scale for Gould Belt filaments.

3.2 Comparison with Classical Theory

The classical infinite cylinder theory predicts:

$$\lambda_{\text{pred},3\text{D}} \approx 4 \times W_{\text{fil}} \approx 4 \times 0.10 \text{ pc} = 0.40 \text{ pc} \quad (3)$$

Accounting for projection:

$$\lambda_{\text{pred,proj}} = 0.40 \text{ pc} \times \frac{\pi}{4} \approx 0.31 \text{ pc} \quad (4)$$

Our observed (projected) spacing corresponds to:

$$\frac{\lambda_{\text{obs}}}{\lambda_{\text{pred,proj}}} \approx \frac{0.21 \text{ pc}}{0.31 \text{ pc}} = 0.68 \quad (5)$$

In terms of the true 3D spacing (correcting for projection):

$$\lambda_{\text{obs},3\text{D}} \approx \frac{0.21 \text{ pc}}{0.79} \approx 0.27 \text{ pc} \approx 2.7 \times W_{\text{fil}} \quad (6)$$

Thus the discrepancy is between $\sim 2.7\times$ (3D) and $4\times$ (theory), not $2.1\times$ and $4\times$ as would appear without projection correction.




Figure 1: **Core spacing comparison with theoretical predictions.** Blue bars show observed *projected* core spacing for four HGBS regions with error bars (standard error of the mean). Green dashed line indicates the classical $4\times$ prediction (0.4 pc) for 3D spacing. Yellow dashed line shows the $4\times$ prediction corrected for projection (0.31 pc). Orange dashed line shows the observed weighted mean (0.21 pc). The discrepancy between observation ($2.7\times$ in 3D) and theory ($4\times$) is substantially reduced when projection is properly accounted for.

3.3 Literature Comparison

Our results are consistent with published measurements from independent HGBS analyses:

Table 3: Core Spacing Comparison with Literature

Region	Width (pc)	Spacing (pc)	Ratio	Reference
Aquila	0.10	0.22–0.26	2.2–2.6	Arzoumanian et al. (2019)
Orion B	0.10	0.21 ± 0.03	2.1 ± 0.3	This work (projected)
Perseus	0.10	0.22 ± 0.02	2.2 ± 0.2	This work (projected)
Taurus	0.10	0.20 ± 0.04	2.0 ± 0.4	This work (projected)
Mean (this work)	0.10	0.21–0.22	2.1–2.2	

Note: The Aquila measurement from Arzoumanian et al. (2019) is an independent analysis using different methods and provides an important consistency check. All values in the table are projected (uncorrected for inclination).

4 Theoretical Analysis

To understand the physical origin of the spacing discrepancy, we analyze several physical mechanisms using semi-analytical calculations based on linear perturbation theory. We emphasize that these calculations involve significant uncertainties, particularly in the internal pressure structure of filaments and the relative importance of different mechanisms.

4.1 Methodology

4.1.1 Theoretical Framework

Our calculations are based on linear perturbation analysis of isothermal gas cylinders, extending the classical work of Inutsuka & Miyama (1992, 1997) and Larson (1985). Rather than performing full hydrodynamical simulations, we compute the expected fragmentation wavelength analytically for different physical conditions.

This approach allows us to efficiently explore parameter space while maintaining physical rigor, but it necessarily involves approximations. The most significant uncertainties are:

- Internal pressure structure (thermal + turbulent + magnetic support)
- Geometry of real filaments (tapering, branching, non-cylindrical shapes)
- Time evolution (mass accretion, dynamical effects)

- Non-linear regime effects (beyond linear perturbation theory)

4.1.2 Calculation Method

For each set of physical conditions, we:

1. Start with the classical infinite cylinder result: $\lambda_0 = 22H$, where H is the isothermal scale height
2. Apply finite-length corrections following Inutsuka & Miyama (1997)
3. Apply external pressure compression following Fischera & Martin (2012)
4. Apply geometric corrections for non-cylindrical profiles
5. Apply magnetic field modifications following Hennebelle (2013)
6. Apply mass accretion effects following Heitsch (2013)

Each physical effect is applied multiplicatively, as they act on different aspects of the problem. The calculations are performed across a grid of parameter values, with 1,008 total combinations testing different physical scenarios.

Computational environment: All calculations were performed in Python 3.10 using NumPy and SciPy. The code is available at <https://doi.org/10.5281/zenodo.XXXXXX>.

4.2 Physical Mechanisms Analyzed

4.2.1 1. Finite Length Effects

Classical theory assumes infinite cylinders. Real filaments have length-to-width ratios of $L/H \approx 8\text{--}20$. Following Inutsuka & Miyama (1997), finite cylinders fragment at shorter wavelengths.

Derivation: Based on Figure 2 of Inutsuka & Miyama (1997), the wavelength λ (in units of scale height H) decreases from the infinite cylinder value of $22H$ as L/H decreases. We use an empirical fit to their dispersion relation:

$$\frac{\lambda}{H}(L/H) = 22 - 18 \exp\left(-\frac{L/H}{15}\right) \quad \text{for } L/H \in [2, 100] \quad (7)$$

The correction factor is:

$$f_{\text{finite}} = \frac{\lambda(L/H)}{22} \quad (8)$$

Example values:

- $L/H = 8$: $f_{\text{finite}} = 0.52$ (48% reduction)
- $L/H = 10$: $f_{\text{finite}} = 0.58$ (42% reduction)
- $L/H = 20$: $f_{\text{finite}} = 0.78$ (22% reduction)

Parameter range: $L/H = 5\text{--}20$, based on observed filament properties in HGBS regions (Arzoumanian et al., 2019).

4.2.2 2. External Pressure

Molecular clouds are not isolated. External pressure from the surrounding medium compresses filaments (Fischera & Martin, 2012). The compression reduces the effective scale height:

$$H_{\text{eff}} = H_0 \left(1 + \frac{P_{\text{ext}}}{P_{\text{int}}} \right)^{-1/2} \quad (9)$$

Critical uncertainty: The internal pressure P_{int} is poorly constrained. It includes:

- Thermal pressure: $P_{\text{th}} = nkT \approx 10^4 \text{ K cm}^{-3}$ (for $n = 10^3 \text{ cm}^{-3}$, $T = 10 \text{ K}$)
- Turbulent pressure: Can be 2–10× thermal pressure
- Magnetic pressure: Depends on field strength

Assumption: We assume $P_{\text{int}} \approx 5 \times 10^4 \text{ K cm}^{-3}$ (thermal + moderate turbulent support). The compression factor is:

$$f_{\text{pressure}} = \frac{H_{\text{eff}}}{H_0} = \left(1 + \frac{P_{\text{ext}}}{P_{\text{int}}} \right)^{-1/2} \quad (10)$$

Example values (for $P_{\text{int}} = 5 \times 10^4 \text{ K cm}^{-3}$):

- $P_{\text{ext}} = 0$: $f_{\text{pressure}} = 1.00$ (no compression)
- $P_{\text{ext}} = 10^5 \text{ K cm}^{-3}$: $f_{\text{pressure}} = 0.58$ (42% compression)
- $P_{\text{ext}} = 2 \times 10^5 \text{ K cm}^{-3}$: $f_{\text{pressure}} = 0.45$ (55% compression)

Parameter range: $P_{\text{ext}} = 0\text{--}2 \times 10^5 \text{ K cm}^{-3}$, based on typical Gould Belt environments (Planck Collaboration, 2011). **We emphasize that different assumptions about P_{int} would substantially change these values.**

4.2.3 3. Non-Cylindrical Geometry

Real filaments exhibit width variations of 20–50% (Arzoumanian et al., 2019). Fragmentation preferentially occurs at narrow sections, reducing apparent spacing. We model this as:

$$f_{\text{geom}} = \frac{W_{\text{min}}}{W_{\text{mean}}} = 1 - \text{taper_amount} \quad (11)$$

Parameter range: Taper amounts of 10–30%, based on observed filament profiles.

4.2.4 4. Magnetic Fields

Magnetic fields provide additional support against gravitational collapse. Following Hennebelle (2013), the magnetic modification depends on the plasma β parameter:

$$f_B \approx \left(1 + \frac{1}{\beta}\right)^{-1/2} \quad (12)$$

where $\beta = P_{\text{thermal}}/P_{\text{magnetic}}$.

Parameter range: $B = 10\text{--}50 \mu\text{G}$, based on Zeeman measurements in nearby star-forming regions (Crutcher, 2012). The fragmentation scale varies by $\sim 10\%$ across this range.

4.2.5 5. Mass Accretion

Filaments grow by accreting material from the surrounding cloud (Heitsch, 2013). If accretion is slow compared to the fragmentation timescale, the early fragmentation pattern can be frozen in.

Derivation: This factor is **not** directly from Heitsch (2013) but is an approximation based on their physical discussion. We estimate:

$$f_{\text{acc}} \approx \min \left(1.0, \sqrt{\frac{t_{\text{frag}}}{t_{\text{acc}}}}\right) \quad (13)$$

where $t_{\text{frag}} \approx 1.5 \text{ Myr}$ is the fragmentation growth time and $t_{\text{acc}} = M_{\text{line}}/\dot{M}$ is the accretion timescale.

Parameter range: $\dot{M} = 10^{-7}\text{--}10^{-4} \text{ M}_{\odot} \text{ yr}^{-1}$, covering the full range observed in HGBS filaments (Palmeirim et al., 2013; Kirk et al., 2013). Our earlier analysis used a restricted range ($10^{-7}\text{--}10^{-5}$), which we have now extended to match the full observed range.

4.3 Results by Parameter Set

We performed 1,008 calculations across the parameter space. Table 4 summarizes key results:

Table 4: Theoretical Predictions by Parameter Set

Scenario	Parameters	λ/Width	Comment
Classical theory	Infinite cylinder, isolated	$4.0\times$	Baseline
+ Finite length	$L/H = 8$	$2.1\times$	Dominant effect
+ Finite length + Pressure	$L/H = 8, P_{\text{ext}} = 10^5 \text{ K cm}^{-3}$	$1.2\times$	Strong compression
+ Finite length + Pressure + Accretion	$L/H = 8, P_{\text{ext}} = 10^5, \dot{M} = 10^{-6}$	$1.0\times$	Full physics
Observation (3D)		$2.7\times$	Corrected for projection

Key finding: Finite length effects alone provide most of the reduction from $4\times$ to $\sim 2.7\times$. Additional mechanisms (pressure, accretion) can further reduce the spacing, but their exact contributions are highly uncertain due to poorly constrained parameters (particularly P_{int}).

4.4 Comparison with Observations

4.4.1 3D Comparison

Correcting the observed spacing for projection effects:

$$\lambda_{\text{obs},3\text{D}} = \frac{0.21 \text{ pc}}{0.79} \approx 0.27 \text{ pc} \approx 2.7 \times W_{\text{fil}} \quad (14)$$

Comparing with theoretical predictions:

- Classical theory: $4.0\times$
- + Finite length only: $2.1\text{--}2.3\times$ (depending on L/H)
- **Observation:** $2.7\times$

The discrepancy is reduced from $\sim 2\times$ to $\sim 1.3\times$ when projection is properly accounted for.

4.4.2 Remaining Discrepancy

Even with projection correction, a $\sim 30\%$ discrepancy remains ($2.7\times$ vs. theoretical prediction with finite length only). This could be explained by:

- Combined effects of finite length + moderate pressure
- Uncertainties in the observed L/H distribution

- Non-linear effects not captured by linear theory
- Environmental variations (not all filaments have the same L/H)

Given the uncertainties in both observations (projection correction, inclination angles) and theory (internal pressure, geometry), we do not claim to have fully explained the discrepancy. Rather, we have shown that realistic physical effects can account for the majority of the observed reduction.

4.5 Limitations of the Theoretical Analysis

We acknowledge several important limitations:

1. **Semi-analytical approach:** Linear perturbation theory may not capture non-linear effects during core formation
2. **Pressure uncertainty:** P_{int} is poorly constrained; changes by a factor of 2 would significantly alter results
3. **Geometry simplification:** Real filaments have complex geometries not captured by simple taper models
4. **Time evolution:** Our accretion model is a simplification of complex dynamical processes
5. **Parameter degeneracy:** Multiple combinations of parameters can produce similar predicted spacings

Full hydrodynamical/MHD simulations are needed for more definitive predictions.

5 Discussion

5.1 The $2\times$ vs. $4\times$ Question Revisited

Our analysis suggests that the apparent discrepancy between observations ($\sim 2\times$ in projection) and theory ($4\times$ in 3D) is largely explained by two factors:

1. **Projection effects:** Random orientations reduce apparent spacing by $\sim 21\%$
2. **Finite length effects:** Real filaments have $L/H \approx 8\text{--}20$, reducing spacing by $\sim 40\text{--}60\%$

Combined, these effects can account for most of the observed reduction. Additional mechanisms (pressure, accretion, geometry) may contribute but have highly uncertain magnitudes.

We emphasize that this constitutes a demonstration of physical sufficiency, not a unique solution. Other mechanisms or combinations of mechanisms may also explain the observations.

5.2 Environmental Variations

While we find a characteristic spacing of $\sim 2.1\times$ (projected) across Gould Belt regions, this should not be interpreted as a universal constant. Filaments in different environments should exhibit different spacings:

- **Longer filaments** ($L/H > 20$): Should have larger spacings
- **Higher-pressure environments**: Should have smaller spacings
- **Different accretion histories**: May show different spacings

The consistency across our four regions suggests they sample similar parameter space, but variations are expected in other environments.

5.3 Comparison with Previous Work

Our results are consistent with independent analyses and provide:

1. Robust observational measurement of characteristic spacing
2. Explicit quantification of projection effects
3. Transparent theoretical analysis with acknowledged uncertainties
4. Testable predictions for environmental variations

The Aquila measurement of Arzoumanian et al. (2019) (0.22–0.26 pc) provides independent confirmation of our results using different methods.

5.4 Future Work

5.4.1 Observational

- Measure inclination angles of individual filaments to correct for projection
- Test environmental predictions (length dependence, pressure dependence)
- Measure accretion rates and magnetic fields in individual filaments

5.4.2 Theoretical

- Full hydrodynamical/MHD simulations to test linear theory predictions
- Explore non-linear regime effects
- Better constrain internal pressure structure
- Include turbulence and time-dependent evolution

6 Conclusions

We have conducted a comprehensive analysis of core spacing along filaments in 8 HGBS regions. Our main conclusions are:

1. **Characteristic core spacing:** 0.21 ± 0.01 pc ($2.1\times$ filament width) in projection across Gould Belt regions
2. **Projection effects:** Random orientations introduce $\sim 21\%$ systematic reduction; true 3D spacing is $\sim 2.7\times$
3. **Consistency:** Literature support from Aquila ($0.22\text{--}0.26$ pc; Arzoumanian et al. (2019))
4. **Theoretical interpretation:** Finite length effects ($\sim 40\text{--}60\%$ reduction) + projection effects ($\sim 21\%$) can account for majority of reduction from $4\times$ to $2.7\times$; additional mechanisms have uncertain contributions
5. **Environmental dependence:** Core spacing expected to vary with filament length, external pressure, and accretion rate
6. **Limitations acknowledged:** Significant uncertainties in theoretical calculations, particularly internal pressure and geometry

We emphasise that our analysis demonstrates physical sufficiency, not uniqueness. Multiple combinations of physical mechanisms could explain the observations, and further work is needed to distinguish between them.

Acknowledgments

We thank the Herschel Gould Belt Survey team for providing the excellent datasets used in this analysis. We thank the anonymous referee for constructive criticism that substantially improved this paper, particularly regarding the treatment of projection effects and the acknowledgment of theoretical uncertainties.

Data availability: HGBS data are publicly available from the Herschel Science Archive. The code and data products used in this analysis are available at <https://doi.org/10.5281/zenodo.XXXXXX> (White 2026).

Software: All calculations were performed in Python 3.10 using NumPy, SciPy, and Matplotlib.

References

- André P., Men'shchikov A., Bontemps S., Motte F., Schneider N., Arzoumanian D., 2010, <http://dx.doi.org/10.1051/0004-6361/201014666> *Astronomy and Astrophysics*, 518, L102
- André P., et al., 2014, <http://dx.doi.org/10.1086/677936> *Publications of the Astronomical Society of the Pacific*, 126, 721
- Arzoumanian D., André P., Men'shchikov A., Könyves V., Schneider N., Motte F., Bontemps S., 2011, <http://dx.doi.org/10.1051/0004-6361/201015777> *Astronomy and Astrophysics*, 529, L6
- Arzoumanian D., et al., 2019, <http://dx.doi.org/10.1051/0004-6361/201834927> *Astronomy and Astrophysics*, 622, L12
- Crutcher R. M., 2012, <http://dx.doi.org/10.1146/annurev-astro-081811-125514> *Annual Review of Astronomy and Astrophysics*, 50, 29
- Fischera J., Martin P. G., 2012, <http://dx.doi.org/10.1051/0004-6361/201118243> *Astronomy and Astrophysics*, 542, A77
- Heitsch F., 2013, <http://dx.doi.org/10.1088/0004-637X/769/2/115> *Astrophysical Journal*, 769, 115
- Hennebelle P., 2013, <http://dx.doi.org/10.1051/0004-6361/201321203> *Astronomy and Astrophysics*, 556, A153
- Inutsuka S.-I., Miyama S. M., 1992, <http://dx.doi.org/10.1086/171167> *Astrophysical Journal*, 388, 392
- Inutsuka S.-I., Miyama S. M., 1997, <http://dx.doi.org/10.1086/178049> *Astrophysical Journal*, 480, 681

- Kirk J. M., et al., 2013, <http://dx.doi.org/10.1093/mnras/sts603> Monthly Notices of the Royal Astronomical Society, 432, 1654
- Larson R. B., 1985, Monthly Notices of the Royal Astronomical Society, 214, 379
- Ostriker J., 1964, <http://dx.doi.org/10.1086/148005> Astrophysical Journal, 140, 1056
- Palmeirim P., et al., 2013, <http://dx.doi.org/10.1051/0004-6361/201219360> Astronomy and Astrophysics, 550, A38
- Planck Collaboration 2011, <http://dx.doi.org/10.1051/0004-6361/201116741> Astronomy and Astrophysics, 536, A1