

ASTRA: Autonomous Scientific Discovery in Astrophysics

Glenn J. White^{1,2*}

¹*Department of Physics and Astronomy, The Open University, Milton Keynes MK7 6AA, UK*

²*RAL Space, STFC Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire, OX11 0QX, UK*

17 June 2026

Key words: methods: data analysis – methods: statistical – ISM: structure – galaxies: general – software: public release

ABSTRACT

We present ASTRA (Autonomous System for Scientific Discovery in Astrophysics), an integrated framework combining numerical data analysis, causal reasoning, and physical validation. ASTRA integrates: dimensional analysis; structural causal models; Bayesian evidence computation; and multi-wavelength data fusion within a unified framework.

We demonstrate capabilities through seven test cases. The first five use real observational data to validate ASTRA’s ability to recover known astrophysical results, including: Herschel filament scaling relations with dimensional analysis; Chandra Deep Field South multi-wavelength cross-matching with Bayesian uncertainty propagation; SDSS galaxy property correlations; Gaia stellar causal inference; and Bayesian model selection for virial scaling relations.

The sixth test case demonstrates discovery-mode operation on realistic galaxy survey data (3,000 galaxies based on SDSS scaling relations). ASTRA recovers the star-forming main sequence and mass-metallicity relation, then applies multi-dimensional outlier detection to identify starbursts, post-starbursts, and transitioning galaxies.

The seventh test case demonstrates autonomous HPC simulation campaign management: ASTRA co-ordinated over 1,200 Athena++ MHD simulations on a remote 224-vCPU server across seven interlinked campaigns to validate the magnetic Jeans fragmentation criterion for ISM filaments, confirming the predicted stability threshold $\beta_{\text{crit}} = 2/3$, calibrating the fragmentation wavelength to $(1.11 \pm 0.12) \lambda_{J,m}(\theta, \beta)$ across 18 field-geometry configurations, establishing 2D power-law scalings for fragmentation timescale ($t_{\text{frag}} \propto f^{-0.54} \beta^{-0.16}$ for longitudinal fields; $t_{\text{frag}} \propto f^{-0.86} \beta^{-0.13}$ for perpendicular fields), measuring the fragmentation wavelength-to-width ratio $\lambda/W = 1.631 \pm 0.034$ (universal for perpendicular-field filaments), and demonstrating that an adiabatic equation of state completely suppresses near-critical fragmentation, validating the isothermal assumption as the conservative lower bound.

Code and documentation: <https://github.com/web3guru888/ASTRA> (White 2026).

1 INTRODUCTION

Modern astronomical surveys generate petabytes of data requiring automated analysis. Multiple computational approaches are available to astronomers, each with distinct strengths and limitations.

Traditional Machine Learning and Statistical Methods: Well-established tools including scikit-learn, emcee, and nested sampling codes (MultiNest, dynesty) can process numerical data, detect patterns, perform statistical inference, and compute Bayesian evidence. These methods are widely used in astronomy with demonstrated success.

Large Language Models with Code Execution: Current frontier models (GPT-4 with Code Interpreter, Claude with code execution) can process numerical data, execute Python code, and perform statistical analysis. These systems provide conversational interfaces to computational tools but require explicit programming and do not natively integrate physical validation or causal interpretation.

The Integration Challenge: While individual tools exist for specific tasks, astronomers typically must manually combine them. This manual integration creates opportunities for inconsistent treatment of data and physical models, leading to errors and inefficiencies. The ASTRA framework addresses these challenges by providing a unified, automated pipeline for data processing, statistical analysis, causal inference, and physical validation. This manual integration creates opportunities for inconsistent treatment of data and physical models, leading to errors and inefficiencies.

quantification are applied consistently throughout multi-step analyses. While individual components use established algorithms (PC algorithm for causal discovery, marginal likelihood estimation for Bayesian evidence, dimensional analysis for physical validation), their integration within a single framework enables analyses that would otherwise require manual combination of multiple separate tools.

In this paper, we focus on seven examples demonstrating how this integrated approach can be applied to astrophysical inference problems:

- (i) **Scaling Relations Analysis** (Section 3): Dimensional analysis and physical validation of filament scaling relations
- (ii) **Multi-Wavelength Data Fusion** (Section 4): Probabilistic cross-matching with Bayesian uncertainty propagation
- (iii) **Pattern Recognition** (Section 5): Identification of established galaxy property relationships
- (iv) **Causal Inference** (Section 6): Structural causal model discovery in stellar data

- (v) **Bayesian Model Selection** (Section 7): Evidence-based model comparison with prior specification
- (vi) **Discovery-Mode Analysis** (Section 8): Galaxy scaling relations and outlier detection on realistic survey data
- (vii) **Autonomous HPC Campaign** (Section 9): Autonomous MHD simulation campaign management for ISM filament fragmentation

Tests 1–5 use real observational data, validating ASTRA’s ability to correctly recover known astrophysical results. Test 6 uses realistic galaxy survey data based on SDSS scaling relations to demonstrate both validation (recovering known relations) and discovery (identifying outliers).

Section 10 discusses why these results demonstrate ASTRA’s unique capabilities, clarifies ASTRA’s role as a tool to assist rather than replace astronomers, and Section 11 concludes.

1.1 Scope and Limitations

What ASTRA Demonstrates: The seven test cases in this paper show ASTRA can: discover physical laws from data with theoretical validation; fuse multi-wavelength observations with proper uncertainty handling; identify established astrophysical relationships through pattern recognition; distinguish causation from correlation; perform rigorous Bayesian model comparison; and operate in discovery mode on realistic survey data to recover known relations while identifying unusual objects for follow-up.

What ASTRA Does Not Claim: ASTRA is not presented as achieving artificial general intelligence (AGI) or AGI-like performance. The system operates within defined astrophysical domains using established algorithms (PC-algorithm causal discovery, Bayesian inference, dimensional analysis) combined through an integrated architecture. Results are validated against known physical theory and observational constraints. The system does not claim general reasoning beyond its training domains or autonomous operation without human oversight. The architecture is designed to support natural language queries that trigger autonomous task completion, though this capability is not demonstrated in the present test cases.

ASTRA’s Role in Scientific Discovery: We should not expect to give ASTRA one of the big contemporary problems like “What is the nature of Dark Matter and Dark Energy,” or “How does gravity connect to quantum physics,” or “What sets the stellar initial mass function (IMF),” and expect to get definitive answers that may be the equivalent of a computational Eureka moment. But ASTRA’s discovery architecture, used alongside and by an experienced astronomer, has capabilities in discovery and inference that go beyond those of straightforward AI (large language models) and/or machine learning. ASTRA’s role in science is to work alongside the experienced astronomer to analyze data and facilitate genuine discovery. ASTRA is therefore a tool to assist the astronomer, rather than a replacement for domain expertise.

Validation Scope: Tests 1–5 use real observational data to validate ASTRA’s ability to correctly recover known astrophysical results. Test 6 uses realistic galaxy survey data based on published SDSS scaling relations to demonstrate both validation (recovering known relations) and discovery (identifying outliers). The data is generated using empirical relations

from [Kauffmann et al. \(2003\)](#), [Tremonti et al. \(2004\)](#), and [Brinchmann et al. \(2004\)](#). The capabilities demonstrated are shown within specific astrophysical domains using particular datasets. Generalization to other domains or datasets requires further validation.

2 METHODS: ASTRA SYSTEM ARCHITECTURE

2.1 Overview

ASTRA implements a modular architecture designed for astrophysical data analysis and inference. The system integrates multiple specialized components through a coordinated framework, enabling complex multi-step analyses that combine data processing, physical reasoning, causal inference, and statistical validation.

2.2 Core Architectural Components

1. Data Processing Pipeline: ASTRA accepts heterogeneous astronomical data formats including catalogs (CSV, FITS), time-series data, images, and spectral data. The pipeline performs ingestion, validation, cleaning, and normalization while preserving measurement uncertainties and metadata.

2. Physics Engine: A unified physics engine implements fundamental physical laws and constraints including conservation laws (mass, energy, momentum), dimensional analysis using Buckingham π theorem, equation solving and numerical simulation, and units consistency checking throughout calculations. This enables ASTRA to validate discoveries against first principles rather than treating correlations as sufficient explanations.

3. Causal Reasoning Module: Implements causal discovery and inference using the PC algorithm ([Spirites et al. 2000](#)) for discovering causal structures from observational data, conditional independence testing for continuous variables (Fisher’s Z-test), V-structure detection for identifying colliders, do-calculus for predicting effects of interventions, and domain-specific adaptations for astrophysical contexts. This module enables ASTRA to distinguish causal relationships from correlations, addressing a fundamental limitation of both traditional ML and LLM approaches.

4. Bayesian Inference Engine: Implements rigorous model comparison and uncertainty quantification through evidence computation via marginal likelihood estimation, Bayes factors for model comparison ([Kass & Raftery 1995](#)), automatic complexity penalty (Occam’s razor), posterior predictive checking, and Monte Carlo methods for uncertainty propagation.

5. Domain Knowledge Systems: Organized astrophysical knowledge through MORK ontology encoding taxonomic and causal relationships, knowledge graphs for semantic relationships, vector-based semantic memory for concept retrieval, and 75 specialized domain modules (ISM, star formation, exoplanets, etc.).

6. Multi-Module Orchestration: Coordinates different analytical perspectives through specialized processing modules for different analysis types, meta-cognitive layer for

context-appropriate module selection, result synthesis and conflict resolution, and quality control and validation checks.

2.3 Unique Capabilities

ASTRA integrates sixteen distinct analytical capabilities that work together to enable comprehensive scientific analysis:

Causal and Statistical Analysis: (1) *Bias Detection*—Identifies and quantifies observational biases like Malmquist bias from survey geometry and selection effects. (2) *Scaling Relations Discovery*—Discovers physical laws from data through dimensional analysis and functional form detection. (3) *Causal Inference*—Discovers causal structures using the PC algorithm and distinguishes physical laws from selection biases. (4) *Model Selection*—Compares competing theories using Bayesian evidence computation with proper complexity penalties.

Data Integration and Analysis: (5) *Multi-Wavelength Fusion*—Cross-matches sources across wavelengths with proper astrometric uncertainty propagation. (6) *Uncertainty Quantification*—Propagates measurement errors through first-order and Monte Carlo methods, separating systematic and statistical components. (7) *Temporal Reasoning*—Detects periodic signals, performs phase folding, and forecasts time-series behavior. (8) *Instrument-Aware Analysis*—Evaluates observational requirements and compatibility across multiple astronomical facilities.

Knowledge Generation: (9) *Hypothesis Generation*—Identifies patterns, detects anomalies, and generates novel testable hypotheses with specific observational requirements. (10) *Analogical Reasoning*—Discovers structural mappings between different astrophysical systems based on shared physics. (11) *Counterfactual Analysis*—Simulates physically-grounded scenarios to predict effects of interventions or observational changes. (12) *Physical Model Discovery*—Automatically identifies functional forms and validates against theoretical predictions.

Meta-Cognitive Capabilities: (13) *Meta-Cognitive Evaluation*—Provides quantitative assessments of data sufficiency, resolution limits, and observational constraints. (14) *Anomaly Detection*—Identifies unusual objects using ensemble methods combining multiple detection approaches. (15) *Ensemble Prediction*—Combines multiple models using Bayesian Model Averaging for robust predictions with uncertainty quantification. (16) *Physical Validation*—Checks all results against dimensional consistency, conservation laws, and established physical principles.

These capabilities are not independent modules but integrated components that work together. For example, discovering a scaling relation (capability 2) involves dimensional analysis (capability 12), physical validation (capability 16), and uncertainty quantification (capability 6). Similarly, multi-wavelength fusion (capability 5) requires coordinate calculations, astrometric error propagation, and physical classification criteria.

Capability Demonstrations in This Paper: Table 1 summarizes which of the 16 capabilities are demonstrated in each of the seven test cases. Capabilities not demonstrated in TCs 1–7 (temporal reasoning, instrument-aware analysis, analogical reasoning, counterfactual analysis, ensemble prediction) are available in ASTRA and demonstrated in ex-

tended test cases in the GitHub repository. TC7 additionally demonstrates a seventeenth capability (HPC simulation management) not listed in the 16-capability inventory.

2.4 Workflow for Scientific Analysis

ASTRA’s analysis workflow follows this general pattern: **Input:** Scientific query or observational data; **Module Selection:** Meta-cognitive layer selects appropriate analysis modules based on query type and data characteristics; **Analysis Execution:** Selected modules perform their specialized analyses including statistical and ML algorithms for pattern detection, causal discovery for identifying relationships, physics engine for theoretical validation, and Bayesian inference for model comparison; **Integration:** Results synthesized from multiple modules, resolving conflicts and identifying consensus conclusions; **Validation:** Results checked against physical constraints (dimensional consistency, conservation laws), statistical significance (p-values, confidence intervals), observational constraints (resolution limits, selection effects), and domain knowledge (established theories, previous results); **Output:** Integrated results with confidence assessments, uncertainty quantification, and physical interpretation.

2.5 Implementation and Availability

ASTRA is implemented in Python with approximately 313,000 lines of code across 620+ files. The system uses established scientific Python libraries (NumPy, SciPy, Pandas, scikit-learn) combined with specialized implementations for causal discovery (causal-learn, dowhy), Bayesian inference, and physics simulation.

The complete system is publicly available at: <https://github.com/web3guru888/ASTRA> (White 2026).

This repository includes: complete source code and installation instructions; system architecture documentation with detailed design diagrams (see also Figure 1); API documentation and user manual; extended validation test cases and example notebooks; and reproducible code for all results presented in this paper. The first five test cases in this paper can be reproduced using the following Jupyter notebooks in the repository: `test02.scaling_relations.ipynb` (Section 3), `test04.multiphysics.fusion.ipynb` (Section 4), `test05.hypothesis_generation.ipynb` (Section 5), `test11.causal_inference.ipynb` (Section 6), and `test12.bayesian_model_selection.ipynb` (Section 7).

2.6 Positioning Relative to Other Approaches

ASTRA occupies a distinct position in the landscape of AI for scientific discovery. **Compared to Pure ML:** Unlike machine learning systems that focus on prediction or classification, ASTRA emphasizes causal understanding, physical interpretation, and theoretical validation. ASTRA can detect correlations like ML systems but goes further by identifying causal structure and validating against physics.

Compared to LLMs: Unlike large language models that require explicit programming and manual tool selection to perform numerical analysis, and do not natively integrate physical validation constraints or causal interpretation, ASTRA provides these capabilities within a unified framework.

Table 1. Capabilities demonstrated in each test case. X = explicitly demonstrated; – = not demonstrated or not applicable. Temporal reasoning, instrument-aware analysis, analogical reasoning, counterfactual analysis, and ensemble prediction are available in ASTRA but demonstrated only in extended test cases on GitHub. TC7 additionally demonstrates HPC Simulation Management (row 17), a qualitatively distinct capability.

Capability	TC1 (Scaling)	TC2 (Multi-λ)	TC3 (Pattern)	TC4 (Causal)	TC5 (Bayesian)	TC6 (Discovery)	TC7 (HPC MHD)
Bias Detection	–	X	–	X	–	–	X
Scaling Relations	X	–	X	–	X	X	X
Causal Inference	–	–	–	X	–	–	–
Model Selection	–	–	–	–	X	–	–
Multi-Wavelength Fusion	–	X	–	–	–	–	–
Uncertainty Quantification	X	X	X	–	X	X	X
Temporal Reasoning	–	–	–	–	–	–	–
Instrument-Aware Analysis	–	–	–	–	–	–	X
Hypothesis Generation	–	–	X	–	–	X	X
Analogical Reasoning	–	–	–	–	–	–	–
Counterfactual Analysis	–	–	–	–	–	–	–
Physical Model Discovery	X	–	–	–	X	–	X
Meta-Cognitive Evaluation	X	X	X	X	X	X	X
Anomaly Detection	–	–	X	–	–	X	–
Ensemble Prediction	–	–	–	–	–	–	–
Physical Validation	X	X	X	X	X	X	X
HPC Simulation Management	–	–	–	–	–	–	X




Figure 1. ASTRA System Architecture. The six core components (Physics Engine, Causal Module, Bayesian Engine, Domain Knowledge, Data Processing Pipeline, HPC Simulation Manager) are co-ordinated through a Meta-Cognitive Orchestration layer that routes natural-language queries and observational data inputs to appropriate analytical modules. Output includes physical interpretations, quantified uncertainties, testable hypotheses, and reproducible workflows. Arrows indicate primary data flow; all components exchange validation signals with the Physics Engine throughout analysis. TC7 (Section 9) exercises the HPC Simulation Manager. The full 16-capability inventory is summarised in Table 1.

Compared to Specialized Tools: Unlike domain-specific tools designed for single tasks (e.g., photometric redshift fitting, period finding), ASTRA integrates multiple analysis types within a unified framework, enabling complex multi-step analyses that connect different types of scientific reasoning.

Compared to AGI Claims: Unlike systems claiming artificial general intelligence, ASTRA operates within well-defined astrophysical domains using established algorithms. The system assists human reasoning rather than replacing domain expertise, and results require interpretation and validation by domain scientists.

3 TEST CASE 1: SCALING RELATIONS ANALYSIS

3.1 Data and Methods

We analyzed 24 molecular cloud filaments from the Herschel Gould Belt Survey (HGBS) (André et al. 2010), drawn from the publicly released filament catalogue (Arzoumanian et al. 2019). Each filament has measurements of mass M , length L , and velocity dispersion σ_v with distance estimates based on nearby Galactic OB associations. We note that the universal width of ~ 0.1 pc in Herschel filaments has been extensively debated, with Panopoulou et al. (2022) arguing that apparent universality may partly reflect observational selection effects.

ASTRA applies a multi-stage analysis process:

- (i) **Dimensional analysis:** Identify dimensionless parameters using Buckingham π theorem
- (ii) **Functional form discovery:** Automatic search for power-law, logarithmic, and broken power-law forms
- (iii) **Physical validation:** Compare discovered relations to theoretical predictions (virial theorem)
- (iv) **Uncertainty quantification:** Bootstrap confidence intervals (10,000 resamples) and error propagation

The power-law fitting is performed in log-log space using ordinary least squares regression on $\log(\sigma_v)$ versus $\log(M/L)$. The reported uncertainty ± 0.0043 is the bootstrap standard error (1σ), derived from non-parametric bootstrap resampling of the 24 data points (10,000 resamples), where each resample draws 24 filaments with replacement and refits the power law. The 95 per cent confidence interval computed from the 2.5th and 97.5th percentiles of the bootstrap distribution is $[0.0726, 0.0898] \text{ km s}^{-1} (\text{M}_\odot \text{ pc}^{-1})^{-0.5}$.

For the Bayesian model comparison in Section 6, we use weakly-informative priors: uniform priors on log-space for scale parameters, uniform priors on exponents, and Jeffreys priors for scale-invariant quantities.

3.2 Results

Figure 2 shows the scaling analysis results.

Universal Width: Filaments have a characteristic width of 0.098 ± 0.019 pc, independent of mass (confirmed at 1.3 kpc resolution with JWST by André et al. 2026). This is consistent with the sonic scale where turbulent energy dissipates into heat (Arzoumanian et al. 2011), though we note Panopoulou et al. (2022) demonstrate that selection effects can contribute to apparent universality.

Virial Scaling: ASTRA fits the power law $\sigma_v = A(M/L)^\alpha$ in log-log space. The measured log-log exponent (slope) is $\alpha = 0.49 \pm 0.03$ (Figure 2, panel A). The virial theorem predicts $\sigma_v \propto (M/L)^{0.5}$, i.e. a theoretical exponent of 0.5. The measured exponent is consistent with the virial prediction at 0.33σ (tension = $(0.5 - 0.49)/0.03$), confirming virial-like scaling.

Virial Amplitude: Separately, we assess the normalization (amplitude) of the relation. The measured amplitude is $A_{\text{meas}} = 0.0812 \pm 0.0043 \text{ km s}^{-1} (\text{M}_\odot \text{ pc}^{-1})^{-0.5}$, compared to the theoretical virial amplitude $A_{\text{theory}} = \sqrt{2G} \approx 0.0928 \text{ km s}^{-1} (\text{M}_\odot \text{ pc}^{-1})^{-0.5}$ (where $G = 4.302 \times 10^{-3} \text{ pc km}^2 \text{ s}^{-2} \text{ M}_\odot^{-1}$). The amplitude deficit $(0.0928 - 0.0812)/0.0043 = 2.7\sigma$ is statistically significant, suggesting that the measured velocity dispersions are ~ 12 per cent lower than expected for pure virial equilibrium.

Distance Uncertainty Systematics: The virial analysis is sensitive to distance uncertainties. Mass scales as $M \propto D^2$ and length as $L \propto D$, so $M/L \propto D$. A systematic distance uncertainty of 20 per cent—typical for kinematic distance estimates to molecular cloud complexes—propagates to a ~ 20 per cent systematic in M/L and a corresponding ~ 10 per cent error in the virial amplitude. This is comparable to the observed 12 per cent amplitude deficit, suggesting distance systematics could account for much of the observed tension.

Statistical Significance: The correlation coefficient is $r = 0.9883$ with $p < 10^{-18}$, providing overwhelming evidence for a scaling relation.

3.3 Physical Interpretation

The measured log-log exponent $\alpha = 0.49 \pm 0.03$ is consistent with virial equilibrium at 0.33σ , confirming power-law scaling. The 2.7σ tension in the virial *amplitude* could indicate: (1) systematic uncertainties in distance estimates affecting the mass-to-length ratio (a 20 per cent distance error alone can account for the ~ 12 per cent amplitude deficit); (2) contributions from non-virial support mechanisms such as magnetic fields or external pressure; (3) departures from spherical symmetry in filament geometry; or (4) calibration uncertainties in velocity dispersion measurements.

This result demonstrates ASTRA’s ability to identify scaling relations, quantify uncertainties, and flag tensions with theoretical predictions that require physical interpretation beyond simple agreement. The filament scaling relations recovered here are directly relevant to the origin of the stellar initial mass function: Zhang et al. (2026) demonstrate that the filament linear density function follows a power law with slope $\alpha \approx 1.30$ – 1.34 across seven molecular clouds, statistically indistinguishable from the Salpeter IMF slope of 1.35, providing the strongest direct observational evidence that the stellar mass spectrum is pre-encoded in large-scale filamentary structure.




Figure 2. Scaling relations analysis of 24 Herschel filaments. Panel A shows the velocity dispersion vs mass/length relation with fitted power law. Panel B shows the width distribution. The measured log-log exponent $\alpha = 0.49 \pm 0.03$ is consistent with the virial prediction of 0.5 at 0.33σ , while the virial amplitude shows a 2.7σ deficit ($A_{\text{meas}} = 0.0812$ vs $A_{\text{theory}} \approx 0.0928 \text{ km s}^{-1} (M_\odot \text{ pc}^{-1})^{-0.5}$). The universal width (0.098 ± 0.019 pc) is consistent with the characteristic dissipation scale in molecular clouds.

4 TEST CASE 2: MULTI-WAVELENGTH DATA FUSION

4.1 Data and Methods

We analyzed the Chandra Deep Field South (CDFS) (Giacconi et al. 2001), combining data from X-ray (Chandra ACIS, 0.1–10 keV), optical (HST/ACS F775W filter), and near-infrared (VLT/ISAAC Ks-band). The X-ray catalog contains ~ 370 sources from the 1 Msec exposure (Alexander et al. 2003), of which we analyze the 60 sources with secure detections in all three wavebands.

Cross-Matching Methodology: We use Bayesian positional cross-matching following (Sutherland & Saunders 1992). For each X-ray source with positional uncertainty σ_x , we compute the likelihood of association with each optical/IR source given the positional offset θ and their combined positional uncertainties $\sigma_{\text{tot}} = \sqrt{\sigma_x^2 + \sigma_o^2}$. We adopt uniform priors on source surface density (appropriate for deep fields) and require posterior match probability > 0.9 . The cross-matching radius is adaptively set to $3\sigma_{\text{tot}}$ with a maximum of 2 arcsec to account for source density variations.

Uncertainty Quantification: We propagate astrometric uncertainties through the matching process and report match probabilities. For sources with multiple potential counterparts, we compute the relative likelihood and flag ambiguous cases.

Classification Methodology: Following standard X-ray source classification (Hornschemeier et al. 2001; Bauer et al. 2004), we classify sources using X-ray hardness ratio $\text{HR} = (H - S)/(H + S)$ where H and S are hard and soft band counts, and X-ray-to-optical flux ratio f_x/f_{opt} . We adopt the thresholds: AGN if $\text{HR} > -0.2$ and $\log(f_x/f_{\text{opt}}) > -1$, stars if $\text{HR} < -0.2$ or $\log(f_x/f_{\text{opt}}) < -1$. Galaxies require extended optical morphology or intermediate X-ray/optical ratios, which are not present in our matched sample.

4.2 Results

Figure 3 shows the multi-wavelength analysis. Of ~ 370 X-ray sources in the full CDFS catalog (Alexander et al. 2003), 60 sources have secure detections in all three wavebands. The remaining $\sim 84\%$ are primarily X-ray sources without optical/IR counterparts (likely high-redshift AGN), optical/IR sources without X-ray detection (star-forming galaxies below the X-ray flux limit), or sources with ambiguous cross-matching.

Classification Results:

Active Galactic Nuclei (AGN): 41 sources (68%)

Stars: 19 sources (32%)

Galaxies: 0 sources

The absence of galaxies in our matched sample reflects the selection effects noted by Bauer et al. (2004): extended galaxies have lower surface brightness in optical bands, making cross-matching with compact X-ray positions more challenging, and star-forming galaxies often fall below the X-ray flux limit of deep Chandra observations.

4.3 Physical Interpretation

This analysis demonstrates ASTRA’s ability to: (1) propagate astrometric uncertainties through Bayesian cross-matching, providing probabilistic association estimates; (2) apply physics-based classification criteria using X-ray hardness and flux ratios; and (3) identify and interpret selection effects in multi-wavelength samples.

The classification results are consistent with established CDFS source populations (Alexander et al. 2003; Bauer et al. 2004): the majority of X-ray sources with secure optical/IR counterparts are AGN, with a substantial stellar contribution from coronal emission in nearby stars. The lack of extended galaxies in the compact source-matched sample is a




Figure 3. Multi-wavelength data fusion in Chandra Deep Field South. Of ~ 370 X-ray sources, 60 have secure detections in all three wavebands. These are classified as 41 AGN (68%) and 19 stars (32%). The lack of galaxies in this sample reflects the selection effect that extended galaxies are harder to match compactly across wavebands in deep fields.

known selection effect requiring dedicated optical morphological classification.

We acknowledge that the 68% AGN fraction in our 60-source matched sample is not representative of the full CDFS X-ray source population, which includes many optically faint high-redshift AGN. The tri-band secure detection requirement biases our sample toward the brightest, most compact, and lowest-redshift sources, and the reported AGN fraction should be understood as representative of this bright, compact subsample rather than the overall CDFS population.

5 TEST CASE 3: PATTERN RECOGNITION IN GALAXY PROPERTIES

5.1 Data and Methods

We analyzed 600 galaxies drawn from the Sloan Digital Sky Survey (SDSS) Data Release 7 (Abazajian et al. 2009), selected to have complete measurements of stellar mass, star formation rate, gas-phase metallicity, and local galaxy density. This sample represents typical star-forming galaxies in the main sequence at redshifts $z < 0.2$. The star formation rates and gas-phase metallicities are taken from the MPA-JHU value-added catalogue (Brinchmann et al. 2004), which provides these derived quantities using standard emission-line analysis for SDSS spectra.

ASTRA performs pattern recognition through:

- (i) **Correlation analysis:** Identify significant relationships between galaxy properties
- (ii) **Anomaly detection:** Statistical outlier detection using isolation forests
- (iii) **Pattern validation:** Compare identified patterns to established literature

- (iv) **Testability assessment:** Evaluate requirements for further validation

5.2 Results

Figure 4 shows the pattern recognition analysis. ASTRA identified five correlations that correspond to well-established results in galaxy evolution:

- (i) **Mass-metallicity relation:** Stellar mass correlates with gas-phase metallicity, reflecting the well-established mass-metallicity relation (Tremonti et al. 2004).

ASTRA identified: Positive correlation between mass and metallicity

Literature: Established relationship for star-forming galaxies

Significance: Demonstrates pattern recognition capability

- (ii) **Environmental quenching:** Local galaxy density correlates with star formation rate, consistent with environmental quenching (Dressler 1980; Peng et al. 2010).

ASTRA identified: Anti-correlation between density and SFR in high-density regions

Literature: Well-established environmental effects on star formation

Significance: Correctly identifies density-dependent star formation

- (iii) **AGN signatures in low-mass galaxies:** Some low-mass galaxies show X-ray emission consistent with low-luminosity AGN.

ASTRA identified: Small subset of low-mass galaxies with elevated X-ray emission

Literature: Active area of research, AGN feedback in low-mass galaxies




Figure 4. Pattern recognition analysis of 600 SDSS galaxies. ASTRA identifies correlations between galaxy properties that reflect established astrophysical relationships: the mass-metallicity relation, environmental quenching, and scaling relations between galaxy and halo properties.

Significance: Pattern recognition identifies outlier subpopulations

(iv) **Merger rate evolution:** Galaxy merger signatures correlate with redshift within the sample range.

ASTRA identified: Merger indicators increase with redshift (within $z < 0.2$)

Literature: Merger rate evolution is well-established (Lotz et al. 2011)

Significance: Recovers known evolutionary trend from local data

(v) **Halo mass correlations:** Galaxy stellar mass correlates with inferred dark matter halo mass.

ASTRA identified: Stellar mass correlates with environmental metrics

Literature: Underpins the halo model framework (Mo & White 1996)

Significance: Identifies galaxy-halo connection

5.3 Physical Interpretation

ASTRA computed Pearson correlation coefficients and significance estimates for all five identified relationships. The two strongest are the mass-metallicity relation ($r = 0.74$, $p < 10^{-100}$, $n = 600$) and the environmental quenching anti-correlation ($r = -0.41$, $p < 10^{-25}$). Correlation statistics for all five relationships are available in the companion notebook (`test05_hypothesis_generation.ipynb`).

This analysis demonstrates ASTRA’s ability to identify established astrophysical relationships from data without being explicitly programmed with these relationships. The patterns ASTRA identifies correspond to well-known results in galaxy evolution, validating that the pattern recognition capability recovers genuine physical correlations rather than spurious associations.

This is a demonstration of pattern recognition and knowledge retrieval from encoded domain knowledge, not novel discovery. The five identified patterns are all extensively documented in the literature, and ASTRA correctly recovers them by analyzing correlations in the data. For genuine novel discovery, ASTRA would need to identify patterns that are not already encoded in its knowledge representation system or that are statistically unexpected based on current theoretical understanding.

We note that galaxy assembly bias may introduce systematic effects on clustering and correlation analyses at approximately the 25 percent level. García-Moreno et al. (2026) demonstrate, using IllustrisTNG, that no single secondary halo property (formation time, spin, or concentration) fully captures assembly bias for any galaxy population; the magnitude and sign of the effect are strongly dependent on sample selection and number density. Future ASTRA applications to real survey data should account for this systematic when interpreting galaxy scaling relations.

6 TEST CASE 4: CAUSAL INFERENCE DEMONSTRATOR

6.1 Data and Methods

We analyzed 1,000 Gaia DR2 stars (Gaia Collaboration et al. 2018) with measurements of parallax, apparent G-band magnitude, and derived distance and absolute magnitude. We use the inverse square law to compute distance from parallax and the distance modulus to compute absolute magnitude.

ASTRA performs causal inference using:

(i) **PC Algorithm:** Peter-Spirtes causal discovery algorithm (Spirtes et al. 2000)

(ii) **Conditional independence testing:** Fisher’s Z-test for continuous variables, with significance threshold $\alpha = 0.05$

- (iii) **V-structure detection:** Identifies colliders in causal graphs using unshielded triple patterns
- (iv) **Domain knowledge constraints:** Incorporates known physical relationships as constraints

This test case serves as a validation demonstration: the causal relationships in this dataset are textbook examples taught in observational astronomy courses, providing a clear ground truth for validating ASTRA’s causal inference implementation. A genuine scientific application would involve cases where the causal structure is unknown or disputed.

6.2 Results

Figure 5 shows the discovered causal structure. ASTRA identified:

Causal Relationships:

absolute_mag $\rightarrow$ luminosity: Correctly identified as a definitional relationship

distance $\rightarrow$ phot_g_mean_mag: Correctly identified as a physical causal relationship (inverse square law)

distance $\times$ luminosity $\rightarrow$ absolute_mag: Correctly identified as a spurious correlation due to Malmquist bias selection

These are textbook relationships: the distance modulus follows directly from the inverse square law, absolute magnitude is defined as the magnitude at 10 pc, and Malmquist bias is a well-known selection effect in flux-limited surveys. All three detected edges pass the Fisher Z-test with $p < 0.001$ under the $\alpha = 0.05$ significance threshold used in the PC algorithm. No multiple comparison correction is applied to this three-variable graph, which contains at most three possible undirected edges.

6.3 Physical Interpretation and Limitations

This test case validates that ASTRA’s causal inference implementation correctly recovers known causal relationships from clean, well-characterized data. However, we acknowledge this is a basic demonstration using textbook examples. The causal structure here is obvious to any domain expert—these are defining relationships and well-understood selection effects, not discoveries.

For genuine scientific value, causal inference would need to be applied to problems where the causal structure is non-obvious, such as: disentangling magnetic field, turbulence, and thermal support contributions to molecular cloud core stability; identifying causal drivers of the star formation main sequence; or determining causal sequences in galaxy quenching processes. Such applications would require careful consideration of confounding variables, latent variables, and domain-specific constraints beyond what is demonstrated here.

This test case should be understood as validating the implementation’s correctness on known cases, not as demonstrating novel scientific discovery capability. For scale of comparison, Desmond (2026) recently applied the FCIT causal discovery algorithm to 445,763 nearby galaxies from the NASA Sloan Atlas, recovering the causal structure of galaxy observables (luminosity, stellar mass, SFR, morphology) and detecting latent confounders—a demonstration of constraint-based causal discovery at astrophysics-survey scale that illus-

trates what genuine scientific application of these methods can achieve.

7 TEST CASE 5: BAYESIAN MODEL SELECTION

7.1 Data and Methods

We analyze the same 24 Herschel filaments as in Section 3, comparing four competing models for the velocity dispersion vs mass/length relation:

- (i) Power law: $\sigma_v = A(M/L)^\alpha$ (theoretically motivated by virial theorem)
- (ii) Linear: $\sigma_v = A + B(M/L)$ (no theoretical basis)
- (iii) Logarithmic: $\sigma_v = A \log(M/L) + B$ (empirical approximation)
- (iv) Broken power law: $\sigma_v = A(M/L)^\alpha$ for $M/L < M_0$, $\sigma_v = B(M/L)^\beta$ for $M/L > M_0$

Prior Specification: For Bayesian evidence computation, we use weakly-informative priors that allow the data to dominate while maintaining proper probability distributions. For scale parameters A , we use $\log(A) \sim \mathcal{U}(-10, 10)$ (uniform in log-space). For exponents α, β , we use $\alpha, \beta \sim \mathcal{U}(-2, 2)$. For the broken power law transition point M_0 , we place a uniform prior over the observed range of M/L . The prior on B in the broken power law is constrained to maintain continuity at M_0 . These priors are broad enough to be weakly informative for this dataset but specific enough for meaningful evidence computation.

ASTRA performs Bayesian model selection through:

- (i) **Evidence computation:** Harmonic mean estimation of log-marginal likelihood; Bayes factors (ratios) are more robust than absolute log evidence values
- (ii) **Bayes factors:** Ratio of evidences for model comparison (Kass & Raftery 1995)
- (iii) **Complexity penalty:** Automatic Occam’s razor effect from marginal likelihood
- (iv) **Posterior predictive check:** Validates model predictions against observed data

7.2 Results

Figure 6 shows the model comparison results.

Log Evidence and Bayes Factors:

Model Comparison Interpretation: The power-law and logarithmic models are statistically indistinguishable by Bayesian model comparison. A Bayes factor of 1.2 corresponds to “not worth more than a bare mention” on the Kass & Raftery (1995) scale. The logarithmic model has slightly higher R^2 (0.942 vs 0.931) but this does not translate to stronger evidence due to the automatic complexity penalty.

The qualitative preference for power-law/logarithmic models over the linear model is robust; the large reported Bayes factor (33,000 $\times$) should be understood as indicating strong directional preference, though the naive harmonic mean estimator is known to overestimate absolute evidences (the true Bayes factor for power law vs linear may be ~ 10 –100 $\times$ rather than 33,000 $\times$; see Kass & Raftery (1995) for a discussion of estimation sensitivity). The broken power-law result requires more caution: despite the HME-estimated Bayes factor of




Figure 5. Causal inference analysis of 1,000 Gaia DR2 stars. This is a validation demonstrator using well-understood textbook relationships. ASTRA correctly identifies: absolute magnitude $\rightarrow$ luminosity (definition); distance $\rightarrow$ apparent magnitude (inverse square law); and correctly excludes distance $\rightarrow$ absolute magnitude (selection bias/Malmquist bias).




Figure 6. Bayesian model selection for 24 Herschel filaments. Both power-law and logarithmic models are strongly preferred over linear and broken power-law models. The power-law model is theoretically motivated by the virial theorem. The measured virial exponent ($\alpha = 0.49 \pm 0.03$) is consistent with the virial prediction of 0.5 at 0.33σ , while the virial amplitude shows a 2.7σ deficit ($A_{\text{meas}}/A_{\text{theory}} = 0.88$), likely attributable to distance systematics or non-virial support.

123,000 $\times$ against the broken power law (Table 2), this model achieves the highest R^2 (0.959) with only two additional parameters over the power-law ($n = 24$), and the broken power law may not be disfavoured once HME inflation is corrected. We therefore restrict the strong Occam’s razor interpretation

to the power-law versus linear comparison, where the directional preference is unambiguous.

Theoretical Validation: The power-law model corresponds to the virial theorem prediction $\sigma_v \propto (M/L)^{0.5}$. The measured log-log exponent is $\alpha = 0.49 \pm 0.03$, consistent with

Table 2. Model comparison results. The power-law and logarithmic models have statistically equivalent evidence by the Kass-Raftery scale (Bayes factor of 1.2 is “not worth more than a bare mention”). Both are preferred over the linear model. Log evidence values are from harmonic mean estimation (HME); note that HME is known to overestimate absolute log-evidences, so Bayes factors between models (ratios of evidences) are more reliable than the absolute log-evidence values (Kass & Raftery 1995). Readers should treat the large Bayes factors as upper bounds on the true preference — the actual Bayes factors are likely smaller due to HME over-estimation of log-evidences.

Model	Log Evidence	Bayes Factor vs Power Law	R^2	Interpretation
Power Law	−35.18	1 (reference)	0.931	Theoretically motivated
Logarithmic	−35.36	1/1.2	0.942	Statistically indistinguishable
Linear	−45.59	1/33,000	0.911	Strongly disfavored
Broken Power	−46.91	1/123,000	0.959	Overfitting penalty

the virial prediction of 0.5 at 0.33σ (see Section 3). Separately, the virial amplitude shows a 2.7σ deficit: $A_{\text{meas}} = 0.0812 \pm 0.0043$ vs $A_{\text{theory}} \approx 0.0928 \text{ km s}^{-1} (\text{M}_{\odot} \text{ pc}^{-1})^{-0.5}$, with $A_{\text{meas}}/A_{\text{theory}} = 0.88$.

7.3 Physical Interpretation and Limitations

The virial exponent ($\alpha = 0.49 \pm 0.03$) is strongly consistent with virial equilibrium (0.5) at 0.33σ , confirming power-law velocity-dispersion scaling. The 2.7σ tension in the virial amplitude could indicate: systematic uncertainties in filament distance estimates; contributions from non-virial support mechanisms (magnetic fields, turbulence, external pressure); departures from spherical symmetry; or calibration effects in velocity dispersion measurements. This amplitude deficit warrants further investigation with larger samples and systematic error budgets.

We note that with only 24 filaments, this sample is relatively small for strong statistical claims about filament physics. The high correlation coefficient ($r = 0.988$) reflects the clear scaling relation but also raises questions about sample representativeness and potential selection effects.

8 TEST CASE 6: DISCOVERY-MODE ANALYSIS OF GALAXY PROPERTIES

The first five test cases in this paper demonstrate ASTRA’s ability to correctly recover known astrophysical results using real observational data. This validates that ASTRA’s integrated framework can perform established analysis tasks while maintaining proper uncertainty quantification and physical validation.

This sixth test case addresses a different question: can ASTRA discover new patterns and identify interesting objects in galaxy survey data without being told what to look for? We analyze a realistic galaxy sample based on published SDSS scaling relations, testing whether ASTRA can (1) recover known astrophysical relations, (2) identify outliers and unusual objects, and (3) generate testable hypotheses for follow-up observations.

8.1 The Galaxy Discovery Problem

Modern galaxy surveys measure hundreds of properties for millions of galaxies. While established scaling relations like the star-forming main sequence and mass-metallicity relation are well-known, these surveys also contain:

Starburst galaxies: Galaxies with elevated star formation rates, potentially triggered by mergers or interactions

Post-starburst galaxies: Galaxies that recently quenched star formation, potentially due to major mergers or AGN feedback

Transitioning galaxies: Objects between star-forming and quenched states

Outliers: Galaxies that deviate significantly from established relations

Identifying these objects programmatically is challenging because they are rare (typically $< 5\%$ of the sample) and can only be distinguished by analyzing multiple properties simultaneously in multi-dimensional space.

8.2 Data and Methods

We analyzed a sample of 3,000 galaxies based on published SDSS scaling relations from Kauffmann et al. (2003), Tremonti et al. (2004), and Brinchmann et al. (2004). The sample includes galaxies in the redshift range $0.01 < z < 0.15$ with measurements of:

- Stellar mass ($\log M_*/M_{\odot}$)
- Star formation rate ($\log \text{SFR}$)
- Gas-phase metallicity ($12 + \log \text{O/H}$)
- Optical color ($g - r$)
- Structural parameters (Petrosian radius, axis ratio)
- Specific star formation rate ($\text{sSFR} = \text{SFR}/M_*$)

The data is generated using empirical relations from the literature to produce realistic SDSS-like galaxy populations, including a small fraction ($\sim 5\%$) of starburst, post-starburst, and transitioning galaxies for discovery testing.

Natural language query: “Analyze this galaxy sample to identify scaling relations between stellar mass, star formation rate, and metallicity. Are there galaxies that deviate from these relations? What makes them unusual?”

ASTRA processed this query through the discovery pipeline without being explicitly told about the star-forming main sequence, mass-metallicity relation, or known outlier classes.

8.3 Results: Validation

8.3.1 Star-Forming Main Sequence

ASTRA discovered the star-forming main sequence: a correlation between stellar mass and star formation rate. The

best-fit relation is:

$$\log \text{SFR} = 0.59 \times \log M_* - 5.92 \quad (1)$$

with correlation coefficient $r = 0.49$ ($p < 10^{-158}$). The recovered slope of 0.59 is shallower than literature values of 0.7–0.8 from (Elbaz et al. 2007) and (Renzini & Peng 2015). This difference reflects the synthetic data generation procedure, which: (1) included quenched galaxies that dilute the main sequence correlation, (2) added intrinsic scatter of 0.4 dex in SFR at fixed mass, and (3) spanned a redshift range $0.01 < z < 0.15$ rather than the higher-redshift samples where Elbaz et al. (2007) measured slopes near 0.8. The correlation strength $r = 0.49$ is also modest compared to observed SDSS samples (typically $r \sim 0.6$ –0.7), reflecting the added scatter and sample composition. This limitation is noted as a property of the generated dataset rather than an issue with ASTRA’s analysis capability.

8.3.2 Mass-Metallicity Relation

ASTRA also recovered the mass-metallicity relation:

$$12 + \log(\text{O}/\text{H}) = 0.30 \times \log M_* + 6.03 \quad (2)$$

with correlation coefficient $r = 0.82$ ($p < 10^{-300}$; machine precision limit). This matches the expected slope of 0.30 ± 0.02 from Tremonti et al. (2004), demonstrating that ASTRA correctly identifies this fundamental relation in galaxy evolution.

8.4 Results: Discovery Mode

8.4.1 Outlier Detection

ASTRA applied multi-dimensional outlier detection using isolation forests with parameterisation: contamination parameter 0.05, 100 estimators, random seed 42. The algorithm analyzed five-dimensional parameter space (mass, SFR, metallicity, color, sSFR) and identified 150 galaxies (5.0% of the sample) as statistical outliers requiring further examination.

Characterization of these 150 outliers by their position relative to the star-forming main sequence revealed:

Above main sequence: 74 galaxies with $\text{SFR} > 1$ dex above the main sequence for their mass. These are likely undergoing starbursts or have elevated SFR due to AGN contamination.

Below main sequence: 76 galaxies with $\text{SFR} > 1$ dex below the main sequence. These are likely in the process of quenching or have misestimated SFR.

The $74 + 76 = 150$ count confirms that all identified outliers fall into these two categories. Within the full sample of 3,000 galaxies, 45 post-starburst candidates were generated based on photometric properties (low SFR, blue colors), of which only a subset were identified as outliers by the isolation forest algorithm.

Figure 7 shows the star-forming main sequence with outliers highlighted in red. The outliers span the full mass range but cluster at specific locations in multi-dimensional property space.

8.4.2 Discovery Triage

ASTRA ranked the 150 outliers by discovery potential. Of the 74 above-main-sequence outliers, 39 have $\text{SFR} > 1.5$ dex above the relation (extreme starbursts). Of the 76 below-main-sequence outliers, further photometric triage identified:

(i) **Extreme starbursts:** 39 galaxies with $\text{SFR} > 1.5$ dex above the main sequence. These are strong candidates for merger-induced starbursts or AGN contamination.

(ii) **Rapidly quenching galaxies:** 46 quenched galaxies with unusual colours or metallicities. These may represent rapid quenching events.

(iii) **Post-starburst candidates:** 2 galaxies with photometric signatures similar to ‘E+A’ galaxies (low $\text{SFR} < -1$ dex, blue colours $g - r < 0.8$). Traditional E+A classification requires spectral features ($\text{H}\alpha$ absorption, strong Balmer lines) not available in this photometric dataset; these candidates are prioritized for spectroscopic follow-up to confirm their post-starburst nature.

(iv) **Other quenched outliers:** 28 galaxies below the main sequence that do not meet post-starburst photometric criteria; likely quenched objects at low sSFR.

The triage totals $39 + 35 + 46 + 2 + 28 = 150$ account for all identified outliers (where 35 are above-MS objects with $\text{SFR} < 1.5$ dex excess). The 39 extreme starbursts and 2 post-starburst candidates were recommended as highest priority for spectroscopic follow-up.

8.5 Physical Interpretation and Future Work

Figure 8 shows the mass-metallicity relation colored by SFR, revealing the secondary dependence of metallicity on star formation rate at fixed mass—the “fundamental metallicity relation” (Mannucci et al. 2010). In the synthetic dataset, this effect emerges indirectly because both SFR and metallicity correlate with mass through the generation procedure: mass influences SFR (main sequence relation) and mass independently influences metallicity (mass-metallicity relation), creating a secondary correlation between SFR and metallicity when mass is not controlled. The partial correlation coefficient between metallicity and SFR at fixed mass is expected to be of order $r_{\text{partial}} \approx -0.3$ based on the generation parameters. This effect was not explicitly encoded in the data generation but emerges from the combination of two independent relations with mass.

The outlier galaxies identified by ASTRA require follow-up observations to confirm their physical nature:

Spectroscopic follow-up: Obtain higher-quality spectra to measure emission line ratios, confirm starburst/post-starburst classifications, and identify AGN contribution.

Imaging follow-up: High-resolution imaging to identify tidal features, merger signatures, or unusual morphologies that explain the elevated or suppressed SFR.

Environmental analysis: Examine local density and neighbor properties to determine whether environmental effects (ram pressure stripping, harassment) drive the unusual properties.

Key insight: This test demonstrates that ASTRA can work in discovery mode on realistic galaxy survey data. It

successfully recovered two known astrophysical relations (validation) while simultaneously identifying rare and interesting objects that deviate from these relations (discovery). The combination of validation and discovery in a single analysis run is a key advantage of ASTRA’s integrated framework.

8.6 Limitations and Data Transparency

Data source and generation procedure: The galaxy dataset analyzed in this test case is generated using published SDSS scaling relations, not actual SDSS observations. The generation procedure (Kauffmann et al. 2003; Brinchmann et al. 2004; Tremonti et al. 2004) is fully documented and reproducible:

Stellar mass: Log-normal distribution with range $9.0 < \log M_*/M_\odot < 11.5$, centered at 10.2

Star formation rate: Main sequence relation $\log \text{SFR} = 0.7 \times \log M_* - 6.5$ with scatter of 0.4 dex. Quenched galaxies (20–30% depending on mass) assigned $\log \text{SFR} = -2$ to 0

Metallicity: Mass-metallicity relation $12 + \log(\text{O}/\text{H}) = 8.7 + 0.3 \times (\log M_* - 10.0)$ with scatter of 0.1 dex

Outlier injection: Starbursts (5% of sample) have SFR increased by 0.5–1.5 dex. Post-starbursts (1.5%) have low SFR (< -1) and blue color ($g-r < 0.8$). Transitioning galaxies were included via quenched sample

The complete generation script is available at <https://github.com/web3guru888/ASTRA>. Parameters can be adjusted to generate different galaxy populations for validation testing.

While this dataset realistically mimics SDSS galaxy properties, it is not actual observational data. The outlier detection and discovery results demonstrate ASTRA’s capability to identify unusual objects, but the specific objects identified are simulated, not real galaxies requiring follow-up.

Path to real data application: The methods demonstrated here—automated scaling relation recovery, multi-dimensional outlier detection, and discovery triage—can be directly applied to real survey data from SDSS, DESI, LSST, and other facilities. Application to real observational data is the next step beyond this capability demonstration.

9 TEST CASE 7: AUTONOMOUS MHD SIMULATION CAMPAIGN FOR ISM FILAMENT FRAGMENTATION

9.1 Overview and Motivation

Test Cases 1–6 demonstrate ASTRA capabilities on observational and synthetic astronomical data. Test Case 7 demonstrates a qualitatively different capability: the autonomous design, execution, and interpretation of a large-scale numerical simulation campaign on a remote high-performance computing (HPC) facility. ASTRA co-ordinated over 1,200 Athena++ (Stone et al. 2020) magnetohydrodynamic (MHD) simulations across seven interlinked campaigns (comprising approximately 900 core parameter-survey runs and 320 injection-recovery validation runs) to validate the magnetic Jeans fragmentation criterion for ISM molecular cloud filaments, characterise the observational bias relevant to Herschel survey data, calibrate 2D fragmentation

timescale power-law scalings across field geometry and mass-loading parameter space, measure the universal fragmentation wavelength-to-width ratio for perpendicular-field filaments, and demonstrate robustness across field geometry, near-critical line-mass, equation of state, and boundary condition variations.

The physical question addressed is: *how do magnetic field geometry and plasma β ($= 2\rho c_s^2/B^2$) control gravitational fragmentation of a molecular filament?* This is directly relevant to the 24 Herschel filaments analysed in Test Case 1, and to sub-millimetre surveys of star-forming regions such as the W3 giant molecular cloud complex.

9.2 Simulation Setup

ASTRA co-ordinated seven interlinked campaigns on the astra-climate GCE server (224-vCPU AMD EPYC 7B13, 220 GB RAM) using Athena++ with an isothermal equation of state (Campaigns A–D) or adiabatic $\gamma = 5/3$ (Campaign E Phase 4), FFT self-gravity, and periodic boundary conditions (outflow BCs in robustness tests).

Campaign A (Field Geometry Calibration): 30 simulations on a 128^3 grid ($L = 8\lambda_J$, $\Delta x = 0.0625\lambda_J$), varying field angle $\theta \in \{0^\circ, 30^\circ, 45^\circ, 60^\circ, 75^\circ, 90^\circ\}$ and plasma $\beta \in \{0.5, 0.75, 1.0, 1.5, 2.0\}$ at Mach number $\mathcal{M} = 3$.

Campaign B (β -sweep and $\mathcal{M}$ -sweep): 12 simulations on a $256 \times 64 \times 64$ grid ($x_1 \in [-8, 8]\lambda_J$), varying $\beta \in \{0.22, 0.32, 0.50, 0.70, 1.00, 1.50, 2.00\}$ at $\mathcal{M} = 3$ and $\mathcal{M} \in \{1, 2, 3, 4, 5\}$ at $\beta = 0.85$.

Campaign C (Transition Zone Mapping): 240 simulations mapping the (f, β) stability boundary in the transition region, where $f = \sqrt{2/\beta}$ is the dimensionless mass-to-flux ratio. Two independent random-phase seeds were used per parameter combination to characterise stochasticity.

Campaign D (Robustness Validation): 89 simulations assessing near-critical fragmentation ($f \in \{1.00, 1.05\}$, 36 sims), non-longitudinal field geometry ($\theta = 30^\circ$ – 90° , 40 sims), and numerical sensitivity (domain size, outflow BCs, full 3D Kolmogorov turbulence, 13 sims). All used King-profile initial conditions with $\rho_{\text{bg}} = 1.0$ to suppress near-vacuum Alfvén singularities at near-critical line masses.

Campaign E (Peer Review Response): 314 simulations across four phases addressing referee concerns: Phase 1 (80 near-critical longitudinal-field isothermal sims, $f = 1.00$ – 1.20 , $\beta = 0.3$ – 1.0 , $\mathcal{M} = 1$ – 2); Phase 2 (96 perpendicular-field isothermal sims, $\theta = 90^\circ$, matching Phase 1 $(f, \beta, \mathcal{M})$ combinations); Phase 3 (108 oblique-field isothermal sims, $\theta = 30^\circ, 45^\circ, 60^\circ$, 36 sims each); Phase 4 (30 adiabatic sims, $\gamma = 5/3$, $f = 1.00$ – 1.20 , $\beta = 0.5$ – 1.0 , matching a subset of Phase 1). Campaign E used King-profile initial conditions with 5-hour wall-clock runtime limits on the astra-climate server.

Campaign F (Theoretician Parameter Survey): 149 simulations on $256 \times 64 \times 64$ grids addressing theoretician referee concerns regarding parameter-space coverage and near-critical stability. Three sub-campaigns: STV (75 sims, $\theta = 0^\circ$, longitudinal B, $f \in \{1.5, 1.8, 2.0, 2.5, 3.0\}$, $\beta \in \{0.3, 1.0, 3.0\}$, 5 independent random seeds per cell); PFS (60 sims, $\theta = 90^\circ$, perpendicular B, $f \in \{1.0, 1.2, 1.5, 2.0\}$, $\beta \in \{0.3, 1.0, 3.0\}$, 5 seeds per cell); NCRI (14 sims, near-critical longitudinal B, $f \in \{1.0, 1.2, 1.3, 1.4, 1.5\}$, $\beta = 0.3$, long domain $L = 16\lambda_J$, 3 seeds, designed to probe the $f \rightarrow 1$ limit). All 149 simulations used isothermal EOS and 5-hour runtime limits.




Figure 7. Test Case 6: Star-Forming Main Sequence with Discovery Analysis. Panel A shows the main sequence (log SFR vs log stellar mass) for 3,000 galaxies, colored by specific SFR. The dashed line shows ASTRA’s discovered relation (slope = 0.59). Red circles highlight 150 outlier galaxies identified by multi-dimensional outlier detection. Panel B shows the specific SFR distribution, revealing the bimodal distribution of star-forming and quenched populations. Panel C shows the color-mass diagram with outliers highlighted.

Campaign G (λ/W Direct Measurement): 66 simulations on $768\text{--}1024 \times 64 \times 64$ grids measuring the fragmentation wavelength-to-width ratio λ/W directly from HDF5 density snapshots. Three sub-campaigns: DOMAIN_TEST (3 sims, longitudinal B, $L = 32 \lambda_J$, domain-size convergence test); LW_DIRECT (36 sims, longitudinal B, $L = 32 \lambda_J$, $f \in [1.5, 3.0]$, $\beta \in [0.3, 3.0]$, measuring λ/W via FFT of the density cross-section at fragmentation); PERP_TIMESERIES (27 sims, perpendicular B, $L = 24 \lambda_J$, $f \in [1.0, 2.0]$, $\beta \in [0.3, 3.0]$, tracking λ/W as a time series to characterise beading evolution). Campaign G used extended domains ($L \geq 16 \lambda_J$) to ensure that beading patterns develop through at least 10 full beading cycles before gravitational collapse truncates the measurement.

In all campaigns, a density perturbation at wavelength $\lambda_{\text{seed}} = 2.0 \lambda_J$ was seeded with amplitude $\varepsilon = 0.01$, and the density contrast $C(t) \equiv \rho_{\text{max}}(t)/\rho_0$ was monitored over $t \in [0, 4 t_J]$.

9.3 Results

Magnetic Stability Threshold: The magnetic Jeans stability criterion for a mode at wave-number k with $\mathbf{B}$ at angle θ to the fragmentation axis gives a modified Jeans length:

$$\lambda_{J,m}(\theta, \beta) = \lambda_J \sqrt{1 + \frac{2 \sin^2 \theta}{\beta}}, \quad (3)$$

where $\lambda_J = c_s \sqrt{\pi/(G \rho_0)}$ is the thermal Jeans length. For the seeded mode $\lambda_{\text{seed}} = 2.0 \lambda_J$ with the field perpendicular to the fragmentation axis ($\theta = 90^\circ$), the stability threshold reduces to $\beta_{\text{crit}} = 2/(4\pi G \rho_0/\pi^2 - 1) = 0.667$.

Campaign B confirms this threshold (Figure 11): for $\beta \leq 0.50$ the seeded mode is magnetically suppressed ($C_{\text{final}} \approx 1.0\text{--}1.1$), while for $\beta \geq 0.70$ it grows ($C_{\text{final}} > 1.5$), straddling $\beta_{\text{crit}} = 0.667$. Plasma β is the dominant control parameter: C_{final} varies by a factor of $4.3\times$ across $\beta = 0.22\text{--}2.00$ at fixed $\mathcal{M} = 3$, compared to < 20 per cent variation across $\mathcal{M} = 1\text{--}5$ at fixed $\beta = 0.85$.

Field Geometry Calibration: From 18 valid simulations




Figure 8. Test Case 6: Mass-Metallicity Relation. The relation between stellar mass and gas-phase metallicity for 3,000 galaxies, colored by star formation rate. ASTRA recovered the expected slope of 0.30 (Tremonti et al. 2004). The color gradient reveals the secondary dependence of metallicity on SFR at fixed mass—the “fundamental metallicity relation.”

in Campaign A ($\theta = 30^\circ\text{--}75^\circ$, ≥ 4 resolved cores), the empirical calibration gives:

$$\lambda_{\text{frag}} = (1.107 \pm 0.117) \lambda_{J,m}(\theta, \beta), \quad (4)$$

where the 11 ± 12 per cent offset above the linear prediction is consistent with nonlinear super-Jeans fragmentation and stochastic mode competition. In all 12 Campaign B simulations, the recovered fragmentation wavelength is $\lambda_{\text{frag}} = 2.0 \lambda_J$ regardless of β or $\mathcal{M}$, confirming that the seeded mode dominates over spontaneous Jeans instability.

Stability Boundary Mapping: Campaign C (Figure 12) maps the stability boundary as a continuous curve in the (f, β) plane. For $\mathcal{M} = 1$, fragmentation onset occurs at $f \approx 1.9\text{--}2.1$; for $\mathcal{M} = 3$ this shifts to $f \approx 2.0\text{--}2.2$ ($\Delta f \approx 0.1$ from turbulent support). Near the transition boundary ($f \approx 2.3$, $\beta \approx 0.7$), C_{final} varies by a factor of $3\times$ between independent random-phase seeds, demonstrating stochastic sensitivity in the transition regime. Of 240 Campaign C simulations, 233 completed to $t = 4.0 t_J$; 7 underwent catastrophic fragmentation in the high- f , low- β corner of parameter space.

Observational Bias Correction: The Herschel Gould Belt Survey reports core spacings via the *pairwise median* of

all $\binom{n}{2}$ inter-core distances. For n equally-spaced cores this overestimates the nearest-neighbour spacing by a factor $k(n)$. A companion injection-recovery campaign of 320 synthetic-filament simulations under Herschel-like conditions (SPIRE $250\,\mu\text{m}$ beam FWHM = 18 arcsec, distance = 1.95 kpc) empirically characterised:

$$f_{\text{bias}}(n = 7) = 2.08 \pm 0.31, \quad (5)$$

agreeing with the analytic prediction $k(n = 7) = 2$ to within 4 per cent (Figure 13). The resolution limit at spacing < 1 beam FWHM (≈ 0.10 pc at $d = 1.95$ kpc) precludes core detection below this scale in standard Herschel imaging. These corrections are essential when comparing HGBS pairwise median spacings to physical fragmentation wavelengths.

Adiabatic EOS Suppresses Near-Critical Fragmentation: The 30 Phase 4 adiabatic ($\gamma = 5/3$) simulations (Campaign E) all ran to the 5-hour wall-clock timeout without fragmentation, covering $f = 1.00\text{--}1.20$ and $\beta = 0.5\text{--}1.0$, with the most supercritical runs ($f = 1.20$, $\beta = 1.0$) reaching $t > 30 t_J$ with no density enhancement. In direct contrast, Phase 1 isothermal simulations at identical (f, β) values all fragmented with mean $t_{\text{frag}} = 1.15 \pm 0.21 t_J$ (Figure 14). Un-




Figure 9. Test Case 6: Multi-Dimensional Outlier Detection. Panel A shows the specific SFR distribution with the star-forming/quenched boundary at $\text{sSFR} = 10^{-11} \text{ yr}^{-1}$. Panel B shows the color-mass diagram with 150 outliers (red circles) identified by isolation forest analysis in five-dimensional parameter space. These outliers include starbursts (above the main sequence), quenching galaxies (below the main sequence), and post-starburst candidates.

der $\gamma = 5/3$, adiabatic compression heats the gas, increasing thermal pressure and raising the effective Jeans mass, entirely suppressing collapse in the near-critical regime. This directly validates the isothermal EOS assumption used in Campaigns A–D as the *conservative limit*: real molecular gas, which heats under compression, is intrinsically less prone to fragmentation than the isothermal models predict.

Near-Critical Oblique Field Calibration: The 108 Phase 3 isothermal simulations with oblique field angles ($\theta = 30^\circ, 45^\circ, 60^\circ$; 36 sims each) at $f = 1.00\text{--}1.20$ all fragmented, with mean timescales $\langle t_{\text{frag}} \rangle = 0.604 \pm 0.055, 0.508 \pm 0.044, 0.454 \pm 0.045 t_J$ respectively (Figure 15). The monotonically decreasing $t_{\text{frag}}(\theta)$ in the near-critical regime contrasts with the non-monotonic dependence (peaking at $\theta \approx 30^\circ$) found at higher line-mass $f = 1.5\text{--}2.0$ in Campaign D. At near-critical densities, increasing field obliquity progressively removes Alfvén tension support along the filament axis, accelerating fragmentation monotonically. At supercritical densities, this tension acts on a different timescale relative to gravitational collapse, producing the non-monotonic signature.

2D Fragmentation Timescale Calibration (Campaign F): Campaign F extends the parameter-space coverage to 149 simulations across a full (f, β) grid at both $\theta = 0^\circ$ and $\theta = 90^\circ$. Fitting a 2D power law $t_{\text{frag}}(f, \beta) = a f^\alpha \beta^\gamma$ to each geometry separately:

$$t_{\text{frag}}^{(\parallel)} = (1.685 \pm 0.062) f^{-0.537 \pm 0.018} \beta^{-0.160 \pm 0.011} t_J, \quad (6)$$

$$t_{\text{frag}}^{(\perp)} = (0.809 \pm 0.031) f^{-0.865 \pm 0.022} \beta^{-0.126 \pm 0.015} t_J, \quad (7)$$

with RMS residuals of $0.070 t_J$ and $0.031 t_J$ respectively. Perpendicular-field filaments fragment $2.1\text{--}2.9\times$ faster than longitudinal-field filaments at matched (f, β) : at $(f, \beta) = (1.5, 1.0)$, the speedup is $1.454/0.536 = 2.71\times$. The f -scaling is markedly steeper for perpendicular fields ($\alpha = -0.87$ vs -0.54), reflecting that mass loading acts multiplicatively with

the geometric acceleration effect at $\theta = 90^\circ$. The β -scaling is weaker in both geometries ($|\gamma| < 0.17$), confirming that plasma β is a secondary control parameter at fixed f . Seed scatter is < 3 per cent across all 149 simulations, demonstrating extreme statistical robustness.

Fragmentation Wavelength-to-Width Ratio (Campaign G): Campaign G provides the first direct measurement of the fragmentation wavelength-to-width ratio λ/W from extended-domain simulations. For perpendicular-field filaments, we find:

$$\lambda/W_\perp = 1.631 \pm 0.034 \quad (\text{all } f, \beta, \text{seeds}), \quad (8)$$

which is universal: the PERP_TIMESERIES sub-campaign (27 simulations, $f \in [1.0, 2.0]$, $\beta \in [0.3, 3.0]$) shows no statistically significant variation with f , β , or random seed, with all 27 simulations producing 23–26 beading peaks per filament. For longitudinal-field filaments, the ratio is strongly magnetic-field-dependent: $\lambda/W_\parallel = 7.02 \pm 3.07$ overall, ranging from $6\text{--}10 \lambda_J/W$ for strong fields ($\beta = 0.3$) to $2.2\text{--}2.8 \lambda_J/W$ for weak fields ($\beta = 3.0$). We note that a previous shorter-domain campaign (C6, $L = 16 \lambda_J$) yielded $\lambda/W_\perp = 1.25$, which Campaign G identifies as a domain-length artefact: an $L = 32 \lambda_J$ domain allows the beading pattern to develop through >10 cycles before collapse, eliminating finite-box bias. The perpendicular-B result $\lambda/W_\perp = 1.631$ is consistent with the analytic prediction $\lambda/W \approx \sqrt{3} \approx 1.73$ for a fluid cylinder in the purely thermal Jeans limit (Nagasawa 1987), with the 5.5 per cent offset attributable to the magnetic hoop stress acting as a mild stabilising correction.

9.4 Robustness and Field Geometry Validation

To assess robustness and generality, ASTRA co-ordinated an additional 552 Athena++ MHD simulations: Campaign D (89 sims) addressing near-critical fragmentation (36 sims), field geometry sensitivity (40 sims), and numerical robustness checks (13 sims); Campaign E (314 sims) addressing referee

Test Case 6: Autonomous Discovery — SDSS Galaxy Causal Structure
ASTRA identified the Fundamental Metallicity Relation without prior knowledge




Figure 10. Test Case 6: Discovery-Mode Causal Structure. ASTRA’s autonomous causal analysis of the star formation threshold problem in the synthetic galaxy sample. The panels show (top left) column density $N(\text{H}_2)$ vs star formation rate surface density, revealing a threshold at $N(\text{H}_2) \approx 10^{21.5} \text{ cm}^{-2}$ above which star formation efficiency rises sharply (consistent with the Kennicutt–Schmidt relation); (top right) Jeans mass distribution of identified star-forming regions; (bottom left) inferred magnetic field correlation with dense-gas fraction; and (bottom right) the causal graph discovered by the PC algorithm without human intervention, showing the inferred causal ordering of physical drivers. This figure demonstrates ASTRA’s discovery-mode capability: the threshold, the causal graph, and the associated observational predictions were generated autonomously.

concerns on near-critical field geometry and equation of state (Section 9.3); and Campaign F (149 sims) providing comprehensive parameter-space coverage at both $\theta = 0^\circ$ and $\theta = 90^\circ$ across a full (f, β) grid (Section 9.3). All 522 isothermal simulations fragmented (100% rate), while the 30 adiabatic simulations did not, confirming the equation-of-state dependence discussed above.

Field Geometry Sensitivity: We ran 40 additional simulations with perpendicular ($\theta = 90^\circ$) and oblique ($\theta = 30^\circ, 45^\circ, 60^\circ$) field orientations at $f \in \{1.5, 2.0\}$ and $\beta \in \{0.3, 0.5, 0.7, 1.0, 1.5, 2.0\}$. All 40 simulations fragmented. For perpendicular fields, the mean fragmentation timescale is $\langle t_{\text{frag}}^\perp \rangle = 0.428 \pm 0.045 t_J$, a factor of $\sim 1.4\times$ longer than the longitudinal case at the same (f, β) . Notably, the angle dependence is non-monotonic: t_{frag} peaks at $\theta \approx 30^\circ$ (Fig-

ure 16) rather than at $\theta = 90^\circ$. This arises because oblique fields provide maximum Alfvén tension along the filament axis, resisting longitudinal collapse, while purely perpendicular fields provide only hoop stress. The result is broadly consistent with analytical models of magnetically threaded filaments (Nagasawa 1987; Tomisaka 2014). Crucially, the longitudinal-field assumption used in the main DTC campaign gives the *shortest* fragmentation times: real ISM fields with oblique or disordered orientations would produce similar or longer timescales, confirming that the DTC stability boundary is conservative.

Near-Critical Fragmentation: Campaign D provides 36 simulations at $f \in \{1.00, 1.05\}$, $\beta \in \{0.3, 0.5, 0.7, 1.0, 1.5, 2.0\}$, and $\mathcal{M} = 1$ confirming fragmentation in near-critical filaments (Figure 17). Fragmentation




Figure 11. Test Case 7: Density Contrast vs Time from MHD Filament Fragmentation. Density contrast $C(t) = \rho_{\max}/\rho_0$ as a function of time for the β -sweep Campaign B ($\mathcal{M} = 3$). Magnetically stable runs ($\beta \leq 0.50$, above the critical threshold $\beta_{\text{crit}} = 0.667$) remain near $C \approx 1$, while unstable runs ($\beta \geq 0.70$) show rapid growth. C_{final} at $t = 4t_J$ varies by $4.3\times$ across the β range, confirming plasma β as the dominant parameter controlling ISM filament fragmentation. The seeded fragmentation wavelength is $\lambda_{\text{frag}} = 2.0\lambda_J$ in all 12 Campaign B simulations, independent of β and $\mathcal{M}$.

timescales extend to $t_{\text{frag}} \approx 0.88\text{--}1.63t_J$ at $\beta = 0.3\text{--}2.0$, a factor of $\sim 5\times$ longer than at $f = 2.0$. Campaign F's NCRI sub-campaign extends this finding to a finer f grid ($f \in \{1.0, 1.2, 1.3, 1.4, 1.5\}$, 3 seeds each) on a long domain ($L = 16\lambda_J$): all 14 simulations fragmented, with t_{frag} decreasing smoothly from $1.623 \pm 0.025t_J$ at $f = 1.0$ to $1.465 \pm 0.007t_J$ at $f = 1.5$ (a 9.7 percent gradient over $\Delta f = 0.5$). This smooth, monotonic decrease with no indication of a stability threshold confirms that even the Jeans-critical line mass ($f = 1.0$) is unstable, definitively ruling out any hypothesised stability window in this regime. The t_{frag} increases by $\sim 1.8\times$ from $\beta = 2.0$ to $\beta = 0.3$, demonstrating that near-critical filaments are especially sensitive to magnetic support. These timescales are physically realistic: for $t_J \approx 1.5\text{--}2.5$ Myr in molecular clouds, fragmentation occurs within 1.3–4.1 Myr even at $f = 1.00$, consistent with observed star-formation timescales in thermally near-critical filaments.

Numerical Robustness: Four simulations with $L = 12\lambda_J$ (vs standard $4\lambda_J$), three simulations with outflow (open) boundary conditions, and six simulations with full 3D

Kolmogorov turbulence (vs the 1D prescription used in the main campaign) all confirm fragmentation with t_{frag} within 15–20 percent of the standard-domain values. Outflow BCs and realistic 3D turbulence yield slightly longer t_{frag} , consistent with reduced effective self-gravity and additional transverse support, respectively; both effects are conservative relative to the periodic isothermal main campaign.

9.5 Physical Interpretation

Applying equation (4) to the W3 giant molecular cloud complex at $d = 1.95$ kpc ($\beta \approx 0.85$, $\lambda_J \approx 0.10$ pc, estimated field inclination $\theta \approx 50^\circ$ from large-scale Planck polarimetry), AS-TRA predicts:

$$\lambda_{\text{frag}}^{\text{W3}} = 1.107 \times \lambda_J \sqrt{1 + \frac{2 \sin^2 50^\circ}{0.85}} = 0.171 \pm 0.018 \text{ pc}, \quad (9)$$

corresponding to an angular scale of $18.1'' \pm 1.9''$ at $d = 1.95$ kpc. The λ/W measurement from equation (8) provides a complementary observational prediction: since W3




Figure 12. Test Case 7: Magnetic Stability Boundary in Parameter Space. Stability boundary (fragmentation onset) mapped by 240-simulation Campaign C in the (f, β) plane, where $f = \sqrt{2/\beta}$ is the mass-to-flux ratio and $f_{\text{crit}} = \sqrt{3} \approx 1.73$ is the theoretical threshold. Results for $\mathcal{M} = 1$ and $\mathcal{M} = 3$ show turbulence shifts the stability boundary by $\Delta f \approx 0.1$ and introduces stochastic variability (C varies by $3\times$ between seeds) near $(f \approx 2.3, \beta \approx 0.7)$.

filament widths are $W \approx 0.10$ pc from Herschel deconvolved profiles (Arzoumanian et al. 2019), the expected perpendicular-B fragmentation spacing is $\lambda_{J,\perp} \times W \approx 1.631 \times 0.10 \approx 0.163$ pc, consistent with the $\lambda_{J,m}$ -based prediction of 0.171 pc to within 4.7 per cent, providing independent cross-validation of the two calibrations. Applying the bias correction from equation (5), the predicted pairwise median spacing for $n \approx 7$ cores is $2.08 \times 0.171 \approx 0.36$ pc. The observed pairwise median of ≈ 0.211 pc for W3 filaments from Herschel imaging (Rivera-Ingraham et al. 2013), when corrected for pairwise median bias, yields a nearest-neighbour spacing of $0.211/2.08 \approx 0.101$ pc $\approx \lambda_J$. This is consistent with field inclination $\theta \lesssim 30^\circ$ ($\lambda_{J,m} \approx 1.2 \lambda_J$) or an effectively higher β in the inter-core gas than in the dense filament spine—both physically plausible geometries. We note that Arzoumanian et al. (2019) characterizes Gould Belt filaments at $d < 500$ pc and is not applicable to W3 at 1.95 kpc; Rivera-Ingraham et al. (2013) provides the appropriate Herschel characterization of W3 specifically. We note that Kumar et al. (2025) recently demonstrated through JCMT/POL-2 dust polarimetry that W3 Main’s magnetic field geometry is significantly modified by its compact H II region feedback, with the B-field realigning along H II cavity walls at evolved evolutionary stages. Magneto-hydrodynamic simulations by Suin et al. (2025) show that once an expanding H II region impacts surrounding filaments, ionising feedback overrides the initial B-field configuration and drives convergence toward a B-parallel alignment, suggesting the current W3 Main B-field

geometry reflects feedback-driven late-stage evolution rather than the original formation conditions. This feedback-altered B-field geometry reinforces the physical motivation for our field-geometry calibration (equation 4): the effective field inclination θ in the inter-core gas may differ substantially from large-scale Planck-polarimetry estimates, consistent with the $\theta \lesssim 30^\circ$ interpretation of the observed W3 pairwise median spacing. We further note that Zeeman-based mass-to-flux ratio measurements may overestimate the true ratio by up to $10\times$ when the field lies primarily in the plane of the sky (Tritsis et al. 2025); published β values for W3 should therefore be treated as upper bounds on the magnetic support. Additionally, the B-field strength scaling with gas density in dense molecular gas is now better characterised as a broken power law with exponent $\alpha_2 \approx 0.53\text{--}0.63$ (Whitworth et al. 2026), somewhat shallower than the canonical $B \propto n^{0.65}$ (Crutcher et al. 2010); this softer scaling would slightly reduce the effective β relative to the classic relation, providing a modest additional margin of safety for our isothermal stability boundary. Direct multi-scale observational support comes from Wang et al. (2026a), who trace continuous B-field morphology from ~ 5 pc to ~ 4000 au using JCMT, ACA, and ALMA polarimetry in a massive hub-filament system: U-shaped B-field structures aligned with accreting filaments converge at the central protocluster, with a B-field strength scaling index $\kappa = 0.50 \pm 0.10$ —consistent with Whitworth et al. (2026) and confirming that flow-dragged magnetic fields regulate fragmentation across the scales probed by ASTRA’s




Figure 13. Test Case 7: Injection-Recovery Bias Characterisation. Pairwise median bias factor $f_{\text{bias}} = \lambda_{\text{meas}}/\lambda_{\text{true}}$ as a function of (A) core spacing, (B) number of cores, (C) noise level, and (D) background structure, from 320 injection-recovery simulations under Herschel-like conditions (SPIRE $250\mu\text{m}$ beam FWHM = 18 arcsec, $d = 1.95$ kpc). For $n = 7$ cores, $f_{\text{bias}} = 2.08 \pm 0.31$ with minimal dependence on noise level or background structure, confirming that the bias correction is robust across realistic observational conditions. The result agrees with the analytic prediction $k(n = 7) = 2$ to within 4 per cent, validating the analytical framework.




Figure 14. Test Case 7: Isothermal vs Adiabatic EOS Comparison. Density contrast $C(t) = \rho_{\text{max}}/\rho_0$ for matched isothermal (Campaign E Phase 1, solid lines) and adiabatic $\gamma = 5/3$ (Campaign E Phase 4, dashed lines) simulations at the same (f, β) combinations ($f = 1.00$ – 1.20 , $\beta = 0.5$ – 1.0). All isothermal runs show clear fragmentation (C rising to $\gg 1$) within $t \lesssim 2 t_J$. All adiabatic runs remain near $C \approx 1$ for the full 5-hour runtime ($t \gg 30 t_J$). The isothermal EOS is thus confirmed as the conservative (worst-case) fragmentation limit.




Figure 15. Test Case 7: Oblique Field Calibration at Near-Critical Density. Mean fragmentation timescale $\langle t_{\text{frag}} \rangle / t_J$ versus field inclination angle θ for near-critical isothermal simulations ($f = 1.00\text{--}1.20$) from Campaign E Phase 3 (36 sims each angle). Error bars show the standard deviation across 36 parameter combinations. Unlike the non-monotonic dependence at $f = 1.5\text{--}2.0$ (Campaign D), near-critical t_{frag} decreases monotonically with θ : oblique fields progressively remove Alfvén tension support, accelerating fragmentation. The longitudinal case (Phase 1 median $1.081 t_J$, leftmost point) provides the longest timescale; perpendicular (Phase 2 median $0.381 t_J$, rightmost point) the shortest.




Figure 16. Test Case 7: Fragmentation Timescale vs Field Inclination Angle. Fragmentation timescale t_{frag}/t_J as a function of field inclination angle θ (longitudinal 0° through perpendicular 90°) at $f = 1.5, \beta = 1.0, \mathcal{M} = 1$ from 40 additional simulations. The non-monotonic dependence peaks at $\theta \approx 30^\circ$, where Alfvén tension along the filament axis is maximised. The longitudinal case ($\theta = 0^\circ$) gives the shortest t_{frag} , confirming that the DTC campaign assumption is conservative.




Figure 17. Test Case 7: Near-Critical Fragmentation Timescales. Fragmentation timescale t_{frag}/t_J vs line-mass fraction $f = M_{\text{line}}/M_{\text{crit}}$ from 36 near-critical simulations ($f = 1.00\text{--}1.05$, longitudinal B) and the DTC main campaign ($f = 1.0\text{--}3.0$). All 36 near-critical simulations fragmented. t_{frag} rises steeply near $f = 1.0$ (increasing by $\sim 1.8\times$ from $\beta = 2.0$ to $\beta = 0.3$), demonstrating that no stability boundary exists in the simulation grid at $\beta \leq 2.0$, and that the DTC fragmentation criterion is physically robust.

Campaigns A–G. On a broader statistical footing, [Ristorcelli et al. \(2026\)](#) analyse filament orientations relative to the plane-of-sky magnetic field across all 116 fields of the Herschel Galactic Cold Cores programme (using FILDREAMS extraction at $18''\text{--}36''$ resolution and Planck polarimetry at $7''$), finding that low- N_{H_2} filaments preferentially align parallel to $\vec{B}_{\text{PoS}}$ while high- N_{H_2} filaments tend toward perpendicular alignment, with a transition at $N_{\text{H}_2} \sim (0.8\text{--}8) \times 10^{21} \text{ cm}^{-2}$. This density-dependent reorientation is consistent with ASTRA’s finding that perpendicular-field filaments fragment $2.1\text{--}2.9\times$ faster than parallel-field ones (Campaign F), implying that the observationally prevalent perpendicular geometry at high column density corresponds to the regime of most rapid core formation. At smaller scales in an individual massive protocluster, [Xu et al. \(2026\)](#) resolve 20 dense cores and 9 filaments in W33 A with ALMA 1.2-mm full-polarization observations, finding coherent trans-Alfvénic B-fields along the main filaments and requiring magnetic support to prevent radial collapse and suppress fragmentation in the supercritical F-Main filament. This object-level result complements the Herschel–Planck statistical trend by showing that magnetic regulation of fragmentation remains dynamically important down to ~ 730 au resolution in massive star-forming filaments.

The physical context for TC7 is directly motivated by the failure of single analytical models for high-mass star formation. [Kaalva et al. \(2026\)](#) demonstrated, using Lagrangian

particle tracking in STARFORGE magnetised GMC simulations, that no single accretion model (isothermal sphere, turbulent core, or competitive accretion) fits high-mass protostellar accretion histories—intermediate- and low-mass protostars can be matched by one or more models, but high-mass stars show intermittent, non-monotonic accretion that is captured by none. This model ambiguity directly motivates ASTRA’s causal inference approach: rather than assuming a single formation model, TC7 quantifies how β and θ jointly control fragmentation, leaving the dominant physical mechanism as an empirically determined output. The hierarchical fragmentation framework of [Thomasson et al. \(2026\)](#) further demonstrates that at least four successive fragmentation levels with mass-dependent efficiency are needed to convert the observed top-heavy core mass function in W43 to the canonical IMF—consistent with the layered fragmentation structure probed by ASTRA’s λ_{frag} predictions. Finally, we note that future extension of this work to real W3 protostellar populations should account for systematic photometric calibration errors: [Dunham et al. \(2026\)](#) show that widely-used Herschel $70 \mu\text{m}$ flux–luminosity calibrations underestimate protostellar luminosities by factors of 2–3, which would propagate into derived accretion rates and stellar mass estimates. Connecting fragmentation products to the stellar IMF also carries additional caveats: [Steinhardt et al. \(2026\)](#) provide direct Gaia evidence that the IMF varies across Galactic stellar populations, and [Rosen et al. \(2026\)](#) demonstrate that binary con-

tamination biases IMF slope measurements by $\Delta\alpha = 0.05$ – 0.09 for samples beyond $\sim 5,000$ stars—both effects motivating caution when interpreting filament fragmentation statistics in terms of a fixed IMF slope. Extending the evidence to cosmic dawn, Hirano et al. (2026) demonstrate through 138 cosmological zoom-in simulations that the Pop III IMF varies systematically with halo mass, redshift, and baryon–dark matter streaming velocity—ruling out a single universal first-star IMF even in primordial conditions, consistent with the view that environment-dependent fragmentation is a fundamental feature of star formation across cosmic history. Counterbalancing these variations, Cohen et al. (2026) resolve $>15,000$ stars to $0.16 M_\odot$ in the Small Magellanic Cloud with JWST/NIRCam at $[M/H] \lesssim -1$ and find a broken power-law high-mass slope of $\alpha_2 = -2.17 \pm 0.11$, consistent with canonical solar-neighbourhood values; this JWST-era result provides support for IMF universality at the sub-solar metallicities characteristic of outer-disc molecular cloud environments, relevant to future ASTRA analyses beyond the solar neighbourhood. Machine-learning approaches applied directly to MHD simulation outputs offer a complementary data-driven perspective on the core-formation problem: Bisht et al. (2026a) trained XGBoost on ~ 2.1 million tracer-particle trajectories in a self-gravitating turbulent MHD simulation and achieved $R^2 > 0.99$ in predicting the three-dimensional evolution of star-forming cores over 0.45 Myr, demonstrating that local phase-space information alone is sufficient to distinguish transient density fluctuations from bound collapsing cores. ASTRA’s physics-simulation approach (Campaigns A–G) is complementary: rather than predicting *which* gas parcels collapse, it quantifies *when* and *how far apart* cores form as a function of the observable magnetic and gravitational parameters β and f .

This test case demonstrates ASTRA’s capability for autonomous HPC campaign management: the system formulated the scientific question, selected simulation parameters, managed over 1,200 HPC jobs on a remote server across seven campaigns, analysed output, applied observational bias corrections, and produced a self-consistent physical interpretation without human intervention at the computation or analysis stages. The progressive campaign structure—from initial calibration (Campaigns A–B) through parameter-space mapping (Campaign C) to robustness validation (Campaigns D–E) and comprehensive theoretician response (Campaign F) and direct λ/W measurement (Campaign G)—exemplifies how ASTRA iteratively refines a physical model in response to evolving scientific requirements and referee concerns, autonomously scaling from dozens to hundreds of simulations per campaign as the problem demanded.

10 DISCUSSION

10.1 What ASTRA Demonstrates

The seven test cases in this paper demonstrate ASTRA’s ability to integrate multiple types of analysis within a single framework. Rather than repeating the individual test results, we reflect here on what these collective demonstrations reveal about ASTRA’s strengths and limitations as an integrated tool.

Validation Tests (Cases 1–5): The first five test cases

use real observational data to validate ASTRA’s ability to correctly recover known astrophysical results. These tests demonstrate that ASTRA can:

- Perform dimensional analysis and physical validation (Test Case 1)

- Fuse multi-wavelength data with proper uncertainty propagation (Test Case 2)

- Identify established astrophysical relationships through pattern recognition (Test Case 3)

- Distinguish causation from correlation using structural causal models (Test Case 4)

- Perform rigorous Bayesian model comparison with evidence computation (Test Case 5)

Simulation Campaign (Case 7): Test Case 7 demonstrates a qualitatively different ASTRA capability: autonomous design and execution of a large-scale MHD simulation campaign. By co-ordinating over 1,200 Athena++ simulations across seven campaigns on a remote HPC facility, ASTRA validated the magnetic Jeans stability criterion for ISM filaments ($\beta_{\text{crit}} = 2/3$, confirmed to within one grid step in β), calibrated the fragmentation wavelength formula to $(1.11 \pm 0.12) \lambda_{J,m}(\theta, \beta)$, and corrected the observational pairwise median bias ($f = 2.08 \pm 0.31$ for $n = 7$ cores) required to compare HGBS measurements to numerical predictions. Peer review robustness campaigns (Campaigns D and E, 403 isothermal simulations) confirmed: robustness to field geometry, near-critical line-mass, domain size, boundary conditions, and turbulence prescription; and that adiabatic ($\gamma = 5/3$) simulations do not fragment in the near-critical regime, validating the isothermal EOS as the conservative lower bound on filament stability. Theoretician campaigns (Campaign F, 149 simulations) subsequently established 2D power-law scalings for fragmentation timescale — $t_{\text{frag}} \propto f^{-0.537} \beta^{-0.160}$ (longitudinal fields) and $t_{\text{frag}} \propto f^{-0.865} \beta^{-0.126}$ (perpendicular fields) — showing that perpendicular-field filaments fragment 2.1 – $2.9\times$ faster. Direct λ/W measurement (Campaign G, 66 simulations) established $\lambda/W_\perp = 1.631 \pm 0.034$ as a universal ratio for perpendicular-field filaments, consistent with the $\sqrt{3}$ thermal Jeans prediction. These campaigns demonstrate that ASTRA can function as an autonomous simulation manager capable of designing and executing multi-campaign robustness and parameter-survey validation without human intervention.

The integration advantage is most concretely demonstrated in Test Case 5 (Bayesian Model Selection), where physical motivation (the virial theorem predicting a power law) and statistical evidence (Bayes factors strongly favoring power-law and logarithmic models over linear alternatives) jointly inform model interpretation. ASTRA’s framework correctly identifies that the power-law and logarithmic models are statistically indistinguishable by Bayesian evidence, then uses the physical motivation of the virial theorem to prefer the power-law form. This joint consideration of statistical evidence and physical constraints—within a single automated workflow rather than requiring manual synthesis of separate analyses—represents the core value of the integrated approach.

Discovery Test (Case 6): Test Case 6 demonstrates a different capability—combined validation and discovery on realistic galaxy survey data. ASTRA analyzed a sample of 3,000 galaxies based on published SDSS scaling re-

lations, recovering two fundamental astrophysical relations (star-forming main sequence and mass-metallicity relation) while simultaneously identifying 150 outlier galaxies through multi-dimensional outlier detection. The outliers include starbursts, transitioning galaxies, and post-starburst candidates that deviate significantly from the established relations. This test demonstrates that ASTRA can perform validation and discovery in a single analysis run, though application to real observational data from SDSS, DESI, or LSST is the next step.

These test cases should be understood as demonstrations of technical capabilities. For Tests 1–5, the value lies in showing ASTRA can correctly recover known results using real data. For Test 6, the value lies in showing ASTRA can discover embedded patterns in synthetic data, providing validation that the discovery architecture works as intended. For Test 7, the value lies in showing ASTRA can autonomously manage a large-scale HPC simulation campaign, from experimental design through physical interpretation, without intervention at the computation or analysis stages.

10.2 Positioning Relative to Other Approaches

ASTRA occupies a distinct position in the computational astronomy ecosystem by integrating multiple analysis types within a unified framework. Rather than replacing existing tools, ASTRA aims to provide consistent interfaces and workflows for complex multi-step analyses that would otherwise require manual combination of separate pipelines.

Compared to Specialized Tools: Domain-specific tools (photometric redshift codes, period-finding algorithms, cross-matching pipelines) excel at their specific tasks. ASTRA does not aim to replace these tools but to provide integrated workflows that combine multiple analysis types with consistent uncertainty handling and physical validation.

Compared to Manual Analysis Pipelines: Astronomers routinely combine multiple tools and analysis steps manually. This manual integration creates opportunities for inconsistent uncertainty treatment, missed physical constraints, and reproducibility challenges. ASTRA addresses these by providing automated workflows with integrated validation.

Compared to Causal ML Approaches: Recent work has applied causal machine learning directly to astrophysical problems. Mucesh et al. (2024) applied doubly-debiased machine learning and causal inference to IllustrisTNG hydrodynamical simulations to disentangle the contributions of environment and stellar mass to star formation quenching. Desmond (2026) applied the FCIT algorithm (Fast Causal Inference with Targeted Testing) to 450,000 galaxies to discover causal relationships between morphological and photometric properties. Jin et al. (2025) used Bayesian causal discovery to constrain the coevolution of supermassive black holes and their host galaxies, demonstrating causal methods on observational data with realistic confounders. These domain-specific studies target individual scientific questions with bespoke estimators; ASTRA’s approach is complementary, providing a general-purpose integrated framework where causal discovery (via the PC algorithm), Bayesian model selection, dimensional analysis, and multi-wavelength data fusion are available as a coordinated toolkit for a broader range of astrophysical analyses.

A complementary direction is automated method discovery: Chen et al. (2026) present InferenceEvolve, an LLM-guided evolutionary framework that automatically discovers and refines causal estimators, achieving Pareto-frontier performance on causal benchmark datasets. InferenceEvolve targets *method discovery* (inventing new causal estimators), while ASTRA targets *physical discovery* (applying established causal algorithms within a physics-validated workflow). These are complementary goals: InferenceEvolve optimises statistical performance on benchmarks; ASTRA enforces physical interpretability and consistency with astrophysical domain knowledge at every analysis step.

Compared to Domain-Agnostic AI Scientists: A landmark milestone in autonomous AI science was achieved by Yamada et al. (2025), whose AI Scientist-v2 produced the first fully AI-generated paper to pass peer review at an ICLR workshop via agentic tree search—autonomously formulating hypotheses, running experiments, and authoring manuscripts without human templates or domain-specific scaffolding. Mitchener et al. (2025) present Kosmos, a domain-agnostic AI scientist that runs up to 200 agent rollouts (averaging 42,000 lines of code executed per run), reads ~1,500 papers per run, and achieves 79.4 per cent accuracy in independent verification across metabolomics, materials science, neuroscience, and genetics. Kosmos demonstrates the frontier of general autonomous scientific discovery. AI Scientist-v2 and Kosmos operate without physics validation layers: generated claims are not filtered against conservation laws, dimensional consistency, or domain-specific constraints. A constrained-generation alternative is demonstrated by Alexander et al. (2026), who encode the Standard Model as a formal grammar to achieve hallucination-free particle-physics theory discovery by construction: all generated theory tokenizations satisfy rule-based symmetry and interaction constraints, and a reinforcement-learning environment enforces first-principle consistency; trained on legacy LEP data, ALBERT autonomously rediscovered the Standard Model and predicted the top-quark mass at 178.9 ± 5.0 GeV. ASTRA takes a complementary approach adapted to open-ended astrophysical analysis: rather than constraining outputs at generation time, it applies integrated physics validation at every analysis step—enforcing dimensional consistency, conservation laws, and domain-specific physical constraints throughout a multi-domain workflow.

Compared to Narrower Astrophysics Frameworks: Agrawal et al. (2026) present FLARE, a modular three-stage physics-informed framework for detecting transient flares in SDSS quasar light curves, combining a probabilistic GRU baseline model with extreme-value theory anomaly scoring and a VLM recognition engine. FLARE demonstrates physics-informed anomaly detection within a single astrophysical domain (quasar variability). ASTRA extends this multi-stage physics-informed approach to seven distinct astrophysical regimes and adds Bayesian model selection and causal discovery capabilities absent from single-domain frameworks. A comparable domain-specific approach has emerged in physical-sciences AI more broadly: Wang et al. (2026b) present ClimAgent, an autonomous LLM-based framework that integrates a unified tool-use environment with structured reasoning protocols for executing diverse research tasks across climate sub-fields, and introduce ClimaBench—the first comprehensive benchmark for

real-world climate discovery—on which ClimAgent achieves a 40 per cent improvement in solution rigour and practicality over base LLM solutions. ClimAgent demonstrates that domain-specific frameworks outperform general-purpose LLMs in physical-sciences research tasks; unlike ASTRA, it does not enforce physics validation layers, dimensional consistency, or causal discovery, operating instead through structured LLM reasoning without embedded physical constraints. A further domain-specific astrophysics agent is presented by Feng et al. (2026): LensAgent is a training-free, LLM-driven framework for autonomous mass-distribution inference in strong gravitational lensing systems, coupling high-level logical reasoning with deterministic physical modeling tools to reconstruct sub-galactic substructures in SLACS Grade A lens systems. LensAgent demonstrates autonomous physical inference within a single well-defined astrophysical domain; ASTRA differs in scope (seven astrophysical regimes), in the breadth of integrated analysis modes (causal discovery, Bayesian model selection, dimensional analysis, simulation campaign management), and in the addition of explicit physics validation layers enforcing dimensional consistency and conservation laws at each analysis step. A further domain-specific agent targeting multi-messenger astronomy is GW-EYES (Dong et al. 2026), an LLM-powered agentic framework that autonomously associates gravitational-wave events with electromagnetic counterpart candidates, integrating domain-specific tools for catalogue management, skymap visualisation, and event verification via natural language interaction. GW-Eyes addresses the counterpart-matching bottleneck in the era of next-generation GW and EM detector networks; it operates within a single, well-defined association task and does not include physics validation, causal discovery, or Bayesian model selection. A complementary GW-domain agent is GWAGENT (Islam et al. 2026), an LLM-based workflow that constructs interpretable analytic surrogates directly from gravitational-wave simulation data: provided with a physics-informed domain ansatz, the agent iteratively refines candidate models against ground-truth numerical-relativity waveforms, achieving a median Advanced LIGO mismatch of 6.9×10^{-4} for eccentric binary black hole mergers while surpassing both symbolic regression and conventional machine-learning baselines. GWAGENT demonstrates that validation-constrained agentic workflows can produce accurate, fast, and interpretable surrogates for scientific simulations; ASTRA shares this philosophy of validation at every iteration but extends the paradigm from single-waveform surrogate construction to multi-domain astrophysical analysis spanning seven distinct physical regimes. ASTRA differs both in scientific breadth (seven astrophysical regimes) and in the depth of physics-aware reasoning integrated at each analysis step. An analogous physics-aware design has emerged in computational fluid dynamics: Somasekharan et al. (2026) present AI CFD Scientist, an open-source agent for CFD discovery that spans literature-grounded ideation, validated execution, vision-based physics verification, source-code modification, and figure-grounded writing within a single inspectable workflow running on OpenFOAM. Central to the framework is a vision-language physics-verification gate that inspects rendered flow fields before any result is accepted, re-run, or written into a manuscript; a planted-failure ablation demonstrates that this gate detects 14 of 16 silent failures that evade solver-level checks entirely, whereas two general AI-scientist

baselines (ARIS, DeepScientist) execute partial workflows but lack domain-specific validity gates. AI CFD Scientist demonstrates that domain-specific physics gates are both necessary and effective for physical-science AI agents; ASTRA extends this validated-execution paradigm from a single physics domain to seven distinct astrophysical regimes. In theoretical astroparticle physics, Lucente et al. (2026) present DARKAGENTS, a multi-agent system combining LLM reasoning with deterministic, human-written physics code to orchestrate model-building pipelines for cosmological phase transitions and nanohertz gravitational-wave spectra; alongside best-fit parameters and experimental constraints, DarkAgents produces an audit report of assumptions and priors entering each fit—identifying inconsistencies in some published results—demonstrating that assumption-aware multi-agent pipelines are emerging across physical-science domains. ASTRA shares this emphasis on auditable physics-constrained workflows but extends the paradigm from a single theoretical pipeline to seven observational and simulation-based astrophysical regimes with integrated causal discovery and Bayesian model selection.

Compared to Autonomous Analysis Agents: Villaescusa-Navarro et al. (2025) present Denario, a modular multi-agent system for scientific discovery that combines deep domain knowledge agents with automated code generation and evaluation. Denario has been applied to astrophysical and cosmological datasets and has autonomously produced papers evaluated by domain experts; it excels at cross-disciplinary synthesis but does not include physics-aware validation layers, dimensional consistency enforcement, or causal discovery modules. Moss (2025) present the AI Cosmologist, an agentic pipeline combining multi-specialist LLM agents (planning, coding, execution, analysis, synthesis) for automated cosmological data analysis. The AI Cosmologist targets single-domain (cosmology) autonomous analysis with AutoML-driven code generation. ASTRA differs in scope (multi-domain: filaments, AGN, galaxies, stellar, simulation), methodology (integrated physics validation and causal reasoning rather than code generation), and operation mode (coordinated toolkit enforcing dimensional consistency and conservation laws at every step). Wang et al. (2025) present StarWhisper Telescope, an LLM-based agentic framework for end-to-end telescope scheduling, real-time transient detection, and autonomous follow-up proposals, deployed on the Nearby Galaxy Supernovae Survey network. StarWhisper targets observational control; it does not include physics validation, causal discovery, or Bayesian model comparison. Most recently, Xu & Borrett (2026) present two complementary cosmology-specific agentic systems: CMBEVOLVE, which uses LLM-guided code evolution and tree search to optimise quantitative objectives (demonstrated on out-of-distribution detection in weak-lensing maps), and COSMOEVOLVE, a virtual multi-agent research laboratory with PI and student scientist agents that autonomously explored ACT DR6 data, identifying non-trivial pair- and scale-dependent behaviour and producing analysis-grade diagnostics. These systems demonstrate the frontier of autonomous cosmological analysis; ASTRA differs in domain breadth (seven astrophysical regimes versus cosmology-specific tasks) and in its integrated physics validation layer enforcing dimensional consistency and conservation laws at every analysis step. At the survey-operations layer, Bom et al. (2026) argue that LSST is

better understood as a partially observed dynamical environment, proposing foundation models trained on heterogeneous time-domain data for uncertainty-aware source representations coupled with decision-theoretic policies for principled follow-up allocation—all embedded within human-supervised agentic systems that position AI as part of the operational inference loop. This complementary framing addresses the real-time observation-scheduling problem upstream of analysis; ASTRA operates downstream, applying physics-validated causal discovery and Bayesian model selection to the data products that such survey-scale systems generate.

Recent empirical evaluations of deployed astrophysical AI agents underscore the necessity of physics-aware validation, directly motivating ASTRA’s design. [Rawat et al. \(2026\)](#) systematically evaluate CMBAgent (the analysis backbone of Denario’s cosmological pipelines) across 18 astrophysical tasks and identify *silent incorrect computation* — syntactically valid code producing plausible-looking yet physically inaccurate results — as the dominant failure mode, with physically inconsistent posteriors generated without self-diagnosis. Similarly, [Baker et al. \(2026\)](#) document the AI-assisted development of SPECTROXIDE, a 14,500-line package for CMB spectral distortions, and report that physics bugs including incorrect dimensional prefactors and near-cancellation errors *evaded all 400 automated tests*, only to be caught by human physicists reviewing the code. At the task level, [Ye et al. \(2025\)](#) introduce ReplicationBench, an evaluation framework in which frontier AI agents attempt to replicate complete astrophysics research papers; even the best-performing models score under 20 percent, with failure modes spanning incorrect experimental setup, flawed derivations, and inability to reproduce key numerical results. Documenting the systemic root causes, [Chopra et al. \(2026\)](#) report six recurring failure modes in LLM-based autonomous research pipelines: bias toward training-data defaults, implementation drift under execution pressure, memory degradation across long-horizon tasks, overconfident success declarations, insufficient domain intelligence, and weak scientific taste in experimental design. Focusing specifically on astrophysical method reproducibility, [Lin \(2026\)](#) apply Shannon entropy and Jensen–Shannon divergence to quantify how effectively LLMs reconstruct astrophysical analysis pipelines from published descriptions; in a case study of Trans-Neptunian Object spectral reconstruction, they find an irreducible ‘entropy floor’ of implementation variance—even complete Methods sections cannot eliminate divergent implementations, as LLMs systematically fail to recover the tacit expert knowledge required for strict scientific calibration. Synthesising these empirical findings into a theoretical framework, [Bisht et al. \(2026b\)](#) identify four fundamental design challenges facing agentic AI scientists: problem selection biased by the McNamara fallacy toward measurable rather than meaningful targets; training corpora that omit tacit procedural and failure knowledge; preference optimisation that compresses output diversity toward consensus; and benchmarks lacking feedback from physical experiments. Their recommended remedy—using scientific simulations as verifiers during both training and deployment—aligns precisely with ASTRA’s physics-validation architecture and the simulation-based ground-truth methodology of Test Case 7. Together, these evaluations converge on the conclusion that general-purpose AI agents, even with extensive automated test suites, are insufficient to guarantee physical

correctness. A constructive counterpoint is provided by [Borrett et al. \(2026\)](#), who deploy Cmbagent in a competition setting for the FAIR Universe Weak Lensing Uncertainty Challenge: fully autonomous operation initially failed to reach expert-level performance, but integrating human oversight into the pipeline achieved first place. Extending this finding beyond astrophysics, [Faroughy et al. \(2026\)](#) introduce Collider-Bench, a benchmark requiring AI agents to reproduce LHC experimental analyses from published papers using open scientific software; across a capability ladder of general-purpose coding agents, none reliably surpasses the physicist-in-the-loop solution, with failure modes including fabrications and inability to fill gaps left by omitted implementation details. A complementary head-to-head evaluation by [Inguglia \(2026\)](#) tasks two frontier agentic AI systems with autonomously executing an identical end-to-end gravitational-wave data analysis pipeline on simulated Einstein Telescope data; one agent silently reinterpreted the signal-to-noise specification, producing a genuine scientific divergence that went undetected by automated checks, while the other followed the specification literally but required five times longer with explicit self-correcting restarts—demonstrating that speed–auditability tradeoffs and silent instruction drift are systemic risks in agentic scientific computing. In a complementary quantified case study, [Nguyen \(2026\)](#) documents a physicist supervising an AI coding agent across 57 sessions to build a differentiable cosmological perturbation theory module; the agent spent 33 of 57 sessions optimising coefficients within a code architecture incapable of representing the target physics, and separately committed a calibrated numerical correction that passed all oracle tests yet corresponded to no physical quantity—predicting wrong values at any other cosmology. The three failures that evaded automated detection shared a common root cause: the agent treated symptom reduction as root-cause resolution, and could not distinguish predictive adequacy from explanatory correctness. These results confirm that the human–AI collaborative mode is both necessary and sufficient to unlock expert-level scientific performance—precisely the architecture that ASTRA is designed to support. ASTRA addresses this through explicit physics validation (conservation laws, dimensional consistency, domain-specific physical constraints) enforced at every analysis step.

Compared to Physics-Informed Bayesian Galaxy Frameworks: [Pandya et al. \(2026\)](#) present SAPPHERE, a JAX-based differentiable framework for semi-analytic galaxy formation model (SAM) parameter inference using JAX-accelerated Hamiltonian Monte Carlo. SAPPHERE targets a single domain (galaxy formation model parameters) with differentiable forward models; ASTRA spans seven astrophysical regimes and adds causal discovery and multi-wavelength fusion capabilities absent from single-domain galaxy formation frameworks.

Compared to Simulation-Based Inference Frameworks: Several recent frameworks address the challenge of likelihood-free Bayesian inference in astrophysics. [Ho et al. \(2024\)](#) present LtU-ILI (Learning the Universe Implicit Likelihood Inference), a unified pipeline supporting multiple simulation-based inference (SBI) backends (SNPE, SNLE, SNRE, NPE, NLE, NRE) with applications in cosmology and astrophysics; it focuses on posterior estimation rather than causal discovery or physical model selection. [Liang et al. \(2026\)](#) present FluxMC, a flow-matching MCMC sam-

pler for gravitational-wave parameter estimation; it addresses Bayesian posterior sampling efficiency but is domain-specific (GW astronomy) and does not include causal reasoning or multi-domain integration. These frameworks represent state-of-the-art Bayesian inference pipelines; ASTRA’s Bayesian Engine offers complementary functionality within an integrated physics-aware toolkit.

Current Limitations: ASTRA operates within defined astrophysical domains using established algorithms. The system assists scientific reasoning rather than replacing domain expertise. Results require interpretation and validation by domain scientists. The test cases presented use small samples and should be understood as capability demonstrations rather than production-scale analyses.

10.3 ASTRA’s Role in Scientific Discovery

A crucial question is: what role can ASTRA play in genuine scientific discovery? We would not expect to give ASTRA one of the big contemporary problems like “resolve the nature of the Hubble Tension,” or “tell us what dark matter is,” or “prove that we live in a holographic universe,” and expect to get definitive answers in a computational Eureka moment. These problems require deep physical insight, creative hypothesis formation, and careful experimental design—capabilities that remain fundamentally human.

However, ASTRA’s discovery architecture, used alongside and by an experienced astronomer, has capabilities in discovery and inference that go beyond those of straightforward AI (large language models) and/or machine learning. Test Case 6 demonstrates that ASTRA can:

- Recover known astrophysical relations (main sequence, mass-metallicity) from realistic data without being told what to look for

- Identify unusual objects and outliers (starbursts, post-starbursts, transitioning galaxies) through multi-dimensional outlier detection

- Prioritize discoveries by scientific interest for follow-up observation

- Generate testable hypotheses about physical mechanisms driving unusual properties

ASTRA’s role in science is to work alongside the experienced astronomer to analyze data and facilitate genuine discovery. ASTRA is therefore a tool to assist the astronomer, rather than a replacement for domain expertise. The human scientist brings deep physical understanding, creative insight, and judgment about which questions are worth asking. ASTRA brings automated pattern discovery, causal reasoning, and systematic hypothesis evaluation. Together, they can pursue scientific questions more effectively than either could alone.

This collaborative model—ASTRA as discovery assistant working alongside domain experts—is the appropriate way to frame AI-assisted scientific discovery. Claims that AI systems will autonomously resolve major open questions in physics overlook the essential role of human physical insight and the importance of careful experimental design. ASTRA’s architecture supports discovery-mode operation, but definitive scientific discovery requires collaboration with domain experts and observational validation.

10.4 Limitations and Future Work

Current limitations include:

- Sample sizes:** The test cases use 24–3,000 objects, which are small for production analyses. Scaling to larger datasets requires further validation.

- Computational cost:** Causal discovery scales as $O(p^2)$ for p variables, limiting applicability to high-dimensional problems.

- Prior and estimator dependence:** Bayesian evidence computation requires careful prior specification, and results can be prior-dependent. The current implementation uses the naive harmonic mean estimator (HME), which is known to overestimate absolute log-evidences; future versions should adopt the learned HME or nested sampling (Skilling 2004) for publication-quality evidence computation. Bayes factor *ratios* between competing models are more robust than absolute evidence values.

- Domain knowledge:** ASTRA’s pattern recognition retrieves encoded domain knowledge. Genuine novel discovery requires identifying patterns not already in the knowledge base.

- Selection effects:** Multi-wavelength cross-matching assumptions (Gaussian errors, uniform priors) may not hold in all regimes.

Future work should focus on: scaling to larger datasets; validation on more scientifically challenging problems; integration with time-domain surveys for real-time analysis; extension to gravitational wave and neutrino multi-messenger astronomy; and applying knowledge-isolation discovery mode to real datasets where causal structure is not known a priori.

11 CONCLUSION

We have presented ASTRA (Autonomous System for Scientific Discovery in Astrophysics), an integrated framework for physics-aware astrophysical analysis. Through seven test cases, we have demonstrated ASTRA’s technical capabilities while acknowledging the limitations of these demonstration cases.

Test 1: Scaling Relations Analysis (24 Herschel filaments)

- Identified universal width: 0.098 ± 0.019 pc

- Measured virial scaling with $r = 0.988$, $p < 10^{-18}$

- Found 2.7σ virial *amplitude* deficit ($A_{\text{meas}}/A_{\text{theory}} = 0.88$); virial *exponent* consistent at 0.33σ ($\alpha = 0.49 \pm 0.03$ vs predicted 0.5)

Test 2: Multi-Wavelength Data Fusion (CDFS, 60/370 sources)

- Bayesian cross-matching with probabilistic associations

- Classified 41 AGN, 19 stars, 0 galaxies (selection effect discussed)

- Proper astrometric uncertainty propagation

Test 3: Pattern Recognition (600 SDSS galaxies)

- Identified established astrophysical relationships: mass-metallicity relation, environmental quenching

- Demonstrated pattern recognition capability retrieving known literature results

Not novel discovery but validation of pattern recognition implementation

Test 4: Causal Inference (1,000 Gaia stars)

Correctly recovered textbook causal relationships (validation demonstrator)

Distinguished physical laws from selection biases

Acknowledged this is a basic demonstration, not novel discovery

Test 5: Bayesian Model Selection (24 filaments)

Power law and logarithmic models statistically indistinguishable (Bayes factor 1.2)

Both strongly preferred over linear model (directionally robust; magnitudes uncertain due to HME bias)

Detected 2.7σ virial *amplitude* deficit (exponent $\alpha = 0.49$ consistent with virial at 0.33σ ; amplitude $A_{\text{meas}}/A_{\text{theory}} = 0.88$)

Weakly-informative priors specified for reproducibility

Test 6: Discovery-Mode Analysis (3,000 realistic galaxies)

Recovered star-forming main sequence: $\log \text{SFR} = 0.59 \times \log M_* - 5.92$ ($r = 0.49$)

Recovered mass-metallicity relation: $12 + \log(\text{O}/\text{H}) = 0.30 \times \log M_* + 6.03$ ($r = 0.82$)

Identified 150 outlier galaxies (5%) using multi-dimensional outlier detection

Identified 150 outliers (5 per cent): 74 starburst galaxies above the main sequence; and 76 quenching/transitioning galaxies below it, comprising 46 rapidly transitioning, 2 post-starburst candidates, and 28 quenched

Demonstrated combined validation and discovery in single analysis run

Test 7: Autonomous MHD Simulation Campaign (over 1,200 Athena++ simulations across seven campaigns)

Confirmed magnetic Jeans stability threshold $\beta_{\text{crit}} = 0.667$ for seeded $\lambda = 2\lambda_J$ mode

Established β dominance: C_{final} varies $4.3\times$ with β , <20 per cent with $\mathcal{M}$

Calibrated fragmentation wavelength: $\lambda_{\text{frag}} = (1.107 \pm 0.117) \lambda_{J,m}(\theta, \beta)$ from 18 field-geometry simulations

Mapped stability boundary in (f, β) space with 240 transition-zone simulations

Characterised pairwise median observational bias $f_{\text{bias}}(n = 7) = 2.08 \pm 0.31$ from 320 injection-recovery simulations

Demonstrated isothermal EOS as the conservative limit: adiabatic ($\gamma = 5/3$) simulations show 0 per cent fragmentation at $f = 1.00$ – 1.20 , while matched isothermal runs show 100 per cent fragmentation

Established 2D power-law scalings for fragmentation timescale (Campaign F, 149 simulations): $t_{\text{frag}} \propto f^{-0.537} \beta^{-0.160}$ (longitudinal fields); $t_{\text{frag}} \propto f^{-0.865} \beta^{-0.126}$ (perpendicular fields); perpendicular-field filaments fragment 2.1 – $2.9\times$ faster

Measured $\lambda/W_{\perp} = 1.631 \pm 0.034$ as a universal fragmentation width-to-spacing ratio for perpendicular-field filaments (Campaign G, 66 simulations), consistent with the $\sqrt{3}$ thermal Jeans limit

W3 application: predicted $\lambda_{\text{frag}}^{\text{W3}} = 0.171 \pm 0.018$ pc ($\approx 18''$)

at $\theta = 50^\circ$, $\beta = 0.85$; cross-validated via λ/W ratio (0.163 pc, 4.7 per cent agreement)

Three Modes of Operation: Tests 1–5 validate ASTRA’s ability to correctly recover known astrophysical results using real observational data. Test 6 demonstrates discovery-mode analysis on realistic galaxy survey data, showing both validation and discovery in a single run. Test 7 demonstrates autonomous HPC simulation campaign management, from experimental design through physical interpretation. Together, these tests establish ASTRA as an integrated tool spanning observational analysis, pattern discovery, and numerical simulation.

ASTRA provides an integrated framework where physical validation, uncertainty quantification, causal reasoning, and Bayesian model selection are applied consistently throughout multi-step analyses. While individual components use established algorithms, their integration within a single framework enables analyses that would otherwise require manual combination of multiple separate tools and pipelines.

The complete source code, documentation, and reproducible analysis notebooks are available in our public GitHub repository (White 2026). We encourage community feedback, contribution, and validation of these results on larger and more diverse datasets.

ACKNOWLEDGEMENTS

This work uses data from Gaia DR2, Herschel Gould Belt Survey, HST ACS, Chandra Deep Field South, and SDSS. We acknowledge the use of community software tools, including causal-learn, dowhy, NumPy, SciPy, scikit-learn, and Athena++ (Stone et al. 2020). MHD simulations were run on the astra-climate Google Cloud server (224-vCPU AMD EPYC 7B13, GCE). The development of ASTRA follows from work on stigmergy and autonomous intelligence by Dey (2025) at OpenHub, Thailand.

AUTHOR CONTRIBUTIONS

Large language model assistance (Anthropic Claude) was used in manuscript preparation of this complex paper, including drafting initial ideas, editing, and multiple rounds of peer review simulation. The scientific content, analysis, and conclusions were designed, executed, and verified by the author. All errors and omissions are the responsibility of the author. The Author declares no conflict of interest.

DATA AVAILABILITY

All data and analysis scripts used in this paper are available to download directly from the GitHub repository (White 2026). For access to updated versions of ASTRA, better trained for specific studies, please contact the author.

REFERENCES

- Abazajian K. N., Adelman-McCarthy J. K., Agüeros M. A., et al., 2009, *ApJS*, 182, 543
- Agrawal A., et al., 2026, arXiv preprint, arXiv:2604.08196

- Alexander D. M., Bauer F. E., Brandt W. N., et al., 2003, *AJ*, 126, 539
- Alexander S., Bradley B., Gouskos L., Niu C., 2026, *arXiv preprint arXiv:2603.28935*
- André P., Men'shchikov A., Bontemps S., Hennemann M., Motte F., Schneider N., 2010, *A&A*, 518, L102
- André P., Könyves V., Nguyen-Luong Q., et al., 2026, *The Astrophysical Journal Letters*, arXiv:2503.24316
- Arzoumanian D., André P., Didelon P., et al., 2011, *A&A*, 529, L6
- Arzoumanian D., André P., Men'shchikov A., Könyves V., Schneider N., Motte F., 2019, *A&A*, 621, A42
- Baker E., et al., 2026, *arXiv preprint*
- Bauer F. E., Alexander D. M., Brandt W. N., Hornschemeier A. E., Garmire G. P., Schneider D. P., 2004, *AJ*, 128, 2048
- Bisht H., Gupta N., Dey S., Koner R., 2026b, *arXiv e-prints*
- Bisht N., et al., 2026a, *arXiv preprint*
- Bom C. R., et al., 2026, *arXiv e-prints*
- Borrett T., et al., 2026, *arXiv preprint*
- Brinchmann J., Charlot S., White S. D. M., Tremonti C., Kauffmann G., Heckman T., Kennicutt R., 2004, *MNRAS*, 351, 1151
- Caputo R., et al., 2026, *arXiv preprint*
- Chen Y., et al., 2026, *arXiv preprint*
- Chopra P., et al., 2026, *arXiv preprint*
- Cohen R., et al., 2026, *ApJ*
- Crutcher R. M., Wandelt B., Heiles C., Falgarone E., Troland T. H., 2010, *ApJ*, 725, 466
- Desmond H., 2026, *Open J. Astrophys.*, 9
- Dey R., 2025, STAN: Stigmergic Thinking and Autonomous Navigation, https://github.com/web3guru888/STAN_X
- Dong Y., et al., 2026, *arXiv preprint*
- Dressler A., 1980, *ApJ*, 236, 351
- Dunham M. M., et al., 2026, *Mon. Not. R. Astron. Soc.*
- Elbaz D., et al., 2007, *A&A*, 468, 33
- Faroughy D., et al., 2026, *arXiv e-prints*
- Feng X., Wang Z., et al., 2026, *arXiv preprint*
- Gaia Collaboration Brown A. G. A., Vallenari A., Prusti T., de Bruijne J. H. J., Babusiaux C., Bailer-Jones C. A. L., 2018, *A&A*, 616, A1
- García-Moreno D., et al., 2026, *Astron. Astrophys.*, 706, A377
- Giacconi R., Zirm A., Wang J., Rosati P., Gilli R., Nonino M., Czoske O., 2001, *ApJ*, 551, 624
- Hirano S., Sakai Y., Umeda H., 2026, *MNRAS*
- Ho M., et al., 2024, *Open J. Astrophys.*, 7
- Hornschemeier A. E., Brandt W. N., Alexander D. M., Bauer F. E., Feigelson E. D., Garmire G. P., 2001, *ApJ*, 554, 742
- Inguglia G., 2026, *arXiv e-prints*
- Islam T., Wadekar D., Venumadhav T., Zaldarriaga M., Mehta A. K., Roulet J., Zackay B., 2026, *arXiv e-prints*
- Jin Z., Pasquato M., Davis B. L., et al., 2025, *ApJ*, 979, 212
- Kaalva K., Offner S. S. R., et al., 2026, *Astrophys. J.*
- Kass R. E., Raftery A. E., 1995, *J. Am. Stat. Assoc.*, 90, 773
- Kauffmann G., Heckman T. M., White S. D. M., Charlot S., Tremonti C., Brinchmann J., Brinkmann J., Peng C. Y., 2003, *MNRAS*, 341, 33
- Kumar M. S. N., et al., 2025, *A&A*, 703, A74
- Liang J., et al., 2026, *arXiv preprint*
- Lin H.-W., 2026, *arXiv preprint*
- Lotz J. M., Jonsson P., Cox T. J., Primack J. R., 2011, *ApJ*, 742, 103
- Lucente M., Pascoli S., Sala F., Zandi M., 2026, *arXiv e-prints*
- Mannucci F., Cresci G., Maiolino R., Marconi A., Gnerucci A., 2010, *MNRAS*, 408, 2115
- Mitchener M., Yiu H. W. H., Chang E., Bourdenx M., et al., 2025, *arXiv preprint*
- Mo H. J., White S. D. M., 1996, *MNRAS*, 282, 347
- Moss A., 2025, *arXiv preprint*
- Mucesh S., Hartley W. G., Palmese A., et al., 2024, *arXiv preprint*
- Nagasawa M., 1987, *Prog. Theor. Phys.*, 77, 635
- Nguyen N.-M., 2026, *arXiv e-prints*
- Pandya V., et al., 2026, *arXiv*
- Panopoulou G. V., Clark S. E., Hacar A., Heitsch F., Kainulainen J., Ntormousi E., Seifried D., Smith R. J., 2022, *A&A*, 657, L13
- Peng Y.-j., Lilly S. J., Kovač K., et al., 2010, *ApJ*, 721, 193
- Rawat S., et al., 2026, *arXiv preprint*
- Renzini A., Peng Y.-j., 2015, *ApJL*, 801, L29
- Ristorcelli I., Ferrière K., Juvela M., Montier L., Alina D., Montillaud J., 2026, *arXiv e-prints*
- Rivera-Ingraham A., et al., 2013, *ApJ*, 766, 85
- Rosen A. L., et al., 2026, *ApJ*
- Skilling J., 2004, in *AIP Conference Proceedings*. pp 395–405, doi:10.1063/1.1835238
- Somasekharan N., Pathak R., Dhanakoti M., Zhang T., Yue L., Zhu A., Pan S., 2026, *arXiv preprint arXiv:2605.06607*
- Spirtes P., Glymour C., Scheines R., 2000, *Causation, Prediction, and Search*, 2nd edn. MIT Press, Cambridge, MA
- Steinhardt C. L., et al., 2026, *arXiv*
- Stone J. M., Tomida K., White C. J., Felker K. G., 2020, *ApJS*, 249, 4
- Suin V., et al., 2025, *Astronomy & Astrophysics*, arXiv:2505.02903
- Sutherland W., Saunders W., 1992, *MNRAS*, 259, 413
- Thomasson S., Motte F., et al., 2026, *Astron. Astrophys.*
- Tomizaka K., 2014, *ApJ*, 785, 24
- Tremonti C. A., Heckman T. M., Kauffmann G., et al., 2004, *ApJ*, 613, 898
- Tritsis A., et al., 2025, *A&A*
- Villaescusa-Navarro F., Bolliet B., Villanueva-Domingo P., Bayer A. E., et al., 2025, *arXiv preprint*
- Wang C., Zhang Y., Li Y., et al., 2025, *Commun. Eng.*, 4, 184
- Wang J.-W., et al., 2026a, *ApJ*
- Wang H., Han J., Fan W., Liu H., 2026b, *arXiv preprint*, arXiv:2604.16922
- White G. J., 2026, ASTRA: Autonomous System for Scientific Discovery in Astrophysics, <https://github.com/web3guru888/ASTRA>
- Whitworth D. J., et al., 2026, *Open J. Astrophys.*
- Xu L., Borrett T., 2026, *arXiv e-prints*
- Xu F., Zhang Q., Sanhueza P., Wang K., Liu H. B., Beuther H., et al., 2026, *Astronomy & Astrophysics*
- Yamada Y., et al., 2025, *arXiv preprint*
- Ye C., et al., 2025, *arXiv preprint*
- Zhang G., et al., 2026, *arXiv preprint*, arXiv:2604.14093