An Archival ALMA Search for Unresolved Spectral-Carrier Technosignatures toward Stars within 40 pc at 114–873 GHz

Glenn J. White[1,2] and Robin Dey[3]

ABSTRACT

Do nearby stars show narrowband, drifting artificial radio carriers in the ALMA archive, and how deep and complete can such a search be? We searched 405 fine-channel windows toward 61 systems within 40 pc, covering 47.7 GHz in 26 separate intervals between 114 and 873 GHz, comparing drift-corrected stellar spectra with nearby control positions; 1258 coarser windows constrain only unresolved emission. Of 49 windows reaching $T_\star \geq 5$, 23 coincide with catalogued molecular lines, 16 of them with further evidence of line emission. Given the adopted line mask and background model, the other 26 are consistent with chance, against 23.8 expected from control positions ($P = 0.36$), which allows up to 12 further crossings. For a carrier narrower than one channel, ninety per cent single-epoch recovery needs an equivalent isotropic radiated power with a median, over systems, of 1.3×10^{15} W, 63 times the Arecibo planetary radar's (Sheikh et al. 2025), and at best 3.7×10^{13} W, still 1.9 times Arecibo's; 25 of 57 systems reach 1×10^{15} W. Confirmation needs a repeat at a covered frequency, which 21 systems lack, so coverage, more than depth, limits the search. No carrier is detected or confirmed: a non-detection, not an exclusion.

Subject headings: Technosignatures, Search for extraterrestrial intelligence, Submillimeter astronomy, Radio spectroscopy, Debris disks

1. INTRODUCTION

The public ALMA archive holds thousands of hours of millimetre and submillimetre observations taken for other reasons, mostly studies of dust discs and planet formation. Many of those fields contain a nearby star. This paper searches them for a narrowband transmitter at the position of such a star. No such signal was found.

Searches for *technosignatures* – detectable imprints of technology, intended or not (Sheikh 2020; Vidal et al. 2026) – have examined many thousands of stars since the first proposals (Cocconi & Morrison 1959; Drake 1961), from Project Phoenix (Tarter 2001; Backus & Project Phoenix Team 2002) to the targeted, commensal and machine-learning-assisted surveys of today (Zhang et al. 2020; Czech et al. 2021; Li et al. 2026; Johnson et al. 2023; Choza et al. 2024; Morrison et al. 2023; Tremblay & Tingay 2024; Manunza et al. 2025; Ma et al. 2023; Jacobson-Bell et al. 2025; Pardo et al. 2025). Even so, the last published accounting of that effort covers only 6×10^{-18} of the space of transmitter properties (Wright et al. 2018), and essentially all of it lies below ∼30 GHz. Above that frequency the record is nearly empty. Steffes & DeBoer (1994) and Mauersberger et al. (1996) observed 40 and 17 nearby targets at 203 GHz, each in no more than 35 MHz of a single line. That is 1363 times less bandwidth than the union searched here, and neither search allowed for drift or tested whether a signal persisted. Of ALMA's receiver bands, only Band 3 has been searched (Mason et al. 2025).

There are good reasons to look at these frequencies. Interstellar dispersion and scintillation are negligible, and an aperture delivers far more gain per watt. The costs are real as well, however. System temperatures are higher, the atmosphere is opaque in places at both ends of the link, and the narrow beam (25 arcsec for the 12 m benchmark of §5.3) would have to be aimed at the Earth deliberately. Since nothing is known about such a transmitter, limits are stated here as an equivalent isotropic radiated power (EIRP), which needs no model of it. The targets were also chosen by other programmes for their own purposes, so the sample cannot bound the fraction of stars that host a transmitter (Wright et al. 2018; Margot et al. 2023; Sheikh et al. 2025) (§2.1).

The signal sought is an *unresolved spectral carrier*: emission confined to a few channels at the stellar position, present at more than one epoch, and allowed to drift in frequency as it must for a transmitter on a rotating, orbiting body (§3.1). Broadband leakage on a planetary scale (Garrett 2026) lies outside the search by construction. What separates an interesting event from a detection is that it recurs (Sheikh et al. 2021), and the survey is built around that test. Mason et al. (2025) identified the two obstacles particular to this band: drift rates scale with frequency, and the spectrum is crowded with molecular lines. The method therefore searches over drift, compares every event with a line catalogue, and looks for the event again in a separated epoch. As ALMA is an interferometer, the star can also be compared with control positions inside its own primary beam, which a single dish cannot resolve, and that comparison takes the place of the usual interference rejection (§3).

Experiments with different channel widths are compared here on the minimum detectable received flux, $F_{\rm min} = S_{\rm min}\,\delta\nu$, each at its own channel width. The comparison has to be made in this way because $F_{\rm min}$ rises as $\delta\nu^{1/2}$, and a limit holds only for a signal narrower than the channel that set it. The channels searched here span 7 orders of magnitude, from 2.79 Hz to 31.2 MHz. Figure 1 draws each published threshold beside its value rescaled to a common 5σ. The rescaling is an arith-

[1] School of Physical Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK; glenn.white@open.ac.uk
[2] RAL Space, STFC Rutherford Appleton Laboratory, Harwell Campus, Didcot, OX11 0QX, UK
[3] VBRL Holdings, 50/5 Huay Kaew Road, Chang Phueak, Chiang Mai, Thailand

metic convention, and it does not match the false-alarm probabilities of the different searches. The plotted values are in `figures/figure_data.csv` of the data archive (White & Dey 2026). Of the limits plotted in Fig. 1, only this survey's EIRP_{90} is calibrated with injected carriers. At a bare 5σ threshold Mason et al. (2025) reach a median $4.9 \times 10^{-23}\,\mathrm{W\,m^{-2}}$ in a 30.52 kHz channel, while the deepest window here outside the line mask reaches $1.2 \times 10^{-23}\,\mathrm{W\,m^{-2}}$ in 122.1 kHz, a factor 4 deeper, and the median window $3.1 \times 10^{-22}\,\mathrm{W\,m^{-2}}$ in 488 kHz, a factor 6 shallower. Channel for channel this survey is therefore the shallower of the two except in its finest windows. It is distinguished instead by its frequency range, and by its 61 systems within 40 pc, where the Band 3 search observed distant stars in 4 calibrator fields over 117 MHz.

2. THE SAMPLE

This search covers 1663 spectral windows toward 90 stars in 83 systems within 40 pc, where a *system* is a bound group, counted once however many of its stars were observed. The windows come from 407 public ALMA execution blocks, all put through the same pipeline: 4.1 TB of raw data, 302 h of telescope time and 1178 h summed over windows. We call this set the *primary census*. Every expectation in the paper, and every limit of §4.1 and Table F1, is computed over it and nothing else. It holds 49 threshold crossings (Table 2), all in quality-passing data, one of them toward HD 139084 B, which lies in a census window but has no catalogue row.

Some data were searched after the census was closed, and these are kept apart. A further 2 crossings lie in an *archival tail* of 8 blocks toward 5 census stars, and they enter no expectation. Nor do an archival extension, a reserved *hold-out* (Appendix B.1) and a control beyond 40 pc (§4.4). Table A1 accounts for every block in every set.

The census divides into two experiments by native channel width, since both the search statistic and the completeness calibration depend on it. *Class A*, the primary experiment, is the 405 fine-channel windows toward 66 stars in 61 systems. Of these, 396 are channelised finely enough for the searched drift to cross a channel (§3.1), and every sensitivity and limit quoted is Class A's unless stated otherwise. *Class B*, the other 1258 windows, supports a secondary search for an unresolved narrowband spectral excess. The channelisation of these windows is bimodal, so the division is unambiguous: the widest fine channel is 1.95 MHz and the narrowest coarse one 7.81 MHz, with nothing in between.

A star enters the sample if its Gaia DR3 parallax is at least 25 mas and measured to better than 20 per cent. In practice the second condition never binds, since the least precise parallax among the stars searched has an uncertainty of 2.5 per cent. Its proper-motion-corrected position must also fall inside the quoted field of view of a public execution block, though not necessarily at the pointing centre, and its distance is taken as $1/\varpi$. A window is kept only where the primary-beam response at the star is at least 0.22. At that level the adopted Gaussian beam and a blocked-aperture response part by 10 per cent, and no window falls below it (Appendix A.2). One serendipitous field star is named throughout after the ALMA source at the centre of its field, ALMA J153702653−33192492. It is 2MASS




FIG. 1.— Where this survey sits among published radio technosignature searches (§1). **(a)** Minimum detectable EIRP against distance; per star, this survey's deepest carrier-search window (blue) or, with coarse channels only, its deepest coarse one (grey); filled, the bare threshold; open, EIRP_{90}. **(b)** Native channel width against sky frequency (filled, Class A; open, Class B).

J15370316−3319273 (Gaia DR3 6015967739554000768), an M5 dwarf at 35.0 pc with no catalogued radial velocity.

The Gaia requirement has a cost at the bright end, since a star with no Gaia DR3 parallax cannot enter the sample. It removes all 22 stars in this volume brighter than second magnitude, and 257 of its 4227 Hipparcos stars. Of these, seven (Sirius A, α Centauri A, Vega, Fomalhaut, α Centauri B, β Leonis and MCC 124) lie in public fields, with 38 fine-channel and 374 coarse-channel windows that a search admitting Hipparcos astrometry could add.

2.1. *The archival selection*

This sample was not designed. It is whatever ALMA programmes with other aims happened to observe among the stars Gaia DR3 catalogues, and 90 per cent of the stars searched were targeted under a debris-disc or planet-formation proposal. Table 1 measures the resulting bias against the volume-limited Gaia DR3 census of which the searched stars are a subset ($\varpi > 0$, $\sigma_\varpi/\varpi \leq 0.1$, $1/\varpi \leq 40$ pc; 17 566 objects). Both columns are classified by BP−RP colour, with absolute G magnitude separating the degenerates.

Distance is the dominant selection effect. Stars within 5 pc are over-represented by a factor 53, and the excess declines monotonically to a factor of two *under*-representation beyond 30 pc. Colour comes second. A stars are over-represented by about 17 and F and G stars by factors of three, because those are the stars with bright debris discs. For the A stars that factor is a floor, since the Gaia requirement itself removes stars as blue as these at 3.7 times its rate among the others. M dwarfs, on the other hand, are depleted. They make up 41 of the 80 non-degenerate stars among the 90 searched, a factor 0.7 of their share of the census, and only 28 of them carry fine-channel coverage. Coverage is also concentrated, with 285 of the 1663 windows coming from 3 systems.

The selection acts hardest on the targets a designed search would pick. Of the 16 sample systems within 15 pc with a confirmed planet, 6 have no Class A coverage and so no drift-resolving limit. No statement in this paper is therefore an occurrence rate for the solar neighbourhood.

TABLE 1
Selection function of the 90 searched stars against the volume-limited Gaia DR3 census of §2.1. "Ratio": the searched over the census column fraction.

Property	Census	Searched	Ratio
Distance			
$d \leq 5$ pc	48 (0.3%)	13 (14.4%)	52.9
5–10 pc	240 (1.4%)	13 (14.4%)	10.6
10–20 pc	2069 (11.8%)	18 (20.0%)	1.7
20–30 pc	5391 (30.7%)	19 (21.1%)	0.7
30–40 pc	9818 (55.9%)	27 (30.0%)	0.5
Colour class[a]			
A	61 (0.4%)	5 (6.2%)	17.2
F	473 (2.8%)	8 (10.0%)	3.6
G	978 (5.8%)	15 (18.8%)	3.2
K	2221 (13.2%)	11 (13.8%)	1.0
M	13 082 (77.8%)	41 (51.2%)	0.7
Other axes			
Confirmed planet hosts[b]	26 (15.5%)	24 (26.7%)	1.7
White dwarfs[a]	463 (2.7%)	8 (9.1%)	3.4
Searched stars / systems	–	90 / 83	–

[a] Colour bins, not MK types: BP−RP < 0.36 for A, 0.36–0.72 for F, 0.72–0.95 for G, 0.95–1.77 for K and > 1.77 for M, where the census's own temperature–colour relation crosses 7500, 6000, 5300 and 3900 K. Degenerate: more than 4 mag below the census main sequence. The five classes are normalised over the 80 non-degenerate searched stars and the 16 815 non-degenerate census objects, the degenerate row over the 88 classified searched stars and the 17 278 classified census objects.

[b] Both columns are the same positional join, of the 168-entry ALMA-covered work list and of its searched subset, to the NASA Exoplanet Archive `pscomppars` table accessed 2026 October 4. The 24 searched hosts carry 51 confirmed planets, 22 of the 24 surviving the deuterium-burning mass limit; the Gaia census carries no planet information, so the census column in this row is the work list and not the 17 566-object Gaia census.

2.2. *Frequency coverage and archival scope*

The Class A experiment covers $\Delta\nu_A = 47.7$ GHz of unique sky frequency, in 26 disjoint islands between 114.0 and 873.1 GHz rather than a continuous band. With the coarse-channel windows the coverage is 118.1 GHz in 46 intervals, of which 113.1 GHz survives the molecular-line mask of §3.2. Fig. 2 shows how that coverage is distributed. A transmitter between the islands was never observable here, whatever its power. Nor is the archive exhausted: 177 further blocks inside the same query were unprocessed when the census was frozen, 18 of them because the archive holds their raw visibilities but never pipeline-calibrated them (Table A1).

3. DATA AND METHODOLOGY

A carrier from an accelerating transmitter drifts in sky frequency, and over a long track its power spreads across many channels. The analysis therefore *de-drifts* the data, shifting each integration's spectrum to follow a trial drift rate before adding the integrations together. This section describes that search, how its output is turned into a power, and the three steps through which every event then passes.

3.1. *The de-drifted search statistic*

The search statistic is the largest stacked signal-to-noise ratio anywhere on the channel×drift grid of a window's dynamic spectrum,

$$T(\mathbf{x}) = \max_{\dot\nu,\,\nu} \frac{\sum_t I(\mathbf{x}, t, \nu + \dot\nu t)\,/\sigma^2(t, \nu + \dot\nu t)}{\left[\sum_t 1/\sigma^2(t, \nu + \dot\nu t)\right]^{1/2}}, \qquad (1)$$

where $I(\mathbf{x}, t, \nu)$ is the spectrum at sky position $\mathbf{x}$ in integration t and channel ν, $\dot\nu$ is the trial drift rate, and $\sigma(t, \nu)$ is the noise scale, both evaluated at the channel the track occupies in that integration. $T_\star$ denotes the value at the star. The time t runs from the first integration, so the frequency that labels a track is the one a carrier would occupy at the start of the block, and a repeat epoch is read at the cell the same convention fixes (§4.4).

Equation 1 is the matched filter, and T an absolute signal-to-noise ratio, only if the noise is independent between integrations and channels. The analysis does not assume that. The trigger is instead set against the distribution T actually takes in the data, measured in every window at 512 control positions on the same visibilities and through the same stack. A value of T is then converted into a power with injected carriers recovered through the unchanged pipeline (§4.1). The controls lie,




FIG. 2.— **Where this survey searched, and where a repeat could test it**: independent systems covering each sky frequency, one panel per populated ALMA band, its width proportional to the band's frequency share; curves step at the searched window edges. *(a)* Every window: blue, the fine-channel carrier search (Class A); grey, the coarse-channel spectral-excess search (Class B); the summed window bandwidth is 24.1× the union. *(b)* Class A: detectable once (grey); with a *separated* block (§3.2) covering the frequency (blue); beyond a year (dark). An accelerating carrier further needs a predicted frequency within the repeat coverage, with adequate sensitivity. *(c)* Class A systems recovering a carrier of that EIRP nine times in ten in one epoch; dashed, 1×10^{15} W.

from a fixed seed, in an annulus set by each block's synthesised and primary beams. The noise scale σ is their median absolute deviation at each integration and channel, scaled to a Gaussian-equivalent standard deviation, and the statistic is evaluated in the same way at the star and at each control. The field read over is the whole primary beam, that is every `FIELD` entry in which the star's primary-beam response reaches 0.05.

Spectra are extracted at the star's apparent position, never from images. For each integration and native channel the pipeline forms the weighted vector average of the phase-shifted visibilities in total intensity, $I = (XX + YY)/2$. That is the dirty-map value at the star, an amplitude and not a deconvolved flux density, so it is not a point-source measurement where the field holds extended emission. The correction for the star's offset from the phase centre is measured with injected carriers (Appendix A.2). The deposited extractor propagates each position to the observation epoch for proper motion and annual parallax. Some products were made by the first forms of the extractor, which propagated proper motion only, and for those windows the parallax loss is corrected window by window in §4.1. Orbital motion is bounded for every component of a multiple in Appendix A.2. The data are the archive's deliveries. Of the 178 blocks that set a quoted limit or hold a crossing, 142 were recalibrated with ALMA's final flags. They are not the same blocks as the 142 that the recurrence test reads. The other 36 keep the delivered calibration, 25 of them setting a limit and 15 holding a crossing. The change that the recalibration makes at the control positions bounds the chance expectation of this mixed calibration at 23.5–24.1.

Every delivered channel is searched, without regridding, deconvolution or edge trimming. A per-integration median along frequency removes the continuum, and none is taken along time, so no feature narrow against the band can be removed. The delivered channels are smoothed, so a correction computed from the correlator's own channel shape turns flux density times channel width into a radiated power (Appendix A).

The search itself runs on the topocentric sky frequencies ALMA delivered, and every drift rate quoted is the apparent one at the telescope. Two later steps need the star's own frame: comparing a crossing with the line catalogue, and reading a repeat epoch at the *predicted cell*, the single channel and drift rate where the discovery's own fit puts a carrier. Both correct each block's frequency axis for the Earth's motion and then for the star's systemic velocity. That is the SIMBAD radial velocity (Wenger et al. 2000), queried on 2026 October 4, or zero where none exists, as it is, among the stars with a crossing, for ALMA J153702653−33192492 alone. Toward β Pic the value of the ALMA CO analysis of its disc (Matrà et al. 2017) is adopted.

Two power quantities follow from the statistic, and both are defined in Table 2. The trigger level $P_{\rm trig}$ is an internal pipeline level. The sensitivity EIRP_{90} is measured by injecting synthetic carriers of known amplitude into the calibrated visibilities of 49 windows and recovering them through the unmodified pipeline. Every limit quoted is the EIRP_{90} of its own window, given

beside its native channel width, except the labelled bare-threshold comparisons of §1, and its error budget is Table 3. The amplitude scale is checked twice. An independent reduction of the same AU Mic visibilities is reproduced to 0.994 ± 0.014, which tests the extraction and not ALMA's flux transfer. The quiescent flux density of ϵ Eridani (Burton et al. 2022) is recovered to 1.10 ± 0.31, which measures the scale to 28 per cent. That is consistent with ALMA's own accuracy specification, ±11 per cent for the sample, which we adopt for the absolute scale (Table 3).

The drift range is bounded by an acceleration and not by a frequency, since drift rate and line-of-sight acceleration are one quantity in two sets of units,

$$\frac{\dot{\nu}}{\nu} = \frac{a_{\rm los}}{c} \qquad \text{and so} \qquad a_{\rm los} = \frac{c\,\dot{\nu}}{\nu}, \tag{2}$$

and a ceiling on $\dot{\nu}/\nu$ is then the same physical limit in every band. The ceiling is $12.0\,\mathrm{Hz\,s^{-1}\,GHz^{-1}}$, or $3.60\,\mathrm{m\,s^{-2}}$ by Equation 2, in 400 of the 405 windows, and $13.3\,\mathrm{Hz\,s^{-1}\,GHz^{-1}}$ ($4.00\,\mathrm{m\,s^{-2}}$) on the 5 TRAPPIST–1 windows, whose innermost planet demands more. The default is roughly the acceleration of a transmitter orbiting a solar-type star at a few hundredths of an au. It is the Mason et al. (2025) fiducial with a threefold margin, and 6 per cent of the 200 nHz that Sheikh et al. (2019) recommend for a general search. The grid is uniform in $\dot{\nu}$ and runs from the negative ceiling to the first trial at or beyond the positive one (Appendix D.5), in steps of half a native channel over the span of the integrations searched. A carrier midway between two trials therefore loses 0.25 of a channel over that span, and at most 0.29 over the longest on-source time.

The survey does not constrain a transmitter whose apparent acceleration exceeds that range, whether it is chirped faster (a chirp is not distinguished from acceleration) or its orbital and rotational accelerations together exceed it. Transiting TRAPPIST-1 b is the most demanding known case. At the default ceiling 29 per cent of its orbital phase would lie out of reach, and the widened grid used on its star matches its maximum acceleration only to within the ±3.1 per cent to which its host's mass and orbit are known.

Searching over drift also only helps if the largest searched drift moves a carrier across about one channel while the star is observed, $\eta_{\rm drift} = |\dot{\nu}_{\rm max}|\tau_{\rm on}/\Delta\nu_{\rm ch} \gtrsim 1$, where $\tau_{\rm on}$ is the time on source. 396 of the 405 Class A windows satisfy it. The other 9 fall below it because their tracks are short, and the per-window catalogue carries $\eta_{\rm drift}$ for every window.

3.2. *From a crossing to a confirmed signal*

Every window passes through three steps in the same order: the trigger, the line attribution and the recurrence test. They yield in turn a *crossing*, an *unattributed crossing* and, had one survived, a *confirmed signal*, each defined once in Table 2. Every count below is read from the ledger of every crossing the search produced, and is of primary crossings, one to a window.

The trigger. The level set on $T_\star$ carries no tail probability. T has not been shown to follow any standard distribution under the null hypothesis, and a window holds between 112 and 6.87×10^6 channel×drift cells, so one level buys very different protection in different windows. Every cell that reaches the trigger is recorded, and not only the largest, because a carrier can be hidden as effectively by a noise excursion as by a line.

The background. With no tail probability attached, a fixed threshold is not a fixed false-alarm rate either. The number of crossings expected by chance is instead computed window by window, from the rate at which that window's own control positions exceed the trigger. This assumes that a star behaves like a control position in its own field, which the reserved hold-out blocks test (§4.2.1). The census's 49 crossings, all of them Class A, are the baseline every expectation is set against. Of these, 1 lies in a window whose control ensemble fails to behave like noise (Appendix D.2), and it is counted but withheld from the amplitude-dependent tests (Appendix D).

The line attribution. At these frequencies many crossings will be molecular lines, so each one is compared with a line catalogue. The comparison is a single database query, made on 2026 October 6 and repeatable as it stands. A crossing within $\pm 50\,\mathrm{km\,s^{-1}}$ of a catalogued transition in the star's rest frame is *attributed*. The half-width follows the Keplerian line-of-sight velocities of gas surviving beyond a few au, with a margin for the systemic velocity. One fixed list of 12 species is searched, whatever a window was tuned to. It was completed at that query, after the crossings had been examined (Appendix C), and the chance comparison of §4.2.1 is therefore conditional on the mask as adopted. Of the 51 crossings of the census and the archival tail, the query attributes 23 and leaves 28. Halving the half-width to $\pm 25\,\mathrm{km\,s^{-1}}$ gives 21 and 30 respectively. An attribution is a coincidence with a catalogue, not an identification or a disposal, so every crossing goes on to the recurrence test whichever side of the mask it falls. The price of the mask is that the survey constrains nothing inside it.

The recurrence test. A repeat counts only if it comes from a *separated* block, which here means a block of another observing session and more than 1 day from the discovery. The session matters as much as the interval, because blocks of one session share calibrators, atmosphere, correlator setup and interference. Two readings are made. The *line-recurrence check* reads the predicted cell, ± 0.11 to $\pm 0.63\,\mathrm{km\,s^{-1}}$ wide, in a separated block, and the *trajectory comparison* asks whether one bounded trajectory inside the drift grid joins two crossings. Table 2 states what each reading can establish, and Fig. 2(b) shows the frequencies at which a separated block exists at all.

Whether the chain finds what it is meant to find is tested with the injected carriers of §4.1, and with β Pictoris CO, a real feature at a known stellar position, which is recovered and localised in two transitions and 10 execution blocks (§4.3).

4. RESULTS

The primary experiment is the Class A search for a drifting, unresolved carrier: 405 fine-channel windows toward 61 systems, 396 of them channelised finely enough to tell a drifting carrier from a stationary one (§3.1). The secondary Class B search looks for an unresolved spectral excess, with no discrimination by drift, in 1258 coarse windows. Together they make up the 1663 windows of

TABLE 2
Operational definitions, used in this sense throughout. A crossing is a single-epoch event; a confirmed signal needs a second, separated epoch, which no sensitivity supplies.

Term	Definition
Crossing	A channel × drift cell at the stellar position reaching $T_\star \geq 5$ in one execution block; neighbouring channels above the trigger belong to one crossing. The largest in a window is its *primary* crossing, any other a *secondary* one. An event requiring investigation; no claimed significance.
Unattributed crossing	A crossing more than $50\,\mathrm{km\,s^{-1}}$ in the star's frame from every transition of the line catalogue. A target for follow-up; not a detection.
Confirmed signal	The same carrier found again in a separated block (§3.2), either at the cell the discovery predicts in the star's frame (the *line-recurrence check*) or in an additional separated epoch at the frequency an independently specified trajectory predicts, with adequate sensitivity there. The check recovers only an emitter at rest in the star's frame, as a line is, so a null there never excludes a drifting or re-tuned carrier. A pair of crossings that one bounded trajectory inside the drift grid joins is *open*, not confirmed: the joining test can reject a pair and cannot accept one (Table 4). For 21 Class A systems this archive holds no separated block at the frequency, so this test can confirm nothing there, whatever the sensitivity.
P_{trig}	$4\pi d^2 S\,\Delta\nu$, the power at distance d of a carrier that makes $T_\star$ reach $T_{\mathrm{trig}} = 5$ in one native channel of width $\Delta\nu$: an internal pipeline level, not a sensitivity.
EIRP_{90}	The power at which nine injected carriers in ten produce a crossing at the star: a measured single-epoch trigger completeness, not a detection probability. One class factor, calibrated by injection in 49 windows, is applied to every window, interpolated within the injected configurations for 299 windows and extrapolated beyond them for 57 (§4.1); each injected window's own point (×0.81–×1.12) is in the data archive. It excludes atmospheric decorrelation (Table 3).
Class A	The primary experiment, defined by channel width: 405 fine-channel windows, median 488.3 kHz, toward 61 systems over 47.7 GHz of union bandwidth, 396 of which satisfy $\eta_{\mathrm{drift}} \gtrsim 1$ (§3.1).
Class B	The secondary experiment: 1258 coarse-channel windows, mode 15.625 MHz, constraining an unresolved narrowband spectral excess. Never a carrier search: in 1216 of them no searched drift leaves one channel, and in the other 42 the whole searched range moves a carrier 1.02–1.43 channels, which registers one but does not discriminate it.

the primary census.

4.1. *Sensitivity*

The median system, on its best window, would have produced a threshold crossing from a carrier of 1.3×10^{15} W. This median, and the counts that follow, are taken over the 57 systems whose limits do not depend on an unchecked position, and the three that do are described at the end of this section. Over these systems the limits run from 3.7×10^{13} to 2.5×10^{16} W (Fig. 3), and 25 of them reach 10^{15} W. A single Class A window is, of course, less deep than the best window of a system, and the median window reaches 2.5×10^{15} W. Each limit is taken over the channels that the line mask leaves free in the star's own rest frame (§3.2), which matters most for the finest windows. These were tuned by the observers to CO, and a 15.26 kHz window spans only 53.9 MHz, less than the 76.9 MHz the mask removes about the line, so 15 of the Class A windows lie wholly inside it and constrain nothing. HIP 10679 has no limit for this reason. Using only the 396 windows fine enough to resolve a drift leaves a system's limit unchanged in the median.

The limits come from injection. Carriers were injected at a random sub-channel phase and with a drift drawn from each window's own grid, and recovered through the unchanged pipeline as far as the localisation at the star. Nine in ten come back at $2.74\,P_{\mathrm{trig}}$ in a Class A window without online spectral averaging, and at $1.90\,P_{\mathrm{trig}}$ with two-channel averaging, where P_{trig} is the bare 5σ of the window itself; that power is EIRP_{90}. The factor shows no trend with the brightness of the control ring, so one value can serve the whole class, and each window's own channel response is carried by injections into the extraction that the search used (Appendix A). The spatial rank against the 512 control positions is not charged against these limits. Charging it would cost a further factor 1.27, and in 3 windows with a bright control ring the rank could not be met at any injected power. Class B has its own factor, $3.65\,P_{\mathrm{trig}}$ (1.71 with two-channel averaging), measured with 5 936 carriers injected into 35 coarse windows. The calibration of the class factor is uncertain by ×0.89–×1.11 (Table 3). That interval applies to the median and to the counts rather than to any one window, and it places the median system between 1.1×10^{15} and 1.4×10^{15} W, so the second figure of the median is set by the calibration.

The largest caveat on these limits is that most of them are transferred rather than measured. Carriers were injected directly into 49 of the 405 Class A windows, 12 per cent, and the result was carried across to the other 356. The injected windows were whole execution blocks, drawn at random and balanced by band and array, and they resemble the population well: their medians match to within 7.1 per cent on five of the 6 properties compared, and differ by ×1.9 on the sixth, the noise level (Appendix F). Only the ninety per cent point, in units of each window's own trigger power, is carried across. Among the injected windows that point ranges from ×0.81 to ×1.12 of the adopted factor, with 40 of the 49 within a tenth of it. This dispersion, and not the calibration interval, measures how uncertain the limit of one uninjected window is. Most of the transferred windows, 299 of the 356, lie inside the injected range on all five of the properties that set the reading, and the other 57 lie outside it on at least one, where that range is not an empirical bound. In all, 6 systems take their headline limit from one of them, and none of them is among the 25 reaching 10^{15} W. Only 7 systems had their best window injected directly, and these are marked in Table F1. For the others the headline limit is an estimate carrying

that dispersion, and so is the median. We have tried to see how much this matters. Drawing each transferred window's factor at random from the measured ones, once per execution block over 16 000 trials, lowers the median by 2 per cent and leaves 25–26 systems at 10^{15} W in nine trials in ten. Even applied at its extreme to every transferred window, the spread could move 3 systems across that level, 1 out and 2 in. Each system's probability of crossing it is given in the data archive.

Two further effects lie outside the injections. The first is atmospheric decorrelation, which a real carrier suffers and an injected one does not, so every limit is nominal in this respect. Estimated from each block's phase-calibrator phase rms (Cortes et al. 2026, §§10.4.11, 11.3.1) with the phase-scatter relation of Richards et al. (2022, Eq. 3), it makes the median system's best window optimistic by 3 per cent. For most systems this is small, but it exceeds a tenth for 13 of them and reaches 200 per cent for HD 172555. Divided out, it would move the median limit by 4.4 per cent and leave 25 systems at 10^{15} W. It is stated rather than applied because only 279 of the 407 blocks record the phase rms, and the others, estimated from blocks matched in baseline, are measurably the harder ones. Table F1 gives it system by system, and the adjusted limits are in the data archive. The channel response, online spectral averaging included, is by contrast measured and applied. The second effect is the position of the star itself. For Barta 161 12, HD 155555C and LHS 1140 the astrometric solutions are disturbed, and there is neither a companion nor a Hipparcos epoch with which to correct them (Appendix A), so the amplitude lost at the adopted position has no bound. Their limits in Table F1 hold only if that position is right, and these three systems are left out of the median, the count at 10^{15} W and Fig. 3.

A carrier that is not always radiating is penalised in proportion. In 3 888 undrifted trials down to a 10 per cent duty cycle the recovered amplitude followed the duty cycle to within 1 per cent, so a transmitter that is on for a tenth of the time would need 10× the power quoted. The drifting, intermittent case has not been measured.

Two of the nearest stars appear only in the secondary search. Barnard's Star and Wolf 359, the second- and third-nearest stellar systems once brown dwarfs are set aside, were not targets of the 203 GHz searches of Steffes & DeBoer (1994) and Mauersberger et al. (1996), and have otherwise been searched at centimetre wavelengths (Enriquez et al. 2017; Price et al. 2020; Margot et al. 2023). The archive holds only coarse Band 6 windows toward them, so their values, 2.5×10^{14}–3.4×10^{14} W over 12 windows and 2.7×10^{14}–3.9×10^{14} W over 12, are limits on an unresolved narrowband spectral excess, and to our knowledge the first at millimetre or submillimetre wavelengths for either star.

No Class B window that passes the quality criterion holds a crossing, which is what chance predicts. Computed exactly as for Class A, the 1254 windows expect 0.11 crossings from their own control positions ($P(0) = 0.89$). The expectation is small because a coarse window holds a median of only 256 independent cells, against 193 200 in Class A.

4.2. *Threshold crossings*




FIG. 3.— **Single-epoch trigger completeness of the carrier search**: the cumulative number of systems whose best window reaches a given EIRP_{90}, decorrelation excluded, the three systems of unbounded position left out and the calibration interval of Table 3 shaded. Confirmation needs a separated repeat, which no power buys and which 21 of the 61 systems lack (Fig. 2b; §5.3).

Of the 1663 windows of the primary census, 49 hold a cell reaching $T \geq 5$ at the stellar position, all in data that pass the quality criterion of Appendix D.2. The window flag of that appendix reaches 5 windows in 2 blocks, and the 1 crossing it reaches, toward HD 48370, is withheld from the tests that depend on amplitude. The $\pm 50\,\mathrm{km\,s^{-1}}$ line mask places 23 of the 49 within that velocity of a catalogued transition in the star's own rest frame (Appendix C). Those 23 are attributed, 16 with corroborating evidence of line emission and 7 on the coincidence alone, and 26 are unattributed. None was identified as interference by the available diagnostics (Appendix E). The archival tail adds two more, both unattributed and both toward HD 14055, in windows that no expectation covers (Table A1). They are in the ledger, but in none of the counts above. The correlator's frame suggests that the attributions reflect line emission, and not merely the width of the mask, since in that frame the crossings crowd onto the transitions these setups were tuned to (Appendix E).

Of the 49, two outrank all 512 control positions of their own window, 1 of them unattributed, and neither is present again at the cell its discovery predicts (§4.4). Behind each window's largest cell, twenty-eight weaker cells also reach the trigger. Of these, six are unattributed, and none reaches the trigger in any covering block the recurrence test reads. No crossing survives the chain, and this survey reports no detection; the deposited crossing ledger lists every one.

4.2.1. *How many crossings chance predicts*

We first ask whether the unattributed crossings are more numerous than noise alone would give. Because the number of searched channel × drift cells in a window spans 4.8 decades, the expectation is built window by window over the 405 Class A windows of the census, from each field's own control positions. The same condition is applied on both sides, to a cell at the position that reaches the trigger and that the line mask does not attribute (Appendix D). The control positions predict 23.8 unattributed Class A crossings, and 26 are found. A count that large or larger arises by chance with

TABLE 3
What sets $EIRP_{90}$, and in which direction. The calibration interval applies to the class-average factor and so to any statistic over many windows; the spread between windows is an observed dispersion and is the statement about one window; the signed terms are biases and are not folded into an error bar (§4.1).

	value	what it does to the limit
Class-average completeness	×2.99	nine carriers in ten recovered at this multiple of a window's own trigger power. *Measured* in the 49 Class A windows injected directly and *estimated*, at this factor, in the other 356
Own channel response	×0.91, ×0.64	the ninety per cent point with the window's own response, online spectral averaging included, over that with the deposited profile, without and with two-channel averaging ($N = 1$, $N = 2$; ×0.68 at $N = 4$). *Measured* on identical carriers in 24 Class A windows and *applied*: a window's factor is ×2.74 or ×1.90
Uncertainty on that calibration (per cent), combined in quadrature		
The factor itself	±2	sampling error of the pooled value over the 49 injected windows
Absolute flux scale	±10.2	the observatory's own band-dependent specification (ALMA Proposer's Guide (Privon et al. 2026), with the Technical Handbook's quality-assurance criterion (Cortes et al. 2026) at Band 8, where the two differ): ±5 in Bands 1–5; ±10 in Bands 6 and 7; ±20 in Bands 8–10, weighted over this sample's bands
Distance and visibility scale	±4.5	the Gaia parallaxes, which enter as d^2, and the block-to-block visibility scale
Combined	$-11/+11$	×0.89–×1.11 on the factor, and so on the median limit and on the system counts; ±21 for the 14 Class A windows in Bands 8 and 10, at 492–493 and 849–873 GHz
Spread between windows. An observed dispersion, not an error bar.		
Window to window	×0.81–×1.12	the 49 injected windows resolve 2.41–3.36 $P_{\rm trig}$ individually, 40 of them within a tenth of the pooled value. This, and not the interval above, is what an estimated limit is uncertain by; it is wider, and it applies to 53 of the 60 systems
Applied window by window where a record allows it and bounded where none does, and so not combined with the above.		
Intra-integration smearing	×1.00–×1.05	a carrier sweeps through part of a channel within one integration and the injected ones do not: $\eta_{\rm smear}$ at the drift that sets a ninety per cent point, from each window's own integration length where it is recorded (1098 windows) and from the longest delivered anywhere where it is not (565)
Primary beam	×1.00–×4.5	the star is not at the pointing centre, so each limit is divided by the response at its own offset, on the full width the observatory documents. Nothing in the median window; the 12 windows furthest off-axis carry the departure of that form from a blocked circular aperture, up to 7.1 per cent, as a one-sided systematic
Annual parallax	×1.00–×2.06	only where a window's product predates the extractor that centres it on the star's apparent position: each such limit (121 Class A windows) is divided by the amplitude a compact source retains at its own displacement; on the 13 windows with no parallax record the loss is bounded (Appendix A.2)
Signed terms. A bias is not an error bar and is not added to one.		
Channel response	×2.286	one over the share of a channel-confined carrier the peak channel keeps. Carried by the injections, so already inside the completeness above, which is this factor times a margin of ×1.309 over the nominal trigger
Decorrelation	$+0/+200$	measured block by block from the phase rms the observatory records on each block's phase calibrator, which an injected carrier does not suffer and a real one does. One-sided, and a floor. A median +3 on the systems' best windows, +8 above 400 GHz, and more than +20 in 5 per cent of windows

probability 0.36.

Two conditions attach to that probability, and to the other probabilities built from the control positions here and in §4.4. The first is that the control positions are a fair model of the background at the stellar position. The searched data cannot test this, and a residual of calibration or imaging that raised the rate of crossings at the star without raising it at the controls would change the probabilities. A calibration error that varied with frequency is one such residual, since it would raise the star and not its continuum-free controls. To make a crossing on its own it would need 21.4 per cent of the star's continuum even in the least demanding window, but a smaller residual adding to the noise could still change the rate. As a sensitivity check, the stellar rate can instead be calibrated at the ratio 1.108 that the star shows over its controls just below the trigger, and the expectation is then 26.4, at $P = 0.6$. The second condition is the mask itself. Its species list was completed after the crossings had been examined, and SO was added after a crossing had been found on one of its lines (Appendix C), so the probabilities, and the allowance given below, are conditional on the mask as adopted. Without SO the same comparison gives 29 observed against 24.7 expected, at $P = 0.23$.

The expectation is not carried by a few noisy fields. Without the ten windows whose control positions trigger most often, the remaining 395 predict 23.7 against 26 observed, and in 12 of 12 subsets by band, array, channel width and field type the observed count lies within the central 95 per cent of its own predicted distribution.

Both sides of the comparison are read over the whole primary beam, the field of §3.1. The extraction's original rule was narrower, keeping only the one field entry whose phase centre lies nearest the star, which drops on-source time and real signal. Neither rule has an adjustable radius. Read over that narrower field on both sides, the comparison gives 29 against 23.2, at $P = 0.15$. Neither geometry was nominated as the primary before the counts were seen, and corrected for the 2 readings the smaller probability is 0.29, so neither departs from chance at the 5 per cent level. That correction allows for the choice of field and for nothing else. It does not cover the choice of mask, or any other choice made after the crossings were seen.

The comparison could not have rejected 12 further crossings against these control positions, or 16 on the narrower reading, and this allowance carries the same two conditions. It is a count of crossings, not of transmitting systems, and not an occurrence rate (Appendix D).

Stellar flares are excluded by a broadband prediction. A millimetre flare raises every channel of a window alike, so one able to make a single-channel crossing would appear as a 53–307σ continuum event in the same integra-

tions. The observed continuum statistic of the crossings, by contrast, has a median of −0.29, and the AU Mic flare of MacGregor et al. (2020), in a block that also holds a crossing, adds only 0.17σ to the single-channel statistic.

4.3. *β Pictoris: the positive control*

A search that finds nothing must show that it could have found something, and β Pictoris provides that test. Without being told the disc is there, the pipeline recovers its circumstellar carbon monoxide (Dent et al. 2014; Matrà et al. 2017) in two rotational transitions, in 10 execution blocks drawn from 6 observing sessions over 8.4 yr. In the star's rest frame, at each block's own epoch, all 10 crossings lie within $0.5\,\mathrm{km\,s^{-1}}$ of the stellar systemic velocity. The same emission is the positive control of the recurrence test of §4.4: every feature recurs in another execution block, and each transition in another session. Emission of this kind is what the line mask removes, including the CO(2→1) toward HD 48370 that Cataldi et al. (2023) report and flag as possible cloud contamination (Appendix C). The mask covers 12 species and costs 6.4 per cent of the Class A union bandwidth, and it is the single most productive test in the chain (§5.3).

4.4. *Candidates and recurrence*

The two field leaders. Two crossings have a stellar statistic above every one of the 512 control positions in their own window: 61 Vir at 346.3232 GHz and CP−72 2713 at 344.2965 GHz, both in Band 7 and in the star's own rest frame (Table 5; Fig. 4). The rank decides which crossings are examined first, and on its own it disposes of nothing (§3.2). If the stellar and control positions were exchangeable, the control ensembles would predict 0.50 unattributed census crossings leading their field by chance, and one or more with probability 0.39. In the reserved blocks 0 of 5 crossings lead their field, and beyond 40 pc 0 of 6. The crossing toward CP−72 2713 lies $-12.3\,\mathrm{km\,s^{-1}}$ from SO(8_8–7_7) in the star's frame, inside the mask, and so is attributed on that coincidence alone. A visibility-domain fit was made for 39 of the 49 census crossings, and for 33 of them it is consistent with a point source at the star. It rejects neither of these two. For comparison, injected sources in the bin they occupy return $\mathrm{Re}/\sigma = 6.58 \pm 1.97$.

Both stars have been observed again at the same stellar-frame frequency in a separated session. For 61 Vir the two separated blocks, the nearest 24.12 h from the discovery, hold a line-like emitter returning to the predicted cell below 0.67 of the discovery amplitude, while for CP−72 2713 the one separated block, 50.50 h from the discovery, is too shallow to test it ($T_{\mathrm{pers}} = 0.49$). Each carrier's own fitted drift would take it out of that half-channel cell (± 0.211 and $\pm 0.213\,\mathrm{km\,s^{-1}}$) in 77 and 119 s, so the repeat reading is the line-recurrence check of Table 2. At the nominal threshold-equivalent amplitude $5S/T_\star$ the separated blocks would recover such an emitter with calculated probability 0.97 and 0.00, conditional on that amplitude, and injected carriers measure 0.94 and 0.00. In Table 5, T_{rep} is one cell at one trial and may be negative, and the recovery chance in brackets, at the discovery amplitude, is biased high. The readings behind each bound are in `candidates/repeat_epoch_readings.csv` of the data archive (White & Dey 2026).

The predicted cell. The same check, applied to every crossing, reads the one cell each discovery predicts at its nearest separated repeat. It works on lines, since 13 of the 19 attributed crossings tested recur in another session, 39–1984 d away, and the 6 that do not are the faintest of them. On the 25 unattributed crossings whose cell could be read it returns nothing. The largest readings are 2.37 at the carried drift and 4.71 at any drift, against a trigger of 5. For 24 of the 25 with a bound, a line-like emitter returning to its cell is held below the discovery amplitude, to a median 0.40 of it, while 5 are too shallow to test at the nominal threshold-equivalent amplitude. A carrier, on the other hand, would not stay in that cell. The bounded crossings accelerate at 0.66 to $3.55\,\mathrm{m\,s^{-2}}$, or 0.63 to 3.52 in the star's frame. At that rate a carrier leaves its half-channel cell in 60 to 471 s, and each crossing's nearest covering block lies at least 221 times that residence time away. Carrying the fitted drift forward does not rescue it, since the drift's own resolution moves the predicted cell by a median 23 channels by the nearest repeat. No bound depends on the cell's registration, the anchor or the threshold (Appendices B.2 and D.3). Three further unattributed crossings have no separated repeat block and are not tested (ALMA J153702653−33192492 at 347.19 GHz and β Pic at 241.77 and 241.86 GHz). No crossing is confirmed in the sense of Table 2.

The widened window. A transmitter in orbit changes its line-of-sight velocity between epochs, so the broadest reading asks whether anything reaches the trigger, at any drift, anywhere in a covering window of a separated session. In the median case that window is $\pm 801\,\mathrm{km\,s^{-1}}$ wide. Of the 93 distinct covering windows, 11 hold a statistic above the trigger, against the 8.2 that each window's own control positions predict ($P = 0.20$; Appendix D). Every one of those windows holds a crossing the ledger already lists, so the comparison re-reads the trigger-level count of §4.2.1 rather than adding a recurrence to it. Of the 11, 8 are toward ALMA J153702653-33192492 and HD 14055, which are also the two most heavily observed objects in the survey: they hold 20 per cent of the Class A census and 49 of the 93 covering windows, so 5.7 of the 8.2 expected exceedances are theirs before anything is looked at. Over the remaining 44 windows the reading is 3 against 2.6, at $P = 0.48$.

The pair test. A carrier that moves might also be caught as two crossings toward one star, and Table 4 asks of the 9 pairs inside the widened window, across 4 stars, whether one transmitter could have produced both. Constant line-of-sight acceleration would require equal drift rates, and the fitted rates differ by more than the grid's resolution in 9 of the 9 pairs, so the question becomes a weaker one: whether any trajectory inside the searched grid joins the two. It separates 0 of the 9. Of the 9 it cannot separate, 9 are *open* and 0 *edge-limited*. An open pair is not a confirmation, since the comparison can reject an association but cannot accept one (§3.2). Chance alone, at each census window's own control rate, would put 3.8 open pairs in separated sessions against the 4 observed, with the pairs that have an archival-tail member set aside. Of random pairs, 91–94 per cent are joinable. As a test of a drifting carrier it is therefore incomplete. Counting archival-tail blocks, one star, HD 14055, has three separated crossings that one trajectory joins pairwise, and in the census alone none

does. However, no constant acceleration or circular orbit inside the grid joins all three. Predicted from the first two, the third misses a constant acceleration by at least 1139 resolutions, and the best circular orbit through all three misses by 56. A line that does not drift behaves as it should under the same test: the CO(2–1) line of HD 285968 is joined across 5.0 d to within 0.6 of the resolution. The accounting for each pair is `candidates/drift_pair_accounting.csv` of the data archive.

HD 14055 and ALMA J153702653−33192492. HD 14055 holds 4 of the exceedances against 4.7 expected ($P = 0.71$), so its concentration is its exposure. Its fields are ordinary, with control rings 1.04 times their class median against 1.00 for the survey, its setups are continuum ones, and its crossings lie 123–1069 km s^{-1} from the nearest masked transition in its own frame. The deepest readings of its crossings come from follow-up blocks outside the census. Over the 9 and 6 of them in separated sessions a line-like emitter at the predicted cell reads -0.7 ± 0.7 and 0.6 ± 1.0 mJy, against discovery amplitudes of 8.0 ± 1.6 and 10.2 ± 1.9 mJy, which bounds it at 0.37 and 0.57 of each.

ALMA J153702653−33192492 is a serendipitous dwarf 5.75 arcsec from the phase centre of fields pointed at the Lupus clouds, observed in 212 s snapshots. It holds 4 exceedances where 1.0 were expected, with $P = 0.013$ for this object alone. That probability, however, belongs to an object picked out because it is the most discrepant of the 16 objects compared, and in Poisson-binomial counts simulated at each object's own per-window control rates some object is that far out in 7 per cent of realisations. Its four exceedances lie on four crossings between 216 and 220 GHz, in two of this object's seven blocks, and their control rings are ordinary. Their fitted drifts lie 15–35 grid steps from zero, against an uncertainty measured by injection of at most 0.7 step. Because the snapshots are spread over a 3613 s track, $\eta_{\rm drift}$ is 19 over the track and 1.13–1.14 over the time on source. A visibility fit places two of the four at the star. A wider list of 225 species places a transition within 15 km s^{-1} of every crossing toward either object, as its 24 976 transitions guarantee, and so identifies nothing. We have no explanation for these crossings. ALMA J153702653−33192492 is a target for follow-up, not evidence of a non-astrophysical source, and a repeat observation should point at it, with a radial velocity and one deeper epoch at 216–220 GHz.

Blocks outside the census. Three sets of execution blocks enter no census statistic (Table A1), and each was searched with the same pipeline. The reserved hold-out, set aside by the identifier rule of Appendix B.1, has 315 windows toward 36 stars and holds 5 crossings, the strongest at $T_\star = 8.64$. Of the 4 tested for recurrence, only a molecular line is present again ($T_{\rm rep} = +8.47$). The archival extension's 463 windows in 115 blocks hold 10 crossings, 7 of them unattributed against 4.5 predicted by chance ($P = 0.17$). The control sample beyond 40 pc holds 6 crossings. Read in the sky frame it supports no limit, and shows only that the pipeline behaves in the same way beyond the sample.

4.5. *A stellar-frame stacked search*

Combining a star's epochs in its own rest frame deepens the limit, provided the emitter holds still in that frame. A frequency fixed in the stellar frame does not stay fixed at the telescope, since the barycentric term alone swings by a median 8.0 km s^{-1} between the epochs of one star, many channels at the 488 kHz Class A width. Each epoch is therefore transformed to the stellar rest frame, divided by its own primary-beam response and combined with the others by inverse variance over the channels of common coverage. The transform is checked on HD 285968's line, which 4 epochs observe 12.2 km s^{-1} apart and place at one stellar-frame frequency to 0.03 km s^{-1}. The criterion is applied in the same way at the stacked control positions, and every stacked limit is measured by injection into the stack.

Of the surveyed stars, 59 can be stacked, 5 of them with blocks beyond the census, and over 316 stacked groups the flux-density limit improves on each star's best single epoch by a median factor 1.50 and by 4.45 at most. A further 10 stars, all control stars beyond 40 pc, are stacked and left out of those figures. The gain is 0.79 of

TABLE 4
Can two crossings be one carrier? The 9 pairs toward one star with an unattributed member, and each HD 14055 triple (§4.4).

Star	blocks	new sess.	Δt (d)	$\dot\nu_{1\star}$ (Hz s^{-1})	$\dot\nu_{2\star}$	$\Delta\nu$ miss (MHz)	one? grid	
61 Vir	X822/X82f	yes	3.00	+1092	+3221	+755.4	15	open
J1537−3319	X10359/X398	yes	123	−2000	+2390	+223.5	3.6	open
HD 14055	X24be0/Xd46	yes	1.92	+1665	−3823	+512.7	100	open
	X2b2ee/X3835	yes	2.09	−2390	+2155	+395.5	55	open
	X3835/Xd526	yes	607	+2155	−2563	−640.4	–	open
	X2b2ee/Xd526	yes	609	−2390	−2563	−244.9	–	open
	X3835/X25c5e	yes	649	+2155	+2611	+338.3	–	open
	X2b2ee/X25c5e	yes	651	−2390	+2611	+733.8	–	open
β Pic	X3a90/Xa9df	no	1.02	−1611	−679	+85.5	132	untest.

Triple	constant acceleration	circular orbit
X2b2ee/X3835/Xd526	misses by 1148	misses by 133
X2b2ee/X3835/X25c5e	misses by 1139	misses by 56

New sess.: *yes* where the two blocks belong to different observing sessions. Starred rates are in the star's frame, the observatory's own acceleration removed (observatory frame: Appendix D.5). *Miss*: departure from constant acceleration, in units of the drift grid's resolution over Δt; a dash, a pair that resolution cannot separate. The verdict is set by the bounded-trajectory test, *yes* at a miss of 3 or less; *edge*, one rate on the grid's outermost trial, a lower bound; *untest.*, untestable: no block of another session covers the window about the pair. Last, each triple's miss of its third crossing, predicted from the first two, in resolutions.

TABLE 5
The two field leaders, the crossings whose stellar statistic exceeds all 512 control positions of their own window (§4.4).

	61 Vir	CP−72 2713
Frequency, stellar frame (GHz)	346.323236	344.296455
Drift rate (Hz s^{-1})	+3182	−2056
$T_\star$	5.947	5.809
Highest control T	5.58	5.68
Fitted amplitude (mJy)	118.6 ± 19.9	11.3 ± 2.0
Peak-channel EIRP (W)	5.0×10^{14}	8.9×10^{14}
Repeat epochs measured	2	1
Nearest separated epoch, separation (h)	−24.12	+50.50
T expected if persistent, $T_{\rm pers}$	8.14	0.49
T measured, matched drift, $T_{\rm rep}$	+0.42	+0.34
Amplitude still bounded, $(T_{\rm rep}+5)/T_{\rm pers}$	0.67 S	untested
Recovery chance Φ if the amplitude is $5S/T_\star$ (S)	0.97 (1.00)	0.00 (0.00)
Nearest masked transition	SO(9_8–8_7)	SO(8_8–7_7)
its offset $\Delta v_\star$ (km s^{-1})	−177.6	−12.3
Disposition	not confirmed	attributed




FIG. 4.— **The strongest unattributed crossing of the quality-passing data, and the control field it leads.** Frequencies are in the star's rest frame. *(a)* The dynamic spectrum at 61 Vir's position around the crossing at 346.3232 GHz, observed 2017 May 20, the fitted drift of $+3182\,\mathrm{Hz\,s^{-1}}$ dashed, 225 integrations binned into 9 and plotted in units of each bin's noise. *(b)* The same rows shifted by that drift. *(c)* The de-drifted stack at the star (black) and at the 8 control positions with retained dynamic spectra (grey), in units of its own noise, the trigger dashed. *(d)* The same statistic at each of the 512 rank-screen control positions: the star (solid) against the highest control (dotted).

what $\sqrt{N}$ equal epochs would give, the shortfall being unequal epoch depth. In the stack, 2 groups pass the full criterion and both are known lines, the β Pictoris carbon monoxide of §4.3 in two transitions, so the positive control passes in the stack as well, and the survey's own $\pm 50\,\mathrm{km\,s^{-1}}$ mask leaves none of them unattributed.

Every stacked limit holds only for an emitter steady in the stellar frame to within one of its own channels. No crossing of §4.4 drifts slowly enough to survive a fine-channel stack, which admits far less stellar-frame excursion than a coarse one, so the single-epoch limits of §4.1 are the ones that apply generally. The deepest stacked limit is Proxima Centauri's, 1.9×10^{13} W from 25 census blocks of coarse 15.625 MHz channels, and like every Class B value it is a limit on an unresolved narrowband spectral excess. It also needs a qualification of its own. A transmitter on Proxima Centauri b would swing over $94 \sin i\,\mathrm{km\,s^{-1}}$ in its orbit (Anglada-Escudé et al. 2016; Kervella et al. 2017; Akeson et al. 2013), some 5 channels at $\sin i = 1$. The stack would cancel such a signal unless the orbit is seen nearly face-on, with $\sin i \leq 0.22$, which holds for 2.5 per cent of random orientations. Within one block, by contrast, the planet moves at most 6 per cent of a channel, so the host's best census single-epoch limit, 7.3×10^{13} W in a Class B window, is the one that applies to it.

5. DISCUSSION

An archival search can only be as good as the observations it inherits, so we ask three questions of the result: what new parameter space the archive allowed us to search, what the absence of a detection actually constrains, and what stops the search from saying more. The answer to the last one also says what a dedicated experiment should do differently.

5.1. *What was searched*

The parameter space opened here is millimetre and submillimetre frequency toward the nearest stars. The archive gives 47.7 GHz of sky frequency between 114 and 873 GHz toward systems within 40 pc, whereas the only other search of nearby stars at these frequencies that covered more than a single line spanned 117 MHz of Band 3 (§1). Most of this range had therefore not been searched at all for narrowband emission from the direction of a nearby star.

The price of using an archive is that the channels were chosen by other observers, for other purposes. They range from 15.3 kHz to 31.25 MHz, while the centimetre-band carrier searches work with channels of about 2.8 Hz (Fig. 1b). A carrier much narrower than one of these channels is measured against the noise of the whole channel, and this is one reason why the limits reached here lie well above those of the centimetre surveys.

Union bandwidth also overstates what was searched, because different frequencies were observed toward different stars, and to very different depths. Figure 2c therefore counts, at each sky frequency and EIRP, the systems in which a carrier would have been recovered nine times in ten. At 1×10^{15} W the covered frequency falls to 19.1 GHz, 40 per cent of the union, and 25 systems are reached at some frequency, 18 of them near 230 GHz. However high the power, no frequency is covered toward more than 38 of the 61 systems, and half of the sample is reached only above 2.9×10^{15} W. The search is thus deep toward a few systems at a few frequencies, and shallow or absent almost everywhere else.

The 1258 coarse-channel windows add a constraint of a weaker kind. Their channels are too wide to follow a drift, so they limit only an unresolved narrowband spectral excess, but they are numerous, and stacked in the star's own rest frame they can be deep. Toward Proxima Centauri, the nearest star, the stacked limit reaches 1.9×10^{13} W for an emitter that stays within one channel in the stellar frame (§4.5).

5.2. *What the non-detection constrains*

Every limit in this paper applies to one kind of signal: an unresolved carrier at the stellar position, confined to a single 15.3–1953 kHz channel, lying outside the molecular-line mask, radiating while an archival block was on sky, and drifting linearly within its window's ceiling of 12.0–13.3 nHz, a fractional drift rate that corresponds to a line-of-sight acceleration of 3.60–$4.00\,\mathrm{m\,s^{-2}}$. That ceiling covers the orbital acceleration of every known planet in the sample except TRAPPIST–1 b, which is reached only as precisely as its host's mass and orbit are known (§3.1). A transmitter on a rapidly

rotating planet, however, uses most of the ceiling on the planet's spin.

A signal that departs from this model is constrained less, or not at all. Emission as wide as the widest line in these fields needs 29 times the power of a one-channel carrier to be found, and a carrier that is switched on for only part of a block is recovered at that fraction of its amplitude. Nothing is constrained inside the $\pm 50\,\mathrm{km\,s^{-1}}$ mask around each listed transition. Nor does the survey say anything about the stars within 40 pc that ALMA has not observed, which are almost all of the 17 566 that this volume holds.

There are also two blind spots of a more specific kind. The analysis forms (XX + YY)/2, which retains the full intensity of a circularly polarised carrier, so what is lost is not sensitivity but the discrimination that polarisation would offer against unpolarised astrophysical emission. ALMA's feeds are linear, and the circular signature lies in the cross-hand products, which this analysis does not extract. The other blind spot is historical. The 203 GHz positronium hyperfine line that Steffes & DeBoer (1994) and Mauersberger et al. (1996) looked for lies in ALMA Band 5, where this survey holds 4 coarse-channel windows, all toward α CMa B, and none of them covers it.

5.3. *What prevents stronger conclusions*

Depth is not the main limit. As an illustration, a 12 m dish radiating 1 MW at 230 GHz with an aperture efficiency of 0.7 has a directive gain $G = \eta(\pi D\nu/c)^2 = 5.9 \times 10^8$, and hence an EIRP of 5.9×10^{14} W. That is above the single-epoch limit of 20 of the 57 Class A systems that retain unmasked bandwidth and have a bounded position, so a transmitter of terrestrial scale would already have been within reach toward those systems. Its beam, however, has a solid angle $\Omega = 4\pi/G = 2.1 \times 10^{-8}$ sr and covers only 1.7×10^{-9} of the sky. EIRP is not transmitter power, and these limits apply to a beamed transmitter only if its beam happens to contain the Solar System. The archive is limited less by its depth than by the astrophysics that can imitate a carrier, and by the repeat coverage that it happens to hold.

At these wavelengths the false positive is line emission. A catalogued molecular transition lies within the mask for 23 of the 49 crossings in the census's quality-passing data. For 16 of them there is support beyond the frequency match: a second transition of the same species, catalogued gas on the sightline, or the crossing's own recurrence. The other seven rest on the coincidence alone, which is not an identification. None was identified as interference by the available diagnostics (Appendix E), which leaves 26 for the comparison with chance of §4.2.1, whose result is therefore conditional on the mask.

Molecular lines also bring a test of their own that has no centimetre analogue. A line that fills the primary beam raises the star and its control annulus together, whereas a carrier from the star raises only the star, and 10 of the 12 windows that reached this spatial stage hold disc or interstellar carbon monoxide (§4.3).

An archive supplies repeat epochs only by accident. A single crossing, however strong, becomes a testable claim only when the same frequency is observed again, and the archive holds such repeats where observers happened to return. We describe this by the same-frequency coverage $C_{\mathrm{conf}}(\nu)$, the number of systems toward which two separated blocks (Table 2) both cover the sky frequency ν. It measures coverage, and is not a completeness for an orbiting carrier, because a drifting carrier must also fall at its predicted frequency in the second block. Its support is 24.0 GHz, 50 per cent of the 47.7 GHz searched, and only 8.5 GHz beyond a year (Fig. 2b). Of the 61 systems, 40 have such a block and 13 have one more than a year away. For the other 21 the archive holds none, so that a crossing there could not have been confirmed at any power. Even where a repeat exists it can be too shallow to test anything, as it was for 5 of the 25 crossings read at a predicted cell.

5.4. *What a dedicated ALMA experiment should do differently*

Each of these limits has a remedy on the same telescope. A dedicated search can choose its targets and its frequencies instead of inheriting them. This archive favours dusty discs, which are the worst case for line confusion, and covers least the M dwarfs that host most of the nearby habitable-zone planets. Fine channels are needed in every window, since the 1258 coarse-channel windows of the archive cannot carry a carrier search at all. Tunings placed away from the bright molecular lines recover most of the 3.05 GHz of Class A coverage that the mask costs here, and probably remove many of the false positives as well. Above all, the recurrence observations have to be scheduled deliberately, at the same tuning and days to years apart, so that every crossing meets a separated epoch at the frequency it predicts. This is chiefly a matter of scheduling, and each additional epoch tests a trajectory predicted independently from the others. Indeed 27 of the 61 systems already carry three epochs at a common frequency, but none of them has a crossing seen in more than one of those epochs. The obvious first target for such a programme is ALMA J153702653−33192492, the most discrepant of the objects whose repeat windows were compared with their own control positions (§4.4).

6. CONCLUSIONS

This paper asked whether any of the nearby stars that ALMA has observed shows a narrowband, drifting radio carrier of the kind an artificial transmitter produces at millimetre and submillimetre wavelengths, and, if none is seen, how deep and how complete a search of the archive for such a carrier can be. To answer it we searched the archival ALMA spectra of 61 star systems within 40 pc. The fine-channel windows cover 47.7 GHz of sky frequency, in 26 separate intervals between 114 and 873 GHz, which is at least 1363 times the bandwidth of either of the millimetre searches of nearby stars that preceded it (Steffes & DeBoer 1994; Mauersberger et al. 1996). We removed the drift from each spectrum and compared it with control positions in the same field, so that the background at the star could be judged from the observation itself.

No transmitter was found. In 49 windows the stellar statistic reaches the threshold $T_\star \geq 5$. Of these, 23 coincide with catalogued molecular lines, 16 of them with further evidence of line emission, and none was identified as interference by the available diagnostics. The other 26 are consistent with chance: the control positions beside

each star predict 23.8, with $P = 0.36$ (0.29 for the alternative extraction field after allowing for the choice, neither significant at the 5 per cent level), which allows up to 12 further crossings. Both that probability and the allowance are conditional on the line mask as adopted, which was completed after the crossings had been examined, and on the control positions being a fair model of the background at the star. Wherever an unattributed crossing was observed again and the repeat could be read, it did not return at its predicted frequency, so no recurrence has been confirmed. This is a non-detection, not an exclusion.

Because nothing was detected, what the search delivers is a set of upper limits. They apply to a carrier narrower than one channel, lying outside the molecular-line mask, radiating while ALMA observed and drifting by no more than 12.0–13.3 nHz, the drift ceiling discussed in §5.2. For such a carrier, recovery nine times in ten in a single epoch needs an equivalent isotropic radiated power, or EIRP, with a median, over systems, of 1.3×10^{15} W in each system's most sensitive window. The deepest limit, 3.7×10^{13} W, is reached toward ϵ Eridani, and 25 of the 57 systems whose limits do not depend on an unchecked position reach 1×10^{15} W.

It helps to set these numbers against something we have built ourselves. The Arecibo planetary radar, one of the most powerful transmitters ever operated on the Earth, had an EIRP of about 2×10^{13} W at 2.38 GHz (Sheikh et al. 2025). The median limit found here is about 63 times that, and even the deepest is 1.9 times it, so no system reaches an Arecibo-like transmitter, although 10 come within a factor of ten of it. Put the other way about, for the median system an Arecibo-like transmitter pointed at the Earth would reach the limit only if the system were 3.4 pc from us, and even in the most favourable case, TWA 3A, only within 7.6 pc. A 12 m dish radiating 1 MW at 230 GHz is, however, a more natural transmitter at these frequencies, and its EIRP lies above the single-epoch limit of 20 of the systems. It could only be seen, of course, if its beam happened to point at us, as discussed in §5.3.

How complete the search can be is set more by the archive's coverage than by its depth. A crossing becomes a testable claim only when its frequency is observed again in a separated session, and 21 of the 61 systems were never re-observed in that way at a covered frequency, so that nothing found toward them could have been confirmed at any power. A programme designed for the purpose, on the same telescope, would use fine channels throughout, tune away from the bright molecular lines and schedule its repeat observations deliberately, so that any crossing it found could be looked for again at the frequency it predicts (§5.4).

ACKNOWLEDGEMENTS

This paper makes use of the following ALMA data: the 67 projects listed in Table A2, each cited as `ADS/JAO.ALMA#` followed by its project code. The 407 primary-census execution blocks searched span 2013 October 6 to 2025 July 11, and we thank the principal investigators and teams who proposed and obtained them. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), NSTC and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (Gaia Collaboration et al. 2023), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC). This research has made use of the SIMBAD database (Wenger et al. 2000) (CDS, Strasbourg, France); of the NASA Exoplanet Archive (Akeson et al. 2013), which is operated by the California Institute of Technology under contract with the National Aeronautics and Space Administration under the Exoplanet Exploration Program; and of the Splatalogue spectral-line database (Remijan et al. 2007).

Use of generative artificial intelligence. Large language model assistance (Anthropic Claude) was used in manuscript preparation of this complex paper. Data analysis was done using the Python based framework ASTRA (White 2026, https://arxiv.org/abs/2610.36144) to assist in the archival retrieval and calibration scripts, search and vetting pipeline, statistical tests, and the generators that produce every number, and carefully cross-checked and validated. The scientific content, analysis, and conclusions were designed, executed, and verified by the author. All errors and omissions are the responsibility of the author.

Testing and validation. No result here is accepted because a model reported it: every number in the text is written by the program that computes it from the data, and a separate audit re-derives the identities that must hold among the published values and refuses the paper where one does not. Every check in the analysis was first made to fail on deliberately wrong input before it was relied on.

Software. The pipeline depends on CASA 6.7.0 (CASA Team 2022), PYTHON 3 with ASTROPY (Astropy Collaboration et al. 2022), NUMPY (Harris et al. 2020) and SCIPY (Virtanen et al. 2020), and the ALMA Science Archive and SIMBAD TAP interfaces through ASTROQUERY (Ginsburg et al. 2019); all are free and open-source, at the versions cited.

Author contributions. G.J.W. conceived the survey, defined the target sample and the selection criteria, and wrote the manuscript; R.D. built and operated the archival retrieval, calibration and search pipeline, implemented the spatial control statistic and the injection–recovery campaigns, and produced the per-window catalogue and its regeneration scripts. Both authors designed the vetting procedure, chose the masked species, carried out the statistical validation reported in §3.1 and Appendix D, and agreed the dispositions of the crossing ledger.

Funding. This work received no dedicated funding. It used public archival data and no observing time was awarded for it.

Competing interests. The authors declare no competing interests. VBRL Holdings is a privately held company through which R.D. conducts independent research, named here solely as his affiliation; it had no role in the design of the survey, the selection of targets, the analysis, the decision to publish or the content of this paper, and provided no funding for the work.

DATA AVAILABILITY

Every observation used here is public in the ALMA Science Archive under the 67 project codes of Table A2, and is retrieved by the execution-block identifier, which the crossing ledger gives for each of the 51 threshold crossings (Chen et al. 2022); the astrometry is Gaia DR3, the planet counts come from the NASA Exoplanet Archive, and the line mask from one Splatalogue query of CDMS and JPL made on 2026-10-06. The data and software are held in a versioned Zenodo record (White & Dey 2026). It contains the per-window catalogue of all 1663 windows, the Class A and Class B limits with the injection trials behind them, the crossing ledger, the control-field statistics, the execution-block and session accounting (every row of Table A1), and the resampling inputs and seeds. From these one program, `code/reproduce.py`, rebuilds the chance expectation and its probability, the completeness factor, the sensitivity summaries and the repeat-window comparison, and checks each against the value printed here; the principal tables also accompany the arXiv version as ancillary files. The record holds the search pipeline as it ran, which takes a public execution block to the statistics of this paper. Some windows, in blocks that were not recalibrated, carry products made by the forms of the extractor and of the search that preceded the final ones. The final extractor places the phase centre at the star's parallax-displaced position and keeps every field entry, which the first forms did not, and the very first also drew its control annulus differently. Those versions are archived with their settings, together with a table that gives, for every window, the versions that made its products; run again on freshly calibrated 7 m and 12 m blocks, they reproduce those products in every window tested. The dynamic spectra at the star and at its 8 retained control positions, around each of the 51 crossings, form a second file of the same record (169 MB). The calibrated visibilities, of order a terabyte, and the per-window spectra were not kept, for lack of storage, so this execution of the search can be run again from the archive with the same code but cannot be repeated exactly.

APPENDIX

A. SPECTRAL RESPONSE, RETENTION AND THE SURVEY LEDGER

A.1. *The response of a channel to a carrier*

A carrier seldom falls at the centre of a channel. ALMA's online Hanning smoothing (Cortes et al. 2026) leaves 0.50 of an unresolved carrier's flux in the peak channel at a channel centre and 0.42 on a boundary, and this loss enters the limits once, through the injections. Each synthetic carrier is deposited at a sub-channel phase drawn afresh and recovered through the unmodified trigger, so the measured completeness already contains it. The Class A factor separates into two parts, $2.99 \simeq 2.286 \times 1.309$. The first is $1/0.4375$, from the phase-averaged share of a carrier's flux that the 0.25/0.5/0.25 kernel places in the peak channel. The second is the margin that a peak channel needs above the nominal trigger for nine carriers in ten to return. $P_{\rm trig}$ and $S_{\rm min}$ therefore stay nominal, comparable with other programmes' thresholds, and the sensitivity carries the correction,

$$\mathrm{EIRP}_{90} = \frac{\kappa\,\rho_N}{R_{\rm px}\,\eta_{\rm smear}}\,4\pi d^2 S_{\rm min}\,\Delta\nu, \qquad S_{\rm min} = \frac{5\sigma}{A}, \tag{A1}$$

where $4\pi d^2 S_{\rm min}\Delta\nu$ is $P_{\rm trig}$, κ is the class factor, ρ_N the response ratio of the window's averaging mode, and A, $R_{\rm px}$ and $\eta_{\rm smear}$ are the primary-beam, parallax and smearing retentions.

The kernel is an approximation. A delivered channel is the mean of N Hann-windowed raw channels ($N = 1$ in 290 Class A windows, online averaging in 115), so carriers were also injected with each window's own response, carrier for carrier against the kernel, into 35 windows of every averaging mode. The ratio ρ_N of the two ninety-per-cent points is 0.91, 0.64 and 0.68 at $N = 1$, 2 and 4 in Class A, and 0.88 and 0.41 at $N = 1$ and 2 in Class B. Smearing within one integration, which the injections do not model, is divided out as $\eta_{\rm smear}$, the mean channel response over the sweep, and is never worse than $\times 1.05$.

A.2. *Primary-beam and positional retention*

The response applied, A in Eq. A1, is a Gaussian primary beam of the documented full width $1.13\,\lambda/D$ (Cortes et al. 2026) at each window's own dish diameter. Archival fields are pointed at catalogue J2000 positions, so these nearby, fast-moving stars sit off axis by a median of $0.082''$ and at most $17.99''$. At the furthest census offset the Gaussian agrees with the blocked-aperture response of CASA (CASA Team 2022) to 7.1 per cent. The statistic, ranks and attribution are ratios within one window, so A enters only $S_{\rm min}$. $R_{\rm px}$ is the modelled share of a carrier's amplitude that survives the annual parallax where the extraction omitted it, and the model errs on the safe side, since carriers injected at the full apparent position of the 8 furthest-out windows retain 0.92 and 0.58 where it predicts 0.89 and 0.49. ϵ Eridani has no parallax record, and its limit may be optimistic by up to 9 per cent.

Propagating a position also assumes straight-line motion. Of the 91 searched positions, 30 have a companion within 400 arcsec (Mason et al. 2001; Matson et al. 2026). The displacement, from a visual orbit, from the separation with masses from Pecaut & Mamajek (2013), or from the proper-motion anomaly (Kervella et al. 2022), moves no system's limit by more than 16 per cent. Barta 161 12, HD 155555C and LHS 1140 are different. Their astrometric solutions are disturbed, yet they have no known companion and no Hipparcos epoch, so the positional loss in their 24 windows has no bound and their limits hold only at the adopted positions. They are marked in Table F1 and left out of the median, the count at 10^{15} W and Fig. 3 (§4.1).

A.3. *The survey ledger*

Every headline count is read from Table A1. The file `catalogues/count_accounting.csv` of the data archive itemises each of its rows under the same number. The planet counts are from the NASA Exoplanet Archive `pscomppars` table (Akeson et al. 2013), accessed 2026 October 4, and Table A2 lists the ALMA proposals the searched blocks belong to.

TABLE A1

The survey ledger: (a) execution blocks and windows, hours being window-hours, so the classes partition the hours as they partition the windows; (b) events, dispositions and confirmation. Each sub-row is a subset of the row its number extends, and a row whose note names sub-rows is their sum.

(a) Execution blocks and windows

	Set of execution blocks	Blocks	Windows	Stars	Systems	Hours
1	Archival scope: blocks the query exposes	656	–	–	–	–
1.1	… processed with the same pipeline	479	–	–	–	–
1.2	… not processed at the freeze	177	–	–	–	–
2	Processed with the same pipeline, 5 of them outside the scope query	484	–	–	–	–
2.1	**Primary census**	407	1663	90	83	1178
2.1.1	Class A, fine channels[a]	–	405	66	61	240
2.1.2	Class B, coarse channels[a]	–	1258	89	82	938
2.2	Reserved hold-out	77	315	36	–	238
3	Not processed at the freeze	177	–	–	–	–
3.1	Calibrated and searched	144	–	–	–	–
3.1.1	**Archival extension**, reported	115	463	16	16	362
3.1.1.1	Class A, fine channels[a]	–	67	11	–	48
3.1.2	**Archival tail**, searched after the extension was frozen	8	32	5	5	23
3.1.2.1	Class A, fine channels[a]	–	14	5	–	10
4	Beyond 40 pc: out-of-sample control	47	198	22	–	90

[a]The two classes partition the census windows, not its blocks: 255 blocks carry a Class A window and 392 a Class B one, and most carry both, so the hours of the two rows partition the windows and not the observing time.

(b) Events, dispositions and confirmation

	Count	Value	What it counts
1	Threshold crossings at $T_\star \geq 5$	51	rows 1.1 and 1.2
1.1 …	in the primary census	49	rows 1.1.1 and 1.1.2
1.1.1	Class A, fine channels	49	the carrier experiment
1.1.2	Class B, coarse channels	0	the released catalogue's 4, all in the one block that fails the quality criterion, fall below the trigger on recalibration; 0.11 expected at the 1254 other coarse windows' own control positions
1.2 …	in the archival tail	2	both unattributed, toward HD 14055, in windows no expectation covers
2	Quality-passing crossings of the census	49	the same rows as 1.1.1
2.1 …	attributed to a transition	23	of 23 coincidences inside the mask in the star's own frame; rows 2.1.1 and 2.1.2
2.1.1 …	with evidence beyond the coincidence	16	recurrence at the predicted cell, a second transition of the same species, or catalogued gas on the sightline
2.1.2 …	on the frequency coincidence alone	7	η Crv, CD−57 1054, CP−72 2713, HD 161868, HIP 10679 and HR 1010: consistent with molecular emission, not identified as it
2.2 …	unattributed	26	rows 2.2.1 and 2.2.2
2.2.1 …	in the primary census	26	against 23.8 expected by chance at the 405 windows' own control positions – a count chance delivers with probability 0.36
2.2.2 …	coincident with an atmospheric transition	0	within 62.5 MHz of one in the topocentric frame (Appendix E), and counted as atmospheric on both sides of that comparison
2.2.3 …	recovered in a later block	0	no confirmed signal; 13 attributed ones do
3	Class A systems	61	of 83 surveyed; rows 3.1 and 3.2
3.1 …	with same-frequency coverage in a separated session	**40**	**the only confirmation count quoted**: a later block in another session, among the census and the repeat blocks the recurrence test reads, covering a frequency the first covered; 13 of them over a year later. For an accelerating carrier, confirmation further requires a predicted frequency within the repeat coverage, with adequate sensitivity
3.2 …	with no separated second block at a covered frequency	21	no confirmation possible by the same-frequency test over the 142 blocks of this archive the recurrence test reads; 36 of all 83
4	Class A union bandwidth	47.7 GHz	26 disjoint islands
4.1 …	removed by the line mask	6.4 %	3.05 GHz: the figure the text quotes, against 4.3 per cent of the 118.1 GHz archival union
5	Threshold crossings in every searched set	72	rows 5.1 to 5.5
5.1 …	in the primary census	49	row 1.1
5.1.1 …	at HD 139084 B	1	in a catalogued census window, at a position the catalogue does not tabulate; on neither side of the comparison of row 2.2.1
5.2 …	in the archival tail	2	row 1.2
5.3 …	in the reserved hold-out	5	enters no census statistic
5.4 …	in the archival extension	10	enters no census statistic
5.5 …	beyond 40 pc	6	the out-of-sample control

TABLE A2
The 67 ALMA proposals the searched blocks belong to, as ADS/JAO.ALMA#*code*. Every census block belongs to one.

2012.1.00142.S, 2012.1.00198.S, 2012.1.00268.S, 2012.1.00385.S, 2012.1.00437.S, 2012.1.00993.S, 2013.1.00359.S, 2013.1.00588.S, 2013.1.00645.S, 2013.1.01147.S, 2013.1.01214.S, 2015.1.00307.S, 2015.1.00633.S, 2015.1.00783.S, 2015.1.01015.S, 2015.1.01260.S, 2016.1.00104.S, 2016.1.00358.S, 2016.1.00515.S, 2016.1.00637.S, 2016.1.00803.S, 2016.1.00817.S, 2016.1.00878.S, 2016.1.00880.S, 2016.2.00015.S, 2016.2.00200.S, 2016.A.00013.S, 2017.1.00167.S, 2017.1.00200.S, 2017.1.00215.S, 2017.1.00561.S, 2017.1.00698.S, 2017.1.00704.S, 2017.1.00786.S, 2017.1.00986.S, 2017.1.01583.S, 2017.1.01644.S, 2018.1.00045.S, 2018.1.00072.S, 2018.1.00126.S, 2018.1.00461.S, 2018.1.00470.S, 2018.1.00947.S, 2018.1.01149.S, 2018.1.01545.S, 2018.1.01833.S, 2019.1.00189.S, 2019.1.00696.S, 2019.1.01443.T, 2019.1.01517.S, 2019.1.01603.S, 2019.1.01681.S, 2019.2.00208.S, 2021.1.00485.S, 2021.1.01209.S, 2021.2.00097.S, 2022.1.00134.S, 2022.1.00338.L, 2022.1.00793.S, 2022.1.01163.S, 2023.1.00390.S, 2023.1.00487.S, 2024.1.00165.S, 2024.1.00595.S, 2024.1.00722.S, 2024.1.01095.S, 2024.A.00040.S

B. THE CONTROL ASSUMPTION, THE HOLD-OUT AND THE RECURRENCE BOUND

The chance expectation of §4.2.1 predicts how often the star should cross the trigger from how often control positions in its own field do. It therefore rests on one assumption the searched data cannot test, that the stellar position behaves like a control position. Below the trigger it does not quite do so. In the band $4 \le T < 5$ the star carries 1.108 times a control position's above-threshold cells (1.05–1.16 at 95 per cent, blocks resampled). Rescaling the expectation by that ratio is a sensitivity check (§4.2.1); the control positions' own rate, the more demanding of the two, is kept for the count under test. For the 26 unattributed crossings to stand above chance at 5 per cent, stars would have to trigger at 0.76 times a control position's rate or less.

The stellar excess is not a primary-beam effect, since a noise rise toward the field edge would put the stellar rank above one half, the opposite of what is seen, and the probes' radial slope on the reserved blocks, -0.027 ± 0.024, is small and of the wrong sign. It appears only in the 7 m-array data, 1.15 (1.1–1.2) against 1.04 (1.0–1.1) on the 12 m array, and its cause has not been identified.

A calibration error that varies with frequency, such as a bandpass residual or a coarse system temperature interpolated across a telluric line (Hunter et al. 2018), would act in the sense observed: it multiplies the continuum, so it acts at the star and not on the empty sky around it. Such a residual matters even if it is too small to make a crossing by itself. A small error coupled to the continuum, added to noise already near the trigger, would still change how often the star crosses it, and the searched data cannot separate that from a real difference between star and controls. The probabilities therefore rest on the stellar-background model (§4.2.1).

B.1. *The reserved hold-out*

Because the statistic and the mask were set on the searched sample, 77 of 484 blocks (315 windows toward 36 stars) were held out by an identifier rule, $\mathrm{sha256}(\mathrm{UID})_{[:8]} \bmod 5 = 0$. Two guards, which also use identifiers alone, keep a block on a named list in the survey and ensure that no system loses every block. Their independence of the statistic and the mask is our assertion, which a reader cannot check. Since the rule hashes the block and not the session, 64 per cent of them also share calibrators, weather and setup with a searched block. They trigger in 5 of 62 windows against 6.5 predicted, and the 28 that share no session hold one trigger event against 1.9, so they neither confirm nor contradict the stellar ratio.

B.2. *The recurrence bound*

A crossing is tested for recurrence by reading its predicted cell, one channel at the carried drift, in the later blocks that cover it. With S the discovery amplitude and $\bar{a}$ and $\bar{\sigma} = (\sum_i \sigma_i^{-2})^{-1/2}$ the inverse-variance combination of the repeat readings, the two signal-to-noise ratios are $T_{\mathrm{pers}} = S/\bar{\sigma}$ and $T_{\mathrm{rep}} = \bar{a}/\bar{\sigma}$. A discovery amplitude selected from 10^2 to 10^7 cells at $T_\star \simeq 5$ is biased high, so it is not compared with the repeat directly. The bound is instead the largest fraction of the discovery amplitude still allowed by the repeat readings,

$$f_{\mathrm{rep}} = \frac{T_{\mathrm{rep}} + 5}{T_{\mathrm{pers}}},$$

and T_{rep} is centred on zero under the null, scattering by 0.994 about -0.016 over 327 single-cell repeat readings. The reading is conditional. Given an assumed true amplitude, the bound recovers a carrier with probability $\Phi(T_{\mathrm{pers}} - 5)$ at the discovery amplitude and $\Phi(5T_{\mathrm{pers}}/T_\star - 5)$ at the nominal threshold-equivalent amplitude $5S/T_\star$, and the 5 crossings where these are smallest are reported as untested.

C. THE LINE MASK

A crossing that falls on a known molecular transition has an astrophysical explanation to hand, and the mask decides which crossings those are.

The species list was assembled by hand. It holds the bright millimetre disc and cloud tracers the pipeline checked from its first search, with [C I] and hydrogen recombination added during the search and SO after a crossing was found on one of its lines. It was completed at the query below, after the crossings had been examined, and it omits other species seen in discs and clouds, such as CH_3OH, DCO^+ and N_2H^+. The mask contains every transition of these 12 species inside the 46 searched intervals with an upper-state energy below 150 K, 111 transitions in all (Table C1). Only main hyperfine components are included, with no strength cut, and all come from one Splatalogue query (Remijan et al. 2007) of CDMS and JPL (Pickett et al. 1998; Müller et al. 2005) dated 2026 October 6.

Attribution is decided in the star's own rest frame. A crossing's topocentric frequency is carried to the barycentre with its own block's pointing and mid-time, and then to the star with its systemic velocity, giving $\nu_\star$. The crossing is attributed if, for any listed transition of rest frequency ν_0,

$$|\Delta v_\star| \le v_{\mathrm{h}}, \qquad \Delta v_\star = c\,\frac{\nu_\star - \nu_0}{\nu_0}, \qquad v_{\mathrm{h}} = 50\,\mathrm{km\,s^{-1}},$$

so that $\Delta v_\star$ is positive for a crossing above the transition in frequency. The same test, in the same frame, decides whether a control maximum is attributed (Appendix D). Of the 49 quality-passing census crossings, 23 meet it, and these are the 23 attributions; the hold-out is treated identically.

TABLE C1
The line mask, by species: the number N of transitions, the frequency range they span inside the searched intervals and their upper-state energies.

Species	N	ν_0 (GHz)	E_u (K)
H	5	92.03 – 231.90	–
SO	15	100.03 – 682.68	16–143
H_2CO	13	101.33 – 491.97	10–106
$C^{18}O$	3	109.78 – 329.33	5–32
CN	50	113.12 – 680.30	5–114
CO	3	115.27 – 345.80	6–33
CS	3	195.95 – 342.88	24–66
SiO	2	217.10 – 347.33	31–75
^{13}CO	8	220.40 – 330.59	16–32
HCN	7	354.50 – 354.51	43–43
HCO^+	1	356.73 – 356.73	43
[C I]	1	492.16 – 492.16	24

The windows were placed by their observers for their own lines, and that limits what "unattributed" means. Of the Class A windows, 39 per cent are centred within v_h of a listed transition, against 1–8 per cent for the same windows displaced, so a crossing is unattributed against a list that these setups helped to shape.

As a check that changes no disposition, every crossing is also read in the local standard of rest. Toward ALMA J153702653−33192492, HD 139084 B, HD 285968 and HD 48370 the CO($1 \rightarrow 0$) survey of Dame & Thaddeus (2022); Dame et al. (2001) shows emission at the crossing's own LSR velocity (within $3\,\mathrm{km\,s^{-1}}$ of its peak, and at least 0.23 K), so 6 of the coincidences are interstellar. Elsewhere the Planck CO map (Planck Collaboration et al. 2016) stays below $0.178\,\mathrm{K\,km\,s^{-1}}$.

Keplerian motion where these molecules survive sets the half-width, $\lesssim 15\,\mathrm{km\,s^{-1}}$ beyond 10 au. It has a cost. Over the union of searched sky frequencies the mask removes 3.05 GHz of the 47.7 GHz Class A union, 6.4 per cent, and 5.04 GHz of 118.1 GHz over both classes, 4.3 per cent, although each limit is formed over its own window's channels outside the mask in its own star's frame (§3.2).

D. THE FALSE-ALARM EXPECTATION AND THE CONTROL ENSEMBLE

D.1. *The expectation and its resampling*

One attribution set serves both sides of the comparison, the mask of Appendix C being applied to a control maximum exactly as to a crossing. Over the 405 Class A windows w of the primary census the expected number of unattributed crossings is

$$\bar{N} = \tau \sum_w r_w\,(1 - a_w), \qquad r_w = \frac{n_w}{N_{\mathrm{ctrl}}},$$

where n_w of the window's $N_{\mathrm{ctrl}} = 512$ control positions reach the trigger, a_w is the share of control exceedances that land inside the mask, and τ is the share the atmospheric screen leaves. The rates alone give 36.5 expected crossings. The attributed share is measured where control maxima actually fall, on the 8 positions per window that keep a dynamic spectrum. Of their 289 exceedances, 92 lie inside the mask, so attribution takes 32 per cent. The screen (Appendix E) then removes 9 of the 197 that remain, leaving $\bar{N} = 23.8$ against 26 observed. The 14 Class A windows of the archival tail and the 1 of HD 139084 B, a position the catalogue does not tabulate, enter neither side. Near a band edge the pooled exceedance rate is $\times 2.464$ the interior rate, but one window, toward HD 48370, holds 59 per cent of the near-edge exceedances; without it the ratio is $\times 1.020$ ($\times 1.181$ on a held-out half, $p = 0.24$), so no edge correction is applied.

Windows share execution blocks, so the count is compared through block resampling. In each of $B = 200\,000$ resamplings the blocks are drawn with replacement, each window's rate is redrawn binomially from its own probes, and the attributed share and the screen are redrawn from their own counts, giving $\bar{N}_b$; then

$$P = \frac{1}{B} \sum_{b=1}^{B} \Pr\!\left(N \geq N_{\mathrm{obs}} \mid \bar{N}_b\right),$$

with N a Poisson count and $N_{\mathrm{obs}} = 26$, gives $P = 0.36$, conditional on the control-position model (Appendix B). With k crossings added to every $\bar{N}_b$, 26 remains credible at 0.06 for $k = 12$ but not for one more (0.04 against 5 per cent). This is the allowance of §4.2.1. As a check, a pseudo-star read at one control position in every window of a block is no more dispersed than one read window by window (variance ratio 1.0) and reaches 26 with probability 0.35. The widened recurrence window of §4.4 is read in the same way over its 93 distinct windows.

D.2. *Execution-block quality*

The comparison assumes that each block's control ensemble behaves like noise, and in one block all 512 probes exceed the trigger. Every block is therefore judged on the median of its windows' control ensembles over its class median, which also flags 5 of the 1663 windows at twice the survey median. The criterion was written after the condition had been seen, but the choice is not a delicate one. The failing block sits at 4.1 and the next at 1.12, and its 4 coarse-channel windows hold no census crossing once ALMA's final flags are restored (§3).

D.3. *Registration*

Between epochs, the predicted cell is registered to 0.75 of a channel in every window. Widening each reading to the worst cell within $0.45\,\mathrm{km\,s^{-1}}$, re-anchoring it along the discovery track, or raising the bar to 3.9 changes no disposition, although registration to a single channel is tested only on the unresolved line of HD 285968, which registers to 0.06 of a channel.

D.4. *The pairs*

Where the epochs of a pair of crossings (§4.4) are close enough for the predicted offset to be tighter than the searched window, all 5 such pairs fail a drift changing at a constant rate, by at least 3.6 times the resolution. The β Pictoris pair, whose crossings lie $106\,\mathrm{km\,s^{-1}}$ apart in the star's frame and 24.4 h apart, cannot be dismissed in this way. Its mean acceleration, $1.21\,\mathrm{m\,s^{-2}}$, is inside the $3.6\,\mathrm{m\,s^{-2}}$ ceiling, and a circular orbit of 1.54 d about the $1.75\,M_\odot$ star (Crifo et al. 1997) reproduces it. Both blocks belong to one session and no block of another session covers the window, so the pair is untestable.

D.5. *The drift grid's edges*

The drift trials run from $-\dot{\nu}_{\max}$ in half-channel steps to the first trial at or beyond $+\dot{\nu}_{\max}$, so the searched interval

is asymmetric by up to one step. Of the 52 crossings of the census, the archival tail and the hold-out with a fitted drift, the 2 toward HD 161868, TRAPPIST−1 lie on an outermost trial of their own window's grid. Their drift rates, in the observatory frame, and the accelerations they imply are lower bounds, and 2 of them carry no amplitude bound.

E. RADIO-FREQUENCY INTERFERENCE SCREENING

Interference is the first alternative explanation for an unattributed crossing, although an interferometer is a poor receiver of it, since a signal not in the far field at the phase centre is suppressed by its rapidly varying geometric phase. Allocations exist across the searched bands (International Telecommunication Union 2024), and the one documented hazard, spaceborne cloud radar at 94.0–94.1 GHz (Stephens et al. 2002; Wehr et al. 2023; Emerson 2004), overlaps 1 of the 1663 census windows, with 0 crossings. The correlator flagging was not retained, so no per-channel interference flag exists. The tests below describe the population. They clear no single crossing, and they do not substitute for the ON/OFF cadence of a dedicated search.

Each test could have failed, and none did. Each p value is computed against the same peaks redrawn on their own channel grids. Crossings cluster in sky frequency only on the survey's own lines, with 15 of 53 within $\pm 50\,\mathrm{km\,s^{-1}}$ of a searched transition against 6 expected ($p < 3 \times 10^{-4}$). The 28 unattributed quality-passing crossings do not crowd on one another between 1 and 300 MHz ($p = 0.6$), whereas the attributed ones do. None of the 56 crossings of the census, the archival tail and the hold-out lies within 62.5 MHz of a telluric line. Peaks are spread uniformly within their windows (median 0.50), which a fixed intermediate frequency would not allow.

Coincidences between different stars within 5 MHz number 569 against 550 expected ($p = 0.1$). Of the 26 unattributed crossings in catalogued blocks, 17 have no catalogued transition in the image band and 4 one their own block saw too faint to leak; for 5 the image frequency was not observed. The one evident concentration is at the carbon monoxide tunings. The 14 crossings between 344.2 and 346.4 GHz include 8 of the 30 unattributed crossings of the census and the hold-out, against 7 expected from where the windows were tuned.

F. PER-SYSTEM SENSITIVITY

Table F1 gives for each system the limit of its best window outside the attribution mask (§3.2). A bound pair appears under its primary's name, and the limits at both ends of the calibration interval are in `sensitivity/system_limits.csv` of the data archive (White & Dey 2026). Beside each limit are that window's centre frequency ν and channel width $\Delta\nu$, the system's merged coverage $\Delta\nu_{\rm cov}$, the percentage *dec.* by which atmospheric decorrelation leaves the limit optimistic, and whether blocks of different sessions cover a common frequency more than a day (d) or a year (yr) apart, over the census and the 142 blocks the recurrence test reads. A dagger marks the 7 systems whose best window was itself injected into, 6 of them among the 57 systems with a bounded position. The other 51 limits of those 57 are transferred from the class average and carry the dispersion between injected windows, $\times 0.81$ to $\times 1.12$. For the 6 systems whose best window lies outside the injected range, that dispersion is itself extrapolated.

REFERENCES

Akeson R. L., et al., 2013, PASP, 125, 989
Anglada-Escudé G., et al., 2016, Nature, 536, 437
Astropy Collaboration, 2022, ApJ, 935, 167
Backus P. R., Project Phoenix Team, 2002, in Stanimirović S., Altschuler D., Goldsmith P., Salter C., eds, ASP Conf. Ser. Vol. 278, Single-Dish Radio Astronomy: Techniques and Applications. Astron. Soc. Pac., San Francisco, p. 525
Burton K., MacGregor M. A., Osten R. A., 2022, ApJ, 939, L6
CASA Team, 2022, PASP, 134, 114501
Cataldi G., et al., 2023, ApJ, 951, 111
Chen T. X., et al., 2022, ApJS, 260, 5
Choza C., et al., 2024, AJ, 167, 10
Cocconi G., Morrison P., 1959, Nature, 184, 844
Cortes P. C., et al., 2026, ALMA Cycle 13 Technical Handbook, ALMA Doc. 13.3, ver. 1.0, doi:10.5281/zenodo.18793803
Crifo F., Vidal-Madjar A., Lallement R., Ferlet R., Gerbaldi M., 1997, A&A, 320, L29
Czech D. J., et al., 2021, PASP, 133, 064502
Dame T. M., Hartmann D., Thaddeus P., 2001, ApJ, 547, 792
Dame T. M., Thaddeus P., 2022, ApJS, 262, 5
Dent W. R. F., et al., 2014, Science, 343, 1490
Drake F. D., 1961, Physics Today, 14, 40
Emerson D., 2004, The CloudSat radar and implications for ALMA, ALMA Memo 504
Enriquez J. E., et al., 2017, ApJ, 849, 104
Gaia Collaboration, Vallenari A., et al., 2023, A&A, 674, A1
Garrett M. A., 2026, MNRAS, 549, stag991
Ginsburg A., et al., 2019, AJ, 157, 98
Harris C. R., et al., 2020, Nature, 585, 357
Hunter T. R., Phillips N. M., Broguière D., González J., 2018, Improving the Calibration of Atmospheric Spectral Features in ALMA Data. NA ALMA 2016 Development Study Final Report, 2018 March 2
International Telecommunication Union, 2024, Radio Regulations, Edition of 2024. ITU, Geneva
Jacobson-Bell B., et al., 2025, AJ, 169, 206
Johnson O. A., et al., 2023, AJ, 166, 193
Kervella P., Thévenin F., Lovis C., 2017, A&A, 598, L7
Kervella P., Arenou F., Thévenin F., 2022, A&A, 657, A7
Li J.-K., Tao Z.-Z., Zhang T.-J., 2026, ApJ, 1006, 163
Ma P. X., et al., 2023, Nature Astronomy, 7, 492
MacGregor M. A., Osten R. A., Hughes A. M., 2020, ApJ, 891, 80
Manunza L., et al., 2025, Acta Astronautica, 233, 155
Margot J.-L., et al., 2023, AJ, 166, 206
Mason B. D., Wycoff G. L., Hartkopf W. I., Douglass G. G., Worley C. E., 2001, AJ, 122, 3466
Mason L. A., Garrett M. A., Wandia K., Siemion A. P. V., 2025, MNRAS, 536, 2127
Matrà L., et al., 2017, MNRAS, 464, 1415
Matson R. A., Williams S. J., Hartkopf W. I., Mason B. D., 2026, Sixth Catalog of Orbits of Visual Binary Stars, U.S. Naval Observatory, accessed 2026 October 14
Mauersberger R., Wilson T. L., Rood R. T., Bania T. M., Hein H., Linhart A., 1996, A&A, 306, 141
Morrison I. S., et al., 2023, PASA, 40, e019
Müller H. S. P., Schlöder F., Stutzki J., Winnewisser G., 2005, J. Mol. Struct., 742, 215
Pardo S., Poznanski D., Croft S., Siemion A. P. V., Lebofsky M., 2025, AJ, 170, 12
Pecaut M. J., Mamajek E. E., 2013, ApJS, 208, 9
Pickett H. M., Poynter R. L., Cohen E. A., Delitsky M. L., Pearson J. C., Müller H. S. P., 1998, J. Quant. Spectrosc. Radiat. Transfer, 60, 883
Planck Collaboration, Adam R., et al., 2016, A&A, 594, A10
Price D. C., et al., 2020, AJ, 159, 86
Privon G., et al., 2026, ALMA Cycle 13 Proposer's Guide, ALMA Doc. 13.2, ver. 1.0, doi:10.5281/zenodo.18793859
Remijan A. J., Markwick-Kemper A., ALMA Working Group on Spectral Line Frequencies, 2007, in American Astronomical Society Meeting Abstracts 211, 132.11
Richards A. M. S., Moravec E., Etoka S., Fomalont E. B., Pérez-Sánchez A. F., Toribio M. C., Laing R. A., 2022, Self-calibration and improving image fidelity for ALMA and other radio interferometers, ALMA Memo 620
Sheikh S. Z., 2020, International Journal of Astrobiology, 19, 237
Sheikh S. Z., Wright J. T., Siemion A. P. V., Enriquez J. E., 2019, ApJ, 884, 14
Sheikh S. Z., et al., 2021, Nature Astronomy, 5, 1153
Sheikh S. Z., Huston M. J., Fan P., Wright J. T., Beatty T., Martini C., Kopparapu R., Frank A., 2025, AJ, 169, 118
Steffes P. G., DeBoer D. R., 1994, Icarus, 107, 215
Stephens G. L., et al., 2002, Bull. Am. Meteorol. Soc., 83, 1771
Tarter J., 2001, ARA&A, 39, 511
Tremblay C. D., Tingay S. J., 2024, ApJ, 972, 76
Vidal C., et al., 2026, arXiv:2605.21093 (preprint)
Virtanen P., et al., 2020, Nature Methods, 17, 261
Wehr T., et al., 2023, Atmos. Meas. Tech., 16, 3581
Wenger M., et al., 2000, A&AS, 143, 9
White G. J., 2026, RAS Techniques and Instruments, in press, arXiv:2610.36144
White G. J., Dey R., 2026, Supplementary Data for an Archival ALMA Search for Technosignatures toward Stars within 40 pc. Zenodo, doi:10.5281/zenodo.0000000
Wright J. T., Kanodia S., Lubar E., 2018, AJ, 156, 260
Zhang Z.-S., et al., 2020, ApJ, 891, 174

TABLE F1

Per-system sensitivity of the carrier search. For each of the 61 systems the fine-channel search covers, the $EIRP_{90}$ of its best window with its own primary-beam, annual-parallax and smearing retentions applied, ordered by the limit. Columns and daggers are defined in Appendix F; the notes below give the other marks.

System	$EIRP_{90}$ (10^{14} W)	ν (GHz)	$\Delta\nu$ (kHz)	$\Delta\nu_{cov}$ (GHz)	dec. (%)	conf.	System	$EIRP_{90}$ (10^{14} W)	ν (GHz)	$\Delta\nu$ (kHz)	$\Delta\nu_{cov}$ (GHz)	dec. (%)	conf.
ϵ Eri	0.37	229.98	977	2.3	–	yr	κ CrB	14†	230.57	977	1.7	2*	–
LP 876-10	0.74	345.83	488	1.7	4*	d	HD 170773	15	230.13	488	7.4	2*	d
τ Cet	0.79	230.55	488	1.8	0.4*	d	HD 61005	17	230.52	244	19	3*	d
HD 69830	0.94†	335.38	244	3.8	0.5	d	LHS 1140	17§	230.00	488	1.8	0.4*	d
AU Mic	1.0	230.56	488	3.5	35*	yr	LP 736-15	17	345.82	977	1.7	4*	–
61 Vir	1.1	345.79	488	1.8	5*	yr	HD 172555	19	340.94	488	5.3	200	–
TRAPPIST-1	1.5	346.58	488	1.7	5	d	HD 155555C	19†§	346.01	488	1.7	43	–
AT Mic AB	1.5	346.04	488	1.7	3	–	L 836-122	19	346.00	488	1.7	1	–
β Pic	1.5‡	226.85	122	4.4	2*	yr	TWA 7	20	345.99	244	0.87	26*	d
HD 38858	2.0	230.50	488	1.8	3*	yr	CP−72 2713	21	345.12	488	5.8	64	d
GJ 3379	2.0	229.96	488	1.9	0.4*	d	HD 14055	22	230.11	488	7.2	3	yr
HD 207129	2.2	230.57	488	1.8	3*	yr	GJ 2006A	22	346.02	488	1.7	1	–
GJ 14	2.8	227.22	488	3.5	19	d	CD−57 1054	22	345.99	488	1.7	20	d
HD 139664	2.9	230.55	488	1.7	3*	–	HD 23484	24	230.53	977	1.8	0.2	d
HD 202628	3.4	230.51	488	1.8	4*	yr	51 Eri	25	230.53	1953	1.7	11*	–
HD 53143	3.4	230.53	977	1.8	2*	yr	TYC 2211-1309-1	26	346.00	488	1.7	5	–
HN Lib	3.5†	229.99	488	1.8	0.4*	d	UCAC3 176-23654	26	346.02	488	1.7	4	–
HD 10647	3.5	230.54	488	6.1	4*	yr	PM J03433+1958	28	230.13	488	1.7	2*	–
η Crv	4.0	343.35	488	7.2	27	yr	UCAC2 20312880	29	345.96	488	1.7	11	–
TWA 3A	4.7†	230.54	122	0.44	35*	d	LP 353-51	30	346.02	488	1.7	26	–
GJ 849	6.1	230.02	488	1.9	0.4*	d	Barta 161 12	31§	346.02	488	1.7	4*	–
HD 92945	7.5	355.11	488	3.7	6*	d	HD 139084	49	346.01	488	1.7	3	–
TX PsA	7.6	346.03	488	1.7	7	–	HD 45184	58	230.53	977	1.7	2	d
HD 285968	8.2	229.97	488	1.8	0.4*	d	WD 0407+179	76	230.13	488	1.7	2*	–
HD 161868	9.0	227.22	488	7.4	5	d	HD 48370	79†	230.73	122	3.3	0.4*	yr
HD 110411	11	219.58	244	4.7	2*	yr	HD 31392	82	230.52	977	1.7	0.9	d
HR 1010	12	230.53	977	2.8	1	d	ALMA J153702653-33192492	89	345.98	488	14	5*	d
HIP 490	12†	230.53	244	0.86	3*	–	HD 89452	160	230.53	977	1.7	0.7	d
HD 107146	13	355.14	488	3.7	6*	d	HD 377	250	230.54	977	1.7	68	d
Wolf 219	13	230.13	488	1.7	2*	d	HIP 10679	–	–	–	0.11	–	–
LP 476-207	13	346.02	488	1.7	4	–							

Notes. A double dagger marks a row keeping less than half its channels outside the mask; HIP 10679, whose 2 windows all lie inside it, has dashes. A section mark (§) flags Barta 161 12, HD 155555C and LHS 1140, whose limits hold only at the adopted position (Appendix A.2) and which are left out of the median and of the counts quoted over systems. *dec.* is a floor, the phase rms being recorded on the calibrator; it is measured where the delivery records one and imputed from blocks matched within a factor of two in baseline length where it does not (asterisk), and a dash marks the one system with neither.