

# A VOLUME-LIMITED SEARCH FOR TECHNOSIGNATURES WITHIN 20 PC USING THE PUBLIC ALMA ARCHIVE: I. BACKGROUND, MOTIVATION AND SURVEY DESIGN

GLENN J. WHITE<sup>1,2</sup> AND ROBIN DEY<sup>3</sup>

Version August 30, 2026

## ABSTRACT

We describe the motivation, background and design of a volume-limited search for radio technosignatures among all stars within 20 pc of the Sun that have public observations in the Atacama Large Millimeter/submillimeter Array (ALMA) science archive. Unlike the great majority of technosignature surveys conducted to date, which are either wide, shallow all-sky surveys or targeted campaigns preferentially selecting exoplanet hosts, young stellar objects, or other astrophysically “interesting” systems, our sample is constructed to be representative of the general stellar population in the immediate solar neighbourhood, irrespective of spectral type, age, multiplicity, or known planetary or circumstellar-disc status. We mine the ALMA archive opportunistically: rather than requesting dedicated observing time, we reprocess existing calibrated (and, where necessary, recalibrated) visibility data to search simultaneously for narrowband Doppler-drifting technosignature candidates and for anomalous broadband continuum emission, at frequencies ( $\sim 90\text{--}950\text{ GHz}$ ) that remain almost entirely unexplored by the technosignature literature. This paper (Paper I of a series) reviews six decades of radio technosignature searches, the small number of millimetre/submillimetre efforts to date, recent applications of machine learning to technosignature detection, and the statistical frameworks used to interpret non-detections, in order to place the present survey in context and to motivate its volume-limited, archival design. The observational results of the survey itself, together with the resulting upper limits on transmitter prevalence in the solar neighbourhood, will be presented in Paper II.

*Subject headings:* extraterrestrial intelligence — radio lines: stars — submillimetre: stars — surveys — astrobiology

## 1. INTRODUCTION

The search for extraterrestrial intelligence (SETI) rests on the premise that a technological civilisation, whether deliberately signalling or simply going about its own affairs, may leave a detectable imprint on the electromagnetic spectrum: a *technosignature* (??). Since the foundational proposal of ? that microwave frequencies near the 21 cm hydrogen line would be an obvious “meeting place” for interstellar communication, and Frank Drake’s first observational test of the idea with Project Ozma (?), the radio band has remained the dominant search space for technosignatures, for reasons that are both historical and physically well motivated: radio photons are cheap to produce in enormous numbers, propagate through the interstellar medium and planetary atmospheres with little attenuation, and (for a sufficiently narrowband signal) can in principle be distinguished from any known natural astrophysical process by their spectral compactness and Doppler dynamics (?).

Six decades on, radio technosignature searches have grown from single-target, single-frequency pointings into surveys of many thousands of stars across broad swathes of the radio spectrum. Yet, as ? demonstrated quantitatively with their “Cosmic Haystack” formalism, even the combined efforts of every major SETI programme to date have sampled only an infinitesimal fraction of the plausible multi-dimensional parameter space of transmitter frequency, bandwidth, power, location, and duty cycle. Persistent, non-trivial gaps remain: almost the entire millimetre and submillimetre regime above  $\sim 50\text{ GHz}$  is essentially unsearched; most surveys have targeted specific classes of star (nearby exoplanet hosts, young stellar objects with protoplanetary discs, stars in the “Earth Transit Zone”) rather than an unbiased, complete sample of the local stellar population; and only a small number of studies have systematically reprocessed *existing* interferometric archives for technosignatures, as opposed to requesting dedicated telescope time.

This paper is the first in a series describing a new survey designed to address several of these gaps simultaneously. We construct a volume-limited sample of every star within 20 pc of the Sun with public ALMA archival coverage, irrespective of spectral type, disc-hosting status, or known planets, and mine that archival data for both narrowband technosignature candidates and anomalous continuum emission, applying corrections for stellar proper motion and for the range of Doppler drift rates plausible for a transmitter on a planet in orbit around its host star. Section ?? reviews the relevant literature: the history of radio SETI (§??), the current generation of large-scale technosignature surveys (§??), the very limited body of work at millimetre/submillimetre frequencies (§??), the growing use of machine learning and anomaly detection in this field (§??), the statistical frameworks used to interpret survey non-detections (§??), the case for a volume-limited, representative sample (§??), and the prospects offered by next-generation facilities (§??). Section ?? summarises our sample selection, Section ?? our data-reduction and search methodology, and Section ??

<sup>1</sup> School of Physical Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK; glenn.white@open.ac.uk

<sup>2</sup> RAL Space, STFC Rutherford Appleton Laboratory, Harwell Campus, Didcot, OX11 0QX, UK

<sup>3</sup> VBRL Holdings Inc.

the scope of the results to be presented in Paper II.

## 2. BACKGROUND

### 2.1. Six decades of radio SETI: a brief history

Project Ozma (?) observed two nearby Sun-like stars,  $\tau$  Ceti and  $\epsilon$  Eridani, with the 26 m Tatel telescope at Green Bank at 1420 MHz for a total of some 150 hours, and found no signal, but established both the methodology and the symbolic starting point for the modern field. Through the 1960s–1980s a steady stream of targeted and sky-survey searches followed, including Ozma II (?), the Ohio State “Big Ear” survey (source of the famous “Wow!” signal), and the META and BETA all-sky surveys conducted from Harvard/Smithsonian facilities. Project Phoenix (?), run by the SETI Institute from 1995–2004, was the most sensitive targeted survey of its era, observing on the order of 800 nearby Sun-like stars using the Arecibo, Parkes and Green Bank telescopes, and pioneered a rigorous methodology for distinguishing candidate signals from terrestrial radio-frequency interference (RFI) through simultaneous multi-site or multi-antenna verification. The Allen Telescope Array (ATA; ?), jointly developed by the SETI Institute and UC Berkeley and first commissioned in 2007, was the first radio telescope purpose-built from the outset for commensal, wide-band SETI observing, and remains in active use, including in recent technosignature searches of interstellar objects (?) and TRAPPIST-1 (?). The distributed-computing project SETI@home (?) processed public-participant-donated CPU cycles to search Arecibo sky-survey data for narrowband and pulsed signals for two decades before its 2020 shutdown, and its final data-analysis summary was only published in 2025.

Two complementary observing strategies have been used throughout this history: *sky surveys*, covering a large fraction of the celestial sphere at modest sensitivity per pointing, and *targeted searches*, concentrating integration time on a preselected list of promising stars (e.g. ?). Both strategies persist in the modern era, but – as we discuss in §§? – the majority of targeted programmes have selected their samples on some presumed indicator of habitability or interest (proximity, Sun-like spectral type, known planets, or membership of the Earth Transit Zone), rather than aiming for a complete, unbiased census of the local stellar population.

### 2.2. Modern large-scale radio technosignature surveys

The current generation of technosignature science is dominated by the Breakthrough Listen initiative, launched in 2015 with a ten-year, \$100 million commitment to survey the one million nearest stars, the full Galactic plane and centre, the 100 nearest galaxies, and a sample of exoplanet and planet-candidate hosts, primarily using the Robert C. Byrd Green Bank Telescope (GBT) and the CSIRO Parkes “Murriyang” telescope, and now increasingly via commensal observing on MeerKAT (?). Its first major published dataset, ?, searched 692 nearby stars over 1.1–1.9 GHz with the GBT and set limits on narrowband transmitters of  $\text{EIRP} > 10^{13}$  W; this was followed by ? (1327 stars, 1.10–3.45 GHz), ?-style searches of the restricted Earth Transit Zone at 3.95–8.00 GHz, and, most recently, a substantial expansion of automated, commensal technosignature searching with MeerKAT that has processed some 1.5 million coherent beams between 2023 and 2026 (?). In parallel, ? conducted a large citizen-science-style search of 11 680 stars with the GBT at 1.15–1.73 GHz, explicitly developing a statistical formalism for interpreting non-detections in terms of the fraction of stars that could host a detectable transmitter, *provided the observed sample is representative of the underlying stellar population of the search volume* – a condition that, as we discuss below, is often not satisfied by targeted, interest-selected samples.

Beyond Breakthrough Listen, the last five years have seen international radio facilities extend technosignature searches to new frequency regimes and observational strategies: the Five-hundred-metre Aperture Spherical Telescope (FAST; ??) in China has conducted commensal searches with unmatched raw sensitivity at L-band, including recent narrowband searches of interstellar objects; LOFAR and the pathfinder NenuFAR have pushed technosignature searches down to 10–190 MHz, with LOFAR alone having surveyed over 1.6 million stars in commensal mode and dual-site LOFAR configurations enabling near-real-time RFI rejection (?); the Sardinia Radio Telescope has conducted one of the first dedicated high-frequency ( $> 18$  GHz) surveys of nearby stars and the Galactic Centre (?); and commensal technosignature back-ends are now operating or in development at the Karl G. Jansky Very Large Array (the COSMIC system; ?), MeerKAT (?), and the Murchison Widefield Array (?). Occultation-based strategies – observing a star during a planet-planet or planet-star occultation, when geometric alignment transiently favours detection of leakage or deliberate signals – have also grown in popularity, exemplified by recent Breakthrough Listen searches of TESS-selected eclipsing systems (?).

### 2.3. Millimetre and submillimetre technosignature searches

In stark contrast to the radio regime below  $\sim 10$  GHz, the millimetre and submillimetre bands remain almost entirely unexplored for technosignatures. ? conducted, to our knowledge, the first technosignature search ever performed with ALMA, searching archival Band 3 data (spectral windows centred at 90.642 and 93.151 GHz) for narrowband signals towards 28 Galactic stars selected from Gaia DR3. Their approach used a “stellar bycatch” method: rather than targeting stars directly, they searched for serendipitous stars falling within the undistorted field of view of four bright ALMA phase calibrators, since these calibrators are observed frequently and are not generally themselves of astrophysical interest. For the nearest star in their sample they placed a limit of  $\text{EIRP}_{\text{min}} > 7 \times 10^{17}$  W, and discussed in detail the specific challenges of high-frequency technosignature work: substantially larger Doppler drift rates at fixed line-of-sight acceleration (since drift rate scales linearly with observing frequency), the resulting sensitivity loss (“smearing”) for a fixed spectral and temporal resolution unless a drift-rate search is performed, and an increased

incidence of spectral confusion from the dense molecular line forest present in protoplanetary discs and star-forming regions at these frequencies. ? extended technosignature work on the TRAPPIST-1 system to the Allen Telescope Array, underlining the continued observational interest in this particular target but also its status as one of very few systems that has now been searched for technosignatures across a wide range of frequencies. Our own earlier work (White & Dey, in prep.; archived at <https://github.com/Tilanthi/SETI>) reproduced and extended the ? methodology directly on ALMA science-target (rather than calibrator-bycatch) fields for TRAPPIST-1 across Bands 3, 6 and 7, finding no detection and placing limits of  $\text{EIRP}_{\min} \sim 5\text{--}10 \times 10^{13} \text{ W}$ , and subsequently built a ranked target list and pilot survey of 100 nearby, ALMA-observed stars, from which the present, volume-limited survey has grown. Beyond these efforts, we are not aware of any other published, systematic technosignature search using ALMA or any comparably sensitive millimetre/submillimetre interferometer, representing a substantial and readily addressable gap in frequency coverage relative to the sub-10 GHz literature.

The near-complete absence of mm/submm technosignature work to date is attributable less to any fundamental unsuitability of these frequencies – Sheikh’s “nine axes of merit” framework (?) notes that higher frequencies suffer less from interstellar scintillation and dispersion, and offer access to much larger instantaneous bandwidths – than to historical circumstance: sensitive mm/submm interferometers with the necessary spectral resolution and archival depth (chiefly ALMA) have only existed, and only accumulated a sufficiently large calibrated science archive, within roughly the last decade. This makes archival reprocessing, rather than dedicated proposal-driven observing, a particularly natural strategy in this regime: ALMA’s archive already contains many thousands of hours of high-spectral-resolution observations of individual, nearby stars, taken for entirely unrelated astrophysical purposes (disc structure, molecular chemistry, stellar activity, debris-disc detection, calibration monitoring), each of which can in principle be re-searched for technosignatures at negligible additional observing cost.

#### 2.4. Machine learning and anomaly detection in technosignature searches

The scale of modern technosignature datasets – hundreds of hours of high-time- and frequency-resolution data per survey, often containing many thousands of candidate “hits” dominated by anthropogenic RFI – has made machine learning an increasingly central tool for both signal detection and RFI rejection. ? presented the first large-scale deep-learning technosignature search, applying a  $\beta$ -convolutional variational autoencoder to 480 hours of Breakthrough Listen GBT data on 820 nearby stars in a semi-unsupervised anomaly-detection mode, recovering candidate signals missed by the standard TURBOSETI narrowband drift-search pipeline (?) and reducing the number of signals requiring human vetting by roughly two orders of magnitude, while returning eight signals of interest for re-observation. Building on this, ? identified and corrected a long-standing sensitivity mischaracterisation in TURBOSETI’s signal-to-noise calibration, prompting a retrospective correction of quoted sensitivities in several earlier Breakthrough Listen publications (?). More recently, unsupervised clustering methods (e.g. the “GLOBULAR” algorithm) have been developed specifically to separate genuine anomalies from RFI in narrowband technosignature searches without requiring labelled training examples (?), and further anomaly-detection searches have been applied to combined Parkes and GBT Breakthrough Listen archives (?). These efforts collectively demonstrate that machine-learning-based anomaly detection can recover classes of signal (broadband, non-linearly drifting, or otherwise morphologically unusual) that narrowband, linear-drift matched-filter searches such as TURBOSETI are, by design, insensitive to (?), and that a purely classical, threshold-based search – of the kind we adopt for the present survey, following ? – necessarily leaves some fraction of the true technosignature parameter space unexamined. We return to this point in §?? as a natural direction for follow-up work once the present survey’s baseline dataset exists.

#### 2.5. Statistical frameworks for interpreting non-detections

A non-detection is not, by itself, a null result: it must be translated into a quantitative statement about which regions of transmitter parameter space have been excluded, and with what confidence. ? formalised this with the “Cosmic Haystack” framework, modelling the technosignature search space as an eight-dimensional volume (spanning, e.g., transmitter frequency, bandwidth, location, distance, and duty cycle) and showing that even the combined efforts of every major SETI programme conducted up to 2018 sampled a small fraction of any astrophysically plausible haystack volume – a result frequently paraphrased as showing that humanity has searched “the equivalent of a large hot tub’s worth of the Earth’s oceans”. ? subsequently proposed a complementary “nine axes of merit” rubric for comparing the relative scientific value of different technosignature search strategies (spanning axes such as detectability, transmitter cost and information content), providing a qualitative framework alongside Wright et al.’s quantitative one. ? developed an explicit formalism for converting a survey’s non-detection rate into an upper bound on the fraction of stars in the search volume hosting a detectable transmitter, but noted explicitly that this inference is only valid if the observed sample is statistically representative of the underlying stellar population of the search volume – a requirement that most targeted, interest-selected samples (including our own earlier 100-star pilot list) do not, by design, satisfy. Most recently, ? inverted the usual question, asking at what distance Earth’s own technosignatures would themselves be detectable by an ALMA- or SKA-like observatory, providing a valuable calibration point (an “ichnoscale” of unity) against which the sensitivity of any given survey, including the present one, can be benchmarked.

#### 2.6. The case for a volume-limited, representative sample

The great majority of the surveys reviewed above – including our own earlier 100-star ALMA pilot programme – select targets by some combination of proximity, Sun-like spectral type, known exoplanets, protoplanetary-disc or debris-disc detection, or membership of specially-constructed target lists such as the Earth Transit Zone. Such selections

are entirely reasonable when the goal is to maximise the a priori probability of detecting a genuine technosignature per unit observing time, but they carry an unavoidable cost: they make it difficult to convert a survey’s results into a statistically clean statement about the prevalence of technosignatures in the general stellar population, precisely because the sample is not representative of that population (?). A disc-hosting or exoplanet-hosting bias is particularly worth flagging in the ALMA context: ALMA-detectable circumstellar and protoplanetary discs are, almost without exception, orders of magnitude brighter than the present-day Solar System’s own zodiacal dust cloud would appear at interstellar distances, so that samples selected *because* they have an ALMA-detected disc are biased towards young, dynamically active systems that are not necessarily more (and could plausibly be less) likely to host a mature, Earth-like technological civilisation than an old, quiescent, disc-poor star of the same spectral type.

The present survey is explicitly designed to avoid this selection effect. We construct our sample by identifying every star within 20 pc of the Sun that happens to fall within the field of view of at least one public ALMA observation – irrespective of whether that observation was proposed to study a disc, a stellar flare, an unrelated background source, or anything else – and irrespective of the target star’s own spectral type, age, multiplicity, or known planet-hosting status. The star need not be at the pointing centre of the observation, provided it lies within the usable primary-beam field of view (see §??). This is, in effect, a volume-limited “bycatch” census of the ALMA archive: rather than sampling stars selected for astrophysical interest in the calibrator fields as in ?, we sample every nearby star that ALMA happens to have observed for any purpose whatsoever, within a fixed physical volume. We recognise, and quantify explicitly in Paper II, that this approach is still subject to a residual, second-order selection effect – ALMA archival coverage of the immediate solar neighbourhood is itself not spatially uniform, since some regions and spectral types are more likely to have been proposed for ALMA time than others – but this bias operates on the presence of *any* ALMA coverage of a star, not on the astrophysical properties that are the actual subject of a technosignature search, and can therefore be characterised and reported as a well-defined completeness function of the 20 pc volume, rather than an implicit and unquantified bias in the target selection itself.

### 2.7. Prospects for next-generation facilities

The coming decade will see a substantial expansion of technosignature search capability. The Square Kilometre Array (SKA; ??), now under construction in Australia (SKA-Low) and South Africa (SKA-Mid), will combine wide instantaneous bandwidth, high angular resolution, and an architecture well suited to commensal, multi-beam technosignature searching, with projected sensitivity sufficient to detect Earth-analogue-level radar and communication leakage from significantly greater distances than is possible today. The next-generation Very Large Array (ngVLA; ?) is specifically being designed with technosignature science as one of its commensal use cases, with proposed antenna configurations optimised to allow of order 64 simultaneous high-spectral-resolution commensal beams. Existing commensal systems at MeerKAT, the VLA (COSMIC), and the Murchison Widefield Array are already providing a preview of the observing modes these larger facilities will employ at scale. None of these facilities, however, offer ALMA’s combination of angular resolution, spectral resolution and raw collecting area above  $\sim 50$  GHz in the near term, so a systematic mm/submm technosignature census – of the kind begun here – is likely to remain a uniquely ALMA-enabled measurement for some years yet, and represents a natural complement to the lower-frequency, larger-sample surveys that will be enabled by SKA and ngVLA.

## 3. SAMPLE SELECTION

Our sample comprises every star within 20 pc of the Sun (based on Gaia DR3 parallaxes) with at least one public ALMA observation, of any band, science category, or proposal type, in which the star’s true (proper-motion-corrected) position at the epoch of observation falls within the field of view of at least one execution block – not necessarily at the observation’s pointing centre, since a substantial fraction of ALMA mosaics and wide single-pointing observations serendipitously cover more than one catalogued star. We deliberately do not weight target selection by the presence of a circumstellar disc, known exoplanets, spectral type, or any other astrophysical property beyond distance and archival data availability; disc presence and planet-hosting status are retained purely as descriptive metadata for the final sample, not as selection or ranking criteria. At the time of writing this identifies 120 unique target stars, spanning spectral types from early A (e.g. Sirius B,  $\alpha$  CMa B) through to the latest, faintest M dwarfs (e.g. Proxima Centauri, TRAPPIST-1, Barnard’s Star, Teegarden’s Star), ranked by increasing distance from the Sun so that the nearest, most sensitively-constrained systems are processed first. The underlying pipeline is built to extend this volume-limited approach to 30, 40 or 50 pc as a natural next step, at the cost of a rapidly increasing target list and a correspondingly reduced fraction with existing ALMA coverage, since the archive’s areal completeness at fixed exposure declines with distance.

## 4. DATA AND METHODOLOGY

For each target we mine the public ALMA science archive for all calibrated (or, where necessary, locally recalibrated from raw data) execution blocks covering the star’s position, apply an efficiency-bounded download and processing budget per target to remain within the resource constraints of the shared compute facility used for this work, and search the resulting visibility data in two complementary ways: (i) a narrowband, Doppler-drift-corrected search for technosignature candidates, following the general methodology of ? and ?, converting any non-detection into an equivalent isotropic radiated power (EIRP) limit via  $\text{EIRP}_{\min} = 4\pi d^2 S_{\min} \Delta\nu$ ; and (ii) a line-excluded continuum search for anomalous broad-band emission at the star’s position, with an accompanying epoch-to-epoch variability check wherever multiple execution blocks exist. Stellar positions are propagated to the epoch of each observation using

Gaia DR3 proper motions and parallaxes. The Doppler-drift search range applied to each target is set to the larger of (a) a generic frequency-scaled default following ?’s adoption of a literature fiducial with a threefold safety margin, and (b) where the target hosts one or more confirmed exoplanets, an explicit calculation of the maximum line-of-sight orbital acceleration (and hence maximum plausible transmitter drift rate) of the innermost known planet, so that a transmitter co-located with a known, short-period planet cannot be missed simply because its orbital motion drifts it out of an insufficiently wide search window. Full details of the pipeline, its validation, and the specific corrections applied for interferometric mosaic fields and for spectral-line contamination of continuum measurements, will be given in Paper II together with the resulting sample-wide detection limits.

## 5. SCOPE OF PAPER II

Paper II of this series will present: the completed narrowband and continuum technosignature search of the full 20 pc sample described in §??; per-target and aggregate EIRP and continuum flux-density upper limits, tabulated together with the exact frequency ranges searched in each case; any candidate signals identified, together with their vetting against known interference and astrophysical contaminants; a discussion of the sample’s completeness as a function of spectral type and distance within 20 pc; and an assessment, following the statistical frameworks of ?, ? and ?, of what fraction of the local stellar population our non-detections allow us to exclude as hosting a detectable millimetre/submillimetre transmitter of Earth-2026-equivalent power. We will also discuss prospects for extending the survey to 30, 40 and 50 pc, and for applying the anomaly-detection and machine-learning techniques reviewed in §?? to the same underlying dataset as a complementary search methodology to the classical narrowband/continuum approach adopted here.

## ACKNOWLEDGEMENTS

This paper makes use of the following ALMA data: [to be completed in Paper II with the full list of ALMA project codes used]. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), MOST and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. This work has made use of data from the European Space Agency (ESA) mission *Gaia*, processed by the *Gaia* Data Processing and Analysis Consortium (DPAC).

## DATA AVAILABILITY

The reduced data products underlying this paper, together with the full target-selection and analysis pipeline, are being made available at <https://github.com/Tilanthi/SETI> as the survey progresses. Raw ALMA visibility data are publicly available from the ALMA Science Archive at <https://almascience.org>.

## REFERENCES

- Backus P. R., Project Phoenix Team, 2002, in *Bulletin of the American Astronomical Society*. p. 1173
- Choza C., et al., 2024, *AJ*, 168, 254
- Cocconi G., Morrison P., 1959, *Nature*, 184, 844
- Czech D., et al., 2021, *PASP*, 133, 064502
- Czech D. J., et al., 2026, *arXiv:2607.23651*
- Drake F. D., 1960, *Physics Today*, 13, 40
- Drake F. D., 1974, in Sagan C., ed., *Communication with Extraterrestrial Intelligence (CETI)*. MIT Press
- Drake F. D., 1986, in *Applications of Digital Image Processing IX*. SPIE
- Enriquez E., Price D., 2019, *turboSETI*, *Astrophysics Source Code Library*, record ascl:1906.006
- Enriquez J. E., et al., 2017, *ApJ*, 849, 104
- Gajjar V., et al., 2019, in *Bulletin of the American Astronomical Society. AAS Meeting #233*
- Garrett M. A., et al., 2026, *MNRAS*, accepted (*arXiv:2605.10212*)
- Isaacson H., et al., 2017, *PASP*, 129, 054501
- Jacobson-Bell K., et al., 2025, *AJ*, 169, 233
- Johnson O. A., et al., 2023, *AJ*, 166, 193
- Lacki B. C., et al., 2021, *arXiv:2101.02244*
- Li J.-K., Tao Z.-Z., Zhang T.-J., 2026, *arXiv:2603.19023*
- Ma P. X., et al., 2023, *Nature Astronomy*, 7, 492
- Margot J.-L., et al., 2023, *AJ*, 166, 189
- Mason L. A., Garrett M. A., Wandia K., Siemion A. P. V., 2024, *MNRAS*, 536, 2127
- Morrison I. S., et al., 2023, *PASA*, 40, e038
- Ng C., et al., 2022, *AJ*, 164, 205
- Poznanski D., et al., 2025, *AJ*, in press (*arXiv:2505.03927*)
- Price D. C., et al., 2020, *AJ*, 159, 86
- Sheikh S. Z., 2020, *International Journal of Astrobiology*, 19, 237
- Sheikh S. Z., et al., 2025, *AJ*, 169, 108
- Sheikh S. Z., et al., 2025, *arXiv:2512.18142* (“A Search for Radio Technosignatures from Interstellar Object 3I/ATLAS with the Allen Telescope Array”)
- Siemion A. P. V., et al., 2015, *arXiv:1412.4867*
- Tarter J., 2001, *ARA&A*, 39, 511
- Tremblay C. D., et al., 2024, *PASA*, 41, e004
- Tremblay C. D., et al., 2026, in *Advancing Astrophysics with the SKA II (AASKAII)*, *arXiv:2606.27565*
- Tusay N., Sheikh S. Z., Sneed E. L., Farah W., Pollak A. W., Cruz L. F., Siemion A., DeBoer D. R., Wright J. T., 2024, *AJ*, 168, 283
- Tusay N., et al., 2025, *arXiv:2506.13459*
- Valente G., et al., 2025, *Acta Astronautica*, in press
- Vidal C., et al., 2026, *arXiv:2605.21093* (“The Search for Technosignatures: a Review of Possibilities”)
- Von Korff J., et al., 2025, *AJ*, 170, 112
- Wright J. T., Kanodia S., Lubar E. G., 2018, *AJ*, 156, 260
- Zhang Z.-S., et al., 2020, *ApJ*, 891, 174

This paper was built using the Open Journal of Astrophysics L<sup>A</sup>T<sub>E</sub>X template. The OJA is a journal which provides fast and easy peer review for new papers in the **astro-ph** section of the arXiv, making the reviewing process simpler for authors and referees alike. Learn more at <http://astro.theoj.org>.