

A First Course in
LINEAR ALGEBRA

Lecture Notes
for Math 1503

**Determinants: Basic Techniques and
Properties**

Creative Commons License (CC BY-NC-SA)

A First Course in Linear Algebra

Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Determinant of a 2×2 Matrix

Definition

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the **determinant** of A is defined as

$$\det A = ad - bc$$

Determinant of a 2×2 Matrix

Definition

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then the **determinant** of A is defined as

$$\det A = ad - bc$$

Notation. For $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we often write $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$, i.e., use **vertical bars** instead of **square brackets**.

How do we find the determinant of an $n \times n$ matrix?

The determinant of an $n \times n$ matrix is defined recursively, using determinants of $(n - 1) \times (n - 1)$ submatrices, and requires some new definitions and notation.

How do we find the determinant of an $n \times n$ matrix?

The determinant of an $n \times n$ matrix is defined recursively, using determinants of $(n - 1) \times (n - 1)$ submatrices, and requires some new definitions and notation.

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix. The **sign** of the (i, j) position is $(-1)^{i+j}$. Thus the sign is 1 if $(i + j)$ is even, and -1 if $(i + j)$ is odd.

The Minor of a Matrix

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix. The ij^{th} minor of A , denoted as $minor(A)_{ij}$, is the determinant of the $n - 1 \times n - 1$ matrix which results from deleting the i^{th} row and the j^{th} column of A .

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1j} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2j} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{i1} & a_{i2} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nj} & \cdots & a_{nn} \end{bmatrix}$$

For any matrix A , $minor(A)_{ij}$ is found by first removing the i^{th} row and j^{th} column, and taking the determinant of the remaining matrix.

The Minor of a Matrix

Example

Let

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

Find $\text{minor}(A)_{12}$.

The Minor of a Matrix

Example

Let

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

Find $\text{minor}(A)_{12}$.

Solution

First, remove the 1st row and 2nd column from A .

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

The Minor of a Matrix

Example

Let

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

Find $\text{minor}(A)_{12}$.

Solution

First, remove the 1st row and 2nd column from A .

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

The resulting matrix is $A = \begin{bmatrix} 2 & 1 \\ 5 & 6 \end{bmatrix}$

Solution (continued)

$$A = \begin{bmatrix} 2 & 1 \\ 5 & 6 \end{bmatrix}$$

Solution (continued)

$$A = \begin{bmatrix} 2 & 1 \\ 5 & 6 \end{bmatrix}$$

Using our previous definition, we can calculate the determinant of this matrix to be

$$(2)(6) - (5)(1) = 12 - 5 = 7$$

Solution (continued)

$$A = \begin{bmatrix} 2 & 1 \\ 5 & 6 \end{bmatrix}$$

Using our previous definition, we can calculate the determinant of this matrix to be

$$(2)(6) - (5)(1) = 12 - 5 = 7$$

Therefore, $\text{minor}(A)_{12} = 7$.

The Cofactors of a Matrix

Definition

The ij^{th} cofactor of A is

$$\text{cof}(A)_{ij} = (-1)^{i+j} \text{minor}(A)_{ij}$$

The Cofactors of a Matrix

Definition

The ij^{th} cofactor of A is

$$\text{cof}(A)_{ij} = (-1)^{i+j} \text{minor}(A)_{ij}$$

Example (continued)

Let

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

Find $\text{cof}(A)_{12}$.

Solution

By the definition, we know that $\text{cof}(A)_{12} = (-1)^{1+2} \text{minor}(A)_{12}$

Solution

By the definition, we know that $\text{cof}(A)_{12} = (-1)^{1+2} \text{minor}(A)_{12}$

From earlier, we know that $\text{minor}(A)_{12} = 7$.

Solution

By the definition, we know that $\text{cof}(A)_{12} = (-1)^{1+2} \text{minor}(A)_{12}$

From earlier, we know that $\text{minor}(A)_{12} = 7$.

Therefore, $\text{cof}(A)_{12} = (-1)^{1+2} \text{minor}(A)_{12} = (-1)^3 7 = -7$

Cofactor Expansion

Using these definitions, we can now define the **determinant of the $n \times n$ matrix A** :

Definition

$$\det A = a_{11}\text{cof}(A)_{11} + a_{12}\text{cof}(A)_{12} + a_{13}\text{cof}(A)_{13} + \cdots + a_{1n}\text{cof}(A)_{1n}$$

This is called the **cofactor expansion of $\det A$ along row 1**.

Cofactor Expansion

Using these definitions, we can now define the **determinant of the $n \times n$ matrix A** :

Definition

$$\det A = a_{11}\text{cof}(A)_{11} + a_{12}\text{cof}(A)_{12} + a_{13}\text{cof}(A)_{13} + \cdots + a_{1n}\text{cof}(A)_{1n}$$

This is called the **cofactor expansion of $\det A$ along row 1**.

We may also write:

$$\det(A) = \sum_{j=1}^n a_{1j}\text{cof}(A)_{1j} = \sum_{i=1}^n a_{ij}\text{cof}(A)_{ij}$$

Cofactor Expansion

Using these definitions, we can now define the **determinant of the $n \times n$ matrix A** :

Definition

$$\det A = a_{11}\text{cof}(A)_{11} + a_{12}\text{cof}(A)_{12} + a_{13}\text{cof}(A)_{13} + \cdots + a_{1n}\text{cof}(A)_{1n}$$

This is called the **cofactor expansion of $\det A$ along row 1**.

We may also write:

$$\det(A) = \sum_{j=1}^n a_{1j}\text{cof}(A)_{1j} = \sum_{i=1}^n a_{ij}\text{cof}(A)_{ij}$$

The first formula consists of expanding the determinant along the i^{th} row and the second expands the determinant along the j^{th} column.

Cofactor Expansion

Using these definitions, we can now define the **determinant of the $n \times n$ matrix A** :

Definition

$$\det A = a_{11}\text{cof}(A)_{11} + a_{12}\text{cof}(A)_{12} + a_{13}\text{cof}(A)_{13} + \cdots + a_{1n}\text{cof}(A)_{1n}$$

This is called the **cofactor expansion of $\det A$ along row 1**.

We may also write:

$$\det(A) = \sum_{j=1}^n a_{1j}\text{cof}(A)_{1j} = \sum_{i=1}^n a_{ij}\text{cof}(A)_{ij}$$

The first formula consists of expanding the determinant along the i^{th} row and the second expands the determinant along the j^{th} column.

Cofactor expansion is also called **Laplace Expansion**.

Cofactor Expansion

Example

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$. Find $\det A$.

Cofactor Expansion

Example

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$. Find $\det A$.

Solution

Using cofactor expansion along row 1,

$$\det A = 1\text{cof}_{11}(A) + 1\text{cof}_{12}(A) + 3\text{cof}_{13}(A)$$

Cofactor Expansion

Example

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$. Find $\det A$.

Solution

Using cofactor expansion along row 1,

$$\begin{aligned}\det A &= 1\text{cof}_{11}(A) + 1\text{cof}_{12}(A) + 3\text{cof}_{13}(A) \\ &= 1(-1)^2 \begin{vmatrix} 4 & 1 \\ 2 & 6 \end{vmatrix} + 1(-1)^3 \begin{vmatrix} 2 & 1 \\ 5 & 6 \end{vmatrix} + 3(-1)^4 \begin{vmatrix} 2 & 4 \\ 5 & 2 \end{vmatrix}\end{aligned}$$

Cofactor Expansion

Example

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$. Find $\det A$.

Solution

Using cofactor expansion along row 1,

$$\begin{aligned}\det A &= 1\text{cof}_{11}(A) + 1\text{cof}_{12}(A) + 3\text{cof}_{13}(A) \\ &= 1(-1)^2 \begin{vmatrix} 4 & 1 \\ 2 & 6 \end{vmatrix} + 1(-1)^3 \begin{vmatrix} 2 & 1 \\ 5 & 6 \end{vmatrix} + 3(-1)^4 \begin{vmatrix} 2 & 4 \\ 5 & 2 \end{vmatrix} \\ &= 1(24 - 2) - 1(12 - 5) + 3(4 - 20)\end{aligned}$$

Cofactor Expansion

Example

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$. Find $\det A$.

Solution

Using cofactor expansion along row 1,

$$\begin{aligned}\det A &= 1\text{cof}_{11}(A) + 1\text{cof}_{12}(A) + 3\text{cof}_{13}(A) \\ &= 1(-1)^2 \begin{vmatrix} 4 & 1 \\ 2 & 6 \end{vmatrix} + 1(-1)^3 \begin{vmatrix} 2 & 1 \\ 5 & 6 \end{vmatrix} + 3(-1)^4 \begin{vmatrix} 2 & 4 \\ 5 & 2 \end{vmatrix} \\ &= 1(24 - 2) - 1(12 - 5) + 3(4 - 20) \\ &= 22 - 7 + 3(-16)\end{aligned}$$

Cofactor Expansion

Example

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$. Find $\det A$.

Solution

Using cofactor expansion along row 1,

$$\begin{aligned}\det A &= 1\text{cof}_{11}(A) + 1\text{cof}_{12}(A) + 3\text{cof}_{13}(A) \\&= 1(-1)^2 \begin{vmatrix} 4 & 1 \\ 2 & 6 \end{vmatrix} + 1(-1)^3 \begin{vmatrix} 2 & 1 \\ 5 & 6 \end{vmatrix} + 3(-1)^4 \begin{vmatrix} 2 & 4 \\ 5 & 2 \end{vmatrix} \\&= 1(24 - 2) - 1(12 - 5) + 3(4 - 20) \\&= 22 - 7 + 3(-16) \\&= 22 - 7 - 48\end{aligned}$$

Cofactor Expansion

Example

Let $A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$. Find $\det A$.

Solution

Using cofactor expansion along row 1,

$$\begin{aligned}\det A &= 1\text{cof}_{11}(A) + 1\text{cof}_{12}(A) + 3\text{cof}_{13}(A) \\&= 1(-1)^2 \begin{vmatrix} 4 & 1 \\ 2 & 6 \end{vmatrix} + 1(-1)^3 \begin{vmatrix} 2 & 1 \\ 5 & 6 \end{vmatrix} + 3(-1)^4 \begin{vmatrix} 2 & 4 \\ 5 & 2 \end{vmatrix} \\&= 1(24 - 2) - 1(12 - 5) + 3(4 - 20) \\&= 22 - 7 + 3(-16) \\&= 22 - 7 - 48 \\&= -33\end{aligned}$$

Solution (continued)

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

Now try cofactor expansion along column 2.

Solution (continued)

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

Now try cofactor expansion along column 2.

$$\begin{aligned} \det A &= 1\text{cof}_{12}(A) + 4\text{cof}_{22}(A) + 2\text{cof}_{32}(A) \\ &= 1(-1)^3 \begin{vmatrix} 2 & 1 \\ 5 & 6 \end{vmatrix} + 4(-1)^4 \begin{vmatrix} 1 & 3 \\ 5 & 6 \end{vmatrix} + 2(-1)^5 \begin{vmatrix} 1 & 3 \\ 2 & 1 \end{vmatrix} \\ &= -1(12 - 5) + 4(6 - 15) - 2(1 - 6) \\ &= -(7) + 4(-9) - 2(-5) \\ &= -7 - 36 + 10 \\ &= -33 \end{aligned}$$

Solution (continued)

$$A = \begin{bmatrix} 1 & 1 & 3 \\ 2 & 4 & 1 \\ 5 & 2 & 6 \end{bmatrix}$$

Now try cofactor expansion along column 2.

$$\begin{aligned} \det A &= 1\text{cof}_{12}(A) + 4\text{cof}_{22}(A) + 2\text{cof}_{32}(A) \\ &= 1(-1)^3 \begin{vmatrix} 2 & 1 \\ 5 & 6 \end{vmatrix} + 4(-1)^4 \begin{vmatrix} 1 & 3 \\ 5 & 6 \end{vmatrix} + 2(-1)^5 \begin{vmatrix} 1 & 3 \\ 2 & 1 \end{vmatrix} \\ &= -1(12 - 5) + 4(6 - 15) - 2(1 - 6) \\ &= -(7) + 4(-9) - 2(-5) \\ &= -7 - 36 + 10 \\ &= -33 \end{aligned}$$

We get the same answer!

The Determinant is Well Defined

Theorem

The determinant of an $n \times n$ matrix A can be computed using cofactor expansion along **any row or column** of A .

The Determinant is Well Defined

Theorem

The determinant of an $n \times n$ matrix A can be computed using cofactor expansion along **any row or column** of A .

What is the significance of this theorem?

This theorem allows us to choose any row or column for computing cofactor expansion, which gives us the opportunity to save ourselves some work!

Problem

Let $A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$. Find $\det A$.

Problem

Let $A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$. Find $\det A$.

Solution

Cofactor expansion along row 1 yields

$$\det A = 0 \times \text{cof}(A)_{11} + 1 \times \text{cof}(A)_{12} + (-2) \times \text{cof}(A)_{13} + 1 \times \text{cof}(A)_{14}$$

Problem

Let $A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$. Find $\det A$.

Solution

Cofactor expansion along row 1 yields

$$\begin{aligned}\det A &= 0 \times \text{cof}(A)_{11} + 1 \times \text{cof}(A)_{12} + (-2) \times \text{cof}(A)_{13} + 1 \times \text{cof}(A)_{14} \\ &= \text{cof}(A)_{12} - 2 \times \text{cof}(A)_{13} + \text{cof}(A)_{14},\end{aligned}$$

Problem

Let $A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$. Find $\det A$.

Solution

Cofactor expansion along row 1 yields

$$\begin{aligned}\det A &= 0 \times \text{cof}(A)_{11} + 1 \times \text{cof}(A)_{12} + (-2) \times \text{cof}(A)_{13} + 1 \times \text{cof}(A)_{14} \\ &= \text{cof}(A)_{12} - 2 \times \text{cof}(A)_{13} + \text{cof}(A)_{14},\end{aligned}$$

whereas cofactor expansion along, row 3 yields

$$\det A = 0 \times \text{cof}(A)_{31} + 1 \times \text{cof}(A)_{32} + (-1) \times \text{cof}(A)_{33} + 0 \times \text{cof}(A)_{34}$$

Problem

$$\text{Let } A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}. \text{ Find } \det A.$$

Solution

Cofactor expansion along row 1 yields

$$\begin{aligned} \det A &= 0 \times \text{cof}(A)_{11} + 1 \times \text{cof}(A)_{12} + (-2) \times \text{cof}(A)_{13} + 1 \times \text{cof}(A)_{14} \\ &= \text{cof}(A)_{12} - 2 \times \text{cof}(A)_{13} + \text{cof}(A)_{14}, \end{aligned}$$

whereas cofactor expansion along, row 3 yields

$$\begin{aligned} \det A &= 0 \times \text{cof}(A)_{31} + 1 \times \text{cof}(A)_{32} + (-1) \times \text{cof}(A)_{33} + 0 \times \text{cof}(A)_{34} \\ &= 1 \text{cof}(A)_{32} + (-1) \text{cof}(A)_{33}, \end{aligned}$$

Problem

Let $A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$. Find $\det A$.

Solution

Cofactor expansion along row 1 yields

$$\begin{aligned}\det A &= 0 \times \text{cof}(A)_{11} + 1 \times \text{cof}(A)_{12} + (-2) \times \text{cof}(A)_{13} + 1 \times \text{cof}(A)_{14} \\ &= \text{cof}(A)_{12} - 2 \times \text{cof}(A)_{13} + \text{cof}(A)_{14},\end{aligned}$$

whereas cofactor expansion along, row 3 yields

$$\begin{aligned}\det A &= 0 \times \text{cof}(A)_{31} + 1 \times \text{cof}(A)_{32} + (-1) \times \text{cof}(A)_{33} + 0 \times \text{cof}(A)_{34} \\ &= 1\text{cof}(A)_{32} + (-1)\text{cof}(A)_{33},\end{aligned}$$

i.e., in the first case we must compute **three** cofactors, but in the second case we need only compute **two** cofactors.

Solution (continued)

Therefore, we can save ourselves some work by using cofactor expansion along row 3 rather than row 1.

$$A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$$

$$\det A = 1 \times \text{cof}(A)_{32} + (-1) \times \text{cof}(A)_{33}$$

Solution (continued)

Therefore, we can save ourselves some work by using cofactor expansion along row 3 rather than row 1.

$$A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$$

$$\begin{aligned} \det A &= 1 \times \text{cof}(A)_{32} + (-1) \times \text{cof}(A)_{33} \\ &= 1 \times (-1)^5 \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} + (-1) \times (-1)^6 \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \end{aligned}$$

Solution (continued)

Therefore, we can save ourselves some work by using cofactor expansion along row 3 rather than row 1.

$$A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$$

$$\begin{aligned} \det A &= 1 \times \text{cof}(A)_{32} + (-1) \times \text{cof}(A)_{33} \\ &= 1 \times (-1)^5 \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} + (-1) \times (-1)^6 \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\ &= - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \end{aligned}$$

Solution (continued)

Therefore, we can save ourselves some work by using cofactor expansion along row 3 rather than row 1.

$$A = \begin{bmatrix} 0 & 1 & -2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$$

$$\begin{aligned} \det A &= 1 \times \text{cof}(A)_{32} + (-1) \times \text{cof}(A)_{33} \\ &= 1 \times (-1)^5 \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} + (-1) \times (-1)^6 \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\ &= - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \end{aligned}$$

Each of the two determinants above can easily be evaluated using **cofactor expansion along column 2**.

Solution (continued)

$$\det A = - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix}$$

Solution (continued)

$$\begin{aligned}\det A &= - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\ &= -(-2)(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} - 1(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix}\end{aligned}$$

Solution (continued)

$$\begin{aligned}\det A &= - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\ &= -(-2)(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} - 1(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} \\ &= -2(10 - 21) + 1(10 - 21)\end{aligned}$$

Solution (continued)

$$\begin{aligned}\det A &= - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\&= -(-2)(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} - 1(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} \\&= -2(10 - 21) + 1(10 - 21) \\&= -2(-11) + (-11)\end{aligned}$$

Solution (continued)

$$\begin{aligned}\det A &= - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\&= -(-2)(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} - 1(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} \\&= -2(10 - 21) + 1(10 - 21) \\&= -2(-11) + (-11) \\&= 22 - 11\end{aligned}$$

Solution (continued)

$$\begin{aligned}\det A &= - \begin{vmatrix} 0 & -2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} - \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\&= -(-2)(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} - 1(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} \\&= -2(10 - 21) + 1(10 - 21) \\&= -2(-11) + (-11) \\&= 22 - 11 \\&= 11.\end{aligned}$$

Therefore, $\det A = 11$.

A Row or Column of Zeros

Example

Let

$$A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}.$$

A Row or Column of Zeros

Example

Let

$$A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}.$$

By choosing column 3 for cofactor expansion, we get $\det A = 0$,

A Row or Column of Zeros

Example

Let

$$A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}.$$

By choosing column 3 for cofactor expansion, we get $\det A = 0$, i.e.,

$$\det A = 0 \times \text{cof}(A)_{13} + 0 \times \text{cof}(A)_{23} + 0 \times \text{cof}(A)_{33} + 0 \times \text{cof}(A)_{43}$$

A Row or Column of Zeros

Example

Let

$$A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}.$$

By choosing column 3 for cofactor expansion, we get $\det A = 0$, i.e.,

$$\det A = 0 \times \text{cof}(A)_{13} + 0 \times \text{cof}(A)_{23} + 0 \times \text{cof}(A)_{33} + 0 \times \text{cof}(A)_{43} = 0.$$

A Row or Column of Zeros

Example

Let

$$A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}.$$

By choosing column 3 for cofactor expansion, we get $\det A = 0$, i.e.,

$$\det A = 0 \times \text{cof}(A)_{13} + 0 \times \text{cof}(A)_{23} + 0 \times \text{cof}(A)_{33} + 0 \times \text{cof}(A)_{43} = 0.$$

Important Fact

If A is an $n \times n$ matrix with a row or column of zeros, then $\det A = 0$.

Determinants of a Triangular Matrices

Definitions

- 1 An $n \times n$ matrix A is called **upper triangular** if all entries **below** the main diagonal are zero.

Determinants of a Triangular Matrices

Definitions

- 1 An $n \times n$ matrix A is called **upper triangular** if all entries **below** the main diagonal are zero.
- 2 An $n \times n$ matrix A is called **lower triangular** if all entries **above** the main diagonal are zero.

Determinants of a Triangular Matrices

Definitions

- 1 An $n \times n$ matrix A is called **upper triangular** if all entries **below** the main diagonal are zero.
- 2 An $n \times n$ matrix A is called **lower triangular** if all entries **above** the main diagonal are zero.
- 3 An $n \times n$ matrix A is called **triangular** if it is upper triangular or lower triangular.

Determinants of a Triangular Matrices

Definitions

- 1 An $n \times n$ matrix A is called **upper triangular** if all entries **below** the main diagonal are zero.
- 2 An $n \times n$ matrix A is called **lower triangular** if all entries **above** the main diagonal are zero.
- 3 An $n \times n$ matrix A is called **triangular** if it is upper triangular or lower triangular.

Theorem

If $A = [a_{ij}]$ is an $n \times n$ triangular matrix, then

$$\det A = a_{11} \times a_{22} \times a_{33} \times \cdots \times a_{nn},$$

i.e., $\det A$ is the product of the entries of the main diagonal of A .

Example

$$\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix}$$

Example

$$\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix} = 1 \times \det \begin{bmatrix} 5 & 6 \\ 0 & 9 \end{bmatrix}$$

Example

$$\begin{aligned}\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix} &= 1 \times \det \begin{bmatrix} 5 & 6 \\ 0 & 9 \end{bmatrix} \\ &= 1 \times 5 \times \det [9]\end{aligned}$$

Example

$$\begin{aligned}\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix} &= 1 \times \det \begin{bmatrix} 5 & 6 \\ 0 & 9 \end{bmatrix} \\ &= 1 \times 5 \times \det [9] \\ &= 1 \times 5 \times 9\end{aligned}$$

Example

$$\begin{aligned}\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix} &= 1 \times \det \begin{bmatrix} 5 & 6 \\ 0 & 9 \end{bmatrix} \\ &= 1 \times 5 \times \det [9] \\ &= 1 \times 5 \times 9 \\ &= 45.\end{aligned}$$

Example

$$\begin{aligned}\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix} &= 1 \times \det \begin{bmatrix} 5 & 6 \\ 0 & 9 \end{bmatrix} \\ &= 1 \times 5 \times \det [9] \\ &= 1 \times 5 \times 9 \\ &= 45.\end{aligned}$$

Notice that 45 is the product of the entries on the main diagonal.

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix}$$

Elementary Row Operations and Determinants

Theorem

Let A be an $n \times n$ matrix and B be an $n \times n$ matrix as defined below.

Elementary Row Operations and Determinants

Theorem

Let A be an $n \times n$ matrix and B be an $n \times n$ matrix as defined below.

- 1 Let B be a matrix which results from switching two rows of A . Then $\det(B) = -\det(A)$.
- 2 Let B be a matrix which results from multiplying some row of A by a scalar k . Then $\det(B) = k \det(A)$.
- 3 Let B be a matrix which results from adding a multiple of a row to another row. Then $\det(A) = \det(B)$.
- 4 If A contains a row which is a multiple of another row of A , then $\det(A) = 0$

Elementary Row Operations and Determinants

Theorem

Let A be an $n \times n$ matrix and B be an $n \times n$ matrix as defined below.

- ① Let B be a matrix which results from switching two rows of A . Then $\det(B) = -\det(A)$.
- ② Let B be a matrix which results from multiplying some row of A by a scalar k . Then $\det(B) = k \det(A)$.
- ③ Let B be a matrix which results from adding a multiple of a row to another row. Then $\det(A) = \det(B)$.
- ④ If A contains a row which is a multiple of another row of A , then $\det(A) = 0$.

An analogous theorem holds for **elementary column operation**. If A is a matrix, then an **elementary column operation** on A is simply the corresponding elementary row operation performed on the transpose of A , A^T .

Computing the Determinant

Example

$$\det \begin{bmatrix} -3 & 5 & -6 \\ 1 & -1 & 3 \\ 2 & -4 & 1 \end{bmatrix}$$

Computing the Determinant

Example

$$\det \begin{bmatrix} -3 & 5 & -6 \\ 1 & -1 & 3 \\ 2 & -4 & 1 \end{bmatrix} =$$

Computing the Determinant

Example

$$\det \begin{bmatrix} -3 & 5 & -6 \\ 1 & -1 & 3 \\ 2 & -4 & 1 \end{bmatrix} =$$

Computing the Determinant

Example

$$\det \begin{bmatrix} -3 & 5 & -6 \\ 1 & -1 & 3 \\ 2 & -4 & 1 \end{bmatrix} =$$

Computing the Determinant

Example

$$\det \begin{bmatrix} -3 & 5 & -6 \\ 1 & -1 & 3 \\ 2 & -4 & 1 \end{bmatrix} =$$

Computing the Determinant

Example

$$\det \begin{bmatrix} -3 & 5 & -6 \\ 1 & -1 & 3 \\ 2 & -4 & 1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix}$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} =$$

Problem

$$\text{If } \det \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} = 4, \text{ find } \det \begin{bmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{bmatrix}.$$

Problem

$$\text{If } \det \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} = 4, \text{ find } \det \begin{bmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{bmatrix}.$$

Solution

$$\begin{vmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{vmatrix}$$

Problem

$$\text{If } \det \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} = 4, \text{ find } \det \begin{bmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{bmatrix}.$$

Solution

$$\begin{aligned} \begin{vmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{vmatrix} &= (-1)(3) \begin{vmatrix} b_1 & b_2 & b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= (-3) \begin{vmatrix} b_1 & b_2 & b_3 \\ a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \end{aligned}$$

Problem

$$\text{If } \det \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} = 4, \text{ find } \det \begin{bmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{bmatrix}.$$

Solution

$$\begin{aligned} \begin{vmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{vmatrix} &= (-1)(3) \begin{vmatrix} b_1 & b_2 & b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= (-3) \begin{vmatrix} b_1 & b_2 & b_3 \\ a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= (-3)(-1) \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= (-3)(-1) \times 4 \end{aligned}$$

Problem

$$\text{If } \det \begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{bmatrix} = 4, \text{ find } \det \begin{bmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{bmatrix}.$$

Solution

$$\begin{aligned} \begin{vmatrix} -b_1 & -b_2 & -b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ 3c_1 & 3c_2 & 3c_3 \end{vmatrix} &= (-1)(3) \begin{vmatrix} b_1 & b_2 & b_3 \\ a_1 + 2b_1 & a_2 + 2b_2 & a_3 + 2b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= (-3) \begin{vmatrix} b_1 & b_2 & b_3 \\ a_1 & a_2 & a_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= (-3)(-1) \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} \\ &= (-3)(-1) \times 4 \\ &= 12. \end{aligned}$$

Lecture II

Problem

Let $A = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{bmatrix}$. Find $\det A$.

Lecture II

Problem

Let $A = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{bmatrix}$. Find $\det A$.

Solution

$$\det A = \begin{vmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix}$$

Lecture II

Problem

Let $A = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{bmatrix}$. Find $\det A$.

Solution

$$\det A = \begin{vmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix} = \begin{vmatrix} 3 & 5 & 9 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix}$$

Lecture II

Problem

Let $A = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{bmatrix}$. Find $\det A$.

Solution

$$\det A = \begin{vmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix} = \begin{vmatrix} 3 & 5 & 9 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix} = \begin{vmatrix} 0 & 0 & 0 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix} = 0.$$

Lecture II

Problem

Let $A = \begin{bmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{bmatrix}$. Find $\det A$.

Solution

$$\det A = \begin{vmatrix} 2 & 3 & 5 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix} = \begin{vmatrix} 3 & 5 & 9 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix} = \begin{vmatrix} 0 & 0 & 0 \\ 3 & 5 & 9 \\ 1 & 2 & 4 \end{vmatrix} = 0.$$

Notice:

$$\text{row2} + \text{row3} - 2(\text{row1}) = \begin{bmatrix} 0 & 0 & 0 \end{bmatrix}$$

Hence the determinant equals 0.

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

Write $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$.

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

$$\det(-2A) = \begin{vmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

$$\det(-2A) = \begin{vmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

$$\det(-2A) = \begin{vmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

$$= (-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

$$\det(-2A) = \begin{vmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

$$= (-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2)(-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

$$\det(-2A) = \begin{vmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

$$= (-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2)(-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$= (-2)^3 \det A$$

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

$$\det(-2A) = \begin{vmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

$$= (-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2)(-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$= (-2)^3 \det A = (-8) \times 7$$

Determinants and Scalar Multiplication

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-2A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -2A = \begin{bmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{bmatrix}.$$

$$\det(-2A) = \begin{vmatrix} -2a_{11} & -2a_{12} & -2a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -2a_{21} & -2a_{22} & -2a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix}$$

$$= (-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ -2a_{31} & -2a_{32} & -2a_{33} \end{vmatrix} = (-2)(-2)(-2) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$= (-2)^3 \det A = (-8) \times 7 = -56.$$

Solution (continued)

Think about the matrix $-2A$ as the matrix obtained from A by multiplying **each of its rows by -2** .

Solution (continued)

Think about the matrix $-2A$ as the matrix obtained from A by multiplying **each of its rows by -2** . This involves **three elementary row operations**, each of which contributes a factor of -2 to the determinant, and thus

Solution (continued)

Think about the matrix $-2A$ as the matrix obtained from A by multiplying **each of its rows by -2** . This involves **three elementary row operations**, each of which contributes a factor of -2 to the determinant, and thus

$$\det A = (-2) \times (-2) \times (-2) \times 7$$

Solution (continued)

Think about the matrix $-2A$ as the matrix obtained from A by multiplying **each of its rows by -2** . This involves **three elementary row operations**, each of which contributes a factor of -2 to the determinant, and thus

$$\det A = (-2) \times (-2) \times (-2) \times 7 = (-2)^3 \times 7.$$

Solution (continued)

Think about the matrix $-2A$ as the matrix obtained from A by multiplying **each of its rows by -2** . This involves **three elementary row operations**, each of which contributes a factor of -2 to the determinant, and thus

$$\det A = (-2) \times (-2) \times (-2) \times 7 = (-2)^3 \times 7.$$

Theorem

If A is an $n \times n$ matrix and k is any scalar, then

$$\det(kA) = k^n \det A.$$

Determinants of Inverses, Products, and Transposes

Theorem

An $n \times n$ matrix A is invertible if and only if $\det A \neq 0$. In this case,

$$\det(A^{-1}) = \frac{1}{\det A}.$$

Determinants of Inverses, Products, and Transposes

Theorem

An $n \times n$ matrix A is invertible if and only if $\det A \neq 0$. In this case,

$$\det(A^{-1}) = \frac{1}{\det A}.$$

Theorem

Let A and B be $n \times n$ matrices. Then

$$\det(AB) = \det A \times \det B.$$

Determinants of Inverses, Products, and Transposes

Theorem

An $n \times n$ matrix A is invertible if and only if $\det A \neq 0$. In this case,

$$\det(A^{-1}) = \frac{1}{\det A}.$$

Theorem

Let A and B be $n \times n$ matrices. Then

$$\det(AB) = \det A \times \det B.$$

Theorem

If A is an $n \times n$ matrix, then the determinant of its transpose is given by

$$\det(A^T) = \det A.$$

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Solution

$$\det A = \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix}$$

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Solution

$$\det A = \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix} = c \begin{vmatrix} 2 & c \\ c & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & c \end{vmatrix}$$

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Solution

$$\begin{aligned} \det A &= \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix} = c \begin{vmatrix} 2 & c \\ c & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & c \end{vmatrix} \\ &= c(10 - c^2) - c \end{aligned}$$

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Solution

$$\begin{aligned} \det A &= \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix} = c \begin{vmatrix} 2 & c \\ c & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & c \end{vmatrix} \\ &= c(10 - c^2) - c \\ &= c(9 - c^2) \end{aligned}$$

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Solution

$$\begin{aligned} \det A &= \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix} = c \begin{vmatrix} 2 & c \\ c & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & c \end{vmatrix} \\ &= c(10 - c^2) - c \\ &= c(9 - c^2) \\ &= c(3 - c)(3 + c). \end{aligned}$$

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Solution

$$\begin{aligned}\det A &= \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix} = c \begin{vmatrix} 2 & c \\ c & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & c \end{vmatrix} \\ &= c(10 - c^2) - c \\ &= c(9 - c^2) \\ &= c(3 - c)(3 + c).\end{aligned}$$

Since A is invertible when $\det(A) \neq 0$,

Problem

Find all values of c for which $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$ is invertible.

Solution

$$\begin{aligned} \det A &= \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix} = c \begin{vmatrix} 2 & c \\ c & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & c \end{vmatrix} \\ &= c(10 - c^2) - c \\ &= c(9 - c^2) \\ &= c(3 - c)(3 + c). \end{aligned}$$

Since A is invertible when $\det(A) \neq 0$, A is invertible for all $c \neq 0, 3, -3$.

Problem

Suppose A , B and C are 4×4 matrices with

$$\det A = -1, \det B = 2, \text{ and } \det C = 1.$$

Find $\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$.

Problem

Suppose A , B and C are 4×4 matrices with

$$\det A = -1, \det B = 2, \text{ and } \det C = 1.$$

Find $\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$.

Solution

$$\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$$

Problem

Suppose A , B and C are 4×4 matrices with

$$\det A = -1, \det B = 2, \text{ and } \det C = 1.$$

Find $\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$.

Solution

$$\det(2A^2(B^{-1})(C^T)^3B(A^{-1})) = 2^4(\det A)^2 \frac{1}{\det B} (\det C)^3 (\det B) \frac{1}{\det A}$$

Problem

Suppose A , B and C are 4×4 matrices with

$$\det A = -1, \det B = 2, \text{ and } \det C = 1.$$

Find $\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$.

Solution

$$\begin{aligned}\det(2A^2(B^{-1})(C^T)^3B(A^{-1})) &= 2^4(\det A)^2 \frac{1}{\det B} (\det C)^3 (\det B) \frac{1}{\det A} \\ &= 16(\det A)(\det C)^3\end{aligned}$$

Problem

Suppose A , B and C are 4×4 matrices with

$$\det A = -1, \det B = 2, \text{ and } \det C = 1.$$

Find $\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$.

Solution

$$\begin{aligned}\det(2A^2(B^{-1})(C^T)^3B(A^{-1})) &= 2^4(\det A)^2 \frac{1}{\det B} (\det C)^3 (\det B) \frac{1}{\det A} \\ &= 16(\det A)(\det C)^3 \\ &= 16 \times (-1) \times 1^3\end{aligned}$$

Problem

Suppose A , B and C are 4×4 matrices with

$$\det A = -1, \det B = 2, \text{ and } \det C = 1.$$

Find $\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$.

Solution

$$\begin{aligned}\det(2A^2(B^{-1})(C^T)^3B(A^{-1})) &= 2^4(\det A)^2 \frac{1}{\det B} (\det C)^3 (\det B) \frac{1}{\det A} \\ &= 16(\det A)(\det C)^3 \\ &= 16 \times (-1) \times 1^3 \\ &= -16.\end{aligned}$$

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Solution

Since $\det(-A) = (-1)^n \det A$,

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Solution

Since $\det(-A) = (-1)^n \det A$, $\det(-A) = \det A$ if and only if

$$(-1)^n \det A = \det A.$$

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Solution

Since $\det(-A) = (-1)^n \det A$, $\det(-A) = \det A$ if and only if

$$(-1)^n \det A = \det A.$$

When is this possible?

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Solution

Since $\det(-A) = (-1)^n \det A$, $\det(-A) = \det A$ if and only if

$$(-1)^n \det A = \det A.$$

When is this possible?

- 1 $(-1)^n \det A = \det A$ whenever $\det A = 0$.

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Solution

Since $\det(-A) = (-1)^n \det A$, $\det(-A) = \det A$ if and only if

$$(-1)^n \det A = \det A.$$

When is this possible?

- 1 $(-1)^n \det A = \det A$ whenever $\det A = 0$.
- 2 If $\det A \neq 0$, then $(-1)^n \det A = \det A$ only if $(-1)^n = 1$,

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Solution

Since $\det(-A) = (-1)^n \det A$, $\det(-A) = \det A$ if and only if

$$(-1)^n \det A = \det A.$$

When is this possible?

- 1 $(-1)^n \det A = \det A$ whenever $\det A = 0$.
- 2 If $\det A \neq 0$, then $(-1)^n \det A = \det A$ only if $(-1)^n = 1$, i.e., only if n is even.

Problem

Let A be an $n \times n$ matrix. Find all conditions that ensure $\det(-A) = \det A$.

Solution

Since $\det(-A) = (-1)^n \det A$, $\det(-A) = \det A$ if and only if

$$(-1)^n \det A = \det A.$$

When is this possible?

- 1 $(-1)^n \det A = \det A$ whenever $\det A = 0$.
- 2 If $\det A \neq 0$, then $(-1)^n \det A = \det A$ only if $(-1)^n = 1$, i.e., only if n is even.

Therefore, $\det(-A) = \det A$ only if **$\det A = 0$ or n is even.**

The Cofactor Matrix

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix. The **cofactor matrix of A** , is the matrix

$$[\text{cof}(A)_{ij}],$$

i.e., the matrix whose (i, j) -entry is the (i, j) -cofactor of A .

The Cofactor Matrix

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix. The **cofactor matrix of A** , is the matrix

$$[\text{cof}(A)_{ij}],$$

i.e., the matrix whose (i, j) -entry is the (i, j) -cofactor of A .

Reminder: the (i, j) -cofactor

$$\text{cof}(A)_{ij} = (-1)^{i+j} \text{minor}(A)_{ij},$$

where $\text{minor}(A)_{ij}$ is the determinant of the matrix obtained from A by deleting row i and column j .

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11}$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix}$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42 = -6.$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42 = -6.$$

$$\text{cof}(A)_{12} = (-1)^{1+2} \det A_{12}$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42 = -6.$$

$$\text{cof}(A)_{12} = (-1)^{1+2} \det A_{12} = - \begin{vmatrix} 1 & 7 \\ 0 & 4 \end{vmatrix}$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42 = -6.$$

$$\text{cof}(A)_{12} = (-1)^{1+2} \det A_{12} = - \begin{vmatrix} 1 & 7 \\ 0 & 4 \end{vmatrix} = -(4 - 0)$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42 = -6.$$

$$\text{cof}(A)_{12} = (-1)^{1+2} \det A_{12} = - \begin{vmatrix} 1 & 7 \\ 0 & 4 \end{vmatrix} = -(4 - 0) = -4.$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42 = -6.$$

$$\text{cof}(A)_{12} = (-1)^{1+2} \det A_{12} = - \begin{vmatrix} 1 & 7 \\ 0 & 4 \end{vmatrix} = -(4 - 0) = -4.$$

$$\text{cof}(A)_{13}$$

Problem

Find the cofactor matrix $[\text{cof}(A)_{ij}]$ of the matrix

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Solution

For each i and j , $1 \leq i, j \leq 3$, we need to compute $\text{cof}(A)_{ij}$, so there are **9 cofactors** to compute.

$$\text{cof}(A)_{11} = (-1)^{1+1} \det A_{11} = \begin{vmatrix} 9 & 7 \\ 6 & 4 \end{vmatrix} = 9 \times 4 - 6 \times 7 = 36 - 42 = -6.$$

$$\text{cof}(A)_{12} = (-1)^{1+2} \det A_{12} = - \begin{vmatrix} 1 & 7 \\ 0 & 4 \end{vmatrix} = -(4 - 0) = -4.$$

$$\text{cof}(A)_{13} = (-1)^{1+3} \det A_{12} = \begin{vmatrix} 1 & 9 \\ 0 & 6 \end{vmatrix} = (6 - 0) = 6.$$

Solution (continued)

Computing the six remaining cofactors results in the cofactor matrix

Solution (continued)

Computing the six remaining cofactors results in the cofactor matrix

$$\begin{bmatrix} -6 & -4 & 6 \\ 18 & 16 & -24 \\ -27 & -25 & 36 \end{bmatrix}.$$

The Adjugate

Definition

If A is an $n \times n$ matrix, then the **adjugate of A** is defined by

$$\text{adj } A = [\text{cof}(A)_{ij}]^T,$$

where $\text{cof}(A)_{ij}$ is the (i,j) -cofactor of A , i.e., **$\text{adj } A$ is the transpose of the cofactor matrix.**

The Adjugate

Definition

If A is an $n \times n$ matrix, then the **adjugate of A** is defined by

$$\text{adj } A = [\text{cof}(A)_{ij}]^T,$$

where $\text{cof}(A)_{ij}$ is the (i,j) -cofactor of A , i.e., **adj A** is the transpose of the cofactor matrix.

Example

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}, \text{ has cofactor matrix } \begin{bmatrix} -6 & -4 & 6 \\ 18 & 16 & -24 \\ -27 & -25 & 36 \end{bmatrix}.$$

The Adjugate

Definition

If A is an $n \times n$ matrix, then the **adjugate of A** is defined by

$$\text{adj } A = [\text{cof}(A)_{ij}]^T,$$

where $\text{cof}(A)_{ij}$ is the (i,j) -cofactor of A , i.e., **$\text{adj } A$ is the transpose of the cofactor matrix.**

Example

$$A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}, \text{ has cofactor matrix } \begin{bmatrix} -6 & -4 & 6 \\ 18 & 16 & -24 \\ -27 & -25 & 36 \end{bmatrix}.$$

Therefore, the adjugate of A is

$$\text{adj } A = \begin{bmatrix} -6 & -4 & 6 \\ 18 & 16 & -24 \\ -27 & -25 & 36 \end{bmatrix}^T = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

Problem

Find $\text{adj } A$ when $A = \begin{bmatrix} 2 & 1 & 3 \\ 5 & -7 & 1 \\ 3 & 0 & -6 \end{bmatrix}$.

Problem

Find $\text{adj } A$ when $A = \begin{bmatrix} 2 & 1 & 3 \\ 5 & -7 & 1 \\ 3 & 0 & -6 \end{bmatrix}$.

Solution

$$\text{adj } A = \begin{bmatrix} 42 & 6 & 22 \\ 33 & -21 & 13 \\ 21 & 3 & -19 \end{bmatrix}.$$

The Adjugate of a 2×2 Matrix

Example

Let A be a 2×2 matrix, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$.

The Adjugate of a 2×2 Matrix

Example

Let A be a 2×2 matrix, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then

$\text{adj } A$

The Adjugate of a 2×2 Matrix

Example

Let A be a 2×2 matrix, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then

$$\text{adj } A = \begin{bmatrix} \text{cof}(A)_{11} & \text{cof}(A)_{12} \\ \text{cof}(A)_{21} & \text{cof}(A)_{22} \end{bmatrix}^T$$

The Adjugate of a 2×2 Matrix

Example

Let A be a 2×2 matrix, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then

$$\text{adj } A = \begin{bmatrix} \text{cof}(A)_{11} & \text{cof}(A)_{12} \\ \text{cof}(A)_{21} & \text{cof}(A)_{22} \end{bmatrix}^T = \begin{bmatrix} (-1)^2 d & (-1)^3 c \\ (-1)^3 b & (-1)^4 a \end{bmatrix}^T$$

The Adjugate of a 2×2 Matrix

Example

Let A be a 2×2 matrix, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then

$$\begin{aligned} \operatorname{adj} A &= \begin{bmatrix} \operatorname{cof}(A)_{11} & \operatorname{cof}(A)_{12} \\ \operatorname{cof}(A)_{21} & \operatorname{cof}(A)_{22} \end{bmatrix}^T = \begin{bmatrix} (-1)^2 d & (-1)^3 c \\ (-1)^3 b & (-1)^4 a \end{bmatrix}^T \\ &= \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}. \end{aligned}$$

The Adjugate of a 2×2 Matrix

Example

Let A be a 2×2 matrix, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then

$$\begin{aligned} \operatorname{adj} A &= \begin{bmatrix} \operatorname{cof}(A)_{11} & \operatorname{cof}(A)_{12} \\ \operatorname{cof}(A)_{21} & \operatorname{cof}(A)_{22} \end{bmatrix}^T = \begin{bmatrix} (-1)^2 d & (-1)^3 c \\ (-1)^3 b & (-1)^4 a \end{bmatrix}^T \\ &= \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}. \end{aligned}$$

We've seen this matrix before: if $\det A \neq 0$, then

$$A^{-1} = \frac{1}{\det A} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

The Adjugate of a 2×2 Matrix

Example

Let A be a 2×2 matrix, say $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. Then

$$\begin{aligned} \operatorname{adj} A &= \begin{bmatrix} \operatorname{cof}(A)_{11} & \operatorname{cof}(A)_{12} \\ \operatorname{cof}(A)_{21} & \operatorname{cof}(A)_{22} \end{bmatrix}^T = \begin{bmatrix} (-1)^2 d & (-1)^3 c \\ (-1)^3 b & (-1)^4 a \end{bmatrix}^T \\ &= \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}. \end{aligned}$$

We've seen this matrix before: if $\det A \neq 0$, then

$$A^{-1} = \frac{1}{\det A} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{1}{\det A} \operatorname{adj} A.$$

Example (continued)

Observe that, regardless of the value of $\det A$,

$$A(\operatorname{adj} A) = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Example (continued)

Observe that, regardless of the value of $\det A$,

$$\begin{aligned} A(\operatorname{adj} A) &= \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \\ &= \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix} \end{aligned}$$

Example (continued)

Observe that, regardless of the value of $\det A$,

$$\begin{aligned} A(\operatorname{adj} A) &= \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \\ &= \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix} \\ &= (ad - bc) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

Example (continued)

Observe that, regardless of the value of $\det A$,

$$\begin{aligned} A(\operatorname{adj} A) &= \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \\ &= \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix} \\ &= (ad - bc) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &= (\det A)I_2. \end{aligned}$$

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$,

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$, has adjugate

$$\text{adj } A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$, has adjugate

$\text{adj } A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$. Computing $A(\text{adj } A)$ we see that

$$A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$$

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$, has adjugate

$\text{adj } A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$. Computing $A(\text{adj } A)$ we see that

$$A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} -6 & 0 & 0 \\ 0 & -6 & 0 \\ 0 & 0 & -6 \end{bmatrix}.$$

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$, has adjugate

$\text{adj } A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$. Computing $A(\text{adj } A)$ we see that

$$A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} -6 & 0 & 0 \\ 0 & -6 & 0 \\ 0 & 0 & -6 \end{bmatrix}.$$

Note that

$$\det A = \begin{vmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{vmatrix}$$

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$, has adjugate

$\text{adj } A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$. Computing $A(\text{adj } A)$ we see that

$$A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} -6 & 0 & 0 \\ 0 & -6 & 0 \\ 0 & 0 & -6 \end{bmatrix}.$$

Note that

$$\det A = \begin{vmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{vmatrix} = \begin{vmatrix} 0 & -36 & -25 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{vmatrix}$$

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$, has adjugate

$\text{adj } A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$. Computing $A(\text{adj } A)$ we see that

$$A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} -6 & 0 & 0 \\ 0 & -6 & 0 \\ 0 & 0 & -6 \end{bmatrix}.$$

Note that

$$\det A = \begin{vmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{vmatrix} = \begin{vmatrix} 0 & -36 & -25 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{vmatrix} = - \begin{vmatrix} -36 & -25 \\ 6 & 4 \end{vmatrix}$$

Example

In an earlier example, we saw that $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$, has adjugate

$\text{adj } A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$. Computing $A(\text{adj } A)$ we see that

$$A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} -6 & 0 & 0 \\ 0 & -6 & 0 \\ 0 & 0 & -6 \end{bmatrix}.$$

Note that

$$\det A = \begin{vmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{vmatrix} = \begin{vmatrix} 0 & -36 & -25 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{vmatrix} = - \begin{vmatrix} -36 & -25 \\ 6 & 4 \end{vmatrix} = -6.$$

Therefore we have $A(\text{adj } A) = (\det A)I$.

The Adjugate Formula

Theorem

If A is an $n \times n$ matrix, then

$$A(\operatorname{adj} A) = (\det A)I = (\operatorname{adj} A)A.$$

Furthermore, if $\det A \neq 0$, then we get a formula for A^{-1} , i.e.,

The Adjugate Formula

Theorem

If A is an $n \times n$ matrix, then

$$A(\operatorname{adj} A) = (\det A)I = (\operatorname{adj} A)A.$$

Furthermore, if $\det A \neq 0$, then we get a formula for A^{-1} , i.e.,

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A.$$

The Adjugate Formula

Theorem

If A is an $n \times n$ matrix, then

$$A(\text{adj } A) = (\det A)I = (\text{adj } A)A.$$

Furthermore, if $\det A \neq 0$, then we get a formula for A^{-1} , i.e.,

$$A^{-1} = \frac{1}{\det A} \text{adj } A.$$

Inverting a matrix using the adjugate

Except in the case of a 2×2 matrix, the adjugate formula is a very inefficient method for computing the inverse of a matrix; the matrix inversion algorithm is much more practical. However, the adjugate formula is of theoretical significance.

Proof of the Adjugate Formula

Example

Recall that the (i, j) -entry of $\text{adj}(A)$ is equal to $\text{cof}(A)_{ji}$.

Proof of the Adjugate Formula

Example

Recall that the (i, j) -entry of $\text{adj}(A)$ is equal to $\text{cof}(A)_{ji}$. Let us compute the (i, j) -entry of $B = A \cdot \text{adj}(A)$:

$$b_{ij} = \sum_{k=1}^n a_{ik} \text{cof}(A)_{ki}$$

Proof of the Adjugate Formula

Example

Recall that the (i, j) -entry of $\text{adj}(A)$ is equal to $\text{cof}(A)_{ji}$. Let us compute the (i, j) -entry of $B = A \cdot \text{adj}(A)$:

$$b_{ij} = \sum_{k=1}^n a_{ik} \text{cof}(A)_{ki}$$

By the cofactor expansion theorem, b_{ij} is equal to the determinant of matrix C obtained from A by replacing its j th column by $a_{i1}, a_{i2}, \dots, a_{in}$ — i.e., its i th column.

Proof of the Adjugate Formula

Example

Recall that the (i, j) -entry of $\text{adj}(A)$ is equal to $\text{cof}(A)_{ji}$. Let us compute the (i, j) -entry of $B = A \cdot \text{adj}(A)$:

$$b_{ij} = \sum_{k=1}^n a_{ik} \text{cof}(A)_{ki}$$

By the cofactor expansion theorem, b_{ij} is equal to the determinant of matrix C obtained from A by replacing its j th column by $a_{i1}, a_{i2}, \dots, a_{in}$ — i.e., its i th column.

If $i = j$ then this matrix is A and therefore

$$a_{ii} = \det A$$

for all i .

Proof of the Adjugate Formula

Example

Recall that the (i, j) -entry of $\text{adj}(A)$ is equal to $\text{cof}(A)_{ji}$. Let us compute the (i, j) -entry of $B = A \cdot \text{adj}(A)$:

$$b_{ij} = \sum_{k=1}^n a_{ik} \text{cof}(A)_{ki}$$

By the cofactor expansion theorem, b_{ij} is equal to the determinant of matrix C obtained from A by replacing its j th column by $a_{i1}, a_{i2}, \dots, a_{in}$ — i.e., its i th column.

If $i = j$ then this matrix is A and therefore

$$a_{ii} = \det A$$

for all i . If $i \neq j$ then this matrix has its i th column equal to its j th column, and therefore

$$a_{ij} = 0 \quad \text{if } i \neq j.$$

Using the Adjugate to Find the Inverse of a Matrix

Example

$$\text{Let } A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}.$$

Using the Adjugate to Find the Inverse of a Matrix

Example

Let $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$. As we saw earlier, $\det A = -6 \neq 0$, so A is invertible, and

Using the Adjugate to Find the Inverse of a Matrix

Example

Let $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$. As we saw earlier, $\det A = -6 \neq 0$, so A is invertible, and

$$\operatorname{adj} A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

Using the Adjugate to Find the Inverse of a Matrix

Example

Let $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$. As we saw earlier, $\det A = -6 \neq 0$, so A is invertible, and

$$\operatorname{adj} A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A$$

Using the Adjugate to Find the Inverse of a Matrix

Example

Let $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$. As we saw earlier, $\det A = -6 \neq 0$, so A is invertible, and

$$\operatorname{adj} A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A = \frac{1}{-6} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}$$

Using the Adjugate to Find the Inverse of a Matrix

Example

Let $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$. As we saw earlier, $\det A = -6 \neq 0$, so A is invertible, and

$$\operatorname{adj} A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A = \frac{1}{-6} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} 1 & -3 & \frac{9}{2} \\ \frac{2}{3} & -\frac{8}{3} & \frac{25}{6} \\ -1 & 4 & -6 \end{bmatrix}.$$

Using the Adjugate to Find the Inverse of a Matrix

Example

Let $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$. As we saw earlier, $\det A = -6 \neq 0$, so A is invertible, and

$$\operatorname{adj} A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A = \frac{1}{-6} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} 1 & -3 & \frac{9}{2} \\ \frac{2}{3} & -\frac{8}{3} & \frac{25}{6} \\ -1 & 4 & -6 \end{bmatrix}.$$

You can check this by computing AA^{-1} .

Using the Adjugate to Find the Inverse of a Matrix

Example

Let $A = \begin{bmatrix} 4 & 0 & 3 \\ 1 & 9 & 7 \\ 0 & 6 & 4 \end{bmatrix}$. As we saw earlier, $\det A = -6 \neq 0$, so A is invertible, and

$$\operatorname{adj} A = \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix}.$$

$$A^{-1} = \frac{1}{\det A} \operatorname{adj} A = \frac{1}{-6} \begin{bmatrix} -6 & 18 & -27 \\ -4 & 16 & -25 \\ 6 & -24 & 36 \end{bmatrix} = \begin{bmatrix} 1 & -3 & \frac{9}{2} \\ \frac{2}{3} & -\frac{8}{3} & \frac{25}{6} \\ -1 & 4 & -6 \end{bmatrix}.$$

You can check this by computing AA^{-1} . You could also check by using the Matrix Inversion Algorithm to find A^{-1} (though this is more work).

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\operatorname{adj} A) = (\det A)^{n-1}$.

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\operatorname{adj} A) = (\det A)^{n-1}$.

Solution

Using the adjugate formula,

$$A(\operatorname{adj} A) = (\det A)I$$

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\operatorname{adj} A) = (\det A)^{n-1}$.

Solution

Using the adjugate formula,

$$\begin{aligned} A(\operatorname{adj} A) &= (\det A)I \\ \det(A(\operatorname{adj} A)) &= \det((\det A)I) \end{aligned}$$

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\operatorname{adj} A) = (\det A)^{n-1}$.

Solution

Using the adjugate formula,

$$\begin{aligned} A(\operatorname{adj} A) &= (\det A)I \\ \det(A(\operatorname{adj} A)) &= \det((\det A)I) \\ (\det A) \times \det(\operatorname{adj} A) &= (\det A)^n (\det I) \end{aligned}$$

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\operatorname{adj} A) = (\det A)^{n-1}$.

Solution

Using the adjugate formula,

$$\begin{aligned} A(\operatorname{adj} A) &= (\det A)I \\ \det(A(\operatorname{adj} A)) &= \det((\det A)I) \\ (\det A) \times \det(\operatorname{adj} A) &= (\det A)^n (\det I) \\ (\det A) \times \det(\operatorname{adj} A) &= (\det A)^n \end{aligned}$$

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\text{adj } A) = (\det A)^{n-1}$.

Solution

Using the adjugate formula,

$$\begin{aligned}A(\text{adj } A) &= (\det A)I \\ \det(A(\text{adj } A)) &= \det((\det A)I) \\ (\det A) \times \det(\text{adj } A) &= (\det A)^n (\det I) \\ (\det A) \times \det(\text{adj } A) &= (\det A)^n\end{aligned}$$

Since A is invertible, $\det A \neq 0$, so we divide both sides of the last equation by $\det A$ to obtain

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\text{adj } A) = (\det A)^{n-1}$.

Solution

Using the adjugate formula,

$$\begin{aligned}A(\text{adj } A) &= (\det A)I \\ \det(A(\text{adj } A)) &= \det((\det A)I) \\ (\det A) \times \det(\text{adj } A) &= (\det A)^n (\det I) \\ (\det A) \times \det(\text{adj } A) &= (\det A)^n\end{aligned}$$

Since A is invertible, $\det A \neq 0$, so we divide both sides of the last equation by $\det A$ to obtain

$$\det(\text{adj } A) = (\det A)^{n-1}.$$

Problem

Let A be an $n \times n$ invertible matrix. Show that $\det(\operatorname{adj} A) = (\det A)^{n-1}$.

Solution

Using the adjugate formula,

$$\begin{aligned}A(\operatorname{adj} A) &= (\det A)I \\ \det(A(\operatorname{adj} A)) &= \det((\det A)I) \\ (\det A) \times \det(\operatorname{adj} A) &= (\det A)^n (\det I) \\ (\det A) \times \det(\operatorname{adj} A) &= (\det A)^n\end{aligned}$$

Since A is invertible, $\det A \neq 0$, so we divide both sides of the last equation by $\det A$ to obtain

$$\det(\operatorname{adj} A) = (\det A)^{n-1}.$$

Even if A is not invertible, $\det(\operatorname{adj} A) = (\det A)^{n-1}$, but the proof is more complicated.

Cramer's Rule

If A is an $n \times n$ invertible matrix, then the solution to $AX = B$ can be given in terms of determinants of matrices.

Cramer's Rule

If A is an $n \times n$ **invertible** matrix, then the solution to $AX = B$ can be given in terms of determinants of matrices.

Theorem (Cramer's Rule)

Let A be an $n \times n$ invertible matrix, and consider the system $AX = B$, where $X = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T$.

Cramer's Rule

If A is an $n \times n$ invertible matrix, then the solution to $AX = B$ can be given in terms of determinants of matrices.

Theorem (Cramer's Rule)

Let A be an $n \times n$ invertible matrix, and consider the system $AX = B$, where $X = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T$. We define A_i to be the matrix obtained from A by replacing column i with B .

Cramer's Rule

If A is an $n \times n$ invertible matrix, then the solution to $AX = B$ can be given in terms of determinants of matrices.

Theorem (Cramer's Rule)

Let A be an $n \times n$ invertible matrix, and consider the system $AX = B$, where $X = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \end{bmatrix}^T$. We define A_i to be the matrix obtained from A by replacing column i with B . Then for each value of i , $1 \leq i \leq n$,

$$x_i = \frac{\det A_i}{\det A}$$

Example (Cramer's Rule)

Solve the following system of linear equations using Cramer's Rule.

$$3x_1 + x_2 - x_3 = -1$$

$$5x_1 + 2x_2 = 2$$

$$x_1 + x_2 - x_3 = 1$$

Example (Cramer's Rule)

Solve the following system of linear equations using Cramer's Rule.

$$\begin{array}{rrcrcl} 3x_1 & + & x_2 & - & x_3 & = & -1 \\ 5x_1 & + & 2x_2 & & & = & 2 \\ x_1 & + & x_2 & - & x_3 & = & 1 \end{array}$$

First, $x_1 = \frac{\det A_1}{\det A}$,

Example (Cramer's Rule)

Solve the following system of linear equations using Cramer's Rule.

$$\begin{array}{rrcrcl} 3x_1 & + & x_2 & - & x_3 & = & -1 \\ 5x_1 & + & 2x_2 & & & = & 2 \\ x_1 & + & x_2 & - & x_3 & = & 1 \end{array}$$

First, $x_1 = \frac{\det A_1}{\det A}$, where

$$A = \begin{bmatrix} 3 & 1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix}$$

Example (Cramer's Rule)

Solve the following system of linear equations using Cramer's Rule.

$$\begin{array}{rrcrcl} 3x_1 & + & x_2 & - & x_3 & = & -1 \\ 5x_1 & + & 2x_2 & & & = & 2 \\ x_1 & + & x_2 & - & x_3 & = & 1 \end{array}$$

First, $x_1 = \frac{\det A_1}{\det A}$, where

$$A = \begin{bmatrix} 3 & 1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix} \text{ and } A_1 = \begin{bmatrix} -1 & 1 & -1 \\ 2 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix}.$$

Example (Cramer's Rule)

Solve the following system of linear equations using Cramer's Rule.

$$\begin{array}{rrcrcl} 3x_1 & + & x_2 & - & x_3 & = & -1 \\ 5x_1 & + & 2x_2 & & & = & 2 \\ x_1 & + & x_2 & - & x_3 & = & 1 \end{array}$$

First, $x_1 = \frac{\det A_1}{\det A}$, where

$$A = \begin{bmatrix} 3 & 1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix} \text{ and } A_1 = \begin{bmatrix} -1 & 1 & -1 \\ 2 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix}.$$

Computing the determinants of these two matrices,

$$\det A = -4 \text{ and } \det A_1 = 4,$$

Example (Cramer's Rule)

Solve the following system of linear equations using Cramer's Rule.

$$\begin{array}{rrcrcl} 3x_1 & + & x_2 & - & x_3 & = & -1 \\ 5x_1 & + & 2x_2 & & & = & 2 \\ x_1 & + & x_2 & - & x_3 & = & 1 \end{array}$$

First, $x_1 = \frac{\det A_1}{\det A}$, where

$$A = \begin{bmatrix} 3 & 1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix} \text{ and } A_1 = \begin{bmatrix} -1 & 1 & -1 \\ 2 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix}.$$

Computing the determinants of these two matrices,

$$\det A = -4 \text{ and } \det A_1 = 4,$$

$$\text{and thus } x_1 = \frac{4}{-4} = -1.$$

Example (continued)

Secondly, $x_2 = \frac{\det A_2}{\det A}$ where $\det A = -4$ and

$$\det A_2 = \begin{vmatrix} 3 & -1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{vmatrix} = -14,$$

Example (continued)

Secondly, $x_2 = \frac{\det A_2}{\det A}$ where $\det A = -4$ and

$$\det A_2 = \begin{vmatrix} 3 & -1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{vmatrix} = -14,$$

and thus $x_2 = \frac{-14}{-4} = \frac{7}{2}$.

Example (continued)

Secondly, $x_2 = \frac{\det A_2}{\det A}$ where $\det A = -4$ and

$$\det A_2 = \begin{vmatrix} 3 & -1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{vmatrix} = -14,$$

and thus $x_2 = \frac{-14}{-4} = \frac{7}{2}$. Finally, $x_3 = \frac{\det A_3}{\det A}$, where $\det A = -4$ and

$$\det A_3 = \begin{vmatrix} 3 & 1 & -1 \\ 5 & 2 & 2 \\ 1 & 1 & 1 \end{vmatrix} = -6,$$

Example (continued)

Secondly, $x_2 = \frac{\det A_2}{\det A}$ where $\det A = -4$ and

$$\det A_2 = \begin{vmatrix} 3 & -1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{vmatrix} = -14,$$

and thus $x_2 = \frac{-14}{-4} = \frac{7}{2}$. Finally, $x_3 = \frac{\det A_3}{\det A}$, where $\det A = -4$ and

$$\det A_3 = \begin{vmatrix} 3 & 1 & -1 \\ 5 & 2 & 2 \\ 1 & 1 & 1 \end{vmatrix} = -6,$$

and thus $x_3 = \frac{-6}{-4} = \frac{3}{2}$.

Example (continued)

Secondly, $x_2 = \frac{\det A_2}{\det A}$ where $\det A = -4$ and

$$\det A_2 = \begin{vmatrix} 3 & -1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{vmatrix} = -14,$$

and thus $x_2 = \frac{-14}{-4} = \frac{7}{2}$. Finally, $x_3 = \frac{\det A_3}{\det A}$, where $\det A = -4$ and

$$\det A_3 = \begin{vmatrix} 3 & 1 & -1 \\ 5 & 2 & 2 \\ 1 & 1 & 1 \end{vmatrix} = -6,$$

and thus $x_3 = \frac{-6}{-4} = \frac{3}{2}$. Therefore, the solution to the system is given by

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -1 \\ \frac{7}{2} \\ \frac{3}{2} \end{bmatrix}.$$

Example (continued)

Secondly, $x_2 = \frac{\det A_2}{\det A}$ where $\det A = -4$ and

$$\det A_2 = \begin{vmatrix} 3 & -1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{vmatrix} = -14,$$

and thus $x_2 = \frac{-14}{-4} = \frac{7}{2}$. Finally, $x_3 = \frac{\det A_3}{\det A}$, where $\det A = -4$ and

$$\det A_3 = \begin{vmatrix} 3 & 1 & -1 \\ 5 & 2 & 2 \\ 1 & 1 & 1 \end{vmatrix} = -6,$$

and thus $x_3 = \frac{-6}{-4} = \frac{3}{2}$. Therefore, the solution to the system is given by

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -1 \\ \frac{7}{2} \\ \frac{3}{2} \end{bmatrix}.$$

You can check this by substituting these values into the original system.

Polynomial Interpolation

Problem

Given data points $(0, 1)$, $(1, 2)$, $(2, 5)$ and $(3, 10)$, find an interpolating polynomial $p(x)$ of degree at most three, and then estimate the value of y corresponding to $x = \frac{3}{2}$.

Polynomial Interpolation

Problem

Given data points $(0, 1)$, $(1, 2)$, $(2, 5)$ and $(3, 10)$, find an interpolating polynomial $p(x)$ of degree at most three, and then estimate the value of y corresponding to $x = \frac{3}{2}$.

Solution

We want to find the coefficients r_0 , r_1 , r_2 and r_3 of

$$p(x) = r_0 + r_1x + r_2x^2 + r_3x^3$$

Polynomial Interpolation

Problem

Given data points $(0, 1)$, $(1, 2)$, $(2, 5)$ and $(3, 10)$, find an interpolating polynomial $p(x)$ of degree at most three, and then estimate the value of y corresponding to $x = \frac{3}{2}$.

Solution

We want to find the coefficients r_0 , r_1 , r_2 and r_3 of

$$p(x) = r_0 + r_1x + r_2x^2 + r_3x^3$$

so that $p(0) = 1$, $p(1) = 2$, $p(2) = 5$, and $p(3) = 10$.

Polynomial Interpolation

Problem

Given data points $(0, 1)$, $(1, 2)$, $(2, 5)$ and $(3, 10)$, find an interpolating polynomial $p(x)$ of degree at most three, and then estimate the value of y corresponding to $x = \frac{3}{2}$.

Solution

We want to find the coefficients r_0 , r_1 , r_2 and r_3 of

$$p(x) = r_0 + r_1x + r_2x^2 + r_3x^3$$

so that $p(0) = 1$, $p(1) = 2$, $p(2) = 5$, and $p(3) = 10$.

$$p(0) = r_0 = 1$$

$$p(1) = r_0 + r_1 + r_2 + r_3 = 2$$

$$p(2) = r_0 + 2r_1 + 4r_2 + 8r_3 = 5$$

$$p(3) = r_0 + 3r_1 + 9r_2 + 27r_3 = 10$$

Solution (continued)

Solve this system of four equations in the four variables r_0 , r_1 , r_2 and r_3 .

Solution (continued)

Solve this system of four equations in the four variables r_0 , r_1 , r_2 and r_3 .

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 2 & 4 & 8 & 5 \\ 1 & 3 & 9 & 27 & 10 \end{array} \right]$$

Solution (continued)

Solve this system of four equations in the four variables r_0 , r_1 , r_2 and r_3 .

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 2 & 4 & 8 & 5 \\ 1 & 3 & 9 & 27 & 10 \end{array} \right] \rightarrow \dots \rightarrow \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

Solution (continued)

Solve this system of four equations in the four variables r_0 , r_1 , r_2 and r_3 .

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 2 & 4 & 8 & 5 \\ 1 & 3 & 9 & 27 & 10 \end{array} \right] \rightarrow \dots \rightarrow \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

Therefore $r_0 = 1$, $r_1 = 0$, $r_2 = 1$, $r_3 = 0$, and so

$$p(x) = 1 + x^2.$$

Solution (continued)

Solve this system of four equations in the four variables r_0 , r_1 , r_2 and r_3 .

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 2 & 4 & 8 & 5 \\ 1 & 3 & 9 & 27 & 10 \end{array} \right] \rightarrow \dots \rightarrow \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

Therefore $r_0 = 1$, $r_1 = 0$, $r_2 = 1$, $r_3 = 0$, and so

$$p(x) = 1 + x^2.$$

The estimate for the value of y corresponding to $x = \frac{3}{2}$ is

Solution (continued)

Solve this system of four equations in the four variables r_0 , r_1 , r_2 and r_3 .

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 2 & 4 & 8 & 5 \\ 1 & 3 & 9 & 27 & 10 \end{array} \right] \rightarrow \cdots \rightarrow \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

Therefore $r_0 = 1$, $r_1 = 0$, $r_2 = 1$, $r_3 = 0$, and so

$$p(x) = 1 + x^2.$$

The estimate for the value of y corresponding to $x = \frac{3}{2}$ is

$$y = p\left(\frac{3}{2}\right) = 1 + \left(\frac{3}{2}\right)^2 = \frac{13}{4},$$

Solution (continued)

Solve this system of four equations in the four variables r_0 , r_1 , r_2 and r_3 .

$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 2 & 4 & 8 & 5 \\ 1 & 3 & 9 & 27 & 10 \end{array} \right] \rightarrow \dots \rightarrow \left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

Therefore $r_0 = 1$, $r_1 = 0$, $r_2 = 1$, $r_3 = 0$, and so

$$p(x) = 1 + x^2.$$

The estimate for the value of y corresponding to $x = \frac{3}{2}$ is

$$y = p\left(\frac{3}{2}\right) = 1 + \left(\frac{3}{2}\right)^2 = \frac{13}{4},$$

resulting in the point $(\frac{3}{2}, \frac{13}{4})$.

Theorem

Given n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with the x_i **distinct**,

Theorem

Given n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with the x_i **distinct**, there is a unique polynomial $p(x) = r_0 + r_1x + r_2x^2 + \dots + r_{n-1}x^{n-1}$ such that $p(x_i) = y_i$ for $i = 1, 2, \dots, n$.

Theorem

Given n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with the x_i **distinct**, there is a unique polynomial $p(x) = r_0 + r_1x + r_2x^2 + \dots + r_{n-1}x^{n-1}$ such that $p(x_i) = y_i$ for $i = 1, 2, \dots, n$. The polynomial $p(x)$ is called the **interpolating polynomial** for the data points.

Theorem

Given n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with the x_i **distinct**, there is a unique polynomial $p(x) = r_0 + r_1x + r_2x^2 + \dots + r_{n-1}x^{n-1}$ such that $p(x_i) = y_i$ for $i = 1, 2, \dots, n$. The polynomial $p(x)$ is called the **interpolating polynomial** for the data points.

To find $p(x)$, set up a system of n linear equations in the n variables $r_0, r_1, r_2, \dots, r_{n-1}$.

Theorem

Given n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with the x_i **distinct**, there is a unique polynomial $p(x) = r_0 + r_1x + r_2x^2 + \dots + r_{n-1}x^{n-1}$ such that $p(x_i) = y_i$ for $i = 1, 2, \dots, n$. The polynomial $p(x)$ is called the **interpolating polynomial** for the data points.

To find $p(x)$, set up a system of n linear equations in the n variables

$$\begin{array}{rcl} r_0, r_1, r_2, \dots, r_{n-1}. & r_0 + r_1x_1 + r_2x_1^2 + \dots + r_{n-1}x_1^{n-1} & = y_1 \\ & r_0 + r_1x_2 + r_2x_2^2 + \dots + r_{n-1}x_2^{n-1} & = y_2 \\ & \vdots & \vdots \\ & r_0 + r_1x_n + r_2x_n^2 + \dots + r_{n-1}x_n^{n-1} & = y_n \end{array}$$

Theorem

Given n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with the x_i **distinct**, there is a unique polynomial $p(x) = r_0 + r_1x + r_2x^2 + \dots + r_{n-1}x^{n-1}$ such that $p(x_i) = y_i$ for $i = 1, 2, \dots, n$. The polynomial $p(x)$ is called the **interpolating polynomial** for the data points.

To find $p(x)$, set up a system of n linear equations in the n variables

$$\begin{array}{rcl} r_0, r_1, r_2, \dots, r_{n-1} \cdot & r_0 + r_1x_1 + r_2x_1^2 + \dots + r_{n-1}x_1^{n-1} & = y_1 \\ & r_0 + r_1x_2 + r_2x_2^2 + \dots + r_{n-1}x_2^{n-1} & = y_2 \\ & \vdots & \vdots \\ & r_0 + r_1x_n + r_2x_n^2 + \dots + r_{n-1}x_n^{n-1} & = y_n \end{array}$$

The fact that the x_i are **distinct** guarantees that the coefficient matrix

$$\begin{bmatrix} 1 & x_1 & x_1^2 & \dots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \dots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_n & x_n^2 & \dots & x_n^{n-1} \end{bmatrix}$$

has determinant **not equal to zero**,

Theorem

Given n data points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with the x_i **distinct**, there is a unique polynomial $p(x) = r_0 + r_1x + r_2x^2 + \dots + r_{n-1}x^{n-1}$ such that $p(x_i) = y_i$ for $i = 1, 2, \dots, n$. The polynomial $p(x)$ is called the **interpolating polynomial** for the data points.

To find $p(x)$, set up a system of n linear equations in the n variables

$$\begin{array}{rcl} r_0, r_1, r_2, \dots, r_{n-1} \cdot & \begin{array}{l} r_0 + r_1x_1 + r_2x_1^2 + \dots + r_{n-1}x_1^{n-1} \\ r_0 + r_1x_2 + r_2x_2^2 + \dots + r_{n-1}x_2^{n-1} \\ \vdots \\ r_0 + r_1x_n + r_2x_n^2 + \dots + r_{n-1}x_n^{n-1} \end{array} & = \begin{array}{l} y_1 \\ y_2 \\ \vdots \\ y_n \end{array} \end{array}$$

The fact that the x_i are **distinct** guarantees that the coefficient matrix

$$\begin{bmatrix} 1 & x_1 & x_1^2 & \dots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \dots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_n & x_n^2 & \dots & x_n^{n-1} \end{bmatrix}$$

has determinant **not equal to zero**, and so the system has a unique solution, i.e., there is a unique interpolating polynomial for the data points.