

Eigenvalues and eigenvectors (review)

Let A be an $n \times n$ matrix

Eigenvalues and eigenvectors (review)

Let A be an $n \times n$ matrix

- 1 Compute the characteristic polynomial of A ,

$$c_A(x) = \det(xI - A).$$

Eigenvalues and eigenvectors (review)

Let A be an $n \times n$ matrix

- 1 Compute the characteristic polynomial of A ,

$$c_A(x) = \det(xI - A).$$

- 2 Factorize $c_A(x)$ and find its roots.

Eigenvalues and eigenvectors (review)

Let A be an $n \times n$ matrix

- 1 Compute the characteristic polynomial of A ,

$$c_A(x) = \det(xI - A).$$

- 2 Factorize $c_A(x)$ and find its roots.
- 3 For each root λ of $c_A(x)$ solve the homogeneous system

$$(\lambda I - A)X = 0.$$

(It always has a nontrivial solution.)

Eigenvalues and eigenvectors (review)

Let A be an $n \times n$ matrix

- 1 Compute the characteristic polynomial of A ,

$$c_A(x) = \det(xI - A).$$

- 2 Factorize $c_A(x)$ and find its roots.
- 3 For each root λ of $c_A(x)$ solve the homogeneous system

$$(\lambda I - A)X = 0.$$

(It always has a nontrivial solution.)

- 4 λ -eigenvectors are the (nontrivial) solutions to this system.

Similar Matrices

Definition

Let A , and B be $n \times n$ matrices. Suppose there exists an invertible matrix P such that

$$A = P^{-1}BP$$

Then A and B are called **similar matrices**.

Similar Matrices

Definition

Let A , and B be $n \times n$ matrices. Suppose there exists an invertible matrix P such that

$$A = P^{-1}BP$$

Then A and B are called **similar matrices**.

How do similar matrices help us in spectral theory?

Similar Matrices

Definition

Let A , and B be $n \times n$ matrices. Suppose there exists an invertible matrix P such that

$$A = P^{-1}BP$$

Then A and B are called **similar matrices**.

How do similar matrices help us in spectral theory?

Theorem

Let A and B be similar matrices, so that $A = P^{-1}BP$ where A, B are $n \times n$ matrices and P is invertible. Then A and B have the same eigenvalues.

Proof

Assume $BX = \lambda X$.

Similar Matrices

Definition

Let A , and B be $n \times n$ matrices. Suppose there exists an invertible matrix P such that

$$A = P^{-1}BP$$

Then A and B are called **similar matrices**.

How do similar matrices help us in spectral theory?

Theorem

Let A and B be similar matrices, so that $A = P^{-1}BP$ where A, B are $n \times n$ matrices and P is invertible. Then A and B have the same eigenvalues.

Proof

Assume $BX = \lambda X$. Let $Y = P^{-1}X$.

Similar Matrices

Definition

Let A , and B be $n \times n$ matrices. Suppose there exists an invertible matrix P such that

$$A = P^{-1}BP$$

Then A and B are called **similar matrices**.

How do similar matrices help us in spectral theory?

Theorem

Let A and B be similar matrices, so that $A = P^{-1}BP$ where A, B are $n \times n$ matrices and P is invertible. Then A and B have the same eigenvalues.

Proof

Assume $BX = \lambda X$. Let $Y = P^{-1}X$. Then

$$AY = (P^{-1}BP)P^{-1}X = P^{-1}BX = P^{-1}\lambda X = \lambda Y.$$

Using Similar and Elementary Matrices

Problem

Find the eigenvalues for the matrix

$$A = \begin{bmatrix} 33 & 105 & 105 \\ 10 & 28 & 30 \\ -20 & -60 & -62 \end{bmatrix}$$

Using Similar and Elementary Matrices

Problem

Find the eigenvalues for the matrix

$$A = \begin{bmatrix} 33 & 105 & 105 \\ 10 & 28 & 30 \\ -20 & -60 & -62 \end{bmatrix}$$

Solution

We will use elementary matrices to simplify A before finding the eigenvalues. Left multiply A by $E(2 \times 2 + 3)$, and right multiply by the inverse of $E(2 \times 2 + 3)$.

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 2 & 1 \end{bmatrix} \begin{bmatrix} 33 & 105 & 105 \\ 10 & 28 & 30 \\ -20 & -60 & -62 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 33 & -105 & 105 \\ 10 & -32 & 30 \\ 0 & 0 & -2 \end{bmatrix}$$

Notice that the resulting matrix and A are similar matrices (with $E(2 \times 2 + 3)$ playing the role of P) so they have the same eigenvalues.

Solution (continued)

We do this step again, on the resulting matrix above.

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 33 & -105 & 105 \\ 10 & -32 & 30 \\ 0 & 0 & -2 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 15 \\ 10 & -2 & 30 \\ 0 & 0 & -2 \end{bmatrix} = B$$

Solution (continued)

We do this step again, on the resulting matrix above.

$$\begin{bmatrix} 1 & -3 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 33 & -105 & 105 \\ 10 & -32 & 30 \\ 0 & 0 & -2 \end{bmatrix} \begin{bmatrix} 1 & 3 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 0 & 15 \\ 10 & -2 & 30 \\ 0 & 0 & -2 \end{bmatrix} = B$$

Again by properties of similar matrices, the resulting matrix here (labeled B) has the same eigenvalues as our original matrix A . The advantage is that it is much simpler to find the eigenvalues of B than A .

Finding these eigenvalues follows the usual procedure and is left as an exercise.

Example (Triangular Matrices)

Consider the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 & 3 \\ 0 & 5 & 1 & -2 \\ 0 & 0 & 0 & 7 \\ 0 & 0 & 0 & -4 \end{bmatrix}.$$

Example (Triangular Matrices)

Consider the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 & 3 \\ 0 & 5 & 1 & -2 \\ 0 & 0 & 0 & 7 \\ 0 & 0 & 0 & -4 \end{bmatrix}.$$

The characteristic polynomial of A is

$$c_A(x) = \det(xI - A)$$

Example (Triangular Matrices)

Consider the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 & 3 \\ 0 & 5 & 1 & -2 \\ 0 & 0 & 0 & 7 \\ 0 & 0 & 0 & -4 \end{bmatrix}.$$

The characteristic polynomial of A is

$$c_A(x) = \det(xI - A) = \det \begin{bmatrix} x-2 & 1 & 0 & -3 \\ 0 & x-5 & -1 & 2 \\ 0 & 0 & x & -7 \\ 0 & 0 & 0 & x+4 \end{bmatrix}$$

Example (Triangular Matrices)

Consider the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 & 3 \\ 0 & 5 & 1 & -2 \\ 0 & 0 & 0 & 7 \\ 0 & 0 & 0 & -4 \end{bmatrix}.$$

The characteristic polynomial of A is

$$c_A(x) = \det(xI - A) = \det \begin{bmatrix} x-2 & 1 & 0 & -3 \\ 0 & x-5 & -1 & 2 \\ 0 & 0 & x & -7 \\ 0 & 0 & 0 & x+4 \end{bmatrix} = (x-2)(x-5)x(x+4).$$

Example (Triangular Matrices)

Consider the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 & 3 \\ 0 & 5 & 1 & -2 \\ 0 & 0 & 0 & 7 \\ 0 & 0 & 0 & -4 \end{bmatrix}.$$

The characteristic polynomial of A is

$$c_A(x) = \det(xI - A) = \det \begin{bmatrix} x-2 & 1 & 0 & -3 \\ 0 & x-5 & -1 & 2 \\ 0 & 0 & x & -7 \\ 0 & 0 & 0 & x+4 \end{bmatrix} = (x-2)(x-5)x(x+4).$$

Therefore the eigenvalues of A are 2, 5, 0 and -4 , **exactly the entries on the main diagonal of A .**

Example (Triangular Matrices)

Consider the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 & 3 \\ 0 & 5 & 1 & -2 \\ 0 & 0 & 0 & 7 \\ 0 & 0 & 0 & -4 \end{bmatrix}.$$

The characteristic polynomial of A is

$$c_A(x) = \det(xI - A) = \det \begin{bmatrix} x-2 & 1 & 0 & -3 \\ 0 & x-5 & -1 & 2 \\ 0 & 0 & x & -7 \\ 0 & 0 & 0 & x+4 \end{bmatrix} = (x-2)(x-5)x(x+4).$$

Therefore the eigenvalues of A are 2, 5, 0 and -4 , **exactly the entries on the main diagonal of A .**

Eigenvalues of Triangular Matrices

If A is an $n \times n$ upper triangular (or lower triangular) matrix, then the eigenvalues of A are the entries on the main diagonal of A .

Geometric Interpretation of Eigenvalues and Eigenvectors


Example

Recall that in \mathbb{R}^2 , **reflection in the x-axis** is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$.

Geometric Interpretation of Eigenvalues and Eigenvectors

Example


Recall that in \mathbb{R}^2 , **reflection in the x-axis** is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$.



Geometric Interpretation of Eigenvalues and Eigenvectors

Example

Recall that in \mathbb{R}^2 , **reflection in the x-axis** is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$.




Let A be the matrix that induces reflection in the x-axis.

Geometric Interpretation of Eigenvalues and Eigenvectors

Example

Recall that in \mathbb{R}^2 , **reflection in the x-axis** is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$.




Let A be the matrix that induces reflection in the x-axis. If λ were an eigenvalue of A and X a corresponding eigenvector, then $AX = \lambda X$ implies that, geometrically, **reflecting X in the x-axis** is the same as changing X to a vector parallel to X .

Geometric Interpretation of Eigenvalues and Eigenvectors

Example

Recall that in \mathbb{R}^2 , **reflection in the x-axis** is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$.




Let A be the matrix that induces reflection in the x -axis. If λ were an eigenvalue of A and X a corresponding eigenvector, then $AX = \lambda X$ implies that, geometrically, **reflecting X in the x -axis** is the same as changing X to a vector parallel to X .

How could this be possible?

Geometric Interpretation of Eigenvalues and Eigenvectors

Example

Recall that in \mathbb{R}^2 , **reflection in the x-axis** is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$.




Let A be the matrix that induces reflection in the x-axis. If λ were an eigenvalue of A and X a corresponding eigenvector, then $AX = \lambda X$ implies that, geometrically, **reflecting X in the x-axis** is the same as changing X to a vector parallel to X .


How could this be possible?

Can you picture what an eigenvector of A would look like?

Example (continued)




Example (continued)




- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the x -axis is $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$.

Example (continued)




- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the x-axis is $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.

Example (continued)




- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the x-axis is $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.
- The reflection of $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the x-axis is $\begin{bmatrix} 0 \\ -1 \end{bmatrix}$,

Example (continued)



- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the x-axis is $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.
- The reflection of $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the x-axis is $\begin{bmatrix} 0 \\ -1 \end{bmatrix}$, so $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.

Example (continued)



- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the x-axis is $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.
- The reflection of $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the x-axis is $\begin{bmatrix} 0 \\ -1 \end{bmatrix}$, so $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.

This makes sense, since we know that reflection in the x-axis is induced by the matrix

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix},$$


which has eigenvalues 1 and -1 .

Example

In \mathbb{R}^2 , reflection in the y -axis is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -a \\ b \end{bmatrix}$.


Example

In \mathbb{R}^2 , reflection in the y -axis is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -a \\ b \end{bmatrix}$.



Example


In \mathbb{R}^2 , reflection in the y -axis is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -a \\ b \end{bmatrix}$.



Let A be the matrix that induces reflection in the y -axis.


Example

In \mathbb{R}^2 , reflection in the y -axis is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -a \\ b \end{bmatrix}$.




Let A be the matrix that induces reflection in the y -axis. If λ were an eigenvalue of A and X a corresponding eigenvector, then $AX = \lambda X$ implies that, geometrically, **reflecting X in the y -axis** is the same as changing X to a vector parallel to X .

Example (continued)




Example (continued)




- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 2 \\ 0 \end{bmatrix}$,

Example (continued)




- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.

Example (continued)




- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.
- The reflection of $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$,

Example (continued)



- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.
- The reflection of $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$, so $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.

Example (continued)



- The reflection of $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 2 \\ 0 \end{bmatrix}$, so $\begin{bmatrix} -2 \\ 0 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.
- The reflection of $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the y-axis is $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$, so $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.


This makes sense, since we know that reflection in the y-axis is induced by the matrix

$$A = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix},$$

which has eigenvalues 1 and -1 .


Example

Reflection in the line $y = x$ is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} b \\ a \end{bmatrix}$.



Example


Reflection in the line $y = x$ is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} b \\ a \end{bmatrix}$.



Let A be the matrix that induces reflection in the line $y = x$.


Example

Reflection in the line $y = x$ is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} b \\ a \end{bmatrix}$.




Let A be the matrix that induces reflection in the line $y = x$. If λ were an eigenvalue of A and X a corresponding eigenvector, then $AX = \lambda X$ implies that, geometrically, **reflecting X in the y -axis** is the same as changing X to a vector parallel to X .

Example (continued)




Example (continued)




- The reflection of $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$,

Example (continued)




- The reflection of $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$, so $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.

Example (continued)




- The reflection of $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$, so $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.
- The reflection of $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$,

Example (continued)



- The reflection of $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$, so $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.
- The reflection of $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$, so $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.

Example (continued)



- The reflection of $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$, so $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = 1$.
- The reflection of $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ in the line $y = x$ is $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$, so $\begin{bmatrix} -1 \\ 1 \end{bmatrix}$ is an eigenvector of A that corresponds to the eigenvalue $\lambda = -1$.

Therefore, 1 and -1 are eigenvalues of A .

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation.

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation. Let X be a vector in \mathbb{R}^2 with tail at the origin.

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation. Let X be a vector in \mathbb{R}^2 with tail at the origin.

If A is the matrix that induces reflection in the line $y = mx$, then

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation. Let X be a vector in \mathbb{R}^2 with tail at the origin.

If A is the matrix that induces reflection in the line $y = mx$, then

- the reflection of a vector X that is parallel to $y = mx$ is simply X ;

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation. Let X be a vector in \mathbb{R}^2 with tail at the origin.

If A is the matrix that induces reflection in the line $y = mx$, then

- the reflection of a vector X that is **parallel** to $y = mx$ is simply X ;
- the reflection of a vector X that is **perpendicular** to $y = mx$ is $-X$.

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation. Let X be a vector in \mathbb{R}^2 with tail at the origin.

If A is the matrix that induces reflection in the line $y = mx$, then

- the reflection of a vector X that is **parallel** to $y = mx$ is simply X ;
- the reflection of a vector X that is **perpendicular** to $y = mx$ is $-X$.

Therefore, 1 and -1 are eigenvalues of A ;

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation. Let X be a vector in \mathbb{R}^2 with tail at the origin.

If A is the matrix that induces reflection in the line $y = mx$, then

- the reflection of a vector X that is **parallel** to $y = mx$ is simply X ;
- the reflection of a vector X that is **perpendicular** to $y = mx$ is $-X$.

Therefore, 1 and -1 are eigenvalues of A ; in fact, these are the only two eigenvalues of A and each has multiplicity one.

\mathbb{R}^2 : Reflections in lines through the origin

In \mathbb{R}^2 , if $y = mx$ is a line through the origin, then reflection in the line $y = mx$ is a linear transformation. Let X be a vector in \mathbb{R}^2 with tail at the origin.

If A is the matrix that induces reflection in the line $y = mx$, then

- the reflection of a vector X that is **parallel** to $y = mx$ is simply X ;
- the reflection of a vector X that is **perpendicular** to $y = mx$ is $-X$.


Therefore, 1 and -1 are eigenvalues of A ; in fact, these are the only two eigenvalues of A and each has multiplicity one. This follows from the fact that A is a 2×2 matrix, so its characteristic polynomial has degree two.

Example (Rotation through π)

We denote by $R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of π .


Example (Rotation through π)

We denote by $R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of π .



Example (Rotation through π)


We denote by $R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of π .



R_π is a linear transformation that transforms X to $-X$.

Example (Rotation through π)

We denote by $R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of π .




R_π is a linear transformation that transforms X to $-X$.

Let A denote the matrix that induces rotation through π .

Example (Rotation through π)

We denote by $R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of π .




R_π is a linear transformation that transforms X to $-X$.

Let A denote the matrix that induces rotation through π . Then $AX = -X$ for every nonzero vector X ,

Example (Rotation through π)

We denote by $R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of π .



R_π is a linear transformation that transforms X to $-X$.


Let A denote the matrix that induces rotation through π . Then $AX = -X$ for every nonzero vector X , meaning that **every nonzero vector of \mathbb{R}^2 is an eigenvector of A corresponding to the eigenvalue $\lambda = -1$.**

Example (Rotation through $\pi/2$)

We denote by $R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of $\pi/2$.


Example (Rotation through $\pi/2$)

We denote by $R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of $\pi/2$.



Example (Rotation through $\pi/2$)


We denote by $R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of $\pi/2$.



R_{π} is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -b \\ a \end{bmatrix}$.

Example (Rotation through $\pi/2$)

We denote by $R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of $\pi/2$.




R_{π} is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -b \\ a \end{bmatrix}$.

Notice that there is no nonzero vector X that can be rotated through an angle of $\pi/2$ to produce a vector parallel to X .

Example (Rotation through $\pi/2$)

We denote by $R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ counterclockwise rotation about the origin through an angle of $\pi/2$.



R_{π} is a linear transformation that transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -b \\ a \end{bmatrix}$.

Notice that there is no nonzero vector X that can be rotated through an angle of $\pi/2$ to produce a vector parallel to X . Therefore, A has no **real** eigenvalues.

Example (continued)

Let A denote the matrix that induces rotation through $\pi/2$.

Example (continued)

Let A denote the matrix that induces rotation through $\pi/2$. Then

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

Example (continued)

Let A denote the matrix that induces rotation through $\pi/2$. Then

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

and $c_A(x) = x^2 + 1$.

Example (continued)

Let A denote the matrix that induces rotation through $\pi/2$. Then

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

and $c_A(x) = x^2 + 1$. Therefore, A has complex eigenvalues i and $-i$.

A First Course in
LINEAR ALGEBRA

Lecture Notes
for Math 1503

Spectral Theory: 7.2 Diagonalization

Creative Commons License (CC BY-NC-SA)

A First Course in Linear Algebra

Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Diagonal Matrices

Definition

An $n \times n$ matrix A is **diagonal** if it is both upper triangular and lower triangular.

Diagonal Matrices

Definition

An $n \times n$ matrix A is **diagonal** if it is both upper triangular and lower triangular. Equivalently, all entries except those on the main diagonal are zeros.

Diagonal Matrices

Definition

An $n \times n$ matrix A is **diagonal** if it is both upper triangular and lower triangular. Equivalently, all entries except those on the main diagonal are zeros.

Notation

An $n \times n$ diagonal matrix

$$D = \begin{bmatrix} a_1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & a_2 & 0 & \cdots & 0 & 0 \\ 0 & 0 & a_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & a_{n-1} & 0 \\ 0 & 0 & 0 & \cdots & 0 & a_n \end{bmatrix}$$

is written $D = \text{diag}(a_1, a_2, a_3, \dots, a_{n-1}, a_n)$.

Diagonalizability

Definition

An $n \times n$ matrix A is said to be **diagonalizable** if there exists an invertible $n \times n$ matrix P such that $A = PDP^{-1}$.

Equivalently: A is **diagonalizable** if $A \sim D$ for some **diagonal matrix** D .

Diagonalizability

Definition

An $n \times n$ matrix A is said to be **diagonalizable** if there exists an invertible $n \times n$ matrix P such that $A = PDP^{-1}$.

Equivalently: A is **diagonalizable** if $A \sim D$ for some **diagonal matrix** D .

Diagonalizing a Matrix

Let A be an $n \times n$ matrix. The process of finding an **invertible** matrix P and a **diagonal** matrix D so that $A = PDP^{-1}$ is referred to as **diagonalizing** the matrix A , and P is called the **diagonalizing** matrix for A .

Diagonalizability

Definition

An $n \times n$ matrix A is said to be **diagonalizable** if there exists an invertible $n \times n$ matrix P such that $A = PDP^{-1}$.

Equivalently: A is **diagonalizable** if $A \sim D$ for some **diagonal matrix** D .

Diagonalizing a Matrix

Let A be an $n \times n$ matrix. The process of finding an **invertible** matrix P and a **diagonal** matrix D so that $A = PDP^{-1}$ is referred to as **diagonalizing** the matrix A , and P is called the **diagonalizing** matrix for A .

The key to diagonalizing a matrix (finding the matrices P and D) lies in the eigenvectors and eigenvalues of the matrix A .

Reminder

Let A be an $n \times n$ matrix and λ a real number. If λ is an eigenvalue of A , then

$$AX = \lambda X$$

for some **nonzero** vector X in \mathbb{R}^n . Such a vector X is called a λ -eigenvector of A or an eigenvector of A corresponding to λ .

Reminder

Let A be an $n \times n$ matrix and λ a real number. If λ is an eigenvalue of A , then

$$AX = \lambda X$$

for some **nonzero** vector X in \mathbb{R}^n . Such a vector X is called a λ -eigenvector of A or an eigenvector of A corresponding to λ .

Theorem

Let A be an $n \times n$ matrix.

Reminder

Let A be an $n \times n$ matrix and λ a real number. If λ is an eigenvalue of A , then

$$AX = \lambda X$$

for some **nonzero** vector X in \mathbb{R}^n . Such a vector X is called a λ -eigenvector of A or an eigenvector of A corresponding to λ .

Theorem

Let A be an $n \times n$ matrix.

- 1 A is diagonalizable if and only if it has eigenvectors X_1, X_2, \dots, X_n so that $P = \begin{bmatrix} X_1 & X_2 & \cdots & X_n \end{bmatrix}$ is invertible.

Reminder

Let A be an $n \times n$ matrix and λ a real number. If λ is an eigenvalue of A , then

$$AX = \lambda X$$

for some **nonzero** vector X in \mathbb{R}^n . Such a vector X is called a **λ -eigenvector of A** or an eigenvector of A corresponding to λ .

Theorem

Let A be an $n \times n$ matrix.

- 1 A is diagonalizable if and only if it has eigenvectors X_1, X_2, \dots, X_n so that $P = \begin{bmatrix} X_1 & X_2 & \cdots & X_n \end{bmatrix}$ is invertible.
- 2 If P is invertible, then

$$P^{-1}AP = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$$

where λ_i is the eigenvalue of A corresponding to the eigenvector X_i , i.e., $AX_i = \lambda_i X_i$.

Example

Diagonalize, if possible, the matrix $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & -3 \end{bmatrix}$.

Example

Diagonalize, if possible, the matrix $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & -3 \end{bmatrix}$.

Solution

$$c_A(x) = \det(xI - A) = \begin{vmatrix} x-1 & 0 & -1 \\ 0 & x-1 & 0 \\ 0 & 0 & x+3 \end{vmatrix} = (x-1)^2(x+3).$$

Example

Diagonalize, if possible, the matrix $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & -3 \end{bmatrix}$.

Solution

$$c_A(x) = \det(xI - A) = \begin{vmatrix} x-1 & 0 & -1 \\ 0 & x-1 & 0 \\ 0 & 0 & x+3 \end{vmatrix} = (x-1)^2(x+3).$$

Therefore, A has eigenvalues $\lambda_1 = 1$ of multiplicity two and $\lambda_2 = -3$ of multiplicity one.

Example

Diagonalize, if possible, the matrix $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & -3 \end{bmatrix}$.

Solution

$$c_A(x) = \det(xI - A) = \begin{vmatrix} x-1 & 0 & -1 \\ 0 & x-1 & 0 \\ 0 & 0 & x+3 \end{vmatrix} = (x-1)^2(x+3).$$

Therefore, A has eigenvalues $\lambda_1 = 1$ of multiplicity two and $\lambda_2 = -3$ of multiplicity one.

Eigenvectors for $\lambda_1 = 1$: solve $(I - A)X = 0$.

$$\left[\begin{array}{ccc|c} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Example

Diagonalize, if possible, the matrix $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & -3 \end{bmatrix}$.

Solution

$$c_A(x) = \det(xI - A) = \begin{vmatrix} x-1 & 0 & -1 \\ 0 & x-1 & 0 \\ 0 & 0 & x+3 \end{vmatrix} = (x-1)^2(x+3).$$

Therefore, A has eigenvalues $\lambda_1 = 1$ of multiplicity two and $\lambda_2 = -3$ of multiplicity one.

Eigenvectors for $\lambda_1 = 1$: solve $(I - A)X = 0$.

$$\left[\begin{array}{ccc|c} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 4 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \text{ so } X = \begin{bmatrix} s \\ t \\ 0 \end{bmatrix}, s, t \in \mathbb{R}.$$

Solution (continued)

Therefore, basic eigenvectors corresponding to $\lambda_1 = 1$ are $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$.

Eigenvectors for $\lambda_2 = -3$: solve $(-3I - A)X = 0$.

$$\left[\begin{array}{ccc|c} -4 & 0 & -1 & 0 \\ 0 & -4 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & \frac{1}{4} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Solution (continued)

Therefore, basic eigenvectors corresponding to $\lambda_1 = 1$ are $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$.

Eigenvectors for $\lambda_2 = -3$: solve $(-3I - A)X = 0$.

$$\left[\begin{array}{ccc|c} -4 & 0 & -1 & 0 \\ 0 & -4 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & \frac{1}{4} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \text{ so } X = \begin{bmatrix} -\frac{1}{4}t \\ 0 \\ t \end{bmatrix}, t \in \mathbb{R}.$$

Solution (continued)

Therefore, basic eigenvectors corresponding to $\lambda_1 = 1$ are $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$.

Eigenvectors for $\lambda_2 = -3$: solve $(-3I - A)X = 0$.

$$\left[\begin{array}{ccc|c} -4 & 0 & -1 & 0 \\ 0 & -4 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & \frac{1}{4} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \text{ so } X = \begin{bmatrix} -\frac{1}{4}t \\ 0 \\ t \end{bmatrix}, t \in \mathbb{R}.$$

Therefore, a basic eigenvector corresponding to $\lambda_2 = -3$ is $\begin{bmatrix} -1 \\ 0 \\ 4 \end{bmatrix}$

Solution (continued)

Therefore, basic eigenvectors corresponding to $\lambda_1 = 1$ are $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$.

Eigenvectors for $\lambda_2 = -3$: solve $(-3I - A)X = 0$.

$$\left[\begin{array}{ccc|c} -4 & 0 & -1 & 0 \\ 0 & -4 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \rightarrow \left[\begin{array}{ccc|c} 1 & 0 & \frac{1}{4} & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right] \text{ so } X = \begin{bmatrix} -\frac{1}{4}t \\ 0 \\ t \end{bmatrix}, t \in \mathbb{R}.$$

Therefore, a basic eigenvector corresponding to $\lambda_2 = -3$ is $\begin{bmatrix} -1 \\ 0 \\ 4 \end{bmatrix}$

Let

$$P = \begin{bmatrix} -1 & 1 & 0 \\ 0 & 0 & 1 \\ 4 & 0 & 0 \end{bmatrix}.$$

Then P is invertible (easily checked by computing $\det P$).

Solution (continued)

Solution (continued)

Furthermore,

$$P^{-1}AP = D = \text{diag}(-3, 1, 1) = \begin{bmatrix} -3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Solution (continued)

Furthermore,

$$P^{-1}AP = D = \text{diag}(-3, 1, 1) = \begin{bmatrix} -3 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

The eigenvalues of A in D (from left to right) occur in the same order as their corresponding eigenvectors as columns of P .

Example

Diagonalize the matrix $A = \begin{bmatrix} 3 & -4 & 2 \\ 1 & -2 & 2 \\ 1 & -5 & 5 \end{bmatrix}$

Example

Diagonalize the matrix $A = \begin{bmatrix} 3 & -4 & 2 \\ 1 & -2 & 2 \\ 1 & -5 & 5 \end{bmatrix}$

Solution

You can check that A has eigenvalues and corresponding basic eigenvectors:

$$\lambda_1 = 3 \text{ and } X_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}; \lambda_2 = 2 \text{ and } X_2 = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}; \lambda_3 = 1 \text{ and } X_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

Example

Diagonalize the matrix $A = \begin{bmatrix} 3 & -4 & 2 \\ 1 & -2 & 2 \\ 1 & -5 & 5 \end{bmatrix}$

Solution

You can check that A has eigenvalues and corresponding basic eigenvectors:

$$\lambda_1 = 3 \text{ and } X_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}; \lambda_2 = 2 \text{ and } X_2 = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}; \lambda_3 = 1 \text{ and } X_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

$$\text{Let } P = [X_1 \quad X_2 \quad X_3] = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 & 1 \\ 2 & 1 & 1 \end{bmatrix}.$$

Example

Diagonalize the matrix $A = \begin{bmatrix} 3 & -4 & 2 \\ 1 & -2 & 2 \\ 1 & -5 & 5 \end{bmatrix}$

Solution

You can check that A has eigenvalues and corresponding basic eigenvectors:

$$\lambda_1 = 3 \text{ and } X_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}; \lambda_2 = 2 \text{ and } X_2 = \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix}; \lambda_3 = 1 \text{ and } X_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

Let $P = [X_1 \ X_2 \ X_3] = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 1 & 1 \\ 2 & 1 & 1 \end{bmatrix}$. Then P is invertible (check this!), so by the previous theorem,

$$P^{-1}AP = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Eigenvalues, Eigenvectors, and Diagonalization

Theorem

Let A be an $n \times n$ matrix, and suppose that A has distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_m$. For each i , let X_i be a λ_i -eigenvector of A . Then $\{X_1, X_2, \dots, X_m\}$ is linearly independent.

Eigenvalues, Eigenvectors, and Diagonalization

Theorem

Let A be an $n \times n$ matrix, and suppose that A has distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_m$. For each i , let X_i be a λ_i -eigenvector of A . Then $\{X_1, X_2, \dots, X_m\}$ is linearly independent.

Diagonalizability

Determining whether or not a square matrix A is diagonalizable can be done using **eigenvalues** and **eigenvectors** of the matrix A .

Theorem

Let A be an $n \times n$ matrix and suppose it has n distinct eigenvalues. Then it follows that A is diagonalizable.

Theorem

Let A be an $n \times n$ matrix and suppose it has n distinct eigenvalues. Then it follows that A is diagonalizable.

Proof.

Let $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$ denote the n (distinct) eigenvalues of A , and let X_i be an eigenvector of A corresponding to λ_i , $1 \leq i \leq n$. Then $\{X_1, X_2, \dots, X_n\}$ is a linearly independent set in \mathbb{R}^n (i.e. a basis).

Theorem

Let A be an $n \times n$ matrix and suppose it has n distinct eigenvalues. Then it follows that A is diagonalizable.

Proof.

Let $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$ denote the n (distinct) eigenvalues of A , and let X_i be an eigenvector of A corresponding to λ_i , $1 \leq i \leq n$. Then $\{X_1, X_2, \dots, X_n\}$ is a linearly independent set in \mathbb{R}^n (i.e. a basis).

It follows that $P = [X_1 \ X_2 \ \cdots \ X_n]$ is invertible, and therefore A is diagonalizable. □

Example

Show that the matrix

$$A = \begin{bmatrix} 0 & -1 & 1 \\ 8 & 6 & -2 \\ 0 & 0 & -3 \end{bmatrix}$$

is diagonalizable.

Example

Show that the matrix

$$A = \begin{bmatrix} 0 & -1 & 1 \\ 8 & 6 & -2 \\ 0 & 0 & -3 \end{bmatrix}$$

is diagonalizable.

Solution

A has characteristic polynomial

$$c_A(x) = (x + 3)(x - 2)(x - 4),$$

and thus A has distinct eigenvalues $-3, 2$ and 4 .

Since A is 3×3 and has three distinct eigenvalues, A is diagonalizable.

Definition

Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$. The eigenspace of A corresponding to λ , written $E_\lambda(A)$ is the subspace spanned by the set of all eigenvectors corresponding to λ .

Definition

Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$. The eigenspace of A corresponding to λ , written $E_\lambda(A)$ is the subspace spanned by the set of all eigenvectors corresponding to λ .

In other words, the eigenspace $E_\lambda(A)$ is all X (including $X = \vec{0}$) such that $AX = \lambda X$. In other words, $E_\lambda(A) = \text{null}(\lambda I - A)$.

Definition (recall)

Let A be an $n \times n$ matrix with characteristic polynomial given by $\det(\lambda I - A)$. Then, the multiplicity of an eigenvalue λ of A is the number of times λ occurs as a root of that characteristic polynomial.

Definition (recall)

Let A be an $n \times n$ matrix with characteristic polynomial given by $\det(\lambda I - A)$. Then, the multiplicity of an eigenvalue λ of A is the number of times λ occurs as a root of that characteristic polynomial.

Lemma

If A is an $n \times n$ matrix, then

$$\dim(E_\lambda(A)) \leq m$$

where λ is an eigenvalue of A of multiplicity m .

Lecture 2

Definition (recall)

Let A be an $n \times n$ matrix with characteristic polynomial given by $\det(\lambda I - A)$. Then, the multiplicity of an eigenvalue λ of A is the number of times λ occurs as a root of that characteristic polynomial.

Lemma

If A is an $n \times n$ matrix, then

$$\dim(E_\lambda(A)) \leq m$$

where λ is an eigenvalue of A of multiplicity m .

This result tells us that if λ is an eigenvalue of A , then the number of linearly independent λ -eigenvectors is never more than the multiplicity of λ .

The crucial consequence of the above Lemma is the characterization of matrices that are diagonalizable.

The crucial consequence of the above Lemma is the characterization of matrices that are diagonalizable.

Theorem

Let A be an $n \times n$ matrix A . Then A is diagonalizable if and only if for each eigenvalue λ of A , $\dim(E_\lambda(A))$ is equal to the multiplicity of λ .

Example

Let

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

Determine if A is diagonalizable

Example

Let

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

Determine if A is diagonalizable

Solution

Through the usual procedure, we find that the eigenvalues of A are $\lambda_1 = 1, \lambda_2 = 1$. Solving as usual, we find that the eigenvectors are given by

$$t \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ and the basic eigenvector is } X_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Example

Let

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

Determine if A is diagonalizable

Solution

Through the usual procedure, we find that the eigenvalues of A are $\lambda_1 = 1, \lambda_2 = 1$. Solving as usual, we find that the eigenvectors are given by

$$t \begin{bmatrix} 1 \\ 0 \end{bmatrix} \text{ and the basic eigenvector is } X_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

This means that $\dim(E_\lambda(A)) = 1$, but the multiplicity of $\lambda = 1$ is 2. Therefore this matrix is not diagonalizable.

Complex Eigenvalues

If a matrix has eigenvalues that have imaginary parts (and aren't simply real numbers), we can still find eigenvectors and possibly diagonalize the matrix.

Complex Eigenvalues

If a matrix has eigenvalues that have imaginary parts (and aren't simply real numbers), we can still find eigenvectors and possibly diagonalize the matrix.

Problem

Diagonalize, if possible, the matrix $A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$.

Complex Eigenvalues

If a matrix has eigenvalues that have imaginary parts (and aren't simply real numbers), we can still find eigenvectors and possibly diagonalize the matrix.

Problem

Diagonalize, if possible, the matrix $A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$.

Solution

$$c_A(x) = \det(xI - A) = \begin{vmatrix} x-1 & -1 \\ 1 & x-1 \end{vmatrix} = x^2 - 2x + 2.$$

The roots of $c_A(x)$ are **distinct complex numbers**: $\lambda_1 = 1 + i$ and $\lambda_2 = 1 - i$, so A is diagonalizable. Corresponding eigenvectors are

$$X_1 = \begin{bmatrix} -i \\ 1 \end{bmatrix} \text{ and } X_2 = \begin{bmatrix} i \\ 1 \end{bmatrix},$$

respectively.

Solution (continued)

A diagonalizing matrix for A is

$$P = \begin{bmatrix} -i & i \\ 1 & 1 \end{bmatrix},$$

and

$$P^{-1}AP = \begin{bmatrix} 1+i & 0 \\ 0 & 1-i \end{bmatrix}.$$

Solution (continued)

A diagonalizing matrix for A is

$$P = \begin{bmatrix} -i & i \\ 1 & 1 \end{bmatrix},$$

and

$$P^{-1}AP = \begin{bmatrix} 1+i & 0 \\ 0 & 1-i \end{bmatrix}.$$

Notice that A is a real matrix, but has complex eigenvalues (and eigenvectors).

Eigenvalues of a Real Symmetric Matrix

Theorem

The eigenvalues of any **real symmetric** matrix are **real**.

Eigenvalues of a Real Symmetric Matrix

Theorem

The eigenvalues of any **real symmetric** matrix are **real**.

Proof.

Let A be an $n \times n$ real symmetric matrix, and let λ be an eigenvalue of A . To prove that λ is real, it is enough to prove that $\bar{\lambda} = \lambda$, i.e., λ is equal to its (complex) conjugate.

We use \bar{A} to denote the matrix obtained from A by replacing each entry by its conjugate. Since A is real, $\bar{A} = A$.

Suppose

$$X = \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_n \end{bmatrix}$$

is a λ -eigenvector of A . Then $AX = \lambda X$.

Proof (continued).

$$\text{Let } y = X^T \overline{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \overline{z}_1 \\ \overline{z}_2 \\ \vdots \\ \overline{z}_n \end{bmatrix}.$$

Proof (continued).

$$\text{Let } y = X^T \overline{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \overline{z}_1 \\ \overline{z}_2 \\ \vdots \\ \overline{z}_n \end{bmatrix}.$$

$$\text{Then } y = z_1 \overline{z}_1 + z_2 \overline{z}_2 + \cdots + z_n \overline{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2;$$

Proof (continued).

$$\text{Let } y = X^T \overline{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \overline{z}_1 \\ \overline{z}_2 \\ \vdots \\ \overline{z}_n \end{bmatrix}.$$

Then $y = z_1 \overline{z}_1 + z_2 \overline{z}_2 + \cdots + z_n \overline{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\lambda y = \lambda(X^T \overline{X}) = (\lambda X^T) \overline{X} = (\lambda X)^T \overline{X}$$

Proof (continued).

$$\text{Let } y = X^T \overline{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \overline{z}_1 \\ \overline{z}_2 \\ \vdots \\ \overline{z}_n \end{bmatrix}.$$

Then $y = z_1 \overline{z}_1 + z_2 \overline{z}_2 + \cdots + z_n \overline{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\begin{aligned} \lambda y &= \lambda(X^T \overline{X}) = (\lambda X^T) \overline{X} = (\lambda X)^T \overline{X} \\ &= (AX)^T \overline{X} = X^T A^T \overline{X} \end{aligned}$$

Proof (continued).

$$\text{Let } y = X^T \bar{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \bar{z}_1 \\ \bar{z}_2 \\ \vdots \\ \bar{z}_n \end{bmatrix}.$$

Then $y = z_1 \bar{z}_1 + z_2 \bar{z}_2 + \cdots + z_n \bar{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\begin{aligned} \lambda y &= \lambda(X^T \bar{X}) = (\lambda X^T) \bar{X} = (\lambda X)^T \bar{X} \\ &= (AX)^T \bar{X} = X^T A^T \bar{X} \\ &= X^T A \bar{X} \quad (\text{since } A \text{ is symmetric}) \end{aligned}$$

Proof (continued).

$$\text{Let } y = X^T \overline{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \overline{z}_1 \\ \overline{z}_2 \\ \vdots \\ \overline{z}_n \end{bmatrix}.$$

Then $y = z_1 \overline{z}_1 + z_2 \overline{z}_2 + \cdots + z_n \overline{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\begin{aligned} \lambda y &= \lambda(X^T \overline{X}) = (\lambda X^T) \overline{X} = (\lambda X)^T \overline{X} \\ &= (AX)^T \overline{X} = X^T A^T \overline{X} \\ &= X^T A \overline{X} \quad (\text{since } A \text{ is symmetric}) \\ &= X^T \overline{AX} \quad (\text{since } A \text{ is real}) \end{aligned}$$

Proof (continued).

$$\text{Let } y = X^T \bar{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \bar{z}_1 \\ \bar{z}_2 \\ \vdots \\ \bar{z}_n \end{bmatrix}.$$

Then $y = z_1 \bar{z}_1 + z_2 \bar{z}_2 + \cdots + z_n \bar{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\begin{aligned} \lambda y &= \lambda(X^T \bar{X}) = (\lambda X^T) \bar{X} = (\lambda X)^T \bar{X} \\ &= (AX)^T \bar{X} = X^T A^T \bar{X} \\ &= X^T A \bar{X} \quad (\text{since } A \text{ is symmetric}) \\ &= X^T \overline{AX} \quad (\text{since } A \text{ is real}) \\ &= X^T (\overline{AX}) = X^T (\overline{\lambda X}) = X^T \overline{\lambda} \bar{X} \end{aligned}$$

Proof (continued).

$$\text{Let } y = X^T \bar{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \bar{z}_1 \\ \bar{z}_2 \\ \vdots \\ \bar{z}_n \end{bmatrix}.$$

Then $y = z_1 \bar{z}_1 + z_2 \bar{z}_2 + \cdots + z_n \bar{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\begin{aligned} \lambda y &= \lambda(X^T \bar{X}) = (\lambda X^T) \bar{X} = (\lambda X)^T \bar{X} \\ &= (AX)^T \bar{X} = X^T A^T \bar{X} \\ &= X^T A \bar{X} \quad (\text{since } A \text{ is symmetric}) \\ &= X^T \overline{AX} \quad (\text{since } A \text{ is real}) \\ &= X^T (\overline{\lambda X}) = X^T (\bar{\lambda} \bar{X}) = X^T \bar{\lambda} \bar{X} \\ &= \bar{\lambda} (X^T \bar{X}) \end{aligned}$$

Proof (continued).

$$\text{Let } y = X^T \bar{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \bar{z}_1 \\ \bar{z}_2 \\ \vdots \\ \bar{z}_n \end{bmatrix}.$$

Then $y = z_1 \bar{z}_1 + z_2 \bar{z}_2 + \cdots + z_n \bar{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\begin{aligned} \lambda y &= \lambda(X^T \bar{X}) = (\lambda X^T) \bar{X} = (\lambda X)^T \bar{X} \\ &= (AX)^T \bar{X} = X^T A^T \bar{X} \\ &= X^T A \bar{X} \quad (\text{since } A \text{ is symmetric}) \\ &= X^T \overline{AX} \quad (\text{since } A \text{ is real}) \\ &= X^T (\overline{AX}) = X^T (\overline{\lambda X}) = X^T \overline{\lambda} \bar{X} \\ &= \overline{\lambda} (X^T \bar{X}) \\ &= \overline{\lambda} y. \end{aligned}$$

Proof (continued).

$$\text{Let } y = X^T \bar{X} = \begin{bmatrix} z_1 & z_2 & \cdots & z_n \end{bmatrix} \begin{bmatrix} \bar{z}_1 \\ \bar{z}_2 \\ \vdots \\ \bar{z}_n \end{bmatrix}.$$

Then $y = z_1 \bar{z}_1 + z_2 \bar{z}_2 + \cdots + z_n \bar{z}_n = |z_1|^2 + |z_2|^2 + \cdots + |z_n|^2$;
since $X \neq 0$, y is a positive real number. So

$$\begin{aligned} \lambda y &= \lambda(X^T \bar{X}) = (\lambda X^T) \bar{X} = (\lambda X)^T \bar{X} \\ &= (AX)^T \bar{X} = X^T A^T \bar{X} \\ &= X^T A \bar{X} \quad (\text{since } A \text{ is symmetric}) \\ &= X^T \overline{AX} \quad (\text{since } A \text{ is real}) \\ &= X^T (\overline{\lambda X}) = X^T (\bar{\lambda} \bar{X}) = X^T \bar{\lambda} \bar{X} \\ &= \bar{\lambda} (X^T \bar{X}) \\ &= \bar{\lambda} y. \end{aligned}$$

Thus, $\lambda y = \bar{\lambda} y$. Since $y \neq 0$, it follows that $\lambda = \bar{\lambda}$, and therefore λ is real. □