

A First Course in
LINEAR ALGEBRA

Lecture Notes
by Karen Seyffarth

\mathbb{R}^n : Subspaces and Basis

Creative Commons License (CC BY-NC-SA)



Champions of Access to Knowledge



OPEN TEXT



ONLINE
ASSESSMENT



SUPPORT



INSTRUCTOR
SUPPLEMENTS

Contact Lyryx Today!

info@lyryx.com

A First Course in Linear Algebra

Lecture Notes

Current Lecture Notes Revision: Version 2017 — Revision A

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Definition

The **vector space** \mathbb{R}^n consists of the set \mathbb{R}^n written as **column matrices**, along with the (matrix) operations of addition and scalar multiplication. Unless stated otherwise, \mathbb{R}^n means the vector space \mathbb{R}^n .

A rigorous definition of an abstract vector space will be given after we have studied properties of the vector space \mathbb{R}^n .

We are not yet ready to formally define the term “subspace”. However, in the context of \mathbb{R}^n , we rely on the **Subspace Test** to determine whether or not a subset of \mathbb{R}^n is a subspace.

Theorem (Subspace Test)

A subset V of \mathbb{R}^n is a subspace of \mathbb{R}^n if

- 1 the zero vector of \mathbb{R}^n , $\vec{0}_n$, is in V ;
- 2 V is closed under addition, i.e., for all $\vec{u}, \vec{w} \in V$, $\vec{u} + \vec{w} \in V$;
- 3 V is closed under scalar multiplication, i.e., for all $\vec{u} \in V$ and $k \in \mathbb{R}$, $k\vec{u} \in V$.

The subset $V = \{\vec{0}_n\}$ is a subspace of \mathbb{R}^n (**verify this**), as is the set \mathbb{R}^n itself. Any other subspace of \mathbb{R}^n is a **proper** subspace of \mathbb{R}^n .

Notation

If V is a subset of \mathbb{R}^n , we write $V \subseteq \mathbb{R}^n$.

Example

In \mathbb{R}^3 , the line L through the origin that is parallel to the vector $\vec{d} = \begin{bmatrix} -5 \\ 1 \\ -4 \end{bmatrix}$ has

(vector) equation $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = t \begin{bmatrix} -5 \\ 1 \\ -4 \end{bmatrix}$, $t \in \mathbb{R}$, so

$$L = \left\{ t\vec{d} \mid t \in \mathbb{R} \right\}.$$

Claim. L is a subspace of \mathbb{R}^3 .

- First: $\vec{0}_3 \in L$ since $0\vec{d} = \vec{0}_3$.
- Suppose $\vec{u}, \vec{v} \in L$. Then by definition, $\vec{u} = s\vec{d}$ and $\vec{v} = t\vec{d}$, for some $s, t \in \mathbb{R}$. Thus

$$\vec{u} + \vec{v} = s\vec{d} + t\vec{d} = (s + t)\vec{d}.$$

Since $s + t \in \mathbb{R}$, $\vec{u} + \vec{v} \in L$; i.e., L is closed under addition.

Example 5 (continued)

- Suppose $\vec{u} \in L$ and $k \in \mathbb{R}$ (k is a scalar). Then $\vec{u} = t\vec{d}$, for some $t \in \mathbb{R}$, so

$$k\vec{u} = k(t\vec{d}) = (kt)\vec{d}.$$

Since $kt \in \mathbb{R}$, $k\vec{u} \in L$; i.e., L is closed under scalar multiplication.

Therefore, L is a subspace of \mathbb{R}^3 .

Note that there is nothing special about the vector \vec{d} used in this example; the same proof works for any **nonzero** vector $\vec{d} \in \mathbb{R}^3$, so any line through the origin is a subspace of \mathbb{R}^3 .

Example

In \mathbb{R}^3 , let M denote the plane through the origin having equation

$3x - 2y + z = 0$; then M has normal vector $\vec{n} = \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$. If $\vec{u} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, then

$$M = \{ \vec{u} \in \mathbb{R}^3 \mid \vec{n} \bullet \vec{u} = 0 \},$$

where $\vec{n} \bullet \vec{u}$ is the dot product of vectors \vec{n} and \vec{u} .

Claim. M is a subspace of \mathbb{R}^3 .

- First: $\vec{0}_3 \in M$ since $\vec{n} \bullet \vec{0}_3 = 0$.
- Suppose $\vec{u}, \vec{v} \in M$. Then by definition, $\vec{n} \bullet \vec{u} = 0$ and $\vec{n} \bullet \vec{v} = 0$, so

$$\vec{n} \bullet (\vec{u} + \vec{v}) = \vec{n} \bullet \vec{u} + \vec{n} \bullet \vec{v} = 0 + 0 = 0,$$

and thus $(\vec{u} + \vec{v}) \in M$; i.e., M is closed under addition.

Example 6 (continued)

- Suppose $\vec{u} \in M$ and $k \in \mathbb{R}$. Then $\vec{n} \bullet \vec{u} = 0$, so

$$\vec{n} \bullet (k\vec{u}) = k(\vec{n} \bullet \vec{u}) = k(0) = 0,$$

and thus $k\vec{u} \in M$; i.e., M is closed under scalar multiplication.

Therefore, M is a subspace of \mathbb{R}^3 .

As in the previous example, there is nothing special about the plane chosen for this example; any plane through the origin is a subspace of \mathbb{R}^3 .

Problem

Is $V = \left\{ \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \mid a, b, c, d \in \mathbb{R} \text{ and } 2a - b = c + 2d \right\}$ a subspace of \mathbb{R}^4 ?

Justify your answer.

Solution 1

The zero vector of \mathbb{R}^4 is the vector $\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$ with $a = b = c = d = 0$.

In this case, $2a - b = 2(0) + 0 = 0$ and $c + 2d = 0 + 2(0) = 0$, so $2a - b = c + 2d$. Therefore, $\vec{0}_4 \in V$.

Solution (continued)

Suppose

$$\vec{v}_1 = \begin{bmatrix} a_1 \\ b_1 \\ c_1 \\ d_1 \end{bmatrix} \text{ and } \vec{v}_2 = \begin{bmatrix} a_2 \\ b_2 \\ c_2 \\ d_2 \end{bmatrix} \text{ are in } V.$$

Then $2a_1 - b_1 = c_1 + 2d_1$ and $2a_2 - b_2 = c_2 + 2d_2$. Now

$$\vec{v}_1 + \vec{v}_2 = \begin{bmatrix} a_1 \\ b_1 \\ c_1 \\ d_1 \end{bmatrix} + \begin{bmatrix} a_2 \\ b_2 \\ c_2 \\ d_2 \end{bmatrix} = \begin{bmatrix} a_1 + a_2 \\ b_1 + b_2 \\ c_1 + c_2 \\ d_1 + d_2 \end{bmatrix},$$

$$\begin{aligned} \text{and} \quad 2(a_1 + a_2) - (b_1 + b_2) &= (2a_1 - b_1) + (2a_2 - b_2) \\ &= (c_1 + 2d_1) + (c_2 + 2d_2) \\ &= (c_1 + c_2) + 2(d_1 + d_2). \end{aligned}$$

Therefore, $\vec{v}_1 + \vec{v}_2 \in V$.

Solution (continued)

Finally, suppose

$$\vec{v} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in V \text{ and } k \in \mathbb{R}.$$

Then $2a - b = c + 2d$. Now

$$k\vec{v} = k \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} ka \\ kb \\ kc \\ kd \end{bmatrix},$$

and

$$2ka - kb = k(2a - b) = k(c + 2d) = kc + 2kd.$$

Therefore, $k\vec{v} \in V$.

It follows from the **Subspace Test** that V is a subspace of \mathbb{R}^4 .

Definition

Let A be an $n \times n$ matrix and $\lambda \in \mathbb{R}$. The **eigenspace of A corresponding to λ** is the set

$$E_\lambda(A) = \{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \lambda\vec{x}\}.$$

Note that

$$\begin{aligned} E_\lambda(A) &= \{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \lambda\vec{x}\}, \\ &= \{\vec{x} \in \mathbb{R}^n \mid \lambda\vec{x} - A\vec{x} = \vec{0}_n\} \\ &= \{\vec{x} \in \mathbb{R}^n \mid (\lambda I - A)\vec{x} = \vec{0}_n\} \end{aligned}$$

showing that

$$E_\lambda(A) = \text{null}(\lambda I - A).$$

It follows that

- if λ is **not** an eigenvalue of A , then $E_\lambda(A) = \{\vec{0}_n\}$;
- the nonzero vectors of $E_\lambda(A)$ are the eigenvectors of A corresponding to λ ;
- the eigenspace of A corresponding to λ is a subspace of \mathbb{R}^n .

Theorem

Let V be a nonempty collection of vectors in \mathbb{R}^n . Then V is a subspace of \mathbb{R}^n if and only if there exist vectors $\{\vec{u}_1, \dots, \vec{u}_k\}$ in V such that

$$V = \text{span} \{ \vec{u}_1, \dots, \vec{u}_k \}$$

Furthermore, let W be another subspace of \mathbb{R}^n and suppose $\{\vec{u}_1, \dots, \vec{u}_k\} \in W$. Then it follows that V is a subset of W .

Proof.

We first show that if V is a subspace, then it can be written as $V = \text{span} \{ \vec{u}_1, \dots, \vec{u}_k \}$.

Pick a vector \vec{u}_1 in V . If $V = \text{span} \{ \vec{u}_1 \}$, then you have found your list of vectors and are done.

Proof (continued).

If $V \neq \text{span}\{\vec{u}_1\}$, then there exists \vec{u}_2 a vector of V which is not in $\text{span}\{\vec{u}_1\}$. Consider $\text{span}\{\vec{u}_1, \vec{u}_2\}$. If $V = \text{span}\{\vec{u}_1, \vec{u}_2\}$, we are done. Otherwise, pick \vec{u}_3 not in $\text{span}\{\vec{u}_1, \vec{u}_2\}$. Continue this way. Note that since V is a subspace, these spans are each contained in V . The process must stop with \vec{u}_k for some $k \leq n$, and thus $V = \text{span}\{\vec{u}_1, \dots, \vec{u}_k\}$.

Now suppose $V = \text{span}\{\vec{u}_1, \dots, \vec{u}_k\}$, we must show this is a subspace. So let $\sum_{i=1}^k c_i \vec{u}_i$ and $\sum_{i=1}^k d_i \vec{u}_i$ be two vectors in V , and let a and b be two scalars. Then

$$a \sum_{i=1}^k c_i \vec{u}_i + b \sum_{i=1}^k d_i \vec{u}_i = \sum_{i=1}^k (ac_i + bd_i) \vec{u}_i$$

which is one of the vectors in $\text{span}\{\vec{u}_1, \dots, \vec{u}_k\}$ and is therefore contained in V . This shows that $\text{span}\{\vec{u}_1, \dots, \vec{u}_k\}$ has the properties of a subspace. □

Proof (continued).

To prove that $V \subseteq W$, we prove that if $\vec{u}_i \in V$, then $\vec{u}_i \in W$.

Suppose $\vec{u} \in V$. Then $\vec{u} = a_1\vec{u}_1 + a_2\vec{u}_2 + \cdots + a_k\vec{u}_k$ for some $a_i \in \mathbb{R}$, $1 \leq i \leq k$.

Since W contain each \vec{u}_i and W is a vector space, it follows that

$a_1\vec{u}_1 + a_2\vec{u}_2 + \cdots + a_k\vec{u}_k \in W$. □

Problem

$$\text{Is } V = \left\{ \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \mid a, b, c, d \in \mathbb{R} \text{ and } 2a - b = c + 2d \right\} \text{ a subspace of } \mathbb{R}^4?$$

Justify your answer.

Solution 2

$$\text{Let } \vec{v} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in V. \text{ Since } 2a - b = c + 2d, c = 2a - b - 2d, \text{ and thus}$$

$$V = \left\{ \begin{bmatrix} a \\ b \\ 2a - b - 2d \\ d \end{bmatrix} \mid a, b, d \in \mathbb{R} \right\} = \text{span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ -2 \\ 1 \end{bmatrix} \right\}.$$

Therefore V is a subspace of \mathbb{R}^4 .

Bases and Dimension

Theorem (Exchange Theorem)

Suppose $\{\vec{u}_1, \dots, \vec{u}_r\}$ is a linearly independent set of vectors in \mathbb{R}^n , and each \vec{u}_k is contained in $\text{span}\{\vec{v}_1, \dots, \vec{v}_s\}$. Then $s \geq r$.

In words, spanning sets have at least as many vectors as linearly independent sets.

Definition

Let V be a subspace of \mathbb{R}^n . Then $\{\vec{u}_1, \dots, \vec{u}_k\}$ is a **basis** for V if the following two conditions hold.

- 1 $\text{span}\{\vec{u}_1, \dots, \vec{u}_k\} = V$
- 2 $\{\vec{u}_1, \dots, \vec{u}_k\}$ is linearly independent

Example

The subset $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ is a basis of \mathbb{R}^n , called the **standard basis** of \mathbb{R}^n . (We've already seen that $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ is linearly independent and that $\mathbb{R}^n = \text{span}\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$.)

Example

In a previous problem, we saw that $\mathbb{R}^4 = \text{span}(S)$ where

$$S = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}.$$

S is also linearly independent (**prove this**). Therefore, S is a basis of \mathbb{R}^4 .

Theorem (Invariance Theorem)

Let V be a subspace of \mathbb{R}^n with two bases B_1 and B_2 . Suppose B_1 contains s vectors and B_2 contains r vectors. Then $s = r$.

Proof.

This follows right away from the Exchange Theorem. Indeed observe that $B_1 = \{\vec{u}_1, \dots, \vec{u}_s\}$ is a spanning set for V while $B_2 = \{\vec{v}_1, \dots, \vec{v}_r\}$ is linearly independent, so $s \geq r$. Similarly $B_2 = \{\vec{v}_1, \dots, \vec{v}_r\}$ is a spanning set for V while $B_1 = \{\vec{u}_1, \dots, \vec{u}_s\}$ is linearly independent, so $r \geq s$. □

Definition

Let V be a subspace of \mathbb{R}^n . Then the **dimension** of V , written $\dim(V)$ is defined to be the number of vectors in a basis.

Problem

In \mathbb{R}^n , what is the dimension of the subspace $\{\vec{0}_n\}$?

Solution

The only basis of the zero subspace is the empty set, \emptyset : (i) the empty set is (trivially) independent, and (ii) any linear combination of no vectors is the zero vector. Therefore, the zero subspace has dimension zero.

Example

Since $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ is a basis of \mathbb{R}^n , \mathbb{R}^n has dimension n .

This is why the Cartesian plane, \mathbb{R}^2 , is called 2-dimensional, and \mathbb{R}^3 is called 3-dimensional.

Problem

Let

$$U = \left\{ \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in \mathbb{R}^4 \mid a - b = d - c \right\}.$$

Show that U is a subspace of \mathbb{R}^4 , find a basis of U , and find $\dim(U)$.

Solution

The condition $a - b = d - c$ is equivalent to the condition $a = b - c + d$, so we may write

$$U = \left\{ \begin{bmatrix} b - c + d \\ b \\ c \\ d \end{bmatrix} \in \mathbb{R}^4 \right\} = \left\{ b \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} + c \begin{bmatrix} -1 \\ 0 \\ 1 \\ 0 \end{bmatrix} + d \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} \mid b, c, d \in \mathbb{R} \right\}$$

This shows that U is a subspace of \mathbb{R}^4 , since $U = \text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$ where

$$\begin{aligned} \vec{x}_1 &= \begin{bmatrix} 1 & 1 & 0 & 0 \end{bmatrix}^T \\ \vec{x}_2 &= \begin{bmatrix} -1 & 0 & 1 & 0 \end{bmatrix}^T \\ \vec{x}_3 &= \begin{bmatrix} 1 & 0 & 0 & 1 \end{bmatrix}^T. \end{aligned}$$

Solution (continued)

Furthermore,

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

is linearly independent, as can be seen by taking the reduced row-echelon form of the matrix whose columns are \vec{x}_1, \vec{x}_2 and \vec{x}_3 .

$$\begin{bmatrix} 1 & -1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Since every column of the reduced row-echelon form matrix has a leading one, the columns are linearly independent.

Therefore $\{\vec{x}_1, \vec{x}_2, \vec{x}_3\}$ is linearly independent and spans U , so is a basis of U , and hence U has dimension three.

Example

Suppose that $B_1 = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is a basis of \mathbb{R}^n and that A is an $n \times n$ invertible matrix. Let $B_2 = \{A\vec{v}_1, A\vec{v}_2, \dots, A\vec{v}_n\}$, and let

$$V = \begin{bmatrix} \vec{v}_1 & \vec{v}_2 & \cdots & \vec{v}_n \end{bmatrix}.$$

Since B is a basis of \mathbb{R}^n , B is independent (also a spanning set of \mathbb{R}^n); thus V is invertible. Now, because A and V are invertible, so is

$$AV = \begin{bmatrix} A\vec{v}_1 & A\vec{v}_2 & \cdots & A\vec{v}_n \end{bmatrix}.$$

Therefore, the columns of AV are independent and span \mathbb{R}^n . Since the columns of AV are the vectors of B_2 , B_2 is a basis of \mathbb{R}^n .

Properties of Bases

Theorem

Let V be a subspace of \mathbb{R}^n . Then there exists a basis of V with $\dim(V) \leq n$.

Example

Previously , we showed that

$$V = \left\{ \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in \mathbb{R}^4 \mid a - b = d - c \right\}$$

is a subspace of \mathbb{R}^4 , and that $\dim(V) = 3$. Also, it is easy to verify that

$$S = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 3 \\ 2 \end{bmatrix} \right\},$$

is an independent subset of V .

S can be extended to a basis of V . To do so, find a vector in V that is **not** in $\text{span}(S)$.

(continued)

$$\begin{bmatrix} 1 & 2 & ? \\ 1 & 3 & ? \\ 1 & 3 & ? \\ 1 & 2 & ? \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 0 \\ 1 & 3 & -1 \\ 1 & 2 & 0 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Therefore, S can be extended to the basis

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 3 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \\ 0 \end{bmatrix} \right\},$$

of V .

Problem

Let W be the subspace

$$\text{span} \left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 8 \\ 19 \\ -8 \\ 8 \end{bmatrix}, \begin{bmatrix} -6 \\ -15 \\ 6 \\ -6 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 5 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Find a basis for W which consists of a subset of the given vectors.

Final Answer

A basis for W is

$$\left\{ \begin{bmatrix} 1 \\ 2 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \\ 0 \\ 1 \end{bmatrix} \right\}$$

Theorem

The following properties hold in \mathbb{R}^n :

- Suppose $\{\vec{u}_1, \dots, \vec{u}_n\}$ is linearly independent. Then $\{\vec{u}_1, \dots, \vec{u}_n\}$ is a basis for \mathbb{R}^n .
- Suppose $\{\vec{u}_1, \dots, \vec{u}_m\}$ spans \mathbb{R}^n . Then $m \geq n$.
- If $\{\vec{u}_1, \dots, \vec{u}_n\}$ spans \mathbb{R}^n , then $\{\vec{u}_1, \dots, \vec{u}_n\}$ is linearly independent.

Question

What is the significance of this result?

Answer

Let V be a subspace of \mathbb{R}^n and suppose $B \subseteq V$.

- If B spans V and $|B| = \dim(V)$, then B is also independent, and hence B is a basis of V .
- If B is independent and $|B| = \dim(V)$, then B also spans V , and hence B is a basis of V .

Therefore if $|B| = \dim(V)$, it is sufficient to prove that B is either independent or spans V in order to prove it is a basis.

Theorem

Let V and W be subspaces of \mathbb{R}^n , and suppose that $W \subseteq V$. Then $\dim(W) \leq \dim(V)$ with equality when $W = V$.

Theorem

Let W be any non-zero subspace \mathbb{R}^n and let $W \subseteq V$ where V is also a subspace of \mathbb{R}^n . Then every basis of W can be extended to a basis for V .