

A First Course in  
**LINEAR ALGEBRA**

**Lecture Notes**  
for Math 1503

**Linear Transformations and Matrices**

Creative Commons License (CC BY-NC-SA)

# A First Course in Linear Algebra

## Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

### License



**Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)**

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

# Transformation by Matrix Multiplication

## Example

Consider the matrix  $A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \end{bmatrix}$ . By matrix multiplication,  $A$  transforms vectors in  $\mathbb{R}^3$  into vectors in  $\mathbb{R}^2$ .

# Transformation by Matrix Multiplication

## Example

Consider the matrix  $A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \end{bmatrix}$ . By matrix multiplication,  $A$  transforms vectors in  $\mathbb{R}^3$  into vectors in  $\mathbb{R}^2$ .

Consider the vector  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ . Transforming this vector by  $A$  looks like:

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x + 2y \\ 2x + y \end{bmatrix}$$

# Transformation by Matrix Multiplication

## Example

Consider the matrix  $A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \end{bmatrix}$ . By matrix multiplication,  $A$  transforms vectors in  $\mathbb{R}^3$  into vectors in  $\mathbb{R}^2$ .

Consider the vector  $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ . Transforming this vector by  $A$  looks like:

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x + 2y \\ 2x + y \end{bmatrix}$$

For example:

$$\begin{bmatrix} 1 & 2 & 0 \\ 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 5 \\ 4 \end{bmatrix}$$

# Transformations

## Definition

A **transformation** is a function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , sometimes written

$$\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m,$$

and is called a **transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$** .

# Transformations

## Definition

A **transformation** is a function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , sometimes written

$$\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m,$$

and is called a **transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$** . If  $m = n$ , then we say  **$T$  is a transformation of  $\mathbb{R}^n$** .

# Transformations

## Definition

A **transformation** is a function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , sometimes written

$$\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m,$$

and is called a **transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$** . If  $m = n$ , then we say  **$T$  is a transformation of  $\mathbb{R}^n$** .

What do we mean by a function?



# Transformations

## Definition

A **transformation** is a function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , sometimes written

$$\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m,$$

and is called a **transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$** . If  $m = n$ , then we say  **$T$  is a transformation of  $\mathbb{R}^n$** .

## What do we mean by a function?

Informally, a function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a rule that assigns exactly one vector of  $\mathbb{R}^m$  to each vector of  $\mathbb{R}^n$ .

We use the notation  $T(\vec{x})$  to mean the transformation  $T$  applied to the vector  $\vec{x}$ .

# Transformations

## Definition

A **transformation** is a function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , sometimes written

$$\mathbb{R}^n \xrightarrow{T} \mathbb{R}^m,$$

and is called a **transformation from  $\mathbb{R}^n$  to  $\mathbb{R}^m$** . If  $m = n$ , then we say  **$T$  is a transformation of  $\mathbb{R}^n$** .

## What do we mean by a function?

Informally, a function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a rule that assigns exactly one vector of  $\mathbb{R}^m$  to each vector of  $\mathbb{R}^n$ .

We use the notation  $T(\vec{x})$  to mean the transformation  $T$  applied to the vector  $\vec{x}$ .

## Definition

If  $T$  acts by matrix multiplication of a matrix  $A$  (such as the previous example), we call  $T$  a **matrix transformation**, and write  $T_A(\vec{x}) = A\vec{x}$ .

# Equality of Transformations

## Definition

Suppose  $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  are transformations. Then  $S = T$  if and only if  $S(\vec{x}) = T(\vec{x})$  for every  $\vec{x} \in \mathbb{R}^n$ .

# Specifying the Action of a Transformation

## Example

$T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  defined by

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

is a transformation

# Specifying the Action of a Transformation

## Example

$T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  defined by

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

is a transformation that **transforms** the vector  $\begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix}$  in  $\mathbb{R}^3$  into the vector

$$T \begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix} = \begin{bmatrix} 1 + 4 \\ 4 + 7 \\ 1 - 7 \\ 7 - 4 \end{bmatrix} = \begin{bmatrix} 5 \\ 11 \\ -6 \\ 3 \end{bmatrix}.$$

# Linear Transformations

## Definition

# Linear Transformations

## Definition

A transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a **linear transformation** if it satisfies the following two properties for all  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and all (scalars)  $a \in \mathbb{R}$ .

# Linear Transformations

## Definition

A transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a **linear transformation** if it satisfies the following two properties for all  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and all (scalars)  $a \in \mathbb{R}$ .

①  $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$  (preservation of addition)



# Linear Transformations

## Definition

A transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a **linear transformation** if it satisfies the following two properties for all  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and all (scalars)  $a \in \mathbb{R}$ .

- 1  $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$  (preservation of addition)
- 2  $T(a\vec{x}) = aT(\vec{x})$  (preservation of scalar multiplication)

## Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, and let  $\vec{x} \in \mathbb{R}^n$ .

## Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, and let  $\vec{x} \in \mathbb{R}^n$ . Since  $T$  preserves scalar multiplication,

## Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, and let  $\vec{x} \in \mathbb{R}^n$ . Since  $T$  preserves scalar multiplication,

- ①  $T(0\vec{x}) = 0T(\vec{x})$  implying  $T(0) = 0$ , so  $T$  preserves the zero vector.

## Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, and let  $\vec{x} \in \mathbb{R}^n$ . Since  $T$  preserves scalar multiplication,

- 1  $T(0\vec{x}) = 0T(\vec{x})$  implying  $T(0) = 0$ , so  $T$  preserves the zero vector.
- 2  $T((-1)\vec{x}) = (-1)T(\vec{x})$ , implying  $T(-\vec{x}) = -T(\vec{x})$ , so  $T$  preserves the negative of a vector.

## Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, and let  $\vec{x} \in \mathbb{R}^n$ . Since  $T$  preserves scalar multiplication,

- ①  $T(0\vec{x}) = 0T(\vec{x})$  implying  $T(0) = 0$ , so  $T$  preserves the zero vector.
- ②  $T((-1)\vec{x}) = (-1)T(\vec{x})$ , implying  $T(-\vec{x}) = -T(\vec{x})$ , so  $T$  preserves the negative of a vector.

Suppose  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$  are vectors in  $\mathbb{R}^n$  and

$$\vec{y} = a_1\vec{x}_1 + a_2\vec{x}_2 + \cdots + a_k\vec{x}_k$$

for some  $a_1, a_2, \dots, a_k \in \mathbb{R}$ .

## Properties of Linear Transformations

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation, and let  $\vec{x} \in \mathbb{R}^n$ . Since  $T$  preserves scalar multiplication,

- ①  $T(0\vec{x}) = 0T(\vec{x})$  implying  $T(0) = 0$ , so  $T$  preserves the zero vector.
- ②  $T((-1)\vec{x}) = (-1)T(\vec{x})$ , implying  $T(-\vec{x}) = -T(\vec{x})$ , so  $T$  preserves the negative of a vector.

Suppose  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$  are vectors in  $\mathbb{R}^n$  and

$$\vec{y} = a_1\vec{x}_1 + a_2\vec{x}_2 + \cdots + a_k\vec{x}_k$$

for some  $a_1, a_2, \dots, a_k \in \mathbb{R}$ . Then

③

$$\begin{aligned} T(\vec{y}) &= T(a_1\vec{x}_1 + a_2\vec{x}_2 + \cdots + a_k\vec{x}_k) \\ &= a_1T(\vec{x}_1) + a_2T(\vec{x}_2) + \cdots + a_kT(\vec{x}_k), \end{aligned}$$

i.e.,  $T$  preserves linear combinations.

## Problem

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}.$$



## Problem

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \text{ and } T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}. \text{ Find } T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix}.$$

## Problem

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \text{ and } T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}. \text{ Find } T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix}.$$

## Solution

## Problem

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \text{ and } T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}. \text{ Find } T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix}.$$

## Solution

The only way it is possible to solve this problem is if

$$\begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} \text{ is a linear combination of } \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \text{ and } \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix},$$

## Problem

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \text{ and } T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}. \text{ Find } T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix}.$$

## Solution

The only way it is possible to solve this problem is if

$$\begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} \text{ is a linear combination of } \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \text{ and } \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix},$$

i.e., if there exist  $a, b \in \mathbb{R}$  so that

$$\begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} = a \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} + b \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix}.$$

## Solution (continued)

To find  $a$  and  $b$ , solve the system of three equations in two variables:

$$\left[ \begin{array}{cc|c} 1 & 4 & -7 \\ 3 & 0 & 3 \\ 1 & 5 & -9 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{cc|c} 1 & 0 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{array} \right]$$

Thus  $a = 1$ ,  $b = -2$ , and

$$\begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix}.$$

## Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} = T \left( \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right)$$

## Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$\begin{aligned} T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} &= T \left( \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right) \\ &= T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \end{aligned}$$

## Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$\begin{aligned} T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} &= T \left( \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right) \\ &= T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \\ &= \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix} = \begin{bmatrix} -4 \\ -6 \\ 2 \\ -12 \end{bmatrix} \end{aligned}$$



## Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$\begin{aligned} T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} &= T \left( \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right) \\ &= T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \\ &= \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix} = \begin{bmatrix} -4 \\ -6 \\ 2 \\ -12 \end{bmatrix} \end{aligned}$$

Therefore,  $T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} = \begin{bmatrix} -4 \\ -6 \\ 2 \\ -12 \end{bmatrix}.$

## Problem

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 1 \\ 0 \\ -2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}. \text{ Find } T \begin{bmatrix} 2 \\ 5 \\ -3 \\ -7 \end{bmatrix}.$$

## Problem

Let  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 1 \\ 0 \\ -2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}. \text{ Find } T \begin{bmatrix} 2 \\ 5 \\ -3 \\ -7 \end{bmatrix}.$$

## Final Answer

$$T \begin{bmatrix} 2 \\ 5 \\ -3 \\ -7 \end{bmatrix} = \begin{bmatrix} -11 \\ 6 \\ -5 \end{bmatrix}.$$

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ ,

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ .

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ . Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and let  $a \in \mathbb{R}$ .

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ . Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and let  $a \in \mathbb{R}$ . Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$



# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ . Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and let  $a \in \mathbb{R}$ . Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that  $T$  preserves addition.

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ . Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and let  $a \in \mathbb{R}$ . Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that  $T$  preserves addition. Also,

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ . Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and let  $a \in \mathbb{R}$ . Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that  $T$  preserves addition. Also,

$$T(a\vec{x}) = A(a\vec{x}) = a(A\vec{x}) = aT(\vec{x}),$$

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ . Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and let  $a \in \mathbb{R}$ . Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that  $T$  preserves addition. Also,

$$T(a\vec{x}) = A(a\vec{x}) = a(A\vec{x}) = aT(\vec{x}),$$

proving that  $T$  preserves scalar multiplication.

# Matrix Transformations

## Theorem

Every matrix transformation is a linear transformation.

## Proof.

Suppose  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a matrix transformation induced by the  $m \times n$  matrix  $A$ , i.e.,  $T(\vec{x}) = A\vec{x}$  for each  $\vec{x} \in \mathbb{R}^n$ . Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and let  $a \in \mathbb{R}$ . Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that  $T$  preserves addition. Also,

$$T(a\vec{x}) = A(a\vec{x}) = a(A\vec{x}) = aT(\vec{x}),$$

proving that  $T$  preserves scalar multiplication.

Since  $T$  preserves addition and scalar multiplication  $T$  is a linear transformation. □

# Some Special Matrix Transformations

## Example (The Zero Transformation)

If  $A$  is the  $m \times n$  matrix of zeros, then the transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  induced by  $A$  is called the **zero transformation** because for every vector  $\vec{x}$  in  $\mathbb{R}^n$

$$T(\vec{x}) = A\vec{x} = 0\vec{x} = \vec{0}.$$

# Some Special Matrix Transformations

## Example (The Zero Transformation)

If  $A$  is the  $m \times n$  matrix of zeros, then the transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  induced by  $A$  is called the **zero transformation** because for every vector  $\vec{x}$  in  $\mathbb{R}^n$

$$T(\vec{x}) = A\vec{x} = 0\vec{x} = \vec{0}.$$

Note that the first zero is the matrix  $A$ , while the second zero is the zero vector of  $\mathbb{R}^m$ .

# Some Special Matrix Transformations

## Example (The Zero Transformation)

If  $A$  is the  $m \times n$  matrix of zeros, then the transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  induced by  $A$  is called the **zero transformation** because for every vector  $\vec{x}$  in  $\mathbb{R}^n$

$$T(\vec{x}) = A\vec{x} = 0\vec{x} = \vec{0}.$$

Note that the first zero is the matrix  $A$ , while the second zero is the zero vector of  $\mathbb{R}^m$ . The zero transformation is usually written as  $T = 0$ .



# Some Special Matrix Transformations

## Example (The Zero Transformation)

If  $A$  is the  $m \times n$  matrix of zeros, then the transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  induced by  $A$  is called the **zero transformation** because for every vector  $\vec{x}$  in  $\mathbb{R}^n$

$$T(\vec{x}) = A\vec{x} = 0\vec{x} = \vec{0}.$$

Note that the first zero is the matrix  $A$ , while the second zero is the zero vector of  $\mathbb{R}^m$ . The zero transformation is usually written as  $T = 0$ .

## Example (The Identity Transformation)

The transformation of  $\mathbb{R}^n$  induced by  $I_n$ , the  $n \times n$  identity matrix, is called the **identity transformation** because for every vector  $\vec{x}$  in  $\mathbb{R}^n$ ,

$$T(\vec{x}) = I_n\vec{x} = \vec{x}.$$

# Some Special Matrix Transformations

## Example (The Zero Transformation)

If  $A$  is the  $m \times n$  matrix of zeros, then the transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  induced by  $A$  is called the **zero transformation** because for every vector  $\vec{x}$  in  $\mathbb{R}^n$

$$T(\vec{x}) = A\vec{x} = 0\vec{x} = \vec{0}.$$

Note that the first zero is the matrix  $A$ , while the second zero is the zero vector of  $\mathbb{R}^m$ . The zero transformation is usually written as  $T = 0$ .

## Example (The Identity Transformation)

The transformation of  $\mathbb{R}^n$  induced by  $I_n$ , the  $n \times n$  identity matrix, is called the **identity transformation** because for every vector  $\vec{x}$  in  $\mathbb{R}^n$ ,

$$T(\vec{x}) = I_n\vec{x} = \vec{x}.$$

The identity transformation on  $\mathbb{R}^n$  is usually written as  $1_{\mathbb{R}^n}$ .

## Example (Revisited)

Recall  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  defined by

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

# Not all transformations are matrix transformations!

## Example

Consider  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

# Not all transformations are matrix transformations!

## Example

Consider  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Why is  $T$  not a matrix transformation?

## Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

### Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation,

### Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider  $T(0)$ , where  $0$  is the zero vector of  $\mathbb{R}^2$ .



### Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider  $T(0)$ , where  $0$  is the zero vector of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

### Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider  $T(0)$ , where  $0$  is the zero vector of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

### Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider  $T(0)$ , where  $0$  is the zero vector of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

### Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider  $T(0)$ , where  $0$  is the zero vector of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

violating one of the properties of a linear transformation.

### Example (continued)

We have  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider  $T(0)$ , where  $0$  is the zero vector of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

violating one of the properties of a linear transformation.

Therefore,  $T$  is not a linear transformation, and hence is not a matrix transformation.

Can you see any other reasons why  $T$  is not a matrix transformation?

# Matrix Transformations

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then we can find an  $n \times m$  matrix  $A$  such that

$$T(\vec{x}) = A\vec{x}$$

# Matrix Transformations

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then we can find an  $n \times m$  matrix  $A$  such that

$$T(\vec{x}) = A\vec{x}$$

In this case, we say that  $T$  is induced, or determined, by  $A$  and we write

$$T_A(\vec{x}) = A\vec{x}$$

# Matrix Transformations

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then we can find an  $n \times m$  matrix  $A$  such that

$$T(\vec{x}) = A\vec{x}$$

In this case, we say that  $T$  is induced, or determined, by  $A$  and we write

$$T_A(\vec{x}) = A\vec{x}$$

## Corollary

A transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a linear transformation if and only if it is a matrix transformation.



# Matrix and Linear Transformations

Good News!

# Matrix and Linear Transformations

Good News!

There is an easy way to find the matrix of  $T$ !

# Matrix and Linear Transformations

## Good News!

There is an easy way to find the matrix of  $T$ !

## Definition

The set of columns  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  of  $I_n$  is called the **standard basis of  $\mathbb{R}^n$** .

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation.

# Matrix and Linear Transformations

## Good News!

There is an easy way to find the matrix of  $T$ !

## Definition

The set of columns  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  of  $I_n$  is called the **standard basis of  $\mathbb{R}^n$** .

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then  $T$  is a matrix transformation.

# Matrix and Linear Transformations

## Good News!

There is an easy way to find the matrix of  $T$ !

## Definition

The set of columns  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  of  $I_n$  is called the **standard basis of  $\mathbb{R}^n$** .

## Theorem

Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then  $T$  is a matrix transformation. Furthermore,  $T$  is induced by the **unique** matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) & \cdots & T(\vec{e}_n) \end{bmatrix},$$

where  $\vec{e}_j$  is the  $j^{\text{th}}$  column of  $I_n$ , and  $T(\vec{e}_j)$  is the  $j^{\text{th}}$  column of  $A$ .

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .



## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix}$$

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix}$$

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

The columns  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$  become the columns of  $A$ ,

## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

The columns  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$  become the columns of  $A$ , so

$$A = \begin{bmatrix} 1 & 2 \\ 1 & -1 \end{bmatrix},$$

and  $T(\vec{x}) = A\vec{x}$  for every  $\vec{x} \in \mathbb{R}^2$ .



## Problem

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each  $\vec{x} \in \mathbb{R}^2$ . Find the matrix,  $A$ , of  $T$ .

## Solution

To find  $A$ , we must find  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$ , where  $\vec{e}_1$  and  $\vec{e}_2$  are the standard basis vectors of  $\mathbb{R}^2$ .

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ and } T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

The columns  $T(\vec{e}_1)$  and  $T(\vec{e}_2)$  become the columns of  $A$ , so

$$A = \begin{bmatrix} 1 & 2 \\ 1 & -1 \end{bmatrix},$$

and  $T(\vec{x}) = A\vec{x}$  for every  $\vec{x} \in \mathbb{R}^2$ . Therefore  $A$  is the matrix for  $T$ .

# Find the Matrix of $T$

## Problem

Sometimes  $T$  is not defined so nicely for us. Suppose  $T$  is given as

$$T \begin{bmatrix} 1 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad T \begin{bmatrix} 1 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

Find the matrix  $A$  of  $T$ .

# Find the Matrix of $T$

## Problem

Sometimes  $T$  is not defined so nicely for us. Suppose  $T$  is given as

$$T \begin{bmatrix} 1 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad T \begin{bmatrix} 1 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

Find the matrix  $A$  of  $T$ .

## Solution

We need to write  $\vec{e}_1$  and  $\vec{e}_2$  as a linear combination of the vectors provided. So we reduce the augmented matrix having  $\vec{e}_1$  and  $\vec{e}_2$  as the third and fourth columns:

$$\left[ \begin{array}{cc|cc} 1 & 1 & \vec{e}_1 & \vec{e}_2 \\ 5 & 4 & & \end{array} \right] = \left[ \begin{array}{cc|cc} 1 & 1 & 1 & 0 \\ 5 & 4 & 0 & 1 \end{array} \right]$$

# Find the Matrix of $T$

## Problem

Sometimes  $T$  is not defined so nicely for us. Suppose  $T$  is given as

$$T \begin{bmatrix} 1 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad T \begin{bmatrix} 1 \\ 4 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

Find the matrix  $A$  of  $T$ .

## Solution

We need to write  $\vec{e}_1$  and  $\vec{e}_2$  as a linear combination of the vectors provided. So we reduce the augmented matrix having  $\vec{e}_1$  and  $\vec{e}_2$  as the third and fourth columns:

$$\left[ \begin{array}{cc|cc} 1 & 1 & \vec{e}_1 & \vec{e}_2 \\ 5 & 4 & & \end{array} \right] = \left[ \begin{array}{cc|cc} 1 & 1 & 1 & 0 \\ 5 & 4 & 0 & 1 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{cc|cc} 1 & 0 & -4 & 1 \\ 0 & 1 & 5 & -1 \end{array} \right]$$

## Solution (continued)

$$\left[ \begin{array}{cc|cc} 1 & 1 & 1 & 0 \\ 5 & 4 & 0 & 1 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{cc|cc} 1 & 0 & -4 & 1 \\ 0 & 1 & 5 & -1 \end{array} \right]$$

From this we can see that

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} = -4 \begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5 \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

and

$$\vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 5 \end{bmatrix} - \begin{bmatrix} 1 \\ 4 \end{bmatrix}$$

## Solution (continued)

So

$$T(\vec{e}_1) = T\left(-4 \begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5 \begin{bmatrix} 1 \\ 4 \end{bmatrix}\right)$$

## Solution (continued)

So

$$\begin{aligned} T(\vec{e}_1) &= T\left(-4\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\ &= -4T\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \end{aligned}$$

## Solution (continued)

So

$$\begin{aligned}T(\vec{e}_1) &= T\left(-4\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= -4T\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= -4\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 5\begin{bmatrix} 3 \\ 2 \end{bmatrix}\end{aligned}$$



## Solution (continued)

So

$$\begin{aligned}T(\vec{e}_1) &= T\left(-4\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= -4T\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= -4\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 5\begin{bmatrix} 3 \\ 2 \end{bmatrix} \\&= \begin{bmatrix} 11 \\ 2 \end{bmatrix}\end{aligned}$$

## Solution (continued)

So

$$\begin{aligned}T(\vec{e}_1) &= T\left(-4\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= -4T\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= -4\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 5\begin{bmatrix} 3 \\ 2 \end{bmatrix} \\&= \begin{bmatrix} 11 \\ 2 \end{bmatrix}\end{aligned}$$

and

$$\begin{aligned}T(\vec{e}_2) &= T\begin{bmatrix} 0 \\ 1 \end{bmatrix} \\&= T\left(\begin{bmatrix} 1 \\ 5 \end{bmatrix} - \begin{bmatrix} 1 \\ 4 \end{bmatrix}\right)\end{aligned}$$

## Solution (continued)

So

$$\begin{aligned}T(\vec{e}_1) &= T\left(-4\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= -4T\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= -4\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 5\begin{bmatrix} 3 \\ 2 \end{bmatrix} \\&= \begin{bmatrix} 11 \\ 2 \end{bmatrix}\end{aligned}$$

and

$$\begin{aligned}T(\vec{e}_2) &= T\begin{bmatrix} 0 \\ 1 \end{bmatrix} \\&= T\left(\begin{bmatrix} 1 \\ 5 \end{bmatrix} - \begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= T\begin{bmatrix} 1 \\ 5 \end{bmatrix} - T\begin{bmatrix} 1 \\ 4 \end{bmatrix}\end{aligned}$$

## Solution (continued)

So

$$\begin{aligned}T(\vec{e}_1) &= T\left(-4\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= -4T\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= -4\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 5\begin{bmatrix} 3 \\ 2 \end{bmatrix} \\&= \begin{bmatrix} 11 \\ 2 \end{bmatrix}\end{aligned}$$

and

$$\begin{aligned}T(\vec{e}_2) &= T\begin{bmatrix} 0 \\ 1 \end{bmatrix} \\&= T\left(\begin{bmatrix} 1 \\ 5 \end{bmatrix} - \begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= T\begin{bmatrix} 1 \\ 5 \end{bmatrix} - T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 3 \\ 2 \end{bmatrix}\end{aligned}$$

## Solution (continued)

So

$$\begin{aligned}T(\vec{e}_1) &= T\left(-4\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= -4T\begin{bmatrix} 1 \\ 5 \end{bmatrix} + 5T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= -4\begin{bmatrix} 1 \\ 2 \end{bmatrix} + 5\begin{bmatrix} 3 \\ 2 \end{bmatrix} \\&= \begin{bmatrix} 11 \\ 2 \end{bmatrix}\end{aligned}$$

and

$$\begin{aligned}T(\vec{e}_2) &= T\begin{bmatrix} 0 \\ 1 \end{bmatrix} \\&= T\left(\begin{bmatrix} 1 \\ 5 \end{bmatrix} - \begin{bmatrix} 1 \\ 4 \end{bmatrix}\right) \\&= T\begin{bmatrix} 1 \\ 5 \end{bmatrix} - T\begin{bmatrix} 1 \\ 4 \end{bmatrix} \\&= \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \begin{bmatrix} 3 \\ 2 \end{bmatrix} \\&= \begin{bmatrix} -2 \\ 0 \end{bmatrix}\end{aligned}$$

### Solution (continued)

The matrix of  $T$  is the matrix whose first column is  $T(\vec{e}_1)$ , and second column is  $T(\vec{e}_2)$ :

### Solution (continued)

The matrix of  $T$  is the matrix whose first column is  $T(\vec{e}_1)$ , and second column is  $T(\vec{e}_2)$ :

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix} = \begin{bmatrix} 11 & -2 \\ 2 & 0 \end{bmatrix}$$

# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .



# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .

One way to show that  $T$  is a linear transformation is to show that it **preserves addition and scalar multiplication**.

# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .

One way to show that  $T$  is a linear transformation is to show that it **preserves addition and scalar multiplication**. However, now that we know that linear transformations are matrix transformations, we can use this to our advantage.

# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .

One way to show that  $T$  is a linear transformation is to show that it **preserves addition and scalar multiplication**. However, now that we know that linear transformations are matrix transformations, we can use this to our advantage.

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .

One way to show that  $T$  is a linear transformation is to show that it **preserves addition and scalar multiplication**. However, now that we know that linear transformations are matrix transformations, we can use this to our advantage.

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix}$$

# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .

One way to show that  $T$  is a linear transformation is to show that it **preserves addition and scalar multiplication**. However, now that we know that linear transformations are matrix transformations, we can use this to our advantage.

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix} = \begin{bmatrix} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} & T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix}$$

# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .

One way to show that  $T$  is a linear transformation is to show that it **preserves addition and scalar multiplication**. However, now that we know that linear transformations are matrix transformations, we can use this to our advantage.

If  $T$  were a linear transformation, then  **$T$  would be induced by the matrix**

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix} = \begin{bmatrix} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} & T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}.$$

# Determining if a Transformation is Linear

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$ .

One way to show that  $T$  is a linear transformation is to show that it **preserves addition and scalar multiplication**. However, now that we know that linear transformations are matrix transformations, we can use this to our advantage.

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 2 & 0 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}.$$

Since

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix},$$

$T$  is a matrix transformation, and is therefore a linear transformation.

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .



## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix}$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right]$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ ,

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix}$$



## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix}$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix} \text{ while } T \begin{bmatrix} x \\ y \end{bmatrix}$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix} \quad \text{while} \quad T \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix} \text{ while } T \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix},$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix} \text{ while } T \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix},$$

$$\text{i.e., } A \begin{bmatrix} 1 \\ 1 \end{bmatrix} \neq T \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

## Example

Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a transformation defined by  $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$ .

If  $T$  were a linear transformation, then  $T$  would be induced by the matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) ] = \left[ T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However,

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix}.$$

We see from this that if  $x = 0$  or  $y = 0$ , then  $xy = 0$ , so  $A \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$ .

But if we take  $x = y = 1$ , then

$$A \begin{bmatrix} x \\ y \end{bmatrix} = A \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 2 \end{bmatrix} \text{ while } T \begin{bmatrix} x \\ y \end{bmatrix} = T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix},$$

i.e.,  $A \begin{bmatrix} 1 \\ 1 \end{bmatrix} \neq T \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . Therefore,  $T$  is **not** a linear transformation.

# Rotations in $\mathbb{R}^2$

## Definition

The transformation

$$R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

denotes counterclockwise rotation about the origin through an angle of  $\theta$ .



# Rotations in $\mathbb{R}^2$

## Definition

The transformation

$$R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

denotes counterclockwise rotation about the origin through an angle of  $\theta$ .

Rotation through an angle of  $\theta$  preserves scalar multiplication.

# Rotations in $\mathbb{R}^2$

## Definition

The transformation

$$R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

denotes counterclockwise rotation about the origin through an angle of  $\theta$ .

Rotation through an angle of  $\theta$  preserves scalar multiplication.

Rotation through an angle of  $\theta$  preserves vector addition.

## $R_\theta$ is a linear transformation

Since  $R_\theta$  preserves addition and scalar multiplication,  $R_\theta$  is a linear transformation, and hence a matrix transformation.

The matrix that induces  $R_\theta$  can be found by computing  $R_\theta(E_1)$  and  $R_\theta(E_2)$ , where

$$E_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad E_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

## $R_\theta$ is a linear transformation

Since  $R_\theta$  preserves addition and scalar multiplication,  $R_\theta$  is a linear transformation, and hence a matrix transformation.

The matrix that induces  $R_\theta$  can be found by computing  $R_\theta(E_1)$  and  $R_\theta(E_2)$ , where

$$E_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad E_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(E_1)$$

## $R_\theta$ is a linear transformation

Since  $R_\theta$  preserves addition and scalar multiplication,  $R_\theta$  is a linear transformation, and hence a matrix transformation.

The matrix that induces  $R_\theta$  can be found by computing  $R_\theta(E_1)$  and  $R_\theta(E_2)$ , where

$$E_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad E_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(E_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

## $R_\theta$ is a linear transformation

Since  $R_\theta$  preserves addition and scalar multiplication,  $R_\theta$  is a linear transformation, and hence a matrix transformation.

The matrix that induces  $R_\theta$  can be found by computing  $R_\theta(E_1)$  and  $R_\theta(E_2)$ , where

$$E_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad E_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(E_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

## $R_\theta$ is a linear transformation

Since  $R_\theta$  preserves addition and scalar multiplication,  $R_\theta$  is a linear transformation, and hence a matrix transformation.

The matrix that induces  $R_\theta$  can be found by computing  $R_\theta(E_1)$  and  $R_\theta(E_2)$ , where

$$E_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad E_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(E_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

and

$$R_\theta(E_2)$$

## $R_\theta$ is a linear transformation

Since  $R_\theta$  preserves addition and scalar multiplication,  $R_\theta$  is a linear transformation, and hence a matrix transformation.

The matrix that induces  $R_\theta$  can be found by computing  $R_\theta(E_1)$  and  $R_\theta(E_2)$ , where

$$E_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad E_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(E_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

and

$$R_\theta(E_2) = R_\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



## $R_\theta$ is a linear transformation

Since  $R_\theta$  preserves addition and scalar multiplication,  $R_\theta$  is a linear transformation, and hence a matrix transformation.

The matrix that induces  $R_\theta$  can be found by computing  $R_\theta(E_1)$  and  $R_\theta(E_2)$ , where

$$E_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad E_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(E_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

and

$$R_\theta(E_2) = R_\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$$

## The Matrix for $R_\theta$

The rotation  $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a linear transformation, and is induced by the matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

## Example (Rotation through $\pi$ )

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$


counterclockwise rotation about the origin through an angle of  $\pi$ .

## Example (Rotation through $\pi$ )

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi$ .




## Example (Rotation through $\pi$ )

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi$ .




## Example (Rotation through $\pi$ )

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi$ .




We see that  $R_\pi \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -a \\ -b \end{bmatrix} =$

## Example (Rotation through $\pi$ )

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi$ .



We see that  $R_\pi \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -a \\ -b \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$ , so  $R_\pi$  is a matrix transformation.

# Rotation

## Problem

The transformation  $R_{\frac{\pi}{2}} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  denotes a **counterclockwise** rotation about the origin through an angle of  $\frac{\pi}{2}$  radians. Find the matrix of  $R_{\frac{\pi}{2}}$ .



# Rotation

## Problem

The transformation  $R_{\frac{\pi}{2}} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  denotes a **counterclockwise** rotation about the origin through an angle of  $\frac{\pi}{2}$  radians. Find the matrix of  $R_{\frac{\pi}{2}}$ .

## Solution

First,

$$R_{\frac{\pi}{2}} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix}$$

# Rotation

## Problem

The transformation  $R_{\frac{\pi}{2}} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  denotes a **counterclockwise** rotation about the origin through an angle of  $\frac{\pi}{2}$  radians. Find the matrix of  $R_{\frac{\pi}{2}}$ .

## Solution

First,

$$R_{\frac{\pi}{2}} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix}$$

Furthermore  $R_{\frac{\pi}{2}}$  is a matrix transformation, and the matrix it is induced by is

$$\begin{bmatrix} -b \\ a \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$

## Example (Rotation through $\pi/2$ )

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$


counterclockwise rotation about the origin through an angle of  $\pi/2$ .

## Example (Rotation through $\pi/2$ )

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi/2$ .




## Example (Rotation through $\pi/2$ )

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi/2$ .




## Example (Rotation through $\pi/2$ )

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi/2$ .




We see that 
$$R_{\pi/2} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} =$$

## Example (Rotation through $\pi/2$ )

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of  $\pi/2$ .



We see that  $R_{\pi/2} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$ , so  $R_{\pi/2}$  is a matrix transformation.

# Reflection in $\mathbb{R}^2$

## Example

In  $\mathbb{R}^2$ , reflection in the  $x$ -axis, which transforms  $\begin{bmatrix} a \\ b \end{bmatrix}$  to  $\begin{bmatrix} a \\ -b \end{bmatrix}$ , is a matrix transformation because

$$\begin{bmatrix} a \\ -b \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$



# Reflection in $\mathbb{R}^2$

## Example

In  $\mathbb{R}^2$ , reflection in the  $x$ -axis, which transforms  $\begin{bmatrix} a \\ b \end{bmatrix}$  to  $\begin{bmatrix} a \\ -b \end{bmatrix}$ , is a matrix transformation because

$$\begin{bmatrix} a \\ -b \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$


## Example

In  $\mathbb{R}^2$ , reflection in the  $y$ -axis transforms  $\begin{bmatrix} a \\ b \end{bmatrix}$  to  $\begin{bmatrix} -a \\ b \end{bmatrix}$ . This is a matrix transformation because

$$\begin{bmatrix} -a \\ b \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$


## Example

Reflection in the line  $y = x$  transforms  $\begin{bmatrix} a \\ b \end{bmatrix}$  to  $\begin{bmatrix} b \\ a \end{bmatrix}$ .



## Example

Reflection in the line  $y = x$  transforms  $\begin{bmatrix} a \\ b \end{bmatrix}$  to  $\begin{bmatrix} b \\ a \end{bmatrix}$ .



This is a matrix transformation because

$$\begin{bmatrix} b \\ a \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$