

A First Course in  
**LINEAR ALGEBRA**

**Lecture Notes**  
for Math 1503

**6.1: Complex Numbers**

Creative Commons License (CC BY-NC-SA)

# A First Course in Linear Algebra

## Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

### License



**Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)**

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

## Why complex numbers?

- Counting numbers:  $1, 2, 3, 4, 5, \dots$

## Why complex numbers?

- Counting numbers:  $1, 2, 3, 4, 5, \dots$
- Integers:  $0, 1, 2, 3, 4 \dots$  but also  $-1, -2, -3 \dots$

## Why complex numbers?

- Counting numbers:  $1, 2, 3, 4, 5, \dots$
- Integers:  $0, 1, 2, 3, 4 \dots$  but also  $-1, -2, -3 \dots$
- To solve  $3x + 2 = 0$ , integers aren't enough, so we have **rational numbers** (fractions), i.e.,  
if  $3x + 2 = 0$ , then  $x = -\frac{2}{3}$ .

## Why complex numbers?

- Counting numbers:  $1, 2, 3, 4, 5, \dots$
- Integers:  $0, 1, 2, 3, 4, \dots$  but also  $-1, -2, -3, \dots$
- To solve  $3x + 2 = 0$ , integers aren't enough, so we have **rational numbers** (fractions), i.e.,  
if  $3x + 2 = 0$ , then  $x = -\frac{2}{3}$ .
- We still can't solve  $x^2 - 2 = 0$  because there are no rational numbers  $x$  with the property that  $x^2 - 2 = 0$ , so we have **irrational numbers**, i.e.,  
if  $x^2 - 2 = 0$ , then  $x = \pm\sqrt{2}$ .

## Why complex numbers?

- Counting numbers:  $1, 2, 3, 4, 5, \dots$
- Integers:  $0, 1, 2, 3, 4, \dots$  but also  $-1, -2, -3, \dots$
- To solve  $3x + 2 = 0$ , integers aren't enough, so we have **rational numbers** (fractions), i.e.,  
if  $3x + 2 = 0$ , then  $x = -\frac{2}{3}$ .
- We still can't solve  $x^2 - 2 = 0$  because there are no rational numbers  $x$  with the property that  $x^2 - 2 = 0$ , so we have **irrational numbers**, i.e.,  
if  $x^2 - 2 = 0$ , then  $x = \pm\sqrt{2}$ .
- The set of **real numbers**,  $\mathbb{R}$ , consists of all rational and irrational numbers (note that integers are rational numbers). However, we still can't solve

$$x^2 + 1 = 0$$

because this requires  $x^2 = -1$ , but any **real** number  $x$  has the property that  $x^2 \geq 0$ .

# Complex Numbers

## Definitions



# Complex Numbers

## Definitions

- The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .

# Complex Numbers

## Definitions

- The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.

# Complex Numbers

## Definitions

- The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

# Complex Numbers

## Definitions

- The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

- ▶  $a$  is called the **real part** of  $z$ .

# Complex Numbers

## Definitions

- The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

- ▶  $a$  is called the **real part** of  $z$ .
- ▶  $b$  is called the **imaginary part** of  $z$ .

# Complex Numbers

## Definitions

- The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

- ▶  $a$  is called the **real part** of  $z$ .
- ▶  $b$  is called the **imaginary part** of  $z$ .
- ▶ If  $b = 0$ , then  $z$  is a real number.

# The Complex Plane (Argand Plane)


## The Complex Plane (Argand Plane)

A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.




## The Complex Plane (Argand Plane)

A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.



## The Complex Plane (Argand Plane)


A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.



- Real numbers:  $a + 0i$  lie on the  $x$ -axis.

## The Complex Plane (Argand Plane)

A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.



- Real numbers:  $a + 0i$  lie on the  $x$ -axis.
- Pure imaginary numbers:  $0 + bi$  ( $b \neq 0$ ) lie on the  $y$ -axis.

## Addition and Subtraction of Complex Numbers

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- **Addition**

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- **Addition**

$$z + w =$$



## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- **Addition**

$$z + w = (a + bi) + (c + di) =$$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

- $(-3 + 6i) + (5 - i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

- $(-3 + 6i) + (5 - i) = 2 + 5i.$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

- $(-3 + 6i) + (5 - i) = 2 + 5i.$
- $(4 - 7i) + (6 - 2i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

### Examples

- $(-3 + 6i) + (5 - i) = 2 + 5i.$
- $(4 - 7i) + (6 - 2i) = 10 - 9i.$



## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

### Examples

- $(-3 + 6i) + (5 - i) = 2 + 5i.$
- $(4 - 7i) + (6 - 2i) = 10 - 9i.$
- $(-3 + 6i) - (5 - i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

### Examples

- $(-3 + 6i) + (5 - i) = 2 + 5i.$
- $(4 - 7i) + (6 - 2i) = 10 - 9i.$
- $(-3 + 6i) - (5 - i) = -8 + 7i.$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

### Examples

- $(-3 + 6i) + (5 - i) = 2 + 5i.$
- $(4 - 7i) + (6 - 2i) = 10 - 9i.$
- $(-3 + 6i) - (5 - i) = -8 + 7i.$
- $(4 - 7i) - (6 - 2i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

- **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

- **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

### Examples

- $(-3 + 6i) + (5 - i) = 2 + 5i.$
- $(4 - 7i) + (6 - 2i) = 10 - 9i.$
- $(-3 + 6i) - (5 - i) = -8 + 7i.$
- $(4 - 7i) - (6 - 2i) = -2 - 5i.$

## Properties of Addition

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

①  $z + w = w + z.$

(addition is commutative)

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

①  $z + w = w + z.$

(addition is commutative)

②  $(z + w) + v = z + (w + v).$

(addition is associative)



## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

①  $z + w = w + z.$

(addition is commutative)

②  $(z + w) + v = z + (w + v).$

(addition is associative)

③  $z + 0 = z.$

(existence of an additive identity)

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

- ①  $z + w = w + z.$  (addition is commutative)
- ②  $(z + w) + v = z + (w + v).$  (addition is associative)
- ③  $z + 0 = z.$  (existence of an additive identity)
- ④ For every  $z = a + bi$  there exists a complex number  $-z = -a - bi$  such that  $z + (-z) = 0.$  (existence of an additive inverse)

## Addition in the Complex Plane


## Addition in the Complex Plane

If  $z = a + bi$  and  $w = c + di$ , then  $z + w = (a + c) + (b + d)i$ .

## Addition in the Complex Plane

If  $z = a + bi$  and  $w = c + di$ , then  $z + w = (a + c) + (b + d)i$ .


Geometrically, we have:



## Addition in the Complex Plane

If  $z = a + bi$  and  $w = c + di$ , then  $z + w = (a + c) + (b + d)i$ .

Geometrically, we have:



$0$ ,  $z$ ,  $w$ , and  $z + w$  are the vertices of a parallelogram.

## Multiplication of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Then the **product** of  $z$  and  $w$  is

$$zw = (a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

## Multiplication of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Then the **product** of  $z$  and  $w$  is

$$zw = (a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

The multiplication is done essentially as the product of two linear polynomials, with  $i^2$  replaced by  $-1$ .



## Multiplication of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Then the **product** of  $z$  and  $w$  is

$$zw = (a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

The multiplication is done essentially as the product of two linear polynomials, with  $i^2$  replaced by  $-1$ .

### Example

$$\begin{aligned}(2 - 3i)(-3 + 4i) &= ((2)(-3) - (-3)(4)) + ((2)(4) + (-3)(-3))i \\ &= (-6 + 12) + (8 + 9)i \\ &= 6 + 17i\end{aligned}$$

## Properties of Multiplication

## Properties of Multiplication

Let  $z$ ,  $w$  and  $v$  be complex numbers.

## Properties of Multiplication

Let  $z, w$  and  $v$  be complex numbers.

- $zw = wz$ .

(multiplication is commutative)

## Properties of Multiplication

Let  $z, w$  and  $v$  be complex numbers.

- $zw = wz$ .

(multiplication is commutative)

- $(zw)v = z(wv)$ .

(multiplication is associative)

## Properties of Multiplication

Let  $z, w$  and  $v$  be complex numbers.

- $zw = wz.$  (multiplication is commutative)
- $(zw)v = z(wv).$  (multiplication is associative)
- $z(w + v) = zw + zv.$  (multiplication distributes over addition)

## Properties of Multiplication

Let  $z, w$  and  $v$  be complex numbers.

- $zw = wz.$  (multiplication is commutative)
- $(zw)v = z(wv).$  (multiplication is associative)
- $z(w + v) = zw + zv.$  (multiplication distributes over addition)
- $1z = z.$  ('1' is the multiplicative identity)

## Properties of Multiplication

Let  $z, w$  and  $v$  be complex numbers.

- $zw = wz.$  (multiplication is commutative)
- $(zw)v = z(wv).$  (multiplication is associative)
- $z(w + v) = zw + zv.$  (multiplication distributes over addition)
- $1z = z.$  ('1' is the multiplicative identity)
- For each  $z \neq 0$ , there exists  $z^{-1}$  such that  $zz^{-1} = 1.$  (existence of a multiplicative inverse)



## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

so

$$a^2 - b^2 = -3 \text{ and } 2ab = 4.$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

so

$$a^2 - b^2 = -3 \text{ and } 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ .

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

so

$$a^2 - b^2 = -3 \text{ and } 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ . Substituting this into the first equation gives us

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

so

$$a^2 - b^2 = -3 \text{ and } 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \end{aligned}$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

so

$$a^2 - b^2 = -3 \text{ and } 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \\ \frac{4}{b^2} - b^2 &= -3 \end{aligned}$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

so

$$a^2 - b^2 = -3 \text{ and } 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \\ \frac{4}{b^2} - b^2 &= -3 \\ 4 - b^4 &= -3b^2 \end{aligned}$$



## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

so

$$a^2 - b^2 = -3 \text{ and } 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \\ \frac{4}{b^2} - b^2 &= -3 \\ 4 - b^4 &= -3b^2 \\ b^4 - 3b^2 - 4 &= 0. \end{aligned}$$

## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored

## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

$$(b - 2)(b + 2)(b^2 + 1) = 0.$$

## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

$$(b - 2)(b + 2)(b^2 + 1) = 0.$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

$$(b - 2)(b + 2)(b^2 + 1) = 0.$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that

## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

$$(b - 2)(b + 2)(b^2 + 1) = 0.$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that

- when  $b = 2$ ,  $a = 1$ , and  $z = a + bi = 1 + 2i$ ;

## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

$$(b - 2)(b + 2)(b^2 + 1) = 0.$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that

- when  $b = 2$ ,  $a = 1$ , and  $z = a + bi = 1 + 2i$ ;
- when  $b = -2$ ,  $a = -1$ , and  $z = a + bi = -1 - 2i$ .



## Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

$$(b - 2)(b + 2)(b^2 + 1) = 0.$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that

- when  $b = 2$ ,  $a = 1$ , and  $z = a + bi = 1 + 2i$ ;
- when  $b = -2$ ,  $a = -1$ , and  $z = a + bi = -1 - 2i$ .

Therefore, if  $z^2 = -3 + 4i$ , then  $z = 1 + 2i$  or  $z = -1 - 2i$ .

## The Conjugate of a Complex Number

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number.

## The Conjugate of a Complex Number


Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ .

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.


## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.



## The Conjugate of a Complex Number


Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.



## Examples

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} =$



## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ ,

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- $\overline{-2 + 5i} =$

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- $\overline{-2 + 5i} = -2 - 5i$ .

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- $\overline{-2 + 5i} = -2 - 5i$ .
- $\bar{i} =$

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- $\overline{-2 + 5i} = -2 - 5i$ .
- $\bar{i} = -i$ .

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- $\overline{-2 + 5i} = -2 - 5i$ .
- $\bar{i} = -i$ .
- $\bar{\bar{z}} = z$

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.



### Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- $\overline{-2 + 5i} = -2 - 5i$ .
- $\bar{i} = -i$ .
- $\bar{7} = 7$ .



## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}.$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}.$
- $\overline{(zw)} = \bar{z} \bar{w}.$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}.$
- $\overline{(zw)} = \bar{z} \bar{w}.$
- $\overline{(\bar{z})} = z.$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}.$

- $\overline{(zw)} = \bar{z} \bar{w}.$

- $\overline{(\bar{z})} = z.$

- $\overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}.$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}$ .
- $\overline{zw} = \bar{z} \bar{w}$ .
- $\overline{(\bar{z})} = z$ .
- $\overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}$ .
- $z$  is real if and only if  $\bar{z} = z$ .

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}$ .
- $\overline{(zw)} = \bar{z} \bar{w}$ .
- $\overline{(\bar{z})} = z$ .
- $\overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}$ .
- $z$  is real if and only if  $\bar{z} = z$ .

### Note

If  $z = a + bi$ , then

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}$ .
- $\overline{(zw)} = \bar{z} \bar{w}$ .
- $\overline{(\bar{z})} = z$ .
- $\overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}$ .
- $z$  is real if and only if  $\bar{z} = z$ .

### Note

If  $z = a + bi$ , then

$$z\bar{z} =$$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}$ .
- $\overline{(zw)} = \bar{z} \bar{w}$ .
- $\overline{(\bar{z})} = z$ .
- $\overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}$ .
- $z$  is real if and only if  $\bar{z} = z$ .

### Note

If  $z = a + bi$ , then

$$z\bar{z} = (a + bi)(a - bi) =$$



## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

- $\overline{z \pm w} = \bar{z} \pm \bar{w}$ .
- $\overline{zw} = \bar{z} \bar{w}$ .
- $\overline{(\bar{z})} = z$ .
- $\overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}$ .
- $z$  is real if and only if  $\bar{z} = z$ .

### Note

If  $z = a + bi$ , then

$$z\bar{z} = (a + bi)(a - bi) = a^2 + b^2.$$

## Lecture 2

## Division of Complex Numbers

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero.

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\frac{z}{w} =$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\frac{z}{w} = \frac{a + bi}{c + di} =$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} &= \frac{a + bi}{c + di} = \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \end{aligned}$$



## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} &= \frac{a + bi}{c + di} = \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2} \\ &= \end{aligned}$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} &= \frac{a + bi}{c + di} = \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2} \\ &= \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i.\end{aligned}$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} &= \frac{a + bi}{c + di} = \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2} \\ &= \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i.\end{aligned}$$

The quotient  $\frac{z}{w}$  is obtained by multiplying both top and bottom of  $\frac{z}{w}$  by  $\overline{w}$  and then simplifying the expression.

## Examples

## Examples



$$\frac{1}{i} =$$

## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$

## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$



$$\frac{2-i}{3+4i} =$$

## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$



$$\frac{2-i}{3+4i} = \frac{2-i}{3+4i} \times \frac{3-4i}{3-4i} = \frac{(6-4) + (-3-8)i}{3^2 + 4^2} = \frac{2-11i}{25} = \frac{2}{25} - \frac{11}{25}i.$$



## Examples

- $$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$

- $$\frac{2-i}{3+4i} = \frac{2-i}{3+4i} \times \frac{3-4i}{3-4i} = \frac{(6-4) + (-3-8)i}{3^2 + 4^2} = \frac{2-11i}{25} = \frac{2}{25} - \frac{11}{25}i.$$

- $$\frac{1-2i}{-2+5i} =$$

## Examples

- $$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$

- $$\frac{2-i}{3+4i} = \frac{2-i}{3+4i} \times \frac{3-4i}{3-4i} = \frac{(6-4) + (-3-8)i}{3^2 + 4^2} = \frac{2-11i}{25} = \frac{2}{25} - \frac{11}{25}i.$$

- $$\frac{1-2i}{-2+5i} = \frac{1-2i}{-2+5i} \times \frac{-2-5i}{-2-5i} = \frac{(-2-10) + (4-5)i}{2^2 + 5^2} = -\frac{12}{29} - \frac{1}{29}i.$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$



## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} = \frac{2 - 6i}{40} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} = \frac{2 - 6i}{40} = \frac{1}{20} - \frac{3}{20}i.$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} = \frac{2 - 6i}{40} = \frac{1}{20} - \frac{3}{20}i.$$

You can always check that  $zz^{-1} = 1$ .



# Modulus

The **absolute value** or **modulus** of a complex number  $z = a + bi$  is

$$|z| = \sqrt{a^2 + b^2}$$

# Modulus

The **absolute value** or **modulus** of a complex number  $z = a + bi$  is

$$|z| = \sqrt{a^2 + b^2}$$

*Note that this is consistent with the definition of the absolute value of a real number.*


# Modulus

The **absolute value** or **modulus** of a complex number  $z = a + bi$  is

$$|z| = \sqrt{a^2 + b^2}$$

*Note that this is consistent with the definition of the absolute value of a real number.*

Geometrically,  $|z| = \sqrt{a^2 + b^2}$  is the distance from  $z$  to the origin.



## Examples

## Examples

①  $|-3 + 4i| =$

## Examples

$$\textcircled{1} \quad |-3 + 4i| = \sqrt{3^2 + 4^2} =$$

## Examples

$$\textcircled{1} \quad |-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} =$$

## Examples

$$\textcircled{1} \quad |-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$$



## Examples

①  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$

②  $|3 - 2i| =$

## Examples

$$\textcircled{1} \quad |-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$$

$$\textcircled{2} \quad |3 - 2i| = \sqrt{3^2 + 2^2} = \sqrt{13}.$$

## Examples

①  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$

②  $|3 - 2i| = \sqrt{3^2 + 2^2} = \sqrt{13}.$

③  $|i| =$


## Examples

$$\textcircled{1} \quad |-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$$

$$\textcircled{2} \quad |3 - 2i| = \sqrt{3^2 + 2^2} = \sqrt{13}.$$

$$\textcircled{3} \quad |i| = \sqrt{1^2} = 1.$$

# Properties of the Modulus



# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

①  $z \cdot \bar{z} = |z|^2.$

# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

①  $z \cdot \bar{z} = |z|^2.$

②  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}.$



# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

- ①  $z \cdot \bar{z} = |z|^2$ .
- ②  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}$ .
- ③  $|z| \geq 0$  for all  $z$ .

# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

①  $z \cdot \bar{z} = |z|^2.$

②  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}.$

③  $|z| \geq 0$  for all  $z$ .

④  $|z| = 0$  if and only if  $z = 0$ .

# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

①  $z \cdot \bar{z} = |z|^2.$

②  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}.$

③  $|z| \geq 0$  for all  $z$ .

④  $|z| = 0$  if and only if  $z = 0$ .

⑤  $|zw| = |z| |w|.$

# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

①  $z \cdot \bar{z} = |z|^2.$

②  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}.$

③  $|z| \geq 0$  for all  $z$ .

④  $|z| = 0$  if and only if  $z = 0$ .

⑤  $|zw| = |z| |w|.$

⑥  $\left| \frac{z}{w} \right| = \frac{|z|}{|w|}.$

# Properties of the Modulus

Let  $z$  and  $w$  be complex numbers.

①  $z \cdot \bar{z} = |z|^2.$

②  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}.$

③  $|z| \geq 0$  for all  $z$ .

④  $|z| = 0$  if and only if  $z = 0$ .

⑤  $|zw| = |z| |w|.$

⑥  $\left| \frac{z}{w} \right| = \frac{|z|}{|w|}.$

⑦ The Triangle Inequality


$$|z + w| \leq |z| + |w|.$$

## Distance in the plane

If  $z = a + bi$  and  $w = c + di$ , then  $|z - w| = \sqrt{(a - c)^2 + (b - d)^2}$ .

## Distance in the plane

If  $z = a + bi$  and  $w = c + di$ , then  $|z - w| = \sqrt{(a - c)^2 + (b - d)^2}$ .



This shows that the **distance** between  $z$  and  $w$  in the complex plane is just the absolute value of their difference.


## The Triangle Inequality: Geometrically

Now consider the points  $z$ ,  $z + w$ , and the origin  $0$  in the complex plane. These points may be considered as vectors in the plane:



## The Triangle Inequality: Geometrically


Now consider the points  $z$ ,  $z + w$ , and the origin  $0$  in the complex plane. These points may be considered as vectors in the plane:



The triangle formed by these points has sides of length  $|z|$ ,  $|z + w|$  and  $|w|$  (the absolute value of the difference between  $z + w$  and  $z$ ).

## The Triangle Inequality: Geometrically

Now consider the points  $z$ ,  $z + w$ , and the origin  $0$  in the complex plane. These points may be considered as vectors in the plane:



The triangle formed by these points has sides of length  $|z|$ ,  $|z + w|$  and  $|w|$  (the absolute value of the difference between  $z + w$  and  $z$ ).

Since the length of **any** side of a triangle is at most the sum of the lengths of the other two sides, we get  $|z + w| \leq |z| + |w|$ .