

A First Course in
LINEAR ALGEBRA

Lecture Notes
for Math 1503

**6.3: Complex Numbers; Roots of Complex
Numbers**

Creative Commons License (CC BY-NC-SA)

A First Course in Linear Algebra

Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Roots of Complex Numbers

Definition

Let z and q be complex numbers, and let n be a positive integer.

Roots of Complex Numbers

Definition

Let z and q be complex numbers, and let n be a positive integer. Then z is called an n^{th} root of q if $z^n = q$.

Roots of Complex Numbers

Definition

Let z and q be complex numbers, and let n be a positive integer. Then z is called **an n^{th} root of q** if $z^n = q$.

De Moivre's Theorem and its implication

If θ is any angle and n is a positive integer, $(e^{i\theta})^n = e^{in\theta}$. This implies that for any real number $r > 0$ and any positive integer n ,

$$(re^{i\theta})^n = r^n e^{in\theta}.$$

This leads to the following result.

Roots of Complex Numbers

Definition

Let z and q be complex numbers, and let n be a positive integer. Then z is called **an n^{th} root of q** if $z^n = q$.

De Moivre's Theorem and its implication

If θ is any angle and n is a positive integer, $(e^{i\theta})^n = e^{in\theta}$. This implies that for any real number $r > 0$ and any positive integer n ,

$$(re^{i\theta})^n = r^n e^{in\theta}.$$

This leads to the following result.

Corollary

Let q be a nonzero complex number and n a positive integer. Then $z^n = q$ has exactly n complex solutions, i.e., q has exactly n complex n^{th} roots.

Example

For any positive real number a , $z^2 = a$ has two complex (in this case, real) solutions, $z = \sqrt{a}$ and $z = -\sqrt{a}$. This is equivalent to the statement that a has two complex (in this case, real) square roots.

Example

For any positive real number a , $z^2 = a$ has two complex (in this case, real) solutions, $z = \sqrt{a}$ and $z = -\sqrt{a}$. This is equivalent to the statement that a has two complex (in this case, real) square roots.

- One particular example: 25 has two square roots, 5 and -5 , and these are the two solutions to $z^2 = 25$.

Example

For any positive real number a , $z^2 = a$ has two complex (in this case, real) solutions, $z = \sqrt{a}$ and $z = -\sqrt{a}$. This is equivalent to the statement that a has two complex (in this case, real) square roots.

- One particular example: 25 has two square roots, 5 and -5 , and these are the two solutions to $z^2 = 25$.
- But we all knew that. A more interesting example is that -1 has no real square roots, but suddenly it has two (complex) square roots, i and $-i$. These are the two (complex) solutions to $z^2 = 1$.

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$.

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 =$$

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1 e^{\pi i/2}$,

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1e^{\pi i/2}$, implying that $r^3 = 1$

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1 e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.
- The statement $3\theta = \frac{\pi}{2}$ is not completely correct.

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1 e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.
- The statement $3\theta = \frac{\pi}{2}$ is **not completely correct**. The problem that arises is that the argument for i , $\frac{\pi}{2}$ is not unique.

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1 e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.
- The statement $3\theta = \frac{\pi}{2}$ is **not completely correct**. The problem that arises is that the argument for i , $\frac{\pi}{2}$ is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.
- The statement $3\theta = \frac{\pi}{2}$ is **not completely correct**. The problem that arises is that the argument for i , $\frac{\pi}{2}$ is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

In fact, there are infinitely many choices for the argument.

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.
- The statement $3\theta = \frac{\pi}{2}$ is **not completely correct**. The problem that arises is that the argument for i , $\frac{\pi}{2}$ is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

In fact, there are infinitely many choices for the argument. The important thing to notice is that any two different arguments differ by a multiple of $2\pi i$,

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.
- The statement $3\theta = \frac{\pi}{2}$ is **not completely correct**. The problem that arises is that the argument for i , $\frac{\pi}{2}$ is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

In fact, there are infinitely many choices for the argument. The important thing to notice is that any two different arguments differ by a multiple of $2\pi i$, and thus we may write

$$3\theta = \frac{\pi}{2} + 2\pi\ell, \ell \in \mathbb{Z}.$$

Cube Roots

Example

To find the (three) cube roots of i , we solve the equation $z^3 = i$. To do so, we express both z and i in polar form: convert i to polar form, and write $z = re^{i\theta}$, giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus $r^3 e^{3i\theta} = 1e^{\pi i/2}$, implying that $r^3 = 1$ and $3\theta = \frac{\pi}{2}$.

- Since r is a non-negative real number, $r^3 = 1$ implies that $r = 1$.
- The statement $3\theta = \frac{\pi}{2}$ is **not completely correct**. The problem that arises is that the argument for i , $\frac{\pi}{2}$ is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

In fact, there are infinitely many choices for the argument. The important thing to notice is that any two different arguments differ by a multiple of $2\pi i$, and thus we may write

$$3\theta = \frac{\pi}{2} + 2\pi\ell, \ell \in \mathbb{Z}.$$

(\mathbb{Z} denotes the set of integers: $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$).

Example (continued)

Dividing both sides of $3\theta = \frac{\pi}{2} + 2\pi\ell$ by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where ℓ is any integer.

Example (continued)

Dividing both sides of $3\theta = \frac{\pi}{2} + 2\pi\ell$ by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where ℓ is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots.

Example (continued)

Dividing both sides of $3\theta = \frac{\pi}{2} + 2\pi\ell$ by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where ℓ is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using $\ell = 0$, $\ell = 1$, and $\ell = 2$, resulting in three values of θ :

Example (continued)

Dividing both sides of $3\theta = \frac{\pi}{2} + 2\pi\ell$ by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where ℓ is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using $\ell = 0$, $\ell = 1$, and $\ell = 2$, resulting in three values of θ :

$$\frac{\pi}{6}, \frac{5\pi}{6}, \text{ and } \frac{9\pi}{6} = \frac{3\pi}{2}.$$

Example (continued)

Dividing both sides of $3\theta = \frac{\pi}{2} + 2\pi\ell$ by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where ℓ is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using $\ell = 0$, $\ell = 1$, and $\ell = 2$, resulting in three values of θ :

$$\frac{\pi}{6}, \frac{5\pi}{6}, \text{ and } \frac{9\pi}{6} = \frac{3\pi}{2}.$$

Thus the cube roots of i are

$$e^{\pi i/6}, e^{5\pi i/6}, \text{ and } e^{3\pi i/2}.$$

Example (continued)

Dividing both sides of $3\theta = \frac{\pi}{2} + 2\pi\ell$ by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where ℓ is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using $\ell = 0$, $\ell = 1$, and $\ell = 2$, resulting in three values of θ :

$$\frac{\pi}{6}, \frac{5\pi}{6}, \text{ and } \frac{9\pi}{6} = \frac{3\pi}{2}.$$

Thus the cube roots of i are

$$e^{\pi i/6}, e^{5\pi i/6}, \text{ and } e^{3\pi i/2}.$$

We now convert these to Cartesian form.

Example (continued)

$$e^{\pi i/6} =$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} =$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} =$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} = -i.$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} = -i.$$

You can check your work by computing the cube of each of these.

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} = -i.$$

You can check your work by computing the cube of each of these.

This process is summarized in the following procedure.

Finding Roots of a Complex Number

Let w be a complex number. We wish to find the n^{th} roots of w , that is all z such that $z^n = w$.

There are n distinct n^{th} roots and they can be found as follows:.

Finding Roots of a Complex Number

Let w be a complex number. We wish to find the n^{th} roots of w , that is all z such that $z^n = w$.

There are n distinct n^{th} roots and they can be found as follows:.

1. Express both z and w in polar form $z = re^{i\theta}$, $w = se^{i\phi}$. Then $z^n = w$ becomes:

$$(re^{i\theta})^n = r^n e^{in\theta} = se^{i\phi}$$

We need to solve for r and θ .

Finding Roots of a Complex Number

Let w be a complex number. We wish to find the n^{th} roots of w , that is all z such that $z^n = w$.

There are n distinct n^{th} roots and they can be found as follows:

1. Express both z and w in polar form $z = re^{i\theta}$, $w = se^{i\phi}$. Then $z^n = w$ becomes:

$$(re^{i\theta})^n = r^n e^{in\theta} = se^{i\phi}$$

We need to solve for r and θ .

2. Solve the following two equations:

$$r^n = s$$

$$e^{in\theta} = e^{i\phi} \tag{1}$$

Continued

3. The solutions to $r^n = s$ are given by $r = \sqrt[n]{s}$.

Continued

3. The solutions to $r^n = s$ are given by $r = \sqrt[n]{s}$.
4. The solutions to $e^{in\theta} = e^{i\phi}$ are given by:

$$n\theta = \phi + 2\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

or

$$\theta = \frac{\phi}{n} + \frac{2}{n}\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

Continued

3. The solutions to $r^n = s$ are given by $r = \sqrt[n]{s}$.

4. The solutions to $e^{in\theta} = e^{i\phi}$ are given by:

$$n\theta = \phi + 2\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

or

$$\theta = \frac{\phi}{n} + \frac{2}{n}\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

5. Using the solutions r, θ to the equations given in (1) construct the n^{th} roots of the form $z = re^{i\theta}$.

Problem

Find all complex numbers z such that $z^4 = 2(\sqrt{3}i - 1)$, and express each in the form $a + bi$.

Problem

Find all complex numbers z such that $z^4 = 2(\sqrt{3}i - 1)$, and express each in the form $a + bi$.

Solution

1. Convert $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$ to polar form:

Problem

Find all complex numbers z such that $z^4 = 2(\sqrt{3}i - 1)$, and express each in the form $a + bi$.

Solution

1. Convert $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$ to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

Problem

Find all complex numbers z such that $z^4 = 2(\sqrt{3}i - 1)$, and express each in the form $a + bi$.

Solution

1. Convert $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$ to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If ϕ is an argument for $-2 + 2\sqrt{3}i$, then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \text{ and } \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \text{ so } \phi = \frac{2\pi}{3}.$$

Problem

Find all complex numbers z such that $z^4 = 2(\sqrt{3}i - 1)$, and express each in the form $a + bi$.

Solution

1. Convert $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$ to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If ϕ is an argument for $-2 + 2\sqrt{3}i$, then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \text{ and } \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \text{ so } \phi = \frac{2\pi}{3}.$$

Thus $z^4 = 4e^{2\pi i/3}$.

Problem

Find all complex numbers z such that $z^4 = 2(\sqrt{3}i - 1)$, and express each in the form $a + bi$.

Solution

1. Convert $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$ to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If ϕ is an argument for $-2 + 2\sqrt{3}i$, then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \text{ and } \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \text{ so } \phi = \frac{2\pi}{3}.$$

Thus $z^4 = 4e^{2\pi i/3}$. Let $z = re^{i\theta}$.

Problem

Find all complex numbers z such that $z^4 = 2(\sqrt{3}i - 1)$, and express each in the form $a + bi$.

Solution

1. Convert $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$ to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If ϕ is an argument for $-2 + 2\sqrt{3}i$, then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \text{ and } \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \text{ so } \phi = \frac{2\pi}{3}.$$

Thus $z^4 = 4e^{2\pi i/3}$. Let $z = re^{i\theta}$.

2. The equation becomes $r^4 e^{i4\theta} = 4e^{2\pi i/3}$, so we need to solve

$$\begin{aligned} r^4 &= 4 \\ e^{i4\theta} &= e^{2\pi i/3} \end{aligned}$$

Solution (continued)

3. Since $r^4 = 4$, $r^2 = \pm 2$. But r is **real**, and so $r^2 = 2$, implying $r = \pm\sqrt{2}$. However $r \geq 0$, and therefore $r = \sqrt{2}$.

Solution (continued)

3. Since $r^4 = 4$, $r^2 = \pm 2$. But r is **real**, and so $r^2 = 2$, implying $r = \pm\sqrt{2}$. However $r \geq 0$, and therefore $r = \sqrt{2}$.
4. The solutions to $e^{i4\theta} = e^{2\pi i/3}$ are given by

$$4\theta = \frac{2}{3}\pi + 2\pi\ell, \ell = 0, 1, 2, 3.$$

Solution (continued)

3. Since $r^4 = 4$, $r^2 = \pm 2$. But r is **real**, and so $r^2 = 2$, implying $r = \pm\sqrt{2}$. However $r \geq 0$, and therefore $r = \sqrt{2}$.
4. The solutions to $e^{i4\theta} = e^{2\pi i/3}$ are given by

$$4\theta = \frac{2}{3}\pi + 2\pi\ell, \ell = 0, 1, 2, 3.$$

Therefore,

$$\theta = \frac{2\pi}{12} + \frac{2\pi\ell}{4} = \frac{\pi}{6} + \frac{\pi\ell}{2} = \frac{\pi(3\ell + 1)}{6}, \text{ for } \ell = 0, 1, 2, 3.$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$.

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\ell = 0 : \quad z = \sqrt{2}e^{\pi i/6} = \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) = \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i\end{aligned}$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i\end{aligned}$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} &= \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) &= \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} = \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) = \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} = \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) = -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} = \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) = -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} = \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of $2(\sqrt{3}i - 1)$ are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i,$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} = \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) = \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} = \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) = -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} = \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) = -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} = \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of $2(\sqrt{3}i - 1)$ are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i, -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i,$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} = \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) = \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} = \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) = -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} = \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) = -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} = \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of $2(\sqrt{3}i - 1)$ are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i, -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i, -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i,$$

Solution (continued)

5. Thus $r = \sqrt{2}$ and $\theta = \left(\frac{3\ell+1}{6}\right)\pi$, $\ell = 0, 1, 2, 3$. Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} = \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) = \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} = \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) = -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} = \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) = -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} = \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of $2(\sqrt{3}i - 1)$ are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i, -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i, -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i, \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i.$$

Roots of Unity

Definition

A complex number z is a **root of unity** if there exists a positive integer n so that $z^n = 1$.

Roots of Unity

Definition

A complex number z is a **root of unity** if there exists a positive integer n so that $z^n = 1$.

Problem

Find the sixth roots of unity, i.e., all solutions to $z^6 = 1$.

Roots of Unity

Definition

A complex number z is a **root of unity** if there exists a positive integer n so that $z^n = 1$.

Problem

Find the sixth roots of unity, i.e., all solutions to $z^6 = 1$.

Solution

Write $z = re^{i\theta}$ and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

Roots of Unity

Definition

A complex number z is a **root of unity** if there exists a positive integer n so that $z^n = 1$.

Problem

Find the sixth roots of unity, i.e., all solutions to $z^6 = 1$.

Solution

Write $z = re^{i\theta}$ and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

Equating the absolute values and arguments,

$$r^6 = 1 \text{ and } 6\theta = 0 + 2\pi\ell, \ell = 0, 1, 2, 3, 4, 5.$$

Roots of Unity

Definition

A complex number z is a **root of unity** if there exists a positive integer n so that $z^n = 1$.

Problem

Find the sixth roots of unity, i.e., all solutions to $z^6 = 1$.

Solution

Write $z = re^{i\theta}$ and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

Equating the absolute values and arguments,

$$r^6 = 1 \text{ and } 6\theta = 0 + 2\pi\ell, \ell = 0, 1, 2, 3, 4, 5.$$

Since r is real, $r = 1$.

Roots of Unity

Definition

A complex number z is a **root of unity** if there exists a positive integer n so that $z^n = 1$.

Problem

Find the sixth roots of unity, i.e., all solutions to $z^6 = 1$.

Solution

Write $z = re^{i\theta}$ and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

Equating the absolute values and arguments,

$$r^6 = 1 \text{ and } 6\theta = 0 + 2\pi\ell, \ell = 0, 1, 2, 3, 4, 5.$$

Since r is real, $r = 1$. The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

ℓ	θ	z
0	0	$e^{0i} = 1$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

ℓ	θ	z
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

ℓ	θ	z
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

ℓ	θ	z
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	π	$e^{\pi i} = -1$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

ℓ	θ	z
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	π	$e^{\pi i} = -1$
4	$\frac{4\pi}{3}$	$e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

ℓ	θ	z
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	π	$e^{\pi i} = -1$
4	$\frac{4\pi}{3}$	$e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$
5	$\frac{5\pi}{3}$	$e^{5\pi i/3} = \frac{1}{2} - \frac{\sqrt{3}}{2}i$

Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

ℓ	θ	z
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	π	$e^{\pi i} = -1$
4	$\frac{4\pi}{3}$	$e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$
5	$\frac{5\pi}{3}$	$e^{5\pi i/3} = \frac{1}{2} - \frac{\sqrt{3}}{2}i$

If you graph these six points in the complex plane, you'll see that they result in six equally spaced points on the unit circle, one of them being $(1, 0)$.

Roots of Unity

For any integer $n \geq 1$, the (complex) solutions to $z^n = 1$ are

$$z = e^{2\pi\ell i/n} \text{ for } \ell = 0, 1, 2, \dots, n-1.$$

Roots of Unity

For any integer $n \geq 1$, the (complex) solutions to $z^n = 1$ are

$$z = e^{2\pi\ell i/n} \text{ for } \ell = 0, 1, 2, \dots, n-1.$$

Furthermore, the n^{th} roots of unity correspond to n equally spaced points on the unit circle, one of them being $(1, 0)$.