

A First Course in
LINEAR ALGEBRA

Lecture Notes
for Math 1503

**5.9: The General Solution to a Linear
System**

Creative Commons License (CC BY-NC-SA)

A First Course in Linear Algebra

Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Definition

Suppose a linear system of equations can be written in the form

$$T(\vec{x}) = \vec{b}$$

where T is a linear transformation determined by $T(\vec{x}) = A\vec{x}$.

Definition

Suppose a linear system of equations can be written in the form

$$T(\vec{x}) = \vec{b}$$

where T is a linear transformation determined by $T(\vec{x}) = A\vec{x}$.

If $T(\vec{x}_p) = \vec{b}$, then \vec{x}_p is called a **particular solution** of the linear system

Definition

Suppose a linear system of equations can be written in the form

$$T(\vec{x}) = \vec{b}$$

where T is a linear transformation determined by $T(\vec{x}) = A\vec{x}$.

If $T(\vec{x}_p) = \vec{b}$, then \vec{x}_p is called a **particular solution** of the linear system

Example

Suppose that a system of linear equations can be written as

$$T\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}\right) = \begin{bmatrix} 5 + 2x_2 - x_4 \\ x_2 \\ -4 - 3x_4 \\ x_4 \end{bmatrix}$$

Definition

Suppose a linear system of equations can be written in the form

$$T(\vec{x}) = \vec{b}$$

where T is a linear transformation determined by $T(\vec{x}) = A\vec{x}$.

If $T(\vec{x}_p) = \vec{b}$, then \vec{x}_p is called a **particular solution** of the linear system

Example

Suppose that a system of linear equations can be written as

$$T\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}\right) = \begin{bmatrix} 5 + 2x_2 - x_4 \\ x_2 \\ -4 - 3x_4 \\ x_4 \end{bmatrix}$$

Setting $x_2 = 2$ and $x_4 = -1$ gives us the **particular solution**

$$\vec{x}_p = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 10 \\ 2 \\ -1 \\ -1 \end{bmatrix}.$$

Null Space and Associated Homogeneous System

Definition

Let T be a linear transformation. Define:

$$\ker(T) = \left\{ \vec{x} : T(\vec{x}) = \vec{0} \right\}$$

Null Space and Associated Homogeneous System

Definition

Let T be a linear transformation. Define:

$$\ker(T) = \left\{ \vec{x} : T(\vec{x}) = \vec{0} \right\}$$

The **kernel**, $\ker(T)$ consists of the set of all vectors \vec{x} such that $T(\vec{x}) = \vec{0}$. This is also called the **null space** of T or the **solution space** of $T(\vec{x}) = \vec{0}$.

Null Space and Associated Homogeneous System

Definition

Let T be a linear transformation. Define:

$$\ker(T) = \left\{ \vec{x} : T(\vec{x}) = \vec{0} \right\}$$

The **kernel**, $\ker(T)$ consists of the set of all vectors \vec{x} such that $T(\vec{x}) = \vec{0}$. This is also called the **null space** of T or the **solution space** of $T(\vec{x}) = \vec{0}$.

Definition

Let $A\vec{x} = \vec{b}$ be a system of equations. Then, the corresponding system given by $A\vec{x} = \vec{0}$, found by replacing \vec{b} with $\vec{0}$ is the **associated homogeneous system**.

Associated Homogeneous System

Example

Consider the system given by

$$\begin{array}{rcccccccl} x_1 & + & x_2 & - & x_3 & + & 3x_4 & = & 2 \\ -x_1 & + & 4x_2 & + & 5x_3 & - & 2x_4 & = & 3 \\ x_1 & + & 6x_2 & + & 3x_3 & + & 4x_4 & = & 4 \end{array}$$

Associated Homogeneous System

Example

Consider the system given by

$$\begin{array}{rrrrrrrrcl} x_1 & + & x_2 & - & x_3 & + & 3x_4 & = & 2 \\ -x_1 & + & 4x_2 & + & 5x_3 & - & 2x_4 & = & 3 \\ x_1 & + & 6x_2 & + & 3x_3 & + & 4x_4 & = & 4 \end{array}$$

We can write this as

$$\begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$$

Associated Homogeneous System

Example

Consider the system given by

$$\begin{array}{rrrrrrrrcl} x_1 & + & x_2 & - & x_3 & + & 3x_4 & = & 2 \\ -x_1 & + & 4x_2 & + & 5x_3 & - & 2x_4 & = & 3 \\ x_1 & + & 6x_2 & + & 3x_3 & + & 4x_4 & = & 4 \end{array}$$

We can write this as

$$\begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}$$

whose **associated homogeneous system** can be written as:

$$\begin{bmatrix} 1 & 1 & -1 & 3 \\ -1 & 4 & 5 & -2 \\ 1 & 6 & 3 & 4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Solutions to Systems as Linear Combinations of Vectors

Solutions to Systems as Linear Combinations of Vectors

Basic Solutions to Homogeneous Systems

In our earlier treatment of homogeneous systems of linear equations, we saw how to write the solution to a homogeneous system of linear equations as a **linear combination** of **basic solutions**.

Solutions to Systems as Linear Combinations of Vectors

Basic Solutions to Homogeneous Systems

In our earlier treatment of homogeneous systems of linear equations, we saw how to write the solution to a homogeneous system of linear equations as a **linear combination** of **basic solutions**.

Example

The system

$$\begin{array}{rcccccccl} x_1 & + & x_2 & - & x_3 & + & 3x_4 & = & 0 \\ -x_1 & + & 4x_2 & + & 5x_3 & - & 2x_4 & = & 0 \\ x_1 & + & 6x_2 & + & 3x_3 & + & 4x_4 & = & 0 \end{array}$$

has general solution

Solutions to Systems as Linear Combinations of Vectors

Basic Solutions to Homogeneous Systems

In our earlier treatment of homogeneous systems of linear equations, we saw how to write the solution to a homogeneous system of linear equations as a **linear combination** of **basic solutions**.

Example

The system
$$\begin{array}{rcccccccl} x_1 & + & x_2 & - & x_3 & + & 3x_4 & = & 0 \\ -x_1 & + & 4x_2 & + & 5x_3 & - & 2x_4 & = & 0 \\ x_1 & + & 6x_2 & + & 3x_3 & + & 4x_4 & = & 0 \end{array}$$
 has general solution

$$x_1 = \frac{9}{5}s - \frac{14}{5}t$$

$$x_2 = -\frac{4}{5}s - \frac{1}{5}t$$

$$x_3 = s$$

$$x_4 = t$$

Solutions to Systems as Linear Combinations of Vectors

Basic Solutions to Homogeneous Systems

In our earlier treatment of homogeneous systems of linear equations, we saw how to write the solution to a homogeneous system of linear equations as a **linear combination** of **basic solutions**.

Example

The system
$$\begin{array}{rcccccccl} x_1 & + & x_2 & - & x_3 & + & 3x_4 & = & 0 \\ -x_1 & + & 4x_2 & + & 5x_3 & - & 2x_4 & = & 0 \\ x_1 & + & 6x_2 & + & 3x_3 & + & 4x_4 & = & 0 \end{array}$$
 has general solution

$$\begin{aligned} x_1 &= \frac{9}{5}s - \frac{14}{5}t \\ x_2 &= -\frac{4}{5}s - \frac{1}{5}t \\ x_3 &= s \\ x_4 &= t \end{aligned} \quad \text{or} \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = s \begin{bmatrix} \frac{9}{5} \\ -\frac{4}{5} \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -\frac{14}{5} \\ -\frac{1}{5} \\ 0 \\ 1 \end{bmatrix}.$$

The General Solution to a Linear System

Theorem

Let T be a linear transformation given by $T(\vec{x}) = A\vec{x}$. Suppose that \vec{x}_p is a particular solution of the system of linear equations given by $T(\vec{x}) = \vec{b}$, i.e., $T(\vec{x}_p) = \vec{b}$. Then if \vec{y} is any other solution of $T(\vec{x}) = \vec{b}$, it can be written in the form $\vec{y} = \vec{x}_p + \vec{x}_0$ for some (particular) solution \vec{x}_0 to the associated homogeneous system $A\vec{x} = 0$.

The General Solution to a Linear System

Theorem

Let T be a linear transformation given by $T(\vec{x}) = A\vec{x}$. Suppose that \vec{x}_p is a particular solution of the system of linear equations given by $T(\vec{x}) = \vec{b}$, i.e., $T(\vec{x}_p) = \vec{b}$. Then if \vec{y} is any other solution of $T(\vec{x}) = \vec{b}$, it can be written in the form $\vec{y} = \vec{x}_p + \vec{x}_0$ for some (particular) solution \vec{x}_0 to the associated homogeneous system $A\vec{x} = 0$.

The theorem implies that if \vec{x}_p is a particular solution to $A\vec{x} = \vec{b}$ and \vec{x}_h is the general solution to the associated homogeneous system $A\vec{x} = 0$, then $\vec{x}_p + \vec{x}_h$ is the general solution to $A\vec{x} = \vec{b}$.

The General Solution to a Linear System

Theorem

Let T be a linear transformation given by $T(\vec{x}) = A\vec{x}$. Suppose that \vec{x}_p is a particular solution of the system of linear equations given by $T(\vec{x}) = \vec{b}$, i.e., $T(\vec{x}_p) = \vec{b}$. Then if \vec{y} is any other solution of $T(\vec{x}) = \vec{b}$, it can be written in the form $\vec{y} = \vec{x}_p + \vec{x}_0$ for some (particular) solution \vec{x}_0 to the associated homogeneous system $A\vec{x} = 0$.

The theorem implies that if \vec{x}_p is a particular solution to $A\vec{x} = \vec{b}$ and \vec{x}_h is the general solution to the associated homogeneous system $A\vec{x} = 0$, then $\vec{x}_p + \vec{x}_h$ is the general solution to $A\vec{x} = \vec{b}$. Furthermore, the general solution to $A\vec{x} = \vec{b}$ can always be written in the form $\vec{x}_p + \vec{x}_h$ where \vec{x}_p is a particular solution to $A\vec{x} = \vec{b}$ and \vec{x}_h is the general solution to $A\vec{x} = 0$.

Example

The system of linear equations $A\vec{x} = \vec{b}$, with

$$A = \begin{bmatrix} 2 & 1 & 3 & 3 \\ 0 & 1 & -1 & 1 \\ -1 & 1 & -3 & 0 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

Example

The system of linear equations $A\vec{x} = \vec{b}$, with

$$A = \begin{bmatrix} 2 & 1 & 3 & 3 \\ 0 & 1 & -1 & 1 \\ -1 & 1 & -3 & 0 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

Notice that $\vec{x}_p = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix}$ is a particular solution to $A\vec{x} = \vec{b}$ (obtained by setting

$s = t = 0$),

Example

The system of linear equations $A\vec{x} = \vec{b}$, with

$$A = \begin{bmatrix} 2 & 1 & 3 & 3 \\ 0 & 1 & -1 & 1 \\ -1 & 1 & -3 & 0 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

Notice that $\vec{x}_p = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix}$ is a particular solution to $A\vec{x} = \vec{b}$ (obtained by setting

$s = t = 0$), while $\vec{x}_h = s \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \end{bmatrix}$, $s, t \in \mathbb{R}$ is the the general solution to the associated homogeneous system $A\vec{x} = 0$.

Example

The system of linear equations $A\vec{x} = \vec{b}$, with

$$A = \begin{bmatrix} 2 & 1 & 3 & 3 \\ 0 & 1 & -1 & 1 \\ -1 & 1 & -3 & 0 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

Notice that $\vec{x}_p = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix}$ is a particular solution to $A\vec{x} = \vec{b}$ (obtained by setting

$s = t = 0$), while $\vec{x}_h = s \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \end{bmatrix}$, $s, t \in \mathbb{R}$ is the the general solution

to the associated homogeneous system $A\vec{x} = 0$. Therefore the general solution is $\vec{y} = \vec{x}_p + \vec{x}_h$:

Example

The system of linear equations $A\vec{x} = \vec{b}$, with

$$A = \begin{bmatrix} 2 & 1 & 3 & 3 \\ 0 & 1 & -1 & 1 \\ -1 & 1 & -3 & 0 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

Notice that $\vec{x}_p = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix}$ is a particular solution to $A\vec{x} = \vec{b}$ (obtained by setting

$s = t = 0$), while $\vec{x}_h = s \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \end{bmatrix}$, $s, t \in \mathbb{R}$ is the the general solution

to the associated homogeneous system $A\vec{x} = 0$. Therefore the general solution is $\vec{y} = \vec{x}_p + \vec{x}_h$:

$$\vec{y} = \begin{bmatrix} 1 - 2s - t \\ 2 + s - t \\ s \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ -1 \\ 0 \\ 1 \end{bmatrix}, \quad s, t \in \mathbb{R}.$$