

A First Course in  
**LINEAR ALGEBRA**

**Lecture Notes**  
for Math 1503

$\mathbb{R}^n$ : Vectors

Creative Commons License (CC BY-NC-SA)

# A First Course in Linear Algebra

## Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

### License



**Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)**

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

# What is $\mathbb{R}^n$ ?

## Notation and Terminology

- $\mathbb{R}$  denotes the set of **real numbers**.
- $\mathbb{R}^2$  denotes the set of all **column vectors with two entries**.
- $\mathbb{R}^3$  denotes the set of all **column vectors with three entries**.
- In general,  $\mathbb{R}^n$  denotes the set of all **column vectors with  $n$  entries**.

# Scalar quantities versus vector quantities

- A **scalar** quantity has only magnitude; e.g. time, temperature.
- A (non-zero) **vector** quantity has both magnitude and direction; e.g. displacement, force, wind velocity.


# Scalar quantities versus vector quantities

- A **scalar** quantity has only magnitude; e.g. time, temperature.
- A (non-zero) **vector** quantity has both magnitude and direction; e.g. displacement, force, wind velocity.


Whereas two scalar quantities are equal if they are represented by the same value, two vector quantities are equal if and only if they have the same **magnitude** and **direction**.

## $\mathbb{R}^2$ and $\mathbb{R}^3$

Vectors in  $\mathbb{R}^2$  and  $\mathbb{R}^3$  have convenient geometric representations as **position vectors** of points in the 2-dimensional (Cartesian) plane and in 3-dimensional space, respectively.

$\mathbb{R}^2$ 

The vector  $\begin{bmatrix} a \\ b \end{bmatrix}$ .

 $\mathbb{R}^3$ 

The vector  $\begin{bmatrix} a \\ b \\ c \end{bmatrix}$ .

## Notation

- If  $P$  is a point in  $\mathbb{R}^n$  with coordinates  $(p_1, p_2, \dots, p_n)$  we denote this by  $P = (p_1, p_2, \dots, p_n)$ .



## Notation

- If  $P$  is a point in  $\mathbb{R}^n$  with coordinates  $(p_1, p_2, \dots, p_n)$  we denote this by  $P = (p_1, p_2, \dots, p_n)$ .
- If  $P = (p_1, p_2, \dots, p_n)$  is a point in  $\mathbb{R}^n$ , then

$$\overrightarrow{OP} = \begin{bmatrix} p_1 \\ p_2 \\ \vdots \\ p_n \end{bmatrix}$$

is often used to denote the position vector of the point.

## Notation

- If  $P$  is a point in  $\mathbb{R}^n$  with coordinates  $(p_1, p_2, \dots, p_n)$  we denote this by  $P = (p_1, p_2, \dots, p_n)$ .
- If  $P = (p_1, p_2, \dots, p_n)$  is a point in  $\mathbb{R}^n$ , then

$$\overrightarrow{OP} = \begin{bmatrix} p_1 \\ p_2 \\ \vdots \\ p_n \end{bmatrix}$$

is often used to denote the position vector of the point.

- Instead of using a capital letter to denote the vector (as we generally do with matrices), we emphasize the importance of the geometry and the direction with an arrow over the name of the vector.

## Notation and Terminology

- The notation  $\overrightarrow{0P}$  emphasizes that this vector goes from the origin  $0$  to the point  $P$ . We can also use lower case letters for names of vectors. In this case, we write  $\overrightarrow{0P} = \vec{p}$ .

## Notation and Terminology

- The notation  $\overrightarrow{0P}$  emphasizes that this vector goes from the origin 0 to the point  $P$ . We can also use lower case letters for names of vectors. In this case, we write  $\overrightarrow{0P} = \vec{p}$ .
- Any vector

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \text{ in } \mathbb{R}^n$$


is associated with the point  $(x_1, x_2, \dots, x_n)$ .

- Often, there is no distinction made between the vector  $\vec{x}$  and the point  $(x_1, x_2, \dots, x_n)$ , and we say that both  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$  and

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n.$$


# Geometric Vectors in $\mathbb{R}^2$ and $\mathbb{R}^3$

Let  $A$  and  $B$  be two points in  $\mathbb{R}^2$  or  $\mathbb{R}^3$ .



## Geometric Vectors in $\mathbb{R}^2$ and $\mathbb{R}^3$


Let  $A$  and  $B$  be two points in  $\mathbb{R}^2$  or  $\mathbb{R}^3$ .



•  $\overrightarrow{AB}$  is the **geometric vector** from  $A$  to  $B$ .


# Geometric Vectors in $\mathbb{R}^2$ and $\mathbb{R}^3$

Let  $A$  and  $B$  be two points in  $\mathbb{R}^2$  or  $\mathbb{R}^3$ .




- $\vec{AB}$  is the **geometric vector** from  $A$  to  $B$ .
- $A$  is the **tail** of  $\vec{AB}$ .
- $B$  is the **tip** of  $\vec{AB}$ .
- the **magnitude** of  $\vec{AB}$  is its length, and is denoted  $||\vec{AB}||$ .

## Equality of geometric vectors






## Equality of geometric vectors




- $\overrightarrow{AB}$  is the vector from  $A = (1, 0)$  to  $B = (2, 2)$ .
- $\overrightarrow{CD}$  is the vector from  $C = (-1, -1)$  to  $D = (0, 1)$ .

## Equality of geometric vectors



- $\overrightarrow{AB}$  is the vector from  $A = (1, 0)$  to  $B = (2, 2)$ .
- $\overrightarrow{CD}$  is the vector from  $C = (-1, -1)$  to  $D = (0, 1)$ .
- $\overrightarrow{AB} = \overrightarrow{CD}$  because the vectors have the same **length** and **direction**.


## Equality of geometric vectors



- $\overrightarrow{AB}$  is the vector from  $A = (1, 0)$  to  $B = (2, 2)$ .
- $\overrightarrow{CD}$  is the vector from  $C = (-1, -1)$  to  $D = (0, 1)$ .
- $\overrightarrow{AB} = \overrightarrow{CD}$  because the vectors have the same **length** and **direction**.


The fact that the points  $A$  and  $B$  are different from the points  $C$  and  $D$  is not important. For geometric vectors, **the location of the vector in the plane (or in 3-dimensional space) is not important**; the important properties are its length and direction.

## Coordinatizing Vectors – Part 1



$\vec{OP}$  is the **position vector** for  $P = (1, 2)$ ,  
and  $\vec{OP} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .


## Coordinatizing Vectors – Part 1



$\vec{OP}$  is the **position vector** for  $P = (1, 2)$ ,  
and  $\vec{OP} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .

Since  $\vec{AB} = \vec{OP}$ , **it should be the case that**  $\vec{AB} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ . This can be seen by moving  $\vec{AB}$  so that its tail is at the origin.

## Coordinatizing Vectors – Part 1



$\vec{OP}$  is the **position vector** for  $P = (1, 2)$ ,  
and  $\vec{OP} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .

Since  $\vec{AB} = \vec{OP}$ , **it should be the case that**  $\vec{AB} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ . This can be seen by moving  $\vec{AB}$  so that its tail is at the origin.

A geometric vector is coordinatized by putting it in **standard position**, meaning with its tail at the origin, and then identifying the vector with its tip.

# Algebra in $\mathbb{R}^n$

## Addition in $\mathbb{R}^n$

Since vectors in  $\mathbb{R}^n$  are  $n \times 1$  matrices, addition in  $\mathbb{R}^n$  is precisely matrix addition using column matrices, i.e.,

- If  $\vec{u}$  and  $\vec{v}$  are in  $\mathbb{R}^n$ , then  $\vec{u} + \vec{v}$  is obtained by adding together corresponding entries of the vectors.
- The zero vector in  $\mathbb{R}^n$  is the  $n \times 1$  zero matrix, and is denoted  $\vec{0}$ .

# Algebra in $\mathbb{R}^n$

## Addition in $\mathbb{R}^n$

Since vectors in  $\mathbb{R}^n$  are  $n \times 1$  matrices, addition in  $\mathbb{R}^n$  is precisely matrix addition using column matrices, i.e.,

- If  $\vec{u}$  and  $\vec{v}$  are in  $\mathbb{R}^n$ , then  $\vec{u} + \vec{v}$  is obtained by adding together corresponding entries of the vectors.
- The zero vector in  $\mathbb{R}^n$  is the  $n \times 1$  zero matrix, and is denoted  $\vec{0}$ .

## Example

Let  $\vec{u} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$ . Then,

$$\vec{u} + \vec{v} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 5 \\ 7 \\ 9 \end{bmatrix}$$



# Properties of Vector Addition

Let  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  be vectors in  $\mathbb{R}^n$ . Then the following properties hold.

# Properties of Vector Addition

Let  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  be vectors in  $\mathbb{R}^n$ . Then the following properties hold.

①  $\vec{u} + \vec{v} = \vec{v} + \vec{u}$  (vector addition is commutative).

# Properties of Vector Addition

Let  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  be vectors in  $\mathbb{R}^n$ . Then the following properties hold.

- ❶  $\vec{u} + \vec{v} = \vec{v} + \vec{u}$  (vector addition is commutative).
- ❷  $(\vec{u} + \vec{v}) + \vec{w} = \vec{u} + (\vec{v} + \vec{w})$  (vector addition is associative).

# Properties of Vector Addition

Let  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  be vectors in  $\mathbb{R}^n$ . Then the following properties hold.

- ①  $\vec{u} + \vec{v} = \vec{v} + \vec{u}$  (vector addition is commutative).
- ②  $(\vec{u} + \vec{v}) + \vec{w} = \vec{u} + (\vec{v} + \vec{w})$  (vector addition is associative).
- ③  $\vec{u} + \vec{0} = \vec{u}$  (existence of an additive identity).

# Properties of Vector Addition

Let  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{w}$  be vectors in  $\mathbb{R}^n$ . Then the following properties hold.

- ①  $\vec{u} + \vec{v} = \vec{v} + \vec{u}$  (vector addition is commutative).
- ②  $(\vec{u} + \vec{v}) + \vec{w} = \vec{u} + (\vec{v} + \vec{w})$  (vector addition is associative).
- ③  $\vec{u} + \vec{0} = \vec{u}$  (existence of an additive identity).
- ④  $\vec{u} + (-\vec{u}) = \vec{0}$  (existence of an additive inverse).

# Scalar Multiplication

Since vectors in  $\mathbb{R}^n$  are  $n \times 1$  matrices, scalar multiplication in  $\mathbb{R}^n$  is precisely matrix scalar multiplication using column matrices, i.e., If  $\vec{u}$  is a vector in  $\mathbb{R}^n$  and  $k \in \mathbb{R}$  is a scalar, then  $k\vec{u}$  is obtained by multiplying every entry of  $\vec{u}$  by  $k$ .

# Scalar Multiplication

Since vectors in  $\mathbb{R}^n$  are  $n \times 1$  matrices, scalar multiplication in  $\mathbb{R}^n$  is precisely matrix scalar multiplication using column matrices, i.e., If  $\vec{u}$  is a vector in  $\mathbb{R}^n$  and  $k \in \mathbb{R}$  is a scalar, then  $k\vec{u}$  is obtained by multiplying every entry of  $\vec{u}$  by  $k$ .

## Example

Let  $\vec{u} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  and  $k = 4$ . Then,

$$k\vec{u} = 4 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 4 \\ 8 \\ 12 \end{bmatrix}$$

# Properties of Scalar Multiplication

Let  $\vec{u}, \vec{v} \in \mathbb{R}^n$  be vectors and  $k, p \in \mathbb{R}$  be scalars. Then the following properties hold.



# Properties of Scalar Multiplication

Let  $\vec{u}, \vec{v} \in \mathbb{R}^n$  be vectors and  $k, p \in \mathbb{R}$  be scalars. Then the following properties hold.

①  $k(\vec{u} + \vec{v}) = k\vec{u} + k\vec{v}$  (scalar multiplication distributes over vector addition).

# Properties of Scalar Multiplication

Let  $\vec{u}, \vec{v} \in \mathbb{R}^n$  be vectors and  $k, p \in \mathbb{R}$  be scalars. Then the following properties hold.

- ①  $k(\vec{u} + \vec{v}) = k\vec{u} + k\vec{v}$  (scalar multiplication distributes over vector addition).
- ②  $(k + p)\vec{u} = k\vec{u} + p\vec{u}$  (addition distributes over scalar multiplication).

# Properties of Scalar Multiplication

Let  $\vec{u}, \vec{v} \in \mathbb{R}^n$  be vectors and  $k, p \in \mathbb{R}$  be scalars. Then the following properties hold.

- ①  $k(\vec{u} + \vec{v}) = k\vec{u} + k\vec{v}$  (scalar multiplication distributes over vector addition).
- ②  $(k + p)\vec{u} = k\vec{u} + p\vec{u}$  (addition distributes over scalar multiplication).
- ③  $k(p\vec{u}) = (kp)\vec{u}$  (scalar multiplication is associative).

# Properties of Scalar Multiplication

Let  $\vec{u}, \vec{v} \in \mathbb{R}^n$  be vectors and  $k, p \in \mathbb{R}$  be scalars. Then the following properties hold.

- ①  $k(\vec{u} + \vec{v}) = k\vec{u} + k\vec{v}$  (scalar multiplication distributes over vector addition).
- ②  $(k + p)\vec{u} = k\vec{u} + p\vec{u}$  (addition distributes over scalar multiplication).
- ③  $k(p\vec{u}) = (kp)\vec{u}$  (scalar multiplication is associative).
- ④  $1\vec{u} = \vec{u}$  (existence of a multiplicative identity).

## Some notation you may encounter

Often, in  $\mathbb{R}^2$  the notation  $\vec{i} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $\vec{j} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$  is used. Whereas in  $\mathbb{R}^3$  the notation is  $\vec{i} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ ,  $\vec{j} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ ,  $\vec{k} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

So we have

$$\begin{bmatrix} a \\ b \end{bmatrix} = a\vec{i} + b\vec{j}$$

and

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} = a\vec{i} + b\vec{j} + c\vec{k}$$

# The Geometry of Vector Addition


- 1 **Vector Equality.** The vectors have the same length and direction.

# The Geometry of Vector Addition

- 1 **Vector Equality.** The vectors have the same length and direction.
- 2 The zero vector,  $\vec{0}$  has length zero and **no direction**.

# The Geometry of Vector Addition

- ① **Vector Equality.** The vectors have the same length and direction.
- ② **The zero vector,  $\vec{0}$**  has length zero and **no direction**.
- ③ **Addition.** Let  $\vec{u}, \vec{v}$  be vectors. Then  $\vec{u} + \vec{v}$  is the diagonal of the parallelogram defined by  $\vec{u}$  and  $\vec{v}$ , and having the same tail as  $\vec{u}$  and  $\vec{v}$ .






## Tip-to-Tail Method for Vector Addition

For points  $A$ ,  $B$  and  $C$ ,

$$\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}.$$




## Example

The diagonals of any parallelogram bisect each other.

## Example

The diagonals of any parallelogram bisect each other.


To see this, denote the parallelogram by its vertices,  $ABCD$ .



## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .




- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .




- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .




- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .




- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .



- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .


$$\overrightarrow{BM} = \overrightarrow{BA} + \overrightarrow{AM}$$



## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .




- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .

$$\overrightarrow{BM} = \overrightarrow{BA} + \overrightarrow{AM} = \overrightarrow{CD} + \overrightarrow{MC}$$

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .




- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .

$$\overrightarrow{BM} = \overrightarrow{BA} + \overrightarrow{AM} = \overrightarrow{CD} + \overrightarrow{MC} = \overrightarrow{MC} + \overrightarrow{CD}$$

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .




- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .

$$\overrightarrow{BM} = \overrightarrow{BA} + \overrightarrow{AM} = \overrightarrow{CD} + \overrightarrow{MC} = \overrightarrow{MC} + \overrightarrow{CD} = \overrightarrow{MD}.$$

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .



- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .


$$\overrightarrow{BM} = \overrightarrow{BA} + \overrightarrow{AM} = \overrightarrow{CD} + \overrightarrow{MC} = \overrightarrow{MC} + \overrightarrow{CD} = \overrightarrow{MD}.$$

Since  $\overrightarrow{BM} = \overrightarrow{MD}$ , these vectors have the same **magnitude** and **direction**,

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .



- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .


$$\overrightarrow{BM} = \overrightarrow{BA} + \overrightarrow{AM} = \overrightarrow{CD} + \overrightarrow{MC} = \overrightarrow{MC} + \overrightarrow{CD} = \overrightarrow{MD}.$$

Since  $\overrightarrow{BM} = \overrightarrow{MD}$ , these vectors have the same **magnitude** and **direction**, implying that  $M$  is the midpoint of  $\overrightarrow{BD}$ .

## Example

The diagonals of any parallelogram bisect each other.

To see this, denote the parallelogram by its vertices,  $ABCD$ .



- Let  $M$  denote the midpoint of  $\overrightarrow{AC}$ .  
Then  $\overrightarrow{AM} = \overrightarrow{MC}$ .
- It now suffices to show that  $\overrightarrow{BM} = \overrightarrow{MD}$ .


$$\overrightarrow{BM} = \overrightarrow{BA} + \overrightarrow{AM} = \overrightarrow{CD} + \overrightarrow{MC} = \overrightarrow{MC} + \overrightarrow{CD} = \overrightarrow{MD}.$$

Since  $\overrightarrow{BM} = \overrightarrow{MD}$ , these vectors have the same **magnitude** and **direction**, implying that  $M$  is the midpoint of  $\overrightarrow{BD}$ .

Therefore, the diagonals of  $ABCD$  bisect each other.


## The Geometry of Vector Subtraction

Let  $\vec{u}$  and  $\vec{v}$  be vectors in  $\mathbb{R}^2$  or  $\mathbb{R}^3$ . The vector  $\vec{u} - \vec{v} = \vec{u} + (-\vec{v})$  is obtained from the parallelogram defined by  $\vec{u}$  and  $\vec{v}$  by taking the vector from the tip of  $\vec{v}$  to the tip of  $\vec{u}$ , i.e., the diagonal of the parallelogram, directed towards the tip of  $\vec{u}$ .



## Coordinatizing Vectors – Part 2


Let  $A = (x_1, y_1, z_1)$  and  $B = (x_2, y_2, z_2)$  be two points in  $\mathbb{R}^3$ .





## Coordinatizing Vectors – Part 2

Let  $A = (x_1, y_1, z_1)$  and  $B = (x_2, y_2, z_2)$  be two points in  $\mathbb{R}^3$ .




We see from the figure that  $\vec{0A} + \vec{AB} = \vec{0B}$ , and hence

$$\vec{AB} = \vec{0B} - \vec{0A} = \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} - \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix} = \begin{bmatrix} x_2 - x_1 \\ y_2 - y_1 \\ z_2 - z_1 \end{bmatrix}.$$

## Length of a Vector, $\mathbb{R}^2$

$$\text{If } \vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^2,$$



then the length of the vector  $\vec{x}$  is the distance from the origin 0 to the point  $X = (x_1, x_2)$  given by  $d(0, X)$ .

The length of  $\vec{x}$ , denoted  $\|\vec{x}\|$ , is given by:

$$d(0, X) = \|\vec{x}\| = \sqrt{x_1^2 + x_2^2}.$$

## Length of a Vector, $\mathbb{R}^3$

This extends clearly to  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \in \mathbb{R}^3$ .

The length of  $\vec{x}$  is the distance from the origin 0 to the point  $X = (x_1, x_2, x_3)$  given by  $d(0, X)$ .

$$d(0, X) = \|\vec{x}\| = \sqrt{x_1^2 + x_2^2 + x_3^2}.$$

## Length of a Vector, $\mathbb{R}^3$

This extends clearly to  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \in \mathbb{R}^3$ .

The length of  $\vec{x}$  is the distance from the origin 0 to the point  $X = (x_1, x_2, x_3)$  given by  $d(0, X)$ .

$$d(0, X) = \|\vec{x}\| = \sqrt{x_1^2 + x_2^2 + x_3^2}.$$

Suppose we want to find the distance between points other than the origin?

## Length of a Vector, $\mathbb{R}^3$

Consider two arbitrary points in  $\mathbb{R}^3$ ,  $A = (x_1, y_1, z_1)$  and  $B = (x_2, y_2, z_2)$ . Then the distance between them is written  $d(A, B)$  and is given by the **distance formula**.

### Distance Formula

$$d(A, B) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$

## Length of a Vector, $\mathbb{R}^3$

Consider two arbitrary points in  $\mathbb{R}^3$ ,  $A = (x_1, y_1, z_1)$  and  $B = (x_2, y_2, z_2)$ . Then the distance between them is written  $d(A, B)$  and is given by the **distance formula**.

### Distance Formula

$$d(A, B) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$

Now let  $P = (x_2 - x_1, y_2 - y_1, z_2 - z_1)$ , and

$$\vec{p} = \begin{bmatrix} x_2 - x_1 \\ y_2 - y_1 \\ z_2 - z_1 \end{bmatrix}$$

## Length of a Vector, $\mathbb{R}^3$

Consider two arbitrary points in  $\mathbb{R}^3$ ,  $A = (x_1, y_1, z_1)$  and  $B = (x_2, y_2, z_2)$ . Then the distance between them is written  $d(A, B)$  and is given by the **distance formula**.

### Distance Formula

$$d(A, B) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$

Now let  $P = (x_2 - x_1, y_2 - y_1, z_2 - z_1)$ , and

$$\vec{p} = \begin{bmatrix} x_2 - x_1 \\ y_2 - y_1 \\ z_2 - z_1 \end{bmatrix}$$

Then the length of  $\vec{p}$  is equal to the distance between the origin and  $P$ , which are both equal to the distance between points  $A$  and  $B$

## Length of a Vector, $\mathbb{R}^3$

Consider two arbitrary points in  $\mathbb{R}^3$ ,  $A = (x_1, y_1, z_1)$  and  $B = (x_2, y_2, z_2)$ . Then the distance between them is written  $d(A, B)$  and is given by the **distance formula**.

### Distance Formula

$$d(A, B) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$

Now let  $P = (x_2 - x_1, y_2 - y_1, z_2 - z_1)$ , and

$$\vec{p} = \begin{bmatrix} x_2 - x_1 \\ y_2 - y_1 \\ z_2 - z_1 \end{bmatrix}$$

Then the length of  $\vec{p}$  is equal to the distance between the origin and  $P$ , which are both equal to the distance between points  $A$  and  $B$

$$\|\vec{p}\| = d(0, P) = d(A, B) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$



## Length of a Vector, $\mathbb{R}^n$

More generally, if  $P = (p_1, p_2, \dots, p_n)$  and  $Q = (q_1, q_2, \dots, q_n)$  are points in  $\mathbb{R}^n$ , then **the distance between  $P$  and  $Q$**  is the length of the vector  $\overrightarrow{PQ}$ , written  $\|\overrightarrow{PQ}\|$ .

$$d(P, Q) = \|\overrightarrow{PQ}\| = \sqrt{(q_1 - p_1)^2 + (q_2 - p_2)^2 + \dots + (q_n - p_n)^2}.$$

## Length of a Vector, $\mathbb{R}^n$

More generally, if  $P = (p_1, p_2, \dots, p_n)$  and  $Q = (q_1, q_2, \dots, q_n)$  are points in  $\mathbb{R}^n$ , then **the distance between  $P$  and  $Q$**  is the length of the vector  $\overrightarrow{PQ}$ , written  $\|\overrightarrow{PQ}\|$ .

$$d(P, Q) = \|\overrightarrow{PQ}\| = \sqrt{(q_1 - p_1)^2 + (q_2 - p_2)^2 + \dots + (q_n - p_n)^2}.$$

The formula for calculating the length of a vector generalizes to  $\mathbb{R}^n$ : if

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n,$$

then

$$\|\vec{x}\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2},$$

which represents the distance from the origin to the point  $(x_1, x_2, \dots, x_n)$ .

# Properties of Distance

Let  $P$  and  $Q$  be two points in  $\mathbb{R}^n$ , and  $d(P, Q)$  the distance between them. Then the following properties hold.

# Properties of Distance

Let  $P$  and  $Q$  be two points in  $\mathbb{R}^n$ , and  $d(P, Q)$  the distance between them. Then the following properties hold.

- 1 The distance between  $P$  and  $Q$  is equal to the distance between  $Q$  and  $P$ , i.e.,  $d(P, Q) = d(Q, P)$ .

# Properties of Distance

Let  $P$  and  $Q$  be two points in  $\mathbb{R}^n$ , and  $d(P, Q)$  the distance between them. Then the following properties hold.

- 1 The distance between  $P$  and  $Q$  is equal to the distance between  $Q$  and  $P$ , i.e.,  $d(P, Q) = d(Q, P)$ .
- 2  $d(P, Q) \geq 0$  with equality if and only if  $P = Q$ .

# Properties of Distance

Let  $P$  and  $Q$  be two points in  $\mathbb{R}^n$ , and  $d(P, Q)$  the distance between them. Then the following properties hold.

- 1 The distance between  $P$  and  $Q$  is equal to the distance between  $Q$  and  $P$ , i.e.,  $d(P, Q) = d(Q, P)$ .
- 2  $d(P, Q) \geq 0$  with equality if and only if  $P = Q$ .

## Example

For  $P = (1, -1, 3)$  and  $Q = (3, 1, 0)$ , the distance between  $P$  and  $Q$  is  $d(P, Q) = \sqrt{2^2 + 2^2 + (-3)^2} = \sqrt{17}$ .

## Example

Let  $\vec{p} = \begin{bmatrix} -3 \\ 4 \end{bmatrix}$  and  $\vec{q} = \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}$ . Then  $-2\vec{q} = (-2)\vec{q} = \begin{bmatrix} -6 \\ 2 \\ 4 \end{bmatrix}$ .

## Example

Let  $\vec{p} = \begin{bmatrix} -3 \\ 4 \end{bmatrix}$  and  $\vec{q} = \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}$ . Then  $-2\vec{q} = (-2)\vec{q} = \begin{bmatrix} -6 \\ 2 \\ 4 \end{bmatrix}$ .

The lengths of these vectors are

$$\|\vec{p}\| = \sqrt{(-3)^2 + 4^2} = \sqrt{9 + 16} = 5,$$



## Example

Let  $\vec{p} = \begin{bmatrix} -3 \\ 4 \end{bmatrix}$  and  $\vec{q} = \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}$ . Then  $-2\vec{q} = (-2)\vec{q} = \begin{bmatrix} -6 \\ 2 \\ 4 \end{bmatrix}$ .

The lengths of these vectors are

$$\|\vec{p}\| = \sqrt{(-3)^2 + 4^2} = \sqrt{9 + 16} = 5,$$

$$\|\vec{q}\| = \sqrt{(3)^2 + (-1)^2 + (-2)^2} = \sqrt{9 + 1 + 4} = \sqrt{14},$$

## Example

Let  $\vec{p} = \begin{bmatrix} -3 \\ 4 \end{bmatrix}$  and  $\vec{q} = \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}$ . Then  $-2\vec{q} = (-2)\vec{q} = \begin{bmatrix} -6 \\ 2 \\ 4 \end{bmatrix}$ .

The lengths of these vectors are

$$\|\vec{p}\| = \sqrt{(-3)^2 + 4^2} = \sqrt{9 + 16} = 5,$$

$$\|\vec{q}\| = \sqrt{(3)^2 + (-1)^2 + (-2)^2} = \sqrt{9 + 1 + 4} = \sqrt{14},$$

and

$$\begin{aligned} \|-2\vec{q}\| &= \sqrt{(-6)^2 + 2^2 + 4^2} \\ &= \sqrt{36 + 4 + 16} \\ &= \sqrt{56} = \sqrt{4 \times 14} \\ &= 2\sqrt{14} = 2\|\vec{q}\|. \end{aligned}$$

# The Geometry of Scalar Multiplication

- **Scalar Multiplication.** If  $\vec{v} \neq \vec{0}$  and  $a \in \mathbb{R}$ ,  $a \neq 0$ , then  $a\vec{v}$  has length  $\|a\vec{v}\| = |a| \cdot \|\vec{v}\|$ , and

# The Geometry of Scalar Multiplication

- **Scalar Multiplication.** If  $\vec{v} \neq \vec{0}$  and  $a \in \mathbb{R}$ ,  $a \neq 0$ , then  $a\vec{v}$  has length  $\|a\vec{v}\| = |a| \cdot \|\vec{v}\|$ , and
  - ▶ has the same direction as  $\vec{v}$  if  $a > 0$ ;

# The Geometry of Scalar Multiplication

- **Scalar Multiplication.** If  $\vec{v} \neq \vec{0}$  and  $a \in \mathbb{R}$ ,  $a \neq 0$ , then  $a\vec{v}$  has length  $\|a\vec{v}\| = |a| \cdot \|\vec{v}\|$ , and
  - ▶ has the same direction as  $\vec{v}$  if  $a > 0$ ;
  - ▶ has direction opposite to  $\vec{v}$  if  $a < 0$ .

# The Geometry of Scalar Multiplication

- **Scalar Multiplication.** If  $\vec{v} \neq \vec{0}$  and  $a \in \mathbb{R}$ ,  $a \neq 0$ , then  $a\vec{v}$  has length  $\|a\vec{v}\| = |a| \cdot \|\vec{v}\|$ , and
  - ▶ has the same direction as  $\vec{v}$  if  $a > 0$ ;
  - ▶ has direction opposite to  $\vec{v}$  if  $a < 0$ .
- **Parallel Vectors.** Two nonzero vectors are called **parallel** if they have the same direction or opposite directions.

# The Geometry of Scalar Multiplication

- **Scalar Multiplication.** If  $\vec{v} \neq \vec{0}$  and  $a \in \mathbb{R}$ ,  $a \neq 0$ , then  $a\vec{v}$  has length  $\|a\vec{v}\| = |a| \cdot \|\vec{v}\|$ , and
  - ▶ has the same direction as  $\vec{v}$  if  $a > 0$ ;
  - ▶ has direction opposite to  $\vec{v}$  if  $a < 0$ .
- **Parallel Vectors.** Two nonzero vectors are called **parallel** if they have the same direction or opposite directions. It follows that nonzero vectors  $\vec{v}$  and  $\vec{w}$  are parallel if and only if one is a scalar multiple of the other.

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?



## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix},$$

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}, \overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix},$$

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ ,

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$$

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

This gives a system of three equations in one variable,

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

This gives a system of three equations in one variable, which is consistent, and has unique solution  $k = -2$ .

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

This gives a system of three equations in one variable, which is consistent, and has unique solution  $k = -2$ . Therefore,  $\overrightarrow{PQ}$  is parallel to  $\overrightarrow{XY}$ .



## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

This gives a system of three equations in one variable, which is consistent, and has unique solution  $k = -2$ . Therefore,  $\overrightarrow{PQ}$  is parallel to  $\overrightarrow{XY}$ .

$$\overrightarrow{PX} = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix},$$

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

This gives a system of three equations in one variable, which is consistent, and has unique solution  $k = -2$ . Therefore,  $\overrightarrow{PQ}$  is parallel to  $\overrightarrow{XY}$ .

$$\overrightarrow{PX} = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix}, \quad \overrightarrow{QY} = \begin{bmatrix} 7 \\ -2 \\ -2 \end{bmatrix},$$

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

This gives a system of three equations in one variable, which is consistent, and has unique solution  $k = -2$ . Therefore,  $\overrightarrow{PQ}$  is parallel to  $\overrightarrow{XY}$ .

$\overrightarrow{PX} = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{QY} = \begin{bmatrix} 7 \\ -2 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PX} = \ell\overrightarrow{QY}$  for some scalar  $\ell$ .

## Problem

Let  $P = (1, -2, 1)$ ,  $Q = (-3, 0, 5)$ ,  $X = (2, -1, 5)$  and  $Y = (4, -2, 3)$  be points in  $\mathbb{R}^3$ . Is  $\overrightarrow{PQ}$  parallel to  $\overrightarrow{XY}$ ? Is  $\overrightarrow{PX}$  parallel to  $\overrightarrow{QY}$ ?

## Solution

$\overrightarrow{PQ} = \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{XY} = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PQ} = k\overrightarrow{XY}$  for some scalar  $k$ , i.e.,

$$\begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = k \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -4 \\ 2 \\ 4 \end{bmatrix} = \begin{bmatrix} 2k \\ -k \\ -2k \end{bmatrix}.$$

This gives a system of three equations in one variable, which is consistent, and has unique solution  $k = -2$ . Therefore,  $\overrightarrow{PQ}$  is parallel to  $\overrightarrow{XY}$ .

$\overrightarrow{PX} = \begin{bmatrix} 1 \\ 1 \\ 4 \end{bmatrix}$ ,  $\overrightarrow{QY} = \begin{bmatrix} 7 \\ -2 \\ -2 \end{bmatrix}$ , and these vectors are parallel if  $\overrightarrow{PX} = \ell\overrightarrow{QY}$  for some scalar  $\ell$ . You will find that no such  $\ell$  exists, so  $\overrightarrow{PX}$  is not parallel to  $\overrightarrow{QY}$ .

# Unit Vectors

## Definition

A **unit vector** is a vector of length one.

# Unit Vectors

## Definition

A **unit vector** is a vector of length one.

## Example

$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} \frac{\sqrt{2}}{2} \\ 0 \\ \frac{\sqrt{2}}{2} \end{bmatrix}$ , are examples of unit vectors.

# Unit Vectors

## Definition

A **unit vector** is a vector of length one.

## Example

$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ ,  $\begin{bmatrix} \frac{\sqrt{2}}{2} \\ 0 \\ \frac{\sqrt{2}}{2} \end{bmatrix}$ , are examples of unit vectors.

## Example

If  $\vec{v} \neq \vec{0}$ , then

$$\frac{1}{\|\vec{v}\|} \vec{v}$$

is a unit vector in the same direction as  $\vec{v}$ .

## Example

$\vec{v} = \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}$  is not a unit vector, since  $\|\vec{v}\| = \sqrt{14}$ .



## Example

$\vec{v} = \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}$  is not a unit vector, since  $\|\vec{v}\| = \sqrt{14}$ . However,

$$\vec{u} = \frac{1}{\sqrt{14}} \vec{v} = \begin{bmatrix} \frac{-1}{\sqrt{14}} \\ \frac{3}{\sqrt{14}} \\ \frac{2}{\sqrt{14}} \end{bmatrix}$$

is a unit vector **in the same direction** as  $\vec{v}$ ,

## Example

$\vec{v} = \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}$  is not a unit vector, since  $\|\vec{v}\| = \sqrt{14}$ . However,

$$\vec{u} = \frac{1}{\sqrt{14}} \vec{v} = \begin{bmatrix} \frac{-1}{\sqrt{14}} \\ \frac{3}{\sqrt{14}} \\ \frac{2}{\sqrt{14}} \end{bmatrix}$$

is a unit vector **in the same direction** as  $\vec{v}$ , i.e.,

$$\|\vec{u}\| = \frac{1}{\sqrt{14}} \|\vec{v}\| = \frac{1}{\sqrt{14}} \sqrt{14} = 1.$$

## Example

$\vec{v} = \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}$  is not a unit vector, since  $\|\vec{v}\| = \sqrt{14}$ . However,

$$\vec{u} = \frac{1}{\sqrt{14}} \vec{v} = \begin{bmatrix} \frac{-1}{\sqrt{14}} \\ \frac{3}{\sqrt{14}} \\ \frac{2}{\sqrt{14}} \end{bmatrix}$$

is a unit vector **in the same direction** as  $\vec{v}$ , i.e.,

$$\|\vec{u}\| = \frac{1}{\sqrt{14}} \|\vec{v}\| = \frac{1}{\sqrt{14}} \sqrt{14} = 1.$$

## Example

If  $\vec{v}$  and  $\vec{w}$  are nonzero that have

- the same direction, then  $\vec{v} = \frac{\|\vec{v}\|}{\|\vec{w}\|} \vec{w}$ ;

## Example

$\vec{v} = \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}$  is not a unit vector, since  $\|\vec{v}\| = \sqrt{14}$ . However,

$$\vec{u} = \frac{1}{\sqrt{14}} \vec{v} = \begin{bmatrix} \frac{-1}{\sqrt{14}} \\ \frac{3}{\sqrt{14}} \\ \frac{2}{\sqrt{14}} \end{bmatrix}$$

is a unit vector **in the same direction** as  $\vec{v}$ , i.e.,

$$\|\vec{u}\| = \frac{1}{\sqrt{14}} \|\vec{v}\| = \frac{1}{\sqrt{14}} \sqrt{14} = 1.$$

## Example

If  $\vec{v}$  and  $\vec{w}$  are nonzero that have

- the same direction, then  $\vec{v} = \frac{\|\vec{v}\|}{\|\vec{w}\|} \vec{w}$ ;
- opposite directions, then  $\vec{v} = -\frac{\|\vec{v}\|}{\|\vec{w}\|} \vec{w}$ .

# Vector problems and examples

## Problem


Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution




# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution




$\vec{OM}$

# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution



$$\overrightarrow{OM} = \overrightarrow{OP_1} + \overrightarrow{P_1M}$$




# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution




$$\overrightarrow{OM} = \overrightarrow{OP_1} + \overrightarrow{P_1M} = \overrightarrow{OP_1} + \frac{1}{2}\overrightarrow{P_1P_2}$$

# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution




$$\overrightarrow{OM} = \overrightarrow{OP_1} + \overrightarrow{P_1M} = \overrightarrow{OP_1} + \frac{1}{2}\overrightarrow{P_1P_2} = \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 6 \\ 4 \\ -6 \end{bmatrix}$$

# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution




$$\begin{aligned}\vec{OM} &= \vec{OP_1} + \vec{P_1M} = \vec{OP_1} + \frac{1}{2}\vec{P_1P_2} = \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 6 \\ 4 \\ -6 \end{bmatrix} \\ &= \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix} + \begin{bmatrix} 3 \\ 2 \\ -3 \end{bmatrix}\end{aligned}$$

# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution




$$\begin{aligned}\vec{OM} &= \vec{OP_1} + \vec{P_1M} = \vec{OP_1} + \frac{1}{2}\vec{P_1P_2} = \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 6 \\ 4 \\ -6 \end{bmatrix} \\ &= \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix} + \begin{bmatrix} 3 \\ 2 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 0 \end{bmatrix}.\end{aligned}$$

# Vector problems and examples

## Problem

Find the point,  $M$ , that is midway between  $P_1 = (-1, -4, 3)$  and  $P_2 = (5, 0, -3)$ .

## Solution



$$\begin{aligned}\vec{OM} &= \vec{OP_1} + \vec{P_1M} = \vec{OP_1} + \frac{1}{2}\vec{P_1P_2} = \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 6 \\ 4 \\ -6 \end{bmatrix} \\ &= \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix} + \begin{bmatrix} 3 \\ 2 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 0 \end{bmatrix}.\end{aligned}$$

Therefore  $M = (2, -2, 0)$ .


## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .

## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .


## Solution



## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .

## Solution




$$\bullet \vec{OA} = \vec{OP} + \frac{1}{3}\vec{PQ}$$



## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .

## Solution




$$\bullet \vec{OA} = \vec{OP} + \frac{1}{3}\vec{PQ}$$

$$\bullet \vec{OB} = \vec{OP} + \frac{2}{3}\vec{PQ}$$

## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .

## Solution




Since  $\vec{PQ} = \begin{bmatrix} 6 \\ -9 \\ -3 \end{bmatrix}$ ,

- $\vec{0A} = \vec{0P} + \frac{1}{3}\vec{PQ}$
- $\vec{0B} = \vec{0P} + \frac{2}{3}\vec{PQ}$

## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .

## Solution



$$\bullet \vec{0A} = \vec{0P} + \frac{1}{3}\vec{PQ}$$

$$\bullet \vec{0B} = \vec{0P} + \frac{2}{3}\vec{PQ}$$


Since  $\vec{PQ} = \begin{bmatrix} 6 \\ -9 \\ -3 \end{bmatrix}$ ,

$$\vec{0A} = \begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix} + \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ 4 \end{bmatrix}$$

## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .

## Solution



$$\bullet \vec{0A} = \vec{0P} + \frac{1}{3}\vec{PQ}$$

$$\bullet \vec{0B} = \vec{0P} + \frac{2}{3}\vec{PQ}$$


Since  $\vec{PQ} = \begin{bmatrix} 6 \\ -9 \\ -3 \end{bmatrix}$ ,

$$\vec{0A} = \begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix} + \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ 4 \end{bmatrix} \quad \text{and} \quad \vec{0B} = \begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix} + \begin{bmatrix} 4 \\ -6 \\ -2 \end{bmatrix} = \begin{bmatrix} 6 \\ -3 \\ 3 \end{bmatrix}.$$

## Problem

Find the two points trisecting the segment between  $P = (2, 3, 5)$  and  $Q = (8, -6, 2)$ .

## Solution



$$\bullet \vec{0A} = \vec{0P} + \frac{1}{3}\vec{PQ}$$

$$\bullet \vec{0B} = \vec{0P} + \frac{2}{3}\vec{PQ}$$

Since  $\vec{PQ} = \begin{bmatrix} 6 \\ -9 \\ -3 \end{bmatrix}$ ,

$$\vec{0A} = \begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix} + \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix} = \begin{bmatrix} 4 \\ 0 \\ 4 \end{bmatrix} \text{ and } \vec{0B} = \begin{bmatrix} 2 \\ 3 \\ 5 \end{bmatrix} + \begin{bmatrix} 4 \\ -6 \\ -2 \end{bmatrix} = \begin{bmatrix} 6 \\ -3 \\ 3 \end{bmatrix}.$$


Therefore, the two points are  $A = (4, 0, 4)$  and  $B = (6, -3, 3)$ .

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.


## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.




Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,



## Example


If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,

## Example


If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and

## Example


If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.




Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .

It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.




Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .

It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .

$$\overrightarrow{M_1M_2} = \overrightarrow{M_1B} + \overrightarrow{BM_2}$$

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.




Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .

It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .

$$\begin{aligned}\overrightarrow{M_1M_2} &= \overrightarrow{M_1B} + \overrightarrow{BM_2} \\ &= \frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC}\end{aligned}$$

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.




Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .

It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .

$$\begin{aligned}\overrightarrow{M_1M_2} &= \overrightarrow{M_1B} + \overrightarrow{BM_2} \\ &= \frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC} \\ &= \frac{1}{2}\overrightarrow{AC}\end{aligned}$$

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .

It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .


$$\begin{aligned}\overrightarrow{M_1M_2} &= \overrightarrow{M_1B} + \overrightarrow{BM_2} \\ &= \frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC} \\ &= \frac{1}{2}\overrightarrow{AC}\end{aligned}$$

$$\overrightarrow{M_4M_3} = \overrightarrow{M_4D} + \overrightarrow{DM_3}$$



## Example

If  $ABCD$  is an arbitrary quadrilateral, then the the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .


It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .

$$\begin{aligned}\overrightarrow{M_1M_2} &= \overrightarrow{M_1B} + \overrightarrow{BM_2} \\ &= \frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC} \\ &= \frac{1}{2}\overrightarrow{AC}\end{aligned}$$

$$\begin{aligned}\overrightarrow{M_4M_3} &= \overrightarrow{M_4D} + \overrightarrow{DM_3} \\ &= \frac{1}{2}\overrightarrow{AD} + \frac{1}{2}\overrightarrow{DC}\end{aligned}$$

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .


It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .

$$\begin{aligned}\overrightarrow{M_1M_2} &= \overrightarrow{M_1B} + \overrightarrow{BM_2} \\ &= \frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC} \\ &= \frac{1}{2}\overrightarrow{AC}\end{aligned}$$

$$\begin{aligned}\overrightarrow{M_4M_3} &= \overrightarrow{M_4D} + \overrightarrow{DM_3} \\ &= \frac{1}{2}\overrightarrow{AD} + \frac{1}{2}\overrightarrow{DC} \\ &= \frac{1}{2}\overrightarrow{AC}\end{aligned}$$

## Example

If  $ABCD$  is an arbitrary quadrilateral, then the midpoints of the four sides of  $ABCD$  are the vertices of a parallelogram.



Let  $M_1$  denote the midpoint of  $\overrightarrow{AB}$ ,  
 $M_2$  the midpoint of  $\overrightarrow{BC}$ ,  
 $M_3$  the midpoint of  $\overrightarrow{CD}$ , and  
 $M_4$  the midpoint of  $\overrightarrow{DA}$ .

It suffices to prove that  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ .

$$\begin{aligned}\overrightarrow{M_1M_2} &= \overrightarrow{M_1B} + \overrightarrow{BM_2} \\ &= \frac{1}{2}\overrightarrow{AB} + \frac{1}{2}\overrightarrow{BC} \\ &= \frac{1}{2}\overrightarrow{AC}\end{aligned}$$

$$\begin{aligned}\overrightarrow{M_4M_3} &= \overrightarrow{M_4D} + \overrightarrow{DM_3} \\ &= \frac{1}{2}\overrightarrow{AD} + \frac{1}{2}\overrightarrow{DC} \\ &= \frac{1}{2}\overrightarrow{AC}\end{aligned}$$

Since  $\overrightarrow{M_1M_2} = \overrightarrow{M_4M_3}$ , the points  $M_1, M_2, M_3, M_4$  are the vertices of a parallelogram.