

A First Course in
LINEAR ALGEBRA

Lecture Notes
for Math 1503

6.2: Complex Numbers; Polar Form

Creative Commons License (CC BY-NC-SA)

A First Course in Linear Algebra

Lecture Slides

These lecture slides were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [A First Course in Linear Algebra](#) based on K. Kuttler's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Tim Alderson, University of New Brunswick
- Ilijas Farah, York University
- Ken Kuttler, Brigham Young University
- Asia Weiss, York University

License




Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.


Complex Numbers in Polar Form

Suppose $z = a + bi$, and let $r = |z| = \sqrt{a^2 + b^2}$. Then r is the distance from z to the origin. Denote by θ the angle that the line through 0 and z makes with the positive x -axis (measured clockwise).



Complex Numbers in Polar Form


Suppose $z = a + bi$, and let $r = |z| = \sqrt{a^2 + b^2}$. Then r is the distance from z to the origin. Denote by θ the angle that the line through 0 and z makes with the positive x -axis (measured clockwise).



Then θ is an angle defined by $\cos \theta = \frac{a}{r}$ and $\sin \theta = \frac{b}{r}$, so
$$z = r \cos \theta + r \sin \theta i = r(\cos \theta + i \sin \theta).$$

Complex Numbers in Polar Form

Suppose $z = a + bi$, and let $r = |z| = \sqrt{a^2 + b^2}$. Then r is the distance from z to the origin. Denote by θ the angle that the line through 0 and z makes with the positive x -axis (measured clockwise).



Then θ is an angle defined by $\cos \theta = \frac{a}{r}$ and $\sin \theta = \frac{b}{r}$, so

$$z = r \cos \theta + r \sin \theta i = r(\cos \theta + i \sin \theta).$$

θ is called an **argument of z** , and is denoted $\arg z$.

The Polar Form

Definition (Polar Form of a Complex Number)

Let z be a complex number with $|z| = r$ and $\arg z = \theta$. Then

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta)$$

is called a **polar form** of z .

The Polar Form

Definition (Polar Form of a Complex Number)

Let z be a complex number with $|z| = r$ and $\arg z = \theta$. Then

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta)$$

is called a **polar form** of z .

Since $\arg z$ is not unique, we do not write **the** polar form of z .

The Polar Form

Definition (Polar Form of a Complex Number)

Let z be a complex number with $|z| = r$ and $\arg z = \theta$. Then

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta)$$

is called a **polar form** of z .

Since $\arg z$ is not unique, we do not write **the** polar form of z .

Definition

Let z be a complex number with $|z| = r$. The **principal argument** of z is the unique angle $\theta = \arg z$ (measured in radians) such that

$$-\pi < \theta \leq \pi.$$

Example

Find the polar form for the number $z = 1$.

Example

Find the polar form for the number $z = 1$.

Solution

To convert z to polar form, we need to find r and θ so that $1 = re^{i\theta}$.

Example

Find the polar form for the number $z = 1$.

Solution

To convert z to polar form, we need to find r and θ so that $1 = re^{i\theta}$. Now $r = |z| = \sqrt{1^2} = 1$,

Example

Find the polar form for the number $z = 1$.

Solution

To convert z to polar form, we need to find r and θ so that $1 = re^{i\theta}$. Now $r = |z| = \sqrt{1^2} = 1$, and $\theta = 0$ is an argument for $z = 1$.

Example

Find the polar form for the number $z = 1$.

Solution

To convert z to polar form, we need to find r and θ so that $1 = re^{i\theta}$. Now $r = |z| = \sqrt{1^2} = 1$, and $\theta = 0$ is an argument for $z = 1$. However, we may also write

$$1 = e^{2\pi i}, 1 = e^{-2\pi i}, e^{4\pi i}, e^{6\pi i}, \dots$$

Example

Find the polar form for the number $z = 1$.

Solution

To convert z to polar form, we need to find r and θ so that $1 = re^{i\theta}$. Now $r = |z| = \sqrt{1^2} = 1$, and $\theta = 0$ is an argument for $z = 1$. However, we may also write

$$1 = e^{2\pi i}, 1 = e^{-2\pi i}, e^{4\pi i}, e^{6\pi i}, \dots$$

Since sine and cosine have periodicity 2π , we may add (or subtract) multiples of 2π to any argument.

Converting to Polar Form

Example

Convert the number $z = -2 + 2\sqrt{3}i$ to polar form.

Converting to Polar Form

Example

Convert the number $z = -2 + 2\sqrt{3}i$ to polar form.

Solution

To convert z to polar form, we need to find r and θ so that $-2 + 2\sqrt{3}i = re^{i\theta}$.

Converting to Polar Form

Example

Convert the number $z = -2 + 2\sqrt{3}i$ to polar form.

Solution

To convert z to polar form, we need to find r and θ so that $-2 + 2\sqrt{3}i = re^{i\theta}$. Since $r = |z|$,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2}$$

Converting to Polar Form

Example

Convert the number $z = -2 + 2\sqrt{3}i$ to polar form.

Solution

To convert z to polar form, we need to find r and θ so that $-2 + 2\sqrt{3}i = re^{i\theta}$. Since $r = |z|$,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)}$$

Converting to Polar Form

Example

Convert the number $z = -2 + 2\sqrt{3}i$ to polar form.

Solution

To convert z to polar form, we need to find r and θ so that $-2 + 2\sqrt{3}i = re^{i\theta}$. Since $r = |z|$,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16}$$

Converting to Polar Form

Example

Convert the number $z = -2 + 2\sqrt{3}i$ to polar form.

Solution

To convert z to polar form, we need to find r and θ so that $-2 + 2\sqrt{3}i = re^{i\theta}$. Since $r = |z|$,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16} = 4.$$

Converting to Polar Form

Example

Convert the number $z = -2 + 2\sqrt{3}i$ to polar form.

Solution

To convert z to polar form, we need to find r and θ so that $-2 + 2\sqrt{3}i = re^{i\theta}$. Since $r = |z|$,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16} = 4.$$

There are two approaches to finding an argument, θ .

Converting to Polar Form

Example


Conver the number $z = -2 + 2\sqrt{3}i$ to polar form.

Solution

To convert z to polar form, we need to find r and θ so that $-2 + 2\sqrt{3}i = re^{i\theta}$. Since $r = |z|$,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16} = 4.$$

There are two approaches to finding an argument, θ . One is to graph $-2 + 2\sqrt{3}i$ in the complex plane.



Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4;

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$.

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Therefore, z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Therefore, z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

The other approach to finding an argument, θ , for $z = -2 + 2\sqrt{3}i$ is as follows. We've already calculated $|z| = r = 4$. By definition, θ is an angle satisfying

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Therefore, z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

The other approach to finding an argument, θ , for $z = -2 + 2\sqrt{3}i$ is as follows. We've already calculated $|z| = r = 4$. By definition, θ is an angle satisfying

$$\cos \theta = \frac{-2}{4}$$

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Therefore, z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

The other approach to finding an argument, θ , for $z = -2 + 2\sqrt{3}i$ is as follows. We've already calculated $|z| = r = 4$. By definition, θ is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2}$$

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Therefore, z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

The other approach to finding an argument, θ , for $z = -2 + 2\sqrt{3}i$ is as follows. We've already calculated $|z| = r = 4$. By definition, θ is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2} \text{ and } \sin \theta = \frac{2\sqrt{3}}{4}$$

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Therefore, z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

The other approach to finding an argument, θ , for $z = -2 + 2\sqrt{3}i$ is as follows. We've already calculated $|z| = r = 4$. By definition, θ is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2} \text{ and } \sin \theta = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}.$$

Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2, $2\sqrt{3}$, and 4; you should recognize this as a right triangle whose other two angles measure $\frac{\pi}{3}$ and $\frac{\pi}{6}$. From this, we see that $\theta = \frac{2\pi}{3}$ is an argument of z .

Therefore, z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

The other approach to finding an argument, θ , for $z = -2 + 2\sqrt{3}i$ is as follows. We've already calculated $|z| = r = 4$. By definition, θ is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2} \text{ and } \sin \theta = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}.$$

By graphing the point $(-\frac{1}{2}, \frac{\sqrt{3}}{2})$, we again determine that $\theta = \frac{2\pi}{3}$, and thus z can be written in polar form as $z = 4e^{i(2\pi/3)}$.

Exercises

Convert each of the following complex numbers to polar form.

① $3i$

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

② $-1 - i$

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

② $-1 - i = \sqrt{2}e^{-(3\pi/4)i}$

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

② $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$.

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

② $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$.

③ $\sqrt{3} - i$

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

② $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$.

③ $\sqrt{3} - i = 2e^{-(\pi/6)i}$.

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

② $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$.

③ $\sqrt{3} - i = 2e^{-(\pi/6)i}$.

④ $\sqrt{3} + 3i$

Exercises

Convert each of the following complex numbers to polar form.

① $3i = 3e^{(\pi/2)i}$.

② $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$.

③ $\sqrt{3} - i = 2e^{-(\pi/6)i}$.

④ $\sqrt{3} + 3i = 2\sqrt{3}e^{(\pi/3)i}$.

Converting from Polar Form to Cartesian form

Problem

Let $z = 2e^{2\pi i/3}$. Write z in the form $z = a + bi$ (this is called **Cartesian form** or **Standard form**).

Converting from Polar Form to Cartesian form

Problem

Let $z = 2e^{2\pi i/3}$. Write z in the form $z = a + bi$ (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that $e^{i\theta} = \cos \theta + i \sin \theta$,

Converting from Polar Form to Cartesian form

Problem

Let $z = 2e^{2\pi i/3}$. Write z in the form $z = a + bi$ (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that $e^{i\theta} = \cos \theta + i \sin \theta$, and thus

$$e^{2\pi i/3} = \cos(2\pi/3) + i \sin(2\pi/3)$$

Converting from Polar Form to Cartesian form

Problem

Let $z = 2e^{2\pi i/3}$. Write z in the form $z = a + bi$ (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that $e^{i\theta} = \cos \theta + i \sin \theta$, and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i\frac{\sqrt{3}}{2}. \end{aligned}$$

Converting from Polar Form to Cartesian form

Problem

Let $z = 2e^{2\pi i/3}$. Write z in the form $z = a + bi$ (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that $e^{i\theta} = \cos \theta + i \sin \theta$, and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i\frac{\sqrt{3}}{2}. \end{aligned}$$

Therefore

$$z = 2e^{2\pi i/3}$$

Converting from Polar Form to Cartesian form

Problem

Let $z = 2e^{2\pi i/3}$. Write z in the form $z = a + bi$ (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that $e^{i\theta} = \cos \theta + i \sin \theta$, and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i\frac{\sqrt{3}}{2}. \end{aligned}$$

Therefore

$$z = 2e^{2\pi i/3} = 2\left(-\frac{1}{2} + i\frac{\sqrt{3}}{2}\right)$$

Converting from Polar Form to Cartesian form

Problem

Let $z = 2e^{2\pi i/3}$. Write z in the form $z = a + bi$ (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that $e^{i\theta} = \cos \theta + i \sin \theta$, and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i\frac{\sqrt{3}}{2}. \end{aligned}$$

Therefore

$$\begin{aligned} z = 2e^{2\pi i/3} &= 2 \left(-\frac{1}{2} + i\frac{\sqrt{3}}{2} \right) \\ &= -1 + \sqrt{3}i. \end{aligned}$$

Exercises

Express each of the following complex numbers in Cartesian form.

① $3e^{-\pi i}$

Exercises

Express each of the following complex numbers in Cartesian form.

① $3e^{-\pi i} = -3$

Exercises

Express each of the following complex numbers in Cartesian form.

① $3e^{-\pi i} = -3$

② $2e^{3\pi i/4}$

Exercises

Express each of the following complex numbers in Cartesian form.

① $3e^{-\pi i} = -3$

② $2e^{3\pi i/4} = -\sqrt{2} + i\sqrt{2}$

Exercises

Express each of the following complex numbers in Cartesian form.

① $3e^{-\pi i} = -3$

② $2e^{3\pi i/4} = -\sqrt{2} + i\sqrt{2}$

③ $2\sqrt{3}e^{-2\pi i/6}$

Exercises

Express each of the following complex numbers in Cartesian form.

① $3e^{-\pi i} = -3$

② $2e^{3\pi i/4} = -\sqrt{2} + i\sqrt{2}$

③ $2\sqrt{3}e^{-2\pi i/6} = \sqrt{3} - 3i$

Using the Polar Form

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

Using the Polar Form

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

Theorem

If $z_1 = r_1 e^{i\theta_1}$ and $z_2 = r_2 e^{i\theta_2}$ are complex numbers, then

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

Using the Polar Form

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

Theorem

If $z_1 = r_1 e^{i\theta_1}$ and $z_2 = r_2 e^{i\theta_2}$ are complex numbers, then

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

Theorem (De Moivre's Formula)

If θ is any angle and n is a positive integer, then $(e^{i\theta})^n = e^{in\theta}$. Equivalently,

$$(\cos \theta + i \sin \theta)^n = (\cos n\theta + i \sin n\theta)$$

Using the Polar Form

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

Theorem

If $z_1 = r_1 e^{i\theta_1}$ and $z_2 = r_2 e^{i\theta_2}$ are complex numbers, then

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

Theorem (De Moivre's Formula)

If θ is any angle and n is a positive integer, then $(e^{i\theta})^n = e^{in\theta}$. Equivalently,

$$(\cos \theta + i \sin \theta)^n = (\cos n\theta + i \sin n\theta)$$

As an immediate consequence of De Moivre's Formula, we have that for any real number $r > 0$ and any positive integer n ,

$$\begin{aligned}(re^{i\theta})^n &= r^n e^{in\theta} \\ (r(\cos \theta + i \sin \theta))^n &= r^n (\cos n\theta + i \sin n\theta)\end{aligned}$$

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$.

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

$$z^6 w^3 = (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3$$

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \end{aligned}$$

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \end{aligned}$$

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \end{aligned}$$

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \\ &= 64e^{\pi i} \end{aligned}$$

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \\ &= 64e^{\pi i} \\ &= 64(\cos \pi + i \sin \pi) \end{aligned}$$

Problem

Express $(1 - i)^6(\sqrt{3} + i)^3$ in the form $a + bi$.

Solution

Let $z = 1 - i = \sqrt{2}e^{-\pi i/4}$ and $w = \sqrt{3} + i = 2e^{\pi i/6}$. We want to compute $z^6 w^3$.

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \\ &= 64e^{\pi i} \\ &= 64(\cos \pi + i \sin \pi) \\ &= -64. \end{aligned}$$

Problem

Express $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$ in the form $a + bi$.

Problem

Express $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$ in the form $a + bi$.

Solution

$$\text{Let } z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}.$$

Problem

Express $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$ in the form $a + bi$.

Solution

Let $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$. Then

$$z^{17} = \left(e^{-\pi i/3}\right)^{17}$$

Problem

Express $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$ in the form $a + bi$.

Solution

Let $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$. Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \end{aligned}$$

Problem

Express $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$ in the form $a + bi$.

Solution

Let $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$. Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \\ &= e^{\pi i/3} \end{aligned}$$

Problem

Express $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$ in the form $a + bi$.

Solution

Let $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$. Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \\ &= e^{\pi i/3} \\ &= \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \end{aligned}$$

Problem

Express $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$ in the form $a + bi$.

Solution

Let $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$. Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \\ &= e^{\pi i/3} \\ &= \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \\ &= \frac{1}{2} + \frac{\sqrt{3}}{2}i. \end{aligned}$$