

# Math 1003

## Functions and Models


Dr. Tim Alderson

University of New Brunswick Saint John

Chapter 1


# Outline

- 1 1.1 Ways to Represent a Function
  - The Definition of a Function
  - The Vertical Line Test
  - Piecewise Defined Functions
  - Symmetry
  - Increasing and Decreasing Functions
  - A Note on Domains of Functions




A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .

A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .



A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .


- Functions are also synonymously called “maps”.



A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .

- Functions are also synonymously called “maps”.

The set  $D$  in the definition of  $f$  is called the domain of  $f$ .




## Definition (Function)

A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .


## Definition (Co-domain)

The set  $E$  in the definition of  $f$  is called the co-domain of  $f$ .



A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .

The number  $f(x)$  is called *the value of  $f$  at  $x$*  and is read “ $f$  of  $x$ ”.




A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .

The number  $f(x)$  is called *the value of  $f$  at  $x$*  and is read “ $f$  of  $x$ ”.

- The value of  $f$  at  $x$  is also called the image of  $x$  under the map  $f$ .






A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .

The number  $f(x)$  is called *the value of  $f$  at  $x$*  and is read “ $f$  of  $x$ ”.

- The value of  $f$  at  $x$  is also called the image of  $x$  under the map  $f$ .
- In the expression  $f(x)$ ,  $x$  is referred to as the *argument* of  $f$ .




A function  $f$  is a rule that assigns to each element  $x$  in a set  $D$  exactly one element, called  $f(x)$ , in a set  $E$ .


The set of all possible values taken by  $f(x)$  as the element  $x$  runs over elements of  $D$  is called the range of  $f$ .

## Question


Given a curve in the plane, is it the graph of a function or not?



?



?



?

# The Vertical Line Test


## Question

*Given a curve in the plane, is it the graph of a function or not?*


The answer is as follows.

## Proposition (The Vertical Line Test)


*A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.*



?



?



?

# The Vertical Line Test


## Question

*Given a curve in the plane, is it the graph of a function or not?*


The answer is as follows.

## Proposition (The Vertical Line Test)


*A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.*



?



?



?

# The Vertical Line Test


## Question

*Given a curve in the plane, is it the graph of a function or not?*


The answer is as follows.

## Proposition (The Vertical Line Test)


*A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.*



Function



?



?

# The Vertical Line Test


## Question

*Given a curve in the plane, is it the graph of a function or not?*


The answer is as follows.

## Proposition (The Vertical Line Test)


*A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.*



Function



?



?

# The Vertical Line Test


## Question

*Given a curve in the plane, is it the graph of a function or not?*


The answer is as follows.

## Proposition (The Vertical Line Test)


*A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.*



Function



Not a function



?



# The Vertical Line Test


## Question

*Given a curve in the plane, is it the graph of a function or not?*


The answer is as follows.

## Proposition (The Vertical Line Test)


*A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.*



Function



Not a function



?

# The Vertical Line Test


## Question

*Given a curve in the plane, is it the graph of a function or not?*


The answer is as follows.

## Proposition (The Vertical Line Test)


*A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.*



Function



Not a function



Function

# Piecewise Defined Functions

## Definition (Piecewise Defined Function)


A piecewise defined function is a function that is defined by different algebraic formulas on different subsets of its domain.

# Piecewise Defined Functions

## Definition (Piecewise Defined Function)

A piecewise defined function is a function that is defined by different algebraic formulas on different subsets of its domain.

## Example



$$f(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ -1 & \text{if } x < 0 \end{cases}$$

The filled red circle means  $(0, 1)$  is on the curve.


The open circle means  $(0, -1)$  is not on the curve.

## Example

The absolute value  $|x|$  of a number  $a$  is defined to be

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

Sketch a graph of the function  $f(x) = |x|$ .




## Example

The absolute value  $|x|$  of a number  $a$  is defined to be

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

Sketch a graph of the function  $f(x) = |x|$ .




## Example

The absolute value  $|x|$  of a number  $a$  is defined to be

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

Sketch a graph of the function  $f(x) = |x|$ .




## Example

The absolute value  $|x|$  of a number  $a$  is defined to be

$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

Sketch a graph of the function  $f(x) = |x|$ .






## Example

The absolute value  $|x|$  of a number  $a$  is defined to be


$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

Sketch a graph of the function  $f(x) = |x|$ .




## Example

Find a formula for the function  $f$  whose graph is given below.



## Example


Find a formula for the function  $f$  whose graph is given below.



Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

## Example

Find a formula for the function  $f$  whose graph is given below.




Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

$$f(x) = \begin{cases} \text{if } 0 \leq x < 1 \\ \text{if } 1 \leq x < 2 \\ \text{if } 2 \leq x < 5 \end{cases}$$

## Example

Find a formula for the function  $f$  whose graph is given below.




Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

$$f(x) = \begin{cases} ? & \text{if } 0 \leq x < 1 \\ & \text{if } 1 \leq x < 2 \\ & \text{if } 2 \leq x < 5 \end{cases}$$

## Example

Find a formula for the function  $f$  whose graph is given below.




Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < 1 \\ 2 - x & \text{if } 1 \leq x < 2 \\ 0 & \text{if } 2 \leq x < 5 \end{cases}$$

## Example

Find a formula for the function  $f$  whose graph is given below.




Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < 1 \\ ? & \text{if } 1 \leq x < 2 \\ & \text{if } 2 \leq x < 5 \end{cases}$$

## Example

Find a formula for the function  $f$  whose graph is given below.




Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < 1 \\ 2 - x & \text{if } 1 \leq x < 2 \\ 0 & \text{if } 2 \leq x < 5 \end{cases}$$



## Example

Find a formula for the function  $f$  whose graph is given below.




Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < 1 \\ 2 - x & \text{if } 1 \leq x < 2 \\ ? & \text{if } 2 \leq x < 5 \end{cases}$$

## Example

Find a formula for the function  $f$  whose graph is given below.




Different formulas on  $[0, 1)$ ,  $[1, 2)$ , and  $[2, 5)$ .

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < 1 \\ 2 - x & \text{if } 1 \leq x < 2 \\ 0 & \text{if } 2 \leq x < 5 \end{cases}$$


## Example

Sketch the function  $f(x) = |2x - 3|$ .



## Example


Sketch the function  $f(x) = |2x - 3|$ .



$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

## Example

Sketch the function  $f(x) = |2x - 3|$ .




$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

$$|2x - 3| = \begin{cases} 2x - 3 & \text{if } 2x - 3 \geq 0 \\ -(2x - 3) & \text{if } 2x - 3 < 0 \end{cases}$$

## Example

Sketch the function  $f(x) = |2x - 3|$ .




$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

$$|2x - 3| = \begin{cases} 2x - 3 & \text{if } 2x - 3 \geq 0 \\ -(2x - 3) & \text{if } 2x - 3 < 0 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } 2x \geq 3 \\ -2x + 3 & \text{if } 2x < 3 \end{cases}$$

## Example

Sketch the function  $f(x) = |2x - 3|$ .



$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$


$$|2x - 3| = \begin{cases} 2x - 3 & \text{if } 2x - 3 \geq 0 \\ -(2x - 3) & \text{if } 2x - 3 < 0 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } 2x \geq 3 \\ -2x + 3 & \text{if } 2x < 3 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } x \geq 3/2 \\ -2x + 3 & \text{if } x < 3/2. \end{cases}$$

## Example

Sketch the function  $f(x) = |2x - 3|$ .



$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

$$|2x - 3| = \begin{cases} 2x - 3 & \text{if } 2x - 3 \geq 0 \\ -(2x - 3) & \text{if } 2x - 3 < 0 \end{cases}$$


$$= \begin{cases} 2x - 3 & \text{if } 2x \geq 3 \\ -2x + 3 & \text{if } 2x < 3 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } x \geq 3/2 \\ -2x + 3 & \text{if } x < 3/2. \end{cases}$$



## Example

Sketch the function  $f(x) = |2x - 3|$ .



$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$


$$|2x - 3| = \begin{cases} 2x - 3 & \text{if } 2x - 3 \geq 0 \\ -(2x - 3) & \text{if } 2x - 3 < 0 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } 2x \geq 3 \\ -2x + 3 & \text{if } 2x < 3 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } x \geq 3/2 \\ -2x + 3 & \text{if } x < 3/2. \end{cases}$$


## Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .



## Example


Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .



$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

## Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .




$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

$$\frac{|4x + 2|}{2x + 1} = \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases}$$

# Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .




$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

$$\begin{aligned} \frac{|4x + 2|}{2x + 1} &= \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases} \\ &= \begin{cases} \frac{?}{2x+1} & \text{if } 4x > -2 \\ \frac{?}{2x+1} & \text{if } 4x < -2 \end{cases} \end{aligned}$$

# Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .




$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

$$\begin{aligned} \frac{|4x + 2|}{2x + 1} &= \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases} \\ &= \begin{cases} \frac{2(2x+1)}{2x+1} & \text{if } 4x > -2 \\ \frac{?}{2x+1} & \text{if } 4x < -2 \end{cases} \end{aligned}$$

## Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .




$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

$$\begin{aligned} \frac{|4x + 2|}{2x + 1} &= \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases} \\ &= \begin{cases} \frac{2(2x+1)}{2x+1} & \text{if } 4x > -2 \\ \frac{?}{2x+1} & \text{if } 4x < -2 \end{cases} \end{aligned}$$

# Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .




$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

$$\begin{aligned} \frac{|4x + 2|}{2x + 1} &= \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases} \\ &= \begin{cases} \frac{2(2x+1)}{2x+1} & \text{if } 4x > -2 \\ \frac{-2(2x+1)}{2x+1} & \text{if } 4x < -2 \end{cases} \end{aligned}$$



## Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .



$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$


$$\frac{|4x + 2|}{2x + 1} = \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases}$$

$$= \begin{cases} \frac{2(\cancel{2x+1})}{\cancel{2x+1}} & \text{if } 4x > -2 \\ \frac{-2(\cancel{2x+1})}{\cancel{2x+1}} & \text{if } 4x < -2 \end{cases}$$

$$= \begin{cases} 2 & \text{if } x > -\frac{1}{2} \\ -2 & \text{if } x < -\frac{1}{2}. \end{cases}$$

# Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .



$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$


$$\frac{|4x + 2|}{2x + 1} = \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases}$$

$$= \begin{cases} \frac{2(2x+1)}{2x+1} & \text{if } 4x > -2 \\ \frac{-2(2x+1)}{2x+1} & \text{if } 4x < -2 \end{cases}$$

$$= \begin{cases} 2 & \text{if } x > -\frac{1}{2} \\ -2 & \text{if } x < -\frac{1}{2}. \end{cases}$$

# Example

Sketch the function  $f(x) = \frac{|4x + 2|}{2x + 1}$ .



$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

$$\frac{|4x + 2|}{2x + 1} = \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases}$$

$$= \begin{cases} \frac{2(2x+1)}{2x+1} & \text{if } 4x > -2 \\ \frac{-2(2x+1)}{2x+1} & \text{if } 4x < -2 \end{cases}$$

$$= \begin{cases} 2 & \text{if } x > -\frac{1}{2} \\ -2 & \text{if } x < -\frac{1}{2}. \end{cases}$$

# Symmetry

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

# Symmetry

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example ( $x^2$ is Even, $x^3$ is Odd)

The function  $f(x) = x^2$  is even:

The function  $g(x) = x^3$  is odd:

# Symmetry

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example ( $x^2$ is Even, $x^3$ is Odd)

The function  $f(x) = x^2$  is even:

$$f(-x) = (-x)^2 = x^2 = f(x).$$

The function  $g(x) = x^3$  is odd:

# Symmetry

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example ( $x^2$ is Even, $x^3$ is Odd)

The function  $f(x) = x^2$  is even:

$$f(-x) = (-x)^2 = x^2 = f(x).$$

The function  $g(x) = x^3$  is odd:

$$g(-x) = (-x)^3 = -x^3 = -g(x).$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$g(x) = 1 - x^4$$

$$h(x) = 2x - 1$$



## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$f(-x) =$$

$$g(x) = 1 - x^4$$

$$g(-x) =$$

$$h(x) = 2x - 1$$

$$h(-x) =$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$f(-x) = (-x)^5 + (-x)$$

$$g(x) = 1 - x^4$$

$$g(-x) =$$

$$h(x) = 2x - 1$$

$$h(-x) =$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \end{aligned}$$

$$g(x) = 1 - x^4$$

$$g(-x) =$$

$$h(x) = 2x - 1$$

$$h(-x) =$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \end{aligned}$$

$$g(x) = 1 - x^4$$

$$g(-x) =$$

$$h(x) = 2x - 1$$

$$h(-x) =$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

$$g(x) = 1 - x^4$$

$$g(-x) =$$

$$h(x) = 2x - 1$$

$$h(-x) =$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$g(x) = 1 - x^4$$

$$h(x) = 2x - 1$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

$$g(-x) =$$

$$h(-x) =$$

Therefore  $f$  is odd.

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$g(x) = 1 - x^4$$

$$h(x) = 2x - 1$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

$$g(-x) = 1 - (-x)^4$$

$$h(-x) =$$

Therefore  $f$  is odd.

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \end{aligned}$$

$$h(x) = 2x - 1$$

$$h(-x) =$$

Therefore  $f$  is odd.



## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \\ &= g(x) \end{aligned}$$

$$h(x) = 2x - 1$$

$$h(-x) =$$

Therefore  $f$  is odd.

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

Therefore  $f$  is odd.

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \\ &= g(x) \end{aligned}$$

Therefore  $g$  is even.

$$h(x) = 2x - 1$$

$$h(-x) =$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

Therefore  $f$  is odd.

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \\ &= g(x) \end{aligned}$$

Therefore  $g$  is even.

$$h(x) = 2x - 1$$

$$h(-x) = 2(-x) - 1$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

Therefore  $f$  is odd.

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \\ &= g(x) \end{aligned}$$

Therefore  $g$  is even.

$$h(x) = 2x - 1$$

$$\begin{aligned} h(-x) &= 2(-x) - 1 \\ &= -2x - 1 \end{aligned}$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

Therefore  $f$  is odd.

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \\ &= g(x) \end{aligned}$$

Therefore  $g$  is even.

$$h(x) = 2x - 1$$

$$\begin{aligned} h(-x) &= 2(-x) - 1 \\ &= -2x - 1 \\ &\neq h(x), -h(x) \end{aligned}$$

## Definition (Even and Odd Functions)

A function  $f$  is called even if  $f(-x) = f(x)$  for all  $x$  in its domain. A function  $f$  is called odd if  $f(-x) = -f(x)$  for all  $x$  in its domain.

## Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

Therefore  $f$  is odd.

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \\ &= g(x) \end{aligned}$$

Therefore  $g$  is even.

$$h(x) = 2x - 1$$

$$\begin{aligned} h(-x) &= 2(-x) - 1 \\ &= -2x - 1 \\ &\neq h(x), -h(x) \end{aligned}$$

Therefore  $h$  is neither even nor odd.

# Increasing and Decreasing Functions

## Definition (Increasing and Decreasing Functions)

A function  $f$  is called increasing on an interval  $I$  if  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

It is called decreasing on the interval  $I$  if  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .


# Increasing and Decreasing Functions

## Definition (Increasing and Decreasing Functions)

A function  $f$  is called increasing on an interval  $I$  if  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

It is called decreasing on the interval  $I$  if  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

## Example (Increasing and Decreasing)






# Increasing and Decreasing Functions

## Definition (Increasing and Decreasing Functions)

A function  $f$  is called increasing on an interval  $I$  if  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

It is called decreasing on the interval  $I$  if  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

## Example (Increasing and Decreasing)



- $f$  is increasing on  $[-1, -\frac{1}{2}]$ .


# Increasing and Decreasing Functions

## Definition (Increasing and Decreasing Functions)

A function  $f$  is called increasing on an interval  $I$  if  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

It is called decreasing on the interval  $I$  if  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

## Example (Increasing and Decreasing)



- $f$  is increasing on  $[-1, -\frac{1}{2}]$ .
- $f$  is decreasing on  $[-\frac{1}{2}, \frac{1}{2}]$ .


# Increasing and Decreasing Functions

## Definition (Increasing and Decreasing Functions)

A function  $f$  is called increasing on an interval  $I$  if  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

It is called decreasing on the interval  $I$  if  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$  in  $I$ .

## Example (Increasing and Decreasing)



- $f$  is increasing on  $[-1, -\frac{1}{2}]$ .
- $f$  is decreasing on  $[-\frac{1}{2}, \frac{1}{2}]$ .
- $f$  is increasing on  $[\frac{1}{2}, 1]$ .

# A Note on Domains of Functions

If the domain of a function isn't specified, it is assumed to be all numbers  $x$  for which the formula  $f(x)$  is defined. There are some restrictions to consider:

# A Note on Domains of Functions

If the domain of a function isn't specified, it is assumed to be all numbers  $x$  for which the formula  $f(x)$  is defined. There are some restrictions to consider:

- Can't divide by 0.

# A Note on Domains of Functions

If the domain of a function isn't specified, it is assumed to be all numbers  $x$  for which the formula  $f(x)$  is defined. There are some restrictions to consider:

- Can't divide by 0.
- Can't take the even root of a negative number ( $\sqrt{-1}$ ,  $\sqrt[4]{-2053}$ ,  $\sqrt[6]{-15} \dots$  not allowed).

# A Note on Domains of Functions

If the domain of a function isn't specified, it is assumed to be all numbers  $x$  for which the formula  $f(x)$  is defined. There are some restrictions to consider:

- Can't divide by 0.
- Can't take the even root of a negative number ( $\sqrt{-1}$ ,  $\sqrt[4]{-2053}$ ,  $\sqrt[6]{-15} \dots$  not allowed).
- Can't take  $\log x$  if  $x \leq 0$ .

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$



## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0?

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number?

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? **Yes.**

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$x - 2 \geq 0$$

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$x - 2 \geq 0$$

$$x \geq 2$$

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$x - 2 \geq 0$$

$$x \geq 2$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .



## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.
- Any risk of dividing by 0?

$$x - 2 \geq 0$$

$$x \geq 2$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.
- Any risk of dividing by 0? **Yes.**

$$x - 2 \geq 0$$

$$x \geq 2$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.
- Any risk of dividing by 0? Yes.
- Any risk of taking the even root of a negative number?

$$x - 2 \geq 0$$

$$x \geq 2$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$x - 2 \geq 0$$

$$x \geq 2$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? Yes.
- Any risk of taking the even root of a negative number? No.

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$x - 2 \geq 0$$

$$x \geq 2$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? Yes.
- Any risk of taking the even root of a negative number? No.
- $x^2 - x - 6$  can't be 0.

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$x - 2 \geq 0$$

$$x \geq 2$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? Yes.
- Any risk of taking the even root of a negative number? No.
- $x^2 - x - 6$  can't be 0.

$$x^2 - x - 6 \neq 0$$

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$\begin{aligned} x - 2 &\geq 0 \\ x &\geq 2 \end{aligned}$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? Yes.
- Any risk of taking the even root of a negative number? No.
- $x^2 - x - 6$  can't be 0.

$$\begin{aligned} x^2 - x - 6 &\neq 0 \\ (x - 3)(x + 2) &\neq 0 \end{aligned}$$

## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$\begin{aligned} x - 2 &\geq 0 \\ x &\geq 2 \end{aligned}$$

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? Yes.
- Any risk of taking the even root of a negative number? No.
- $x^2 - x - 6$  can't be 0.

$$\begin{aligned} x^2 - x - 6 &\neq 0 \\ (x - 3)(x + 2) &\neq 0 \\ x &\neq 3 \text{ or } -2 \end{aligned}$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .



## Example (Two Functions and Their Domains)

Find the domains of the following two functions:

$$f(x) = \sqrt[4]{x-2} + \sqrt[3]{6-x}$$

- Any risk of dividing by 0? No.
- Any risk of taking the even root of a negative number? Yes.
- $x - 2$  can't be negative.

$$\begin{aligned} x - 2 &\geq 0 \\ x &\geq 2 \end{aligned}$$

Domain is all real numbers bigger than or equal to 2; that is,  $[2, \infty)$ .

$$g(x) = \frac{x^2 - 9}{x^2 - x - 6}$$

- Any risk of dividing by 0? Yes.
- Any risk of taking the even root of a negative number? No.
- $x^2 - x - 6$  can't be 0.

$$\begin{aligned} x^2 - x - 6 &\neq 0 \\ (x - 3)(x + 2) &\neq 0 \\ x &\neq 3 \text{ or } -2 \end{aligned}$$

Domain is all real numbers except 3 and  $-2$ ; that is,  $(-\infty, -2) \cup (-2, 3) \cup (3, \infty)$ .