

# Math 1013

## Improper Integrals and Arc Length

Dr. Tim Alderson

University of New Brunswick Saint John

Week 9

# Outline

## 1 Improper Integrals

# Outline

1 Improper Integrals

2 Arc Length

# Improper Integrals and Arc Length

## Differentiation..WT\*?

The derivative of  $x^2$ , with respect to  $x$ , is  $2x$ . However, suppose we write  $x^2$  as the sum of  $x$ 's, and then take the derivative:

$$\text{Let } f(x) = x + x + \cdots + x \text{ (} x \text{ times)}$$

Then

$$\begin{aligned} f'(x) &= \frac{d}{dx}[x + x + \cdots + x] \text{ (} x \text{ times)} \\ &= \frac{d}{dx}(x) + \frac{d}{dx}(x) + \cdots + \frac{d}{dx}(x) \text{ (} x \text{ times)} \\ &= 1 + 1 + \cdots + 1 \text{ (} x \text{ times)} \\ &= x \end{aligned}$$

This argument appears to show that the derivative of  $x^2$ , with respect to  $x$ , is actually  $x$ . Where is the fallacy?





# Improper Integrals

- The definition of  $\int_a^b f(x)dx$ , where  $f$  is defined on  $[a, b]$ , has two requirements:
  - 1  $[a, b]$  is a finite interval.
  - 2  $f$  has no infinite discontinuities in  $[a, b]$ .

# Improper Integrals

- The definition of  $\int_a^b f(x)dx$ , where  $f$  is defined on  $[a, b]$ , has two requirements:
  - 1  $[a, b]$  is a finite interval.
  - 2  $f$  has no infinite discontinuities in  $[a, b]$ .
- We are now going to relax these requirements.
  - 1 We allow infinite intervals, such as  $(a, \infty)$ ,  $(-\infty, b)$ , and  $(-\infty, \infty)$ .
  - 2  $f$  might have infinite discontinuities in  $[a, b]$ .

# Improper Integrals

- The definition of  $\int_a^b f(x)dx$ , where  $f$  is defined on  $[a, b]$ , has two requirements:
  - 1  $[a, b]$  is a finite interval.
  - 2  $f$  has no infinite discontinuities in  $[a, b]$ .
- We are now going to relax these requirements.
  - 1 We allow infinite intervals, such as  $(a, \infty)$ ,  $(-\infty, b)$ , and  $(-\infty, \infty)$ .
  - 2  $f$  might have infinite discontinuities in  $[a, b]$ .
- Such integrals are called improper integrals.



# Improper Integrals

- The definition of  $\int_a^b f(x)dx$ , where  $f$  is defined on  $[a, b]$ , has two requirements:
  - 1  $[a, b]$  is a finite interval.
  - 2  $f$  has no infinite discontinuities in  $[a, b]$ .
- We are now going to relax these requirements.
  - 1 We allow infinite intervals, such as  $(a, \infty)$ ,  $(-\infty, b)$ , and  $(-\infty, \infty)$ .
  - 2  $f$  might have infinite discontinuities in  $[a, b]$ .
- Such integrals are called improper integrals.

## Definition (Improper Integral)


The integral

$$\int_a^b f(x)dx$$

is called improper if one or more of the endpoints  $a$  and  $b$  is infinite, or if  $f$  has an infinite discontinuity on  $[a, b]$ .

# Type I: Infinite Intervals


- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .



# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .


$$A(t) = \int_1^t \frac{dx}{x^2} =$$



# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .


$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t =$$



# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .


$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$



# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) =$

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) =$



# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) = 1$ .

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) = 1$ .

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) = 1$ .

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) = 1$ .

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) = 1$ .

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$




- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) = 1$ .

# Type I: Infinite Intervals

- Consider the region  $A$  that lies under  $y = 1/x^2$ , above the  $x$ -axis, and to the right of  $x = 1$ .
- To find its area, approximate with  $A(t)$ , the area of the region under  $1/x^2$ , above the  $x$ -axis, right of  $x = 1$ , and left of  $x = t$ .

$$A(t) = \int_1^t \frac{dx}{x^2} = \left[ -\frac{1}{x} \right]_1^t = 1 - \frac{1}{t}$$



- Notice  $A(t) < 1$  no matter how big  $t$  is.
- Also notice  $\lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left( 1 - \frac{1}{t} \right) = 1$ .
- We say that the area  $A$  is equal to 1 and write

$$\int_1^{\infty} \frac{1}{x^2} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2} dx = 1.$$

## Definition (Improper Integral of Type I)

- 1 If  $\int_a^t f(x)dx$  exists for every  $t \geq a$ , then

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$$

if the limit exists.



## Definition (Improper Integral of Type I)

- 1 If  $\int_a^t f(x)dx$  exists for every  $t \geq a$ , then

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$$

if the limit exists.

- 2 If  $\int_t^b f(x)dx$  exists for every  $t \leq b$ , then

$$\int_{-\infty}^b f(x)dx = \lim_{t \rightarrow -\infty} \int_t^b f(x)dx$$

if the limit exists.

## Definition (Improper Integral of Type I)

- ❶ If  $\int_a^t f(x)dx$  exists for every  $t \geq a$ , then

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$$

if the limit exists.

- ❷ If  $\int_t^b f(x)dx$  exists for every  $t \leq b$ , then

$$\int_{-\infty}^b f(x)dx = \lim_{t \rightarrow -\infty} \int_t^b f(x)dx$$

if the limit exists.

$\int_a^\infty f(x)dx$  and  $\int_{-\infty}^b f(x)dx$  are called **convergent** if the corresponding limit exists and **divergent** if it doesn't exist.

## Definition (Improper Integral of Type I)

- ❶ If  $\int_a^t f(x)dx$  exists for every  $t \geq a$ , then

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$$

if the limit exists.

- ❷ If  $\int_t^b f(x)dx$  exists for every  $t \leq b$ , then

$$\int_{-\infty}^b f(x)dx = \lim_{t \rightarrow -\infty} \int_t^b f(x)dx$$

if the limit exists.


$\int_a^\infty f(x)dx$  and  $\int_{-\infty}^b f(x)dx$  are called **convergent** if the corresponding limit exists and **divergent** if it doesn't exist.

- ❸ If both  $\int_a^\infty f(x)dx$  and  $\int_{-\infty}^a f(x)dx$  are convergent, then we define

$$\int_{-\infty}^\infty f(x)dx = \int_{-\infty}^a f(x)dx + \int_a^\infty f(x)dx.$$


## Example

Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.



## Example


Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.



$$\int_1^{\infty} \frac{1}{x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx$$

## Example


Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.



$$\begin{aligned}\int_1^{\infty} \frac{1}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx \\ &= \lim_{t \rightarrow \infty} [\ln |x|]_1^t\end{aligned}$$

## Example


Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.



$$\begin{aligned}\int_1^{\infty} \frac{1}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx \\ &= \lim_{t \rightarrow \infty} [\ln |x|]_1^t \\ &= \lim_{t \rightarrow \infty} (\ln t - \ln 1)\end{aligned}$$

## Example

Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.




$$\begin{aligned}\int_1^{\infty} \frac{1}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx \\&= \lim_{t \rightarrow \infty} [\ln |x|]_1^t \\&= \lim_{t \rightarrow \infty} (\ln t - \ln 1) \\&= \lim_{t \rightarrow \infty} \ln t\end{aligned}$$



## Example


Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.



$$\begin{aligned}\int_1^{\infty} \frac{1}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx \\ &= \lim_{t \rightarrow \infty} [\ln |x|]_1^t \\ &= \lim_{t \rightarrow \infty} (\ln t - \ln 1) \\ &= \lim_{t \rightarrow \infty} \ln t = \infty\end{aligned}$$

## Example

Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.




Infinite area

$$\begin{aligned}\int_1^{\infty} \frac{1}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx \\ &= \lim_{t \rightarrow \infty} [\ln |x|]_1^t \\ &= \lim_{t \rightarrow \infty} (\ln t - \ln 1) \\ &= \lim_{t \rightarrow \infty} \ln t = \infty\end{aligned}$$


Therefore the improper integral is divergent.

## Example

Determine whether  $\int_1^{\infty} \frac{1}{x} dx$  is convergent or divergent.



Infinite area



Finite area

$$\begin{aligned}
 \int_1^{\infty} \frac{1}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx \\
 &= \lim_{t \rightarrow \infty} [\ln |x|]_1^t \\
 &= \lim_{t \rightarrow \infty} (\ln t - \ln 1) \\
 &= \lim_{t \rightarrow \infty} \ln t = \infty
 \end{aligned}$$

Therefore the improper integral is divergent.

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\int_{-\infty}^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx$$

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\int_{-\infty}^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0$$

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx$$



## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) \end{aligned}$$

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) = \lim_{t \rightarrow -\infty} (0 - \tan^{-1} t) \end{aligned}$$

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) = \lim_{t \rightarrow -\infty} (0 - \tan^{-1} t) \\ &= 0 - \left(-\frac{\pi}{2}\right) = \frac{\pi}{2} \end{aligned}$$

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) = \lim_{t \rightarrow -\infty} (0 - \tan^{-1} t) \\ &= 0 - \left(-\frac{\pi}{2}\right) = \frac{\pi}{2} \end{aligned}$$

$$\int_0^{\infty} \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} [\tan^{-1} x]_0^t$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) = \lim_{t \rightarrow -\infty} (0 - \tan^{-1} t) \\ &= 0 - \left(-\frac{\pi}{2}\right) = \frac{\pi}{2} \end{aligned}$$

$$\begin{aligned} \int_0^{\infty} \frac{1}{1+x^2} dx &= \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} [\tan^{-1} x]_0^t \\ &= \lim_{t \rightarrow \infty} (\tan^{-1} t - \tan^{-1} 0) \end{aligned}$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) = \lim_{t \rightarrow -\infty} (0 - \tan^{-1} t) \\ &= 0 - \left(-\frac{\pi}{2}\right) = \frac{\pi}{2} \end{aligned}$$

$$\begin{aligned} \int_0^{\infty} \frac{1}{1+x^2} dx &= \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} [\tan^{-1} x]_0^t \\ &= \lim_{t \rightarrow \infty} (\tan^{-1} t - \tan^{-1} 0) = \lim_{t \rightarrow \infty} \tan^{-1} t \end{aligned}$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) = \lim_{t \rightarrow -\infty} (0 - \tan^{-1} t) \\ &= 0 - \left(-\frac{\pi}{2}\right) = \frac{\pi}{2} \end{aligned}$$

$$\begin{aligned} \int_0^{\infty} \frac{1}{1+x^2} dx &= \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} [\tan^{-1} x]_0^t \\ &= \lim_{t \rightarrow \infty} (\tan^{-1} t - \tan^{-1} 0) = \lim_{t \rightarrow \infty} \tan^{-1} t = \frac{\pi}{2} \end{aligned}$$

## Example

Evaluate  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

$$\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \int_{-\infty}^0 \frac{1}{1+x^2} dx + \int_0^{\infty} \frac{1}{1+x^2} dx$$

Evaluate the two integrals separately:

$$\begin{aligned} \int_{-\infty}^0 \frac{1}{1+x^2} dx &= \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{1+x^2} dx = \lim_{t \rightarrow -\infty} [\tan^{-1} x]_t^0 \\ &= \lim_{t \rightarrow -\infty} (\tan^{-1} 0 - \tan^{-1} t) = \lim_{t \rightarrow -\infty} (0 - \tan^{-1} t) \\ &= 0 - \left(-\frac{\pi}{2}\right) = \frac{\pi}{2} \end{aligned}$$

$$\begin{aligned} \int_0^{\infty} \frac{1}{1+x^2} dx &= \lim_{t \rightarrow \infty} \int_0^t \frac{1}{1+x^2} dx = \lim_{t \rightarrow \infty} [\tan^{-1} x]_0^t \\ &= \lim_{t \rightarrow \infty} (\tan^{-1} t - \tan^{-1} 0) = \lim_{t \rightarrow \infty} \tan^{-1} t = \frac{\pi}{2} \end{aligned}$$

Therefore  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = \frac{\pi}{2} + \frac{\pi}{2} = \pi$ .



## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx$$

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \left[ \frac{x^{-p+1}}{-p+1} \right]_1^t$$

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \left[ \frac{x^{-p+1}}{-p+1} \right]_1^t = \lim_{t \rightarrow \infty} \frac{\frac{1}{t^{p-1}} - 1}{1-p}$$

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \left[ \frac{x^{-p+1}}{-p+1} \right]_1^t = \lim_{t \rightarrow \infty} \frac{\frac{1}{t^{p-1}} - 1}{1-p}$$

- If  $p > 1$ , then  $p - 1 > 0$ , so as  $t \rightarrow \infty$ ,  $t^{p-1} \rightarrow \infty$  and  $\frac{1}{t^{p-1}} \rightarrow 0$ .

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \left[ \frac{x^{-p+1}}{-p+1} \right]_1^t = \lim_{t \rightarrow \infty} \frac{\frac{1}{t^{p-1}} - 1}{1-p}$$

- If  $p > 1$ , then  $p - 1 > 0$ , so as  $t \rightarrow \infty$ ,  $t^{p-1} \rightarrow \infty$  and  $\frac{1}{t^{p-1}} \rightarrow 0$ .
- Therefore  $\int_1^{\infty} \frac{1}{x^p} dx = \frac{1}{p-1}$  if  $p > 1$ , so the integral is convergent.



## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \left[ \frac{x^{-p+1}}{-p+1} \right]_1^t = \lim_{t \rightarrow \infty} \frac{\frac{1}{t^{p-1}} - 1}{1-p}$$

- If  $p > 1$ , then  $p - 1 > 0$ , so as  $t \rightarrow \infty$ ,  $t^{p-1} \rightarrow \infty$  and  $\frac{1}{t^{p-1}} \rightarrow 0$ .
- Therefore  $\int_1^{\infty} \frac{1}{x^p} dx = \frac{1}{p-1}$  if  $p > 1$ , so the integral is convergent.
- If  $p < 1$ , then  $p - 1 < 0$ , so  $\frac{1}{t^{p-1}} = t^{1-p} \rightarrow \infty$  as  $t \rightarrow \infty$ .

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \left[ \frac{x^{-p+1}}{-p+1} \right]_1^t = \lim_{t \rightarrow \infty} \frac{\frac{1}{t^{p-1}} - 1}{1-p}$$

- If  $p > 1$ , then  $p - 1 > 0$ , so as  $t \rightarrow \infty$ ,  $t^{p-1} \rightarrow \infty$  and  $\frac{1}{t^{p-1}} \rightarrow 0$ .
- Therefore  $\int_1^{\infty} \frac{1}{x^p} dx = \frac{1}{p-1}$  if  $p > 1$ , so the integral is convergent.
- If  $p < 1$ , then  $p - 1 < 0$ , so  $\frac{1}{t^{p-1}} = t^{1-p} \rightarrow \infty$  as  $t \rightarrow \infty$ .
- Therefore  $\int_1^{\infty} \frac{1}{x^p} dx$  is divergent if  $p < 1$ .

## Example

For what values of  $p$  is the integral  $\int_1^{\infty} \frac{1}{x^p} dx$  convergent?

- If  $p = 1$ , the integral is divergent (Example 1), so assume  $p \neq 1$ .

$$\int_1^{\infty} \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^p} dx = \lim_{t \rightarrow \infty} \left[ \frac{x^{-p+1}}{-p+1} \right]_1^t = \lim_{t \rightarrow \infty} \frac{\frac{1}{t^{p-1}} - 1}{1-p}$$

- If  $p > 1$ , then  $p - 1 > 0$ , so as  $t \rightarrow \infty$ ,  $t^{p-1} \rightarrow \infty$  and  $\frac{1}{t^{p-1}} \rightarrow 0$ .
- Therefore  $\int_1^{\infty} \frac{1}{x^p} dx = \frac{1}{p-1}$  if  $p > 1$ , so the integral is convergent.
- If  $p < 1$ , then  $p - 1 < 0$ , so  $\frac{1}{t^{p-1}} = t^{1-p} \rightarrow \infty$  as  $t \rightarrow \infty$ .
- Therefore  $\int_1^{\infty} \frac{1}{x^p} dx$  is divergent if  $p < 1$ .

## Theorem

$\int_1^{\infty} \frac{1}{x^p} dx$  converges if  $p > 1$  and diverges if  $p \leq 1$ .


## Type II: Discontinuous Integrands

We can use the same approach if the function  $f$  is discontinuous at one of the endpoints  $a$  and  $b$  in the integral  $\int_a^b f(x)dx$ .

For example,  $\frac{1}{\sqrt{x-2}}$  is discontinuous at 2, so we might wonder if the integral

$$\int_2^5 \frac{1}{\sqrt{x-2}} dx$$

exists.



## Definition (Improper Integral of Type II)

- 1 If  $f$  is continuous on  $[a, b)$  and discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

if the limit exists.

## Definition (Improper Integral of Type II)

- ❶ If  $f$  is continuous on  $[a, b)$  and discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

if the limit exists.

- ❷ If  $f$  is continuous on  $(a, b]$  and discontinuous at  $a$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$$

if the limit exists.

## Definition (Improper Integral of Type II)

- ❶ If  $f$  is continuous on  $[a, b)$  and discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

if the limit exists.

- ❷ If  $f$  is continuous on  $(a, b]$  and discontinuous at  $a$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$$

if the limit exists.

$\int_a^b f(x) dx$  is called **convergent** if the corresponding limit exists and **divergent** if it doesn't exist.

## Definition (Improper Integral of Type II)

- ❶ If  $f$  is continuous on  $[a, b)$  and discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$$

if the limit exists.

- ❷ If  $f$  is continuous on  $(a, b]$  and discontinuous at  $a$ , then

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$$

if the limit exists.

$\int_a^b f(x) dx$  is called **convergent** if the corresponding limit exists and **divergent** if it doesn't exist.


- ❸ If  $f$  has a discontinuity at  $c$ , where  $a < c < b$ , and both  $\int_a^c f(x) dx$  and  $\int_c^b f(x) dx$  are convergent, then we define

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$



## Example


Find  $\int_2^5 \frac{1}{\sqrt{x-2}} dx$ .



## Example

Find  $\int_2^5 \frac{1}{\sqrt{x-2}} dx$ .


Observe that  $x = 2$  is a vertical asymptote for the integrand.



## Example

Find  $\int_2^5 \frac{1}{\sqrt{x-2}} dx$ .

Observe that  $x = 2$  is a vertical asymptote for the integrand.




$$\begin{aligned} & \int_2^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \int_t^5 \frac{1}{\sqrt{x-2}} dx \end{aligned}$$

## Example

Find  $\int_2^5 \frac{1}{\sqrt{x-2}} dx$ .

Observe that  $x = 2$  is a vertical asymptote for the integrand.




$$\begin{aligned} & \int_2^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \int_t^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \left[ 2\sqrt{x-2} \right]_t^5 \end{aligned}$$

## Example

Find  $\int_2^5 \frac{1}{\sqrt{x-2}} dx$ .

Observe that  $x = 2$  is a vertical asymptote for the integrand.




$$\begin{aligned} & \int_2^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \int_t^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \left[ 2\sqrt{x-2} \right]_t^5 \\ &= \lim_{t \rightarrow 2^+} 2(\sqrt{5-2} - \sqrt{t-2}) \end{aligned}$$

## Example

Find  $\int_2^5 \frac{1}{\sqrt{x-2}} dx$ .

Observe that  $x = 2$  is a vertical asymptote for the integrand.




$$\begin{aligned} & \int_2^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \int_t^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \left[ 2\sqrt{x-2} \right]_t^5 \\ &= \lim_{t \rightarrow 2^+} 2(\sqrt{5-2} - \sqrt{t-2}) \\ &= 2\sqrt{3} \end{aligned}$$

## Example

Find  $\int_2^5 \frac{1}{\sqrt{x-2}} dx$ .

Observe that  $x = 2$  is a vertical asymptote for the integrand.



Area =  $2\sqrt{3}$

$$\begin{aligned} & \int_2^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \int_t^5 \frac{1}{\sqrt{x-2}} dx \\ &= \lim_{t \rightarrow 2^+} \left[ 2\sqrt{x-2} \right]_t^5 \\ &= \lim_{t \rightarrow 2^+} 2(\sqrt{5-2} - \sqrt{t-2}) \\ &= 2\sqrt{3} \end{aligned}$$

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .



## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

$$\int_0^3 \frac{1}{x-1} dx = \int_0^1 \frac{1}{x-1} dx + \int_1^3 \frac{1}{x-1} dx$$

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

$$\int_0^3 \frac{1}{x-1} dx = \int_0^1 \frac{1}{x-1} dx + \int_1^3 \frac{1}{x-1} dx$$

$$\int_0^1 \frac{dx}{x-1} = \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{x-1}$$

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

$$\int_0^3 \frac{1}{x-1} dx = \int_0^1 \frac{1}{x-1} dx + \int_1^3 \frac{1}{x-1} dx$$

$$\int_0^1 \frac{dx}{x-1} = \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{x-1} = \lim_{t \rightarrow 1^-} [\ln |x-1|]_0^t$$

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

$$\int_0^3 \frac{1}{x-1} dx = \int_0^1 \frac{1}{x-1} dx + \int_1^3 \frac{1}{x-1} dx$$

$$\begin{aligned} \int_0^1 \frac{dx}{x-1} &= \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{x-1} = \lim_{t \rightarrow 1^-} [\ln |x-1|]_0^t \\ &= \lim_{t \rightarrow 1^-} \ln |t-1| - \ln 1 \end{aligned}$$

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

$$\int_0^3 \frac{1}{x-1} dx = \int_0^1 \frac{1}{x-1} dx + \int_1^3 \frac{1}{x-1} dx$$

$$\begin{aligned} \int_0^1 \frac{dx}{x-1} &= \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{x-1} = \lim_{t \rightarrow 1^-} [\ln |x-1|]_0^t \\ &= \lim_{t \rightarrow 1^-} \ln |t-1| - \ln 1 = -\infty \end{aligned}$$

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

$$\int_0^3 \frac{1}{x-1} dx = \int_0^1 \frac{1}{x-1} dx + \int_1^3 \frac{1}{x-1} dx$$

$$\begin{aligned} \int_0^1 \frac{dx}{x-1} &= \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{x-1} = \lim_{t \rightarrow 1^-} [\ln |x-1|]_0^t \\ &= \lim_{t \rightarrow 1^-} \ln |t-1| - \ln 1 = -\infty \end{aligned}$$

- Therefore the integral diverges.

## Example

Evaluate  $\int_0^3 \frac{1}{x-1} dx$ .

Observe that  $x = 1$  is a vertical asymptote for the integrand.

$$\int_0^3 \frac{1}{x-1} dx = \int_0^1 \frac{1}{x-1} dx + \int_1^3 \frac{1}{x-1} dx$$

$$\begin{aligned} \int_0^1 \frac{dx}{x-1} &= \lim_{t \rightarrow 1^-} \int_0^t \frac{dx}{x-1} = \lim_{t \rightarrow 1^-} [\ln |x-1|]_0^t \\ &= \lim_{t \rightarrow 1^-} \ln |t-1| - \ln 1 = -\infty \end{aligned}$$

- Therefore the integral diverges.
- If we had not noticed the vertical asymptote, we might have made the following **mistake**:

$$\int_0^3 \frac{dx}{x-1} = [\ln |x-1|]_0^3 = \ln 2 - \ln 1 = \ln 2.$$




# A Comparison Test for Improper Integrals

Sometimes it's impossible to find the exact value of an integral, but we still want to know if it's convergent or divergent. For such cases, we can sometimes use the following theorem.

## Theorem (Comparison Theorem)

*Suppose  $f$  and  $g$  are continuous and  $f(x) \geq g(x) \geq 0$  for  $x \geq a$ .*

- 1 If  $\int_a^{\infty} f(x) dx$  is convergent, then  $\int_a^{\infty} g(x) dx$  is convergent.
- 2 If  $\int_a^{\infty} g(x) dx$  is divergent, then  $\int_a^{\infty} f(x) dx$  is divergent.




# A Comparison Test for Improper Integrals

Sometimes it's impossible to find the exact value of an integral, but we still want to know if it's convergent or divergent. For such cases, we can sometimes use the following theorem.

## Theorem (Comparison Theorem)

*Suppose  $f$  and  $g$  are continuous and  $f(x) \geq g(x) \geq 0$  for  $x \geq a$ .*

- 1 *If  $\int_a^\infty f(x)dx$  is convergent, then  $\int_a^\infty g(x)dx$  is convergent.*
- 2 *If  $\int_a^\infty g(x)dx$  is divergent, then  $\int_a^\infty f(x)dx$  is divergent.*



A similar theorem holds for Type II improper integrals.

## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.

## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.

- $$\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx.$$

## Example


Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.
- $\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx.$
- The first integral on the right hand side is a proper integral.

## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.


- The antiderivative isn't an elementary function.
- $\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx$ .
- The first integral on the right hand side is a proper integral.
- Notice that  $e^{-x^2} \leq e^{-x}$  for  $x \geq 1$ .



## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.
- $\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx$ .
- The first integral on the right hand side is a proper integral.
- Notice that  $e^{-x^2} \leq e^{-x}$  for  $x \geq 1$ .




$$\int_1^{\infty} e^{-x} dx = \lim_{t \rightarrow \infty} \int_1^t e^{-x} dx$$



## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.
- $\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx$ .
- The first integral on the right hand side is a proper integral.
- Notice that  $e^{-x^2} \leq e^{-x}$  for  $x \geq 1$ .




$$\begin{aligned}\int_1^{\infty} e^{-x} dx &= \lim_{t \rightarrow \infty} \int_1^t e^{-x} dx \\ &= \lim_{t \rightarrow \infty} [-e^{-x}]_1^t\end{aligned}$$

## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.
- $\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx$ .
- The first integral on the right hand side is a proper integral.
- Notice that  $e^{-x^2} \leq e^{-x}$  for  $x \geq 1$ .




$$\begin{aligned}\int_1^{\infty} e^{-x} dx &= \lim_{t \rightarrow \infty} \int_1^t e^{-x} dx \\ &= \lim_{t \rightarrow \infty} [-e^{-x}]_1^t \\ &= \lim_{t \rightarrow \infty} (e^{-1} - e^{-t})\end{aligned}$$

## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.
- $\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx$ .
- The first integral on the right hand side is a proper integral.
- Notice that  $e^{-x^2} \leq e^{-x}$  for  $x \geq 1$ .




$$\begin{aligned}\int_1^{\infty} e^{-x} dx &= \lim_{t \rightarrow \infty} \int_1^t e^{-x} dx \\ &= \lim_{t \rightarrow \infty} [-e^{-x}]_1^t \\ &= \lim_{t \rightarrow \infty} (e^{-1} - e^{-t}) \\ &= e^{-1}\end{aligned}$$

## Example

Show that  $\int_0^{\infty} e^{-x^2} dx$  is convergent.

- The antiderivative isn't an elementary function.
- $\int_0^{\infty} e^{-x^2} dx = \int_0^1 e^{-x^2} dx + \int_1^{\infty} e^{-x^2} dx.$
- The first integral on the right hand side is a proper integral.
- Notice that  $e^{-x^2} \leq e^{-x}$  for  $x \geq 1$ .



$$\begin{aligned}
 \int_1^{\infty} e^{-x} dx &= \lim_{t \rightarrow \infty} \int_1^t e^{-x} dx \\
 &= \lim_{t \rightarrow \infty} [-e^{-x}]_1^t \\
 &= \lim_{t \rightarrow \infty} (e^{-1} - e^{-t}) \\
 &= e^{-1}
 \end{aligned}$$

Therefore by the Comparison Theorem,  $\int_0^{\infty} e^{-x^2} dx$  converges.

## Example

Is  $\int_1^{\infty} \frac{1 + e^{-x}}{x} dx$  convergent or divergent?

## Example

Is  $\int_1^{\infty} \frac{1 + e^{-x}}{x} dx$  convergent or divergent?

- Notice that for  $x \geq 1$  we have  $\frac{1 + e^{-x}}{x} > \frac{1}{x}$ .

## Example

Is  $\int_1^{\infty} \frac{1 + e^{-x}}{x} dx$  convergent or divergent?

- Notice that for  $x \geq 1$  we have  $\frac{1 + e^{-x}}{x} > \frac{1}{x}$ .
- By a previously studied example,  $\int_1^{\infty} \frac{dx}{x}$  is divergent.


## Example

Is  $\int_1^{\infty} \frac{1 + e^{-x}}{x} dx$  convergent or divergent?

- Notice that for  $x \geq 1$  we have  $\frac{1 + e^{-x}}{x} > \frac{1}{x}$ .
- By a previously studied example,  $\int_1^{\infty} \frac{dx}{x}$  is divergent.
- Therefore  $\int_1^{\infty} \frac{1 + e^{-x}}{x} dx$  is divergent by the Comparison Theorem.




# Arc Length




- What do we mean by the length of a curve?

# Arc Length




- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.

# Arc Length




- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.

# Arc Length




- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.
- Then take the limit as the number of segments of the polygon goes to  $\infty$ .

# Arc Length




- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.
- Then take the limit as the number of segments of the polygon goes to  $\infty$ .

# Arc Length




- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.
- Then take the limit as the number of segments of the polygon goes to  $\infty$ .

# Arc Length



- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.
- Then take the limit as the number of segments of the polygon goes to  $\infty$ .


# Arc Length



- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.
- Then take the limit as the number of segments of the polygon goes to  $\infty$ .




# Arc Length




- What do we mean by the length of a curve?
- The length of a polygon is easy to compute: add up the length of the line segments that form the polygon.
- If the curve is a circle, approximate it by a polygon.
- Then take the limit as the number of segments of the polygon goes to  $\infty$ .


- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?




- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .




- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .




- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .
- The length  $L$  of the curve  $y = f(x)$  is the limit of the lengths of these segments as  $n \rightarrow \infty$ .




- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .
- The length  $L$  of the curve  $y = f(x)$  is the limit of the lengths of these segments as  $n \rightarrow \infty$ .



- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .
- The length  $L$  of the curve  $y = f(x)$  is the limit of the lengths of these segments as  $n \rightarrow \infty$ .




- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .
- The length  $L$  of the curve  $y = f(x)$  is the limit of the lengths of these segments as  $n \rightarrow \infty$ .







- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .
- The length  $L$  of the curve  $y = f(x)$  is the limit of the lengths of these segments as  $n \rightarrow \infty$ .



- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .
- The length  $L$  of the curve  $y = f(x)$  is the limit of the lengths of these segments as  $n \rightarrow \infty$ .



- What about  $y = f(x)$  for a continuous function  $f$  on  $[a, b]$ ?
- Divide  $[a, b]$  into  $n$  subintervals with endpoints  $x_0, x_1, \dots, x_n$  and equal width  $\Delta x$ .
- The points  $P_i = (x_i, f(x_i))$  lie on  $y = f(x)$ , and the segments with vertices  $P_0, P_1, \dots, P_n$  are an approximation to  $y = f(x)$ .
- The length  $L$  of the curve  $y = f(x)$  is the limit of the lengths of these segments as  $n \rightarrow \infty$ .



$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- Mean Value Theorem: there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- Mean Value Theorem: there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .



$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- **Then**  $|P_i P_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- **Mean Value Theorem:** there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$|P_{i-1}P_i| = \sqrt{(\Delta x)^2 + (\Delta y)^2}$$

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- Mean Value Theorem: there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$|P_{i-1}P_i| = \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2}$$

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- **Mean Value Theorem:** there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$\begin{aligned}|P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2} \\ &= \sqrt{1 + (f'(x_i^*))^2} \sqrt{(\Delta x)^2}\end{aligned}$$

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i|$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- **Mean Value Theorem:** there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$\begin{aligned} |P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2} \\ &= \sqrt{1 + (f'(x_i^*))^2} \sqrt{(\Delta x)^2} = \sqrt{1 + (f'(x_i^*))^2} \Delta x \end{aligned}$$

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i| = \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{1 + (f'(x_i^*))^2} \Delta x$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_i P_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- Mean Value Theorem: there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$\begin{aligned} |P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2} \\ &= \sqrt{1 + (f'(x_i^*))^2} \sqrt{(\Delta x)^2} = \sqrt{1 + (f'(x_i^*))^2} \Delta x \end{aligned}$$

$$\begin{aligned} L &= \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i| &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{1 + (f'(x_i^*))^2} \Delta x \\ & &= \int_a^b \sqrt{1 + (f'(x))^2} dx \end{aligned}$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- **Mean Value Theorem:** there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$\begin{aligned} |P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2} \\ &= \sqrt{1 + (f'(x_i^*))^2} \sqrt{(\Delta x)^2} = \sqrt{1 + (f'(x_i^*))^2} \Delta x \end{aligned}$$

$$\begin{aligned}
 L &= \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i| &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{1 + (f'(x_i^*))^2} \Delta x \\
 & &= \int_a^b \sqrt{1 + (f'(x))^2} dx
 \end{aligned}$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- Mean Value Theorem: there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$\begin{aligned}
 |P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2} \\
 &= \sqrt{1 + (f'(x_i^*))^2} \sqrt{(\Delta x)^2} = \sqrt{1 + (f'(x_i^*))^2} \Delta x
 \end{aligned}$$

$$\begin{aligned}
 L &= \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i| &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{1 + (f'(x_i^*))^2} \Delta x \\
 & &= \int_a^b \sqrt{1 + (f'(x))^2} dx
 \end{aligned}$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_iP_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- Mean Value Theorem: there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$\begin{aligned}
 |P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2} \\
 &= \sqrt{1 + (f'(x_i^*))^2} \sqrt{(\Delta x)^2} = \sqrt{1 + (f'(x_i^*))^2} \Delta x
 \end{aligned}$$



$$\begin{aligned}
 L &= \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}P_i| &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{1 + (f'(x_i^*))^2} \, \Delta x \\
 & &= \int_a^b \sqrt{1 + (f'(x))^2} \, dx
 \end{aligned}$$

- This formula is not useful for computational purposes.
- We can find a better formula when  $f$  has a continuous derivative.
- Let  $y_i = f(x_i)$ , and  $\Delta y = y_i - y_{i-1} = f(x_i) - f(x_{i-1})$ .
- Then  $|P_i P_{i-1}| = \sqrt{(x_i - x_{i-1})^2 + (y_i - y_{i-1})^2} = \sqrt{(\Delta x)^2 + (\Delta y)^2}$ .
- **Mean Value Theorem:** there exists  $x_i^*$  between  $x_{i-1}$  and  $x_i$  such that  $f(x_i) - f(x_{i-1}) = f'(x_i^*)(x_i - x_{i-1})$ .
- $\Delta y = f'(x_i^*)\Delta x$ .

$$\begin{aligned}
 |P_{i-1}P_i| &= \sqrt{(\Delta x)^2 + (\Delta y)^2} = \sqrt{(\Delta x)^2 + (f'(x_i^*)\Delta x)^2} \\
 &= \sqrt{1 + (f'(x_i^*))^2} \sqrt{(\Delta x)^2} = \sqrt{1 + (f'(x_i^*))^2} \, \Delta x
 \end{aligned}$$

# The Arc Length Formula


## Definition

Suppose  $f'$  exists and is continuous on  $[a, b]$ . Then the length of the curve  $y = f(x)$ ,  $a \leq x \leq b$ , is

$$\begin{aligned} L &= \int_a^b \sqrt{1 + (f'(x))^2} \, dx \\ &= \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx \quad (\text{in Leibniz notation}) \quad . \end{aligned}$$


## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .



## Example


Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .



- For the top half of the curve we have:
- $y =$       and  $y' =$       .

## Example


Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .



- For the top half of the curve we have:
- $y =$       and  $y' =$       .

## Example


Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .



- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' =$  .

## Example


Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .



- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' =$  .

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .



## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .

$$L = \int_1^4 \sqrt{1 + (y')^2} dx$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u =$                       and  $dx =$                       .
- When  $x = 1$ ,  $u =$                       .
- When  $x = 4$ ,  $u =$                       .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u =$  and  $dx =$  .
- When  $x = 1$ ,  $u =$  .
- When  $x = 4$ ,  $u =$  .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx =$  .
- When  $x = 1$ ,  $u =$  .
- When  $x = 4$ ,  $u =$  .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int \sqrt{u} \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx =$  .
- When  $x = 1$ ,  $u =$  .
- When  $x = 4$ ,  $u =$  .

$$\begin{aligned}
 L &= \int_1^4 \sqrt{1 + (y')^2} dx \\
 &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int \sqrt{u}
 \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $\textcolor{red}{dx} = \frac{4}{9}\textcolor{red}{du}$ .
- When  $x = 1$ ,  $u =$  .
- When  $x = 4$ ,  $u =$  .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} \textcolor{red}{dx} = \int \textcolor{red}{\frac{4}{9}} \sqrt{u} \textcolor{red}{du} \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx = \frac{4}{9}du$ .
- When  $x = 1$ ,  $u =$  .
- When  $x = 4$ ,  $u =$  .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int \frac{4}{9} \sqrt{u} du \end{aligned}$$



## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx = \frac{4}{9}du$ .
- When  $x = 1$ ,  $u = \frac{13}{4}$ .
- When  $x = 4$ ,  $u =$  .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int_{13/4} \frac{4}{9} \sqrt{u} du \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx = \frac{4}{9}du$ .
- When  $x = 1$ ,  $u = \frac{13}{4}$ .
- When  $x = 4$ ,  $u = \quad$ .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int_{13/4}^{\quad} \frac{4}{9} \sqrt{u} du \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx = \frac{4}{9}du$ .
- When  $x = 1$ ,  $u = \frac{13}{4}$ .
- **When  $x = 4$ ,  $u = 10$ .**

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int_{13/4}^{10} \frac{4}{9} \sqrt{u} du \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .




- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx = \frac{4}{9}du$ .
- When  $x = 1$ ,  $u = \frac{13}{4}$ .
- When  $x = 4$ ,  $u = 10$ .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int_{13/4}^{10} \frac{4}{9} \sqrt{u} du \\ &= \frac{4}{9} \left[ \frac{2}{3} u^{3/2} \right]_{13/4}^{10} \end{aligned}$$

## Example

Find the length of the arc of  $y^2 = x^3$  between  $(1, 1)$  and  $(4, 8)$ .



- For the top half of the curve we have:
- $y = x^{3/2}$  and  $y' = \frac{3}{2}x^{1/2}$ .
- $u = 1 + \frac{9}{4}x$  and  $dx = \frac{4}{9}du$ .
- When  $x = 1$ ,  $u = \frac{13}{4}$ .
- When  $x = 4$ ,  $u = 10$ .

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + (y')^2} dx \\ &= \int_1^4 \sqrt{1 + \frac{9}{4}x} dx = \int_{13/4}^{10} \frac{4}{9} \sqrt{u} du \\ &= \frac{4}{9} \left[ \frac{2}{3} u^{3/2} \right]_{13/4}^{10} = \frac{8}{27} \left( 10^{3/2} - \left( \frac{13}{4} \right)^{3/2} \right) \end{aligned}$$

If a curve has equation  $x = g(y)$ ,  $c \leq y \leq d$ , and  $g'(y)$  is continuous, then we can get the length of the curve by interchanging the roles of  $x$  and  $y$  in the arc length formula:

$$L = \int_c^d \sqrt{1 + (g'(y))^2} \, dy = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} \, dy$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

$$L = \int_0^1 \sqrt{1 + (dx/dy)^2} dy$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy =$  .

$$L = \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + } dy$$



## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy =$  .

$$L = \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + } dy$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .

$$L = \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y =$  , so  $dy =$  , and  $\sqrt{1 + 4y^2} =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} \, dy = \int_0^1 \sqrt{1 + 4y^2} \, dy \\ &= \int \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y =$  , so  $dy =$  , and  $\sqrt{1 + 4y^2} =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} \, dy = \int_0^1 \sqrt{1 + 4y^2} \, dy \\ &= \int \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy =$  , and  $\sqrt{1 + 4y^2} =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} \, dy = \int_0^1 \sqrt{1 + 4y^2} \, dy \\ &= \int \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy =$  , and  $\sqrt{1 + 4y^2} =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$



## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 1$ , so  $\theta = \frac{\pi}{4}$ .

$$\begin{aligned}
 L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\
 &= \int \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta
 \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta =$  .
- When  $y = 1$ ,  $\tan \theta =$  , so  $\theta =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 1$ , so  $\theta = \frac{\pi}{4}$ .

$$\begin{aligned}
 L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\
 &= \int \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta
 \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \arctan 2$ .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^{\arctan 2} \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \arctan 2$ .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^{\arctan 2} \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta \\ &= \frac{1}{2} \int_0^{\arctan 2} \sec^3 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta =$  .

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^{\theta} \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta =$

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^{\theta} \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$



## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^{\alpha} \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^\alpha \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^\alpha \sec^3 \theta d\theta \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^\alpha \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^\alpha \sec^3 \theta d\theta \\ &= \frac{1}{2} \cdot \frac{1}{2} [\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|]_0^\alpha \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^\alpha \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^\alpha \sec^3 \theta d\theta \\ &= \frac{1}{2} \cdot \frac{1}{2} [\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|]_0^\alpha \\ &= \frac{1}{4} (\sec \alpha \tan \alpha + \ln |\sec \alpha + \tan \alpha|) \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned}
 L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\
 &= \int_0^\alpha \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^\alpha \sec^3 \theta d\theta \\
 &= \frac{1}{2} \cdot \frac{1}{2} [\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|]_0^\alpha \\
 &= \frac{1}{4} (\sec \alpha \tan \alpha + \ln |\sec \alpha + \tan \alpha|) \\
 &= \frac{1}{4} \left( \quad + \ln | \quad + | \right)
 \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned}
 L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\
 &= \int_0^\alpha \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^\alpha \sec^3 \theta d\theta \\
 &= \frac{1}{2} \cdot \frac{1}{2} [\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|]_0^\alpha \\
 &= \frac{1}{4} (\sec \alpha \tan \alpha + \ln |\sec \alpha + \tan \alpha|) \\
 &= \frac{1}{4} (2 + \ln |2 + 2|)
 \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned}
 L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\
 &= \int_0^\alpha \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^\alpha \sec^3 \theta d\theta \\
 &= \frac{1}{2} \cdot \frac{1}{2} [\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|]_0^\alpha \\
 &= \frac{1}{4} (\sec \alpha \tan \alpha + \ln |\sec \alpha + \tan \alpha|) \\
 &= \frac{1}{4} (2 + \ln |2 + 2|)
 \end{aligned}$$

## Example

Find the length of the arc of  $x = y^2$  from  $(0, 0)$  to  $(1, 1)$ .

- $x = y^2$ , so  $dx/dy = 2y$ .
- Substitute  $y = \frac{1}{2} \tan \theta$ , so  $dy = \frac{1}{2} \sec^2 \theta d\theta$ , and  $\sqrt{1 + 4y^2} = \sec \theta$ .
- When  $y = 0$ ,  $\tan \theta = 0$ , so  $\theta = 0$ .
- When  $y = 1$ ,  $\tan \theta = 2$ , so  $\theta = \tan^{-1}(2)$  (call this  $\alpha$ ).

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + (dx/dy)^2} dy = \int_0^1 \sqrt{1 + 4y^2} dy \\ &= \int_0^\alpha \sec \theta \cdot \frac{1}{2} \sec^2 \theta d\theta = \frac{1}{2} \int_0^\alpha \sec^3 \theta d\theta \\ &= \frac{1}{2} \cdot \frac{1}{2} [\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|]_0^\alpha \\ &= \frac{1}{4} (\sec \alpha \tan \alpha + \ln |\sec \alpha + \tan \alpha|) \\ &= \frac{1}{4} (2\sqrt{5} + \ln |\sqrt{5} + 2|) \end{aligned}$$



Sometimes an arc length problem can be simplified significantly if the function  $y = f(x)$  has a certain form.

In particular, if  $y' = a - b$ , where  $2ab = \frac{1}{2}$ , then  $1 + (y')^2$  is a perfect square, simplifying the arc length formula.

$$\begin{aligned}1 + (y')^2 &= 1 + (a - b)^2 \\&= 1 + \left(a^2 - \frac{1}{2} + b^2\right) \\&= a^2 + \frac{1}{2} + b^2 \\&= a^2 + 2ab + b^2 \\&= (a + b)^2.\end{aligned}$$

There are several exercises like this in the textbook (such as numbers 9 and 10) and on WebWork (question 2 from this section) of this type.

Consider the following examples:

Example: Find the arc length of  $y = \frac{x^7}{14} + \frac{1}{10x^5}$   
between  $x = 1$  and  $x = 2$ .

**Solution:** Arc length formula:  $L = \int_1^2 \sqrt{1 + (y')^2} dx$ , so first find  $y'$ .

$$\begin{aligned} y' &= 7\frac{x^6}{14} + (-5)\frac{1}{10x^6} \\ &= \frac{1}{2}x^6 - \frac{1}{2}x^{-6} \end{aligned}$$

Example: Find the arc length of  $y = \frac{x^7}{14} + \frac{1}{10x^5}$   
between  $x = 1$  and  $x = 2$ .

**Solution:** Arc length formula:  $L = \int_1^2 \sqrt{1 + (y')^2} dx$ , so first find  $y'$ .

$$\begin{aligned}y' &= 7\frac{x^6}{14} + (-5)\frac{1}{10x^6} \\ &= \frac{1}{2}x^6 - \frac{1}{2}x^{-6}\end{aligned}$$

Observe that if  $a = \frac{1}{2}x^6$  and  $b = \frac{1}{2}x^{-6}$ , then  $y' = a - b$ , and  
 $2ab = 2 \left(\frac{1}{2}x^6\right) \left(\frac{1}{2}x^{-6}\right) = \frac{1}{2}$ , so

$$1 + (y')^2 = \left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right)^2$$

Now apply this to the arc length formula:

$$L = \int_1^2 \sqrt{1 + (y')^2} dx = \int_1^2 \sqrt{\left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right)^2} dx$$

$$\begin{aligned} L &= \int_1^2 \sqrt{1 + (y')^2} dx = \int_1^2 \sqrt{\left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right)^2} dx \\ &= \int_1^2 \left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right) dx \end{aligned}$$

$$\begin{aligned} L &= \int_1^2 \sqrt{1 + (y')^2} \, dx = \int_1^2 \sqrt{\left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right)^2} \, dx \\ &= \int_1^2 \left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right) \, dx \\ &= \left[\frac{1}{2} \cdot \frac{1}{7}x^7 + \frac{1}{2} \cdot \frac{1}{-5}x^{-5}\right]_1^2 \end{aligned}$$


$$\begin{aligned} L &= \int_1^2 \sqrt{1 + (y')^2} dx = \int_1^2 \sqrt{\left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right)^2} dx \\ &= \int_1^2 \left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right) dx \\ &= \left[\frac{1}{2} \cdot \frac{1}{7}x^7 + \frac{1}{2} \cdot \frac{1}{-5}x^{-5}\right]_1^2 \\ &= \left[\frac{1}{14}x^7 - \frac{1}{10}x^{-5}\right]_1^2 \end{aligned}$$

$$\begin{aligned} L &= \int_1^2 \sqrt{1 + (y')^2} \, dx = \int_1^2 \sqrt{\left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right)^2} \, dx \\ &= \int_1^2 \left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right) \, dx \\ &= \left[\frac{1}{2} \cdot \frac{1}{7}x^7 + \frac{1}{2} \cdot \frac{1}{-5}x^{-5}\right]_1^2 \\ &= \left[\frac{1}{14}x^7 - \frac{1}{10}x^{-5}\right]_1^2 \\ &= \left(\frac{128}{14} - \frac{1}{10} \cdot \frac{1}{32}\right) - \left(\frac{1}{14} - \frac{1}{10}\right) \end{aligned}$$



$$\begin{aligned} L &= \int_1^2 \sqrt{1 + (y')^2} \, dx = \int_1^2 \sqrt{\left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right)^2} \, dx \\ &= \int_1^2 \left(\frac{1}{2}x^6 + \frac{1}{2}x^{-6}\right) \, dx \\ &= \left[\frac{1}{2} \cdot \frac{1}{7}x^7 + \frac{1}{2} \cdot \frac{1}{-5}x^{-5}\right]_1^2 \\ &= \left[\frac{1}{14}x^7 - \frac{1}{10}x^{-5}\right]_1^2 \\ &= \left(\frac{128}{14} - \frac{1}{10} \cdot \frac{1}{32}\right) - \left(\frac{1}{14} - \frac{1}{10}\right) = \frac{20537}{2240}. \end{aligned}$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)




Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' =$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)




$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)




$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx$$

$$= \int_0^1 \left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right) dx$$



Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx$$

$$= \int_0^1 \left( \frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x} \right) dx = \left[ \quad \quad \quad \right]_0^1$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx$$

$$= \int_0^1 \left( \frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x} \right) dx = \left[ \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x} \right]_0^1$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)




$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx$$

$$= \int_0^1 \left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right) dx = \left[\frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}\right]_0^1$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx$$

$$= \int_0^1 \left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right) dx = \left[\frac{1}{6}e^{3x} - \frac{1}{6}e^{-3x}\right]_0^1$$

Example (Find the length of the arc of  $y = \frac{1}{6}e^{3x} + \frac{1}{6}e^{-3x}$  from  $x = 0$  to  $x = 1$ .)



$$y' = \frac{1}{2}e^{3x} - \frac{1}{2}e^{-3x}.$$

$$= a - b \text{ where } a = \frac{1}{2}e^{3x}, b = \frac{1}{2}e^{-3x}, \text{ and } 2ab = \frac{1}{2}$$

$$L = \int_0^1 \sqrt{1 + (y')^2} dx = \int_0^1 \sqrt{\left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right)^2} dx$$

$$= \int_0^1 \left(\frac{1}{2}e^{3x} + \frac{1}{2}e^{-3x}\right) dx = \left[\frac{1}{6}e^{3x} - \frac{1}{6}e^{-3x}\right]_0^1 = \frac{e^3 - e^{-3}}{6}.$$