


Starter



Find the general solution to:

1.
$$\frac{dy}{dx} + 2y = x$$

2.
$$\ddot{x} + \dot{x} - 6x = 0$$

3. Find the particular solution to

$$\frac{d^2z}{dt^2} - 9z = 0, \text{ given } \left. \frac{dz}{dt} \right|_{t=0} = 16, z|_{t=0} = 4$$

Solutions

1. IF = e^{2x} , $\therefore e^{2x} \frac{dy}{dx} + 2e^{2x}y = xe^{2x}$

$$e^{2x}y = \int xe^{2x} dx = \frac{1}{2}xe^{2x} - \int \frac{1}{2}e^{2x} dx$$

$$= \frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} + c$$

$$\therefore y = \frac{1}{2}x - \frac{1}{4} + ce^{-2x}$$

Solutions

$$2. \text{ AE} = m^2 + m - 6 = 0, m \in \{2, -3\},$$

$$\therefore x = Ae^{2t} + Be^{-3t}$$

Solutions

$$3. \quad AE = m^2 - 9 = 0, \quad m = 3, \quad \therefore x = Ae^{3t} + Bte^{3t}$$

$$\dot{x} = 3Ae^{3t} + Be^{3t} + 3Bte^{3t}$$

$$x|_{t=0} = 4 = A + 0 \Rightarrow A = 4$$

$$\dot{x}|_{t=0} = 12 = 3A + B + 0 \Rightarrow B = 4$$

$$\therefore x = 4e^{3t} + 4te^{3t}$$


Differential Equations 2

2nd Order Homogeneous 2

Purely Imaginary Roots

Find the general solution to

$$\frac{d^2y}{dx^2} + 9y = 0$$




Purely Imaginary Roots

Find the general solution to

$$\frac{d^2y}{dx^2} + 9y = 0$$

$$AE = m^2 + 9 = 0 \Rightarrow m = \pm 3i$$



Purely Imaginary Roots

Find the general solution to

$$\frac{d^2y}{dx^2} + 9y = 0$$

$$AE = m^2 + 9 = 0 \Rightarrow m = \pm 3i$$


$$y = Ae^{3ix} + Be^{-3ix}$$

$$= A(\cos 3x + i \sin 3x) + B(\cos -3x + i \sin -3x)$$

$$= A(\cos 3x + i \sin 3x) + B(\cos 3x - i \sin 3x)$$

$$= (A + B) \cos 3x + (A - B) i \sin 3x$$

$$= C \cos 3x + D \sin 3x$$



Purely Imaginary roots in General


Show that the general solution to

$$\frac{d^2y}{dx^2} + \omega^2 y = 0$$

can be written in the form


$$y = A \cos \omega x + B \sin \omega x$$

where A, B, ω are real numbers.



Complex roots


Can we extend this to a complex root, $\lambda \pm \mu i$?



Complex roots

Can we extend this to a complex root, $\lambda \pm \mu i$?


$$\begin{aligned} y &= Ae^{(\lambda + \mu i)x} + Be^{(\lambda - \mu i)x} \\ &= Ae^{\lambda x} e^{\mu i x} + Be^{\lambda x} e^{-\mu i x} \\ &= e^{\lambda x} (Ae^{\mu i x} + Be^{-\mu i x}) \\ &= e^{\lambda x} (C \cos \mu x + D \sin \mu x) \end{aligned}$$



Sketching the solutions

It's natural that when we look at either of the two solution forms, we want to know what they look like!

Harmonic forms from A Level Maths can give us an answer.





Sketching the solutions

It's natural that when we look at either of the two solution forms, we want to know what they look like!

Harmonic forms from A Level Maths can give us an answer.

$$y = A \cos \mu x + B \sin \mu x \equiv R \cos(x + \alpha)$$





Sketching the solutions

It's natural that when we look at either of the two solution forms, we want to know what they look like!

Harmonic forms from A Level Maths can give us an answer.

$$y = e^{\lambda x} (A \cos \mu x + B \sin \mu x) \equiv R e^{\lambda x} \cos(x + \alpha)$$




Finding a Particular Solution

Find and sketch the particular solution to

$$\ddot{x} - 2\dot{x} + 2x = 0$$

$$\text{Initially } x = 1, \dot{x} = 2$$



Finding a Particular Solution

Find and sketch the particular solution to

$$\ddot{x} - 2\dot{x} + 2x = 0$$

$$\text{Initially } x = 1, \dot{x} = 2$$

$$\text{AE: } m^2 - 2m + 2 = 0 \Rightarrow m = 1 \pm i$$


$$\therefore x = e^t (A \cos t + B \sin t)$$

$$\therefore \dot{x} = e^t (A \cos t + B \sin t) + e^t (B \cos t - A \sin t)$$

$$x|_{t=0} = 1 = e^0 (A \cos 0 + B \sin 0) \Rightarrow A = 1$$

$$\dot{x}|_{t=0} = 2 = e^0 (A \cos 0 + B \sin 0) + e^0 (B \cos 0 - A \sin 0) \Rightarrow B = 1$$

$$\therefore x = e^t (\cos t + \sin t) = \sqrt{2}e^t \cos\left(t + \frac{\pi}{4}\right)$$





Finding a Particular Solution


Find and sketch the particular solution to

$$\ddot{x} - 2\dot{x} + 2x = 0$$

Initially $x = 1, \dot{x} = 2$



Recall



Given the equation

$$a \frac{d^2 y}{dx^2} + b \frac{dy}{dx} + cy = 0$$

state the four possible forms the general solution may take, and the conditions under which it will take them.

General Solutions

If the auxiliary equation, $am^2 + bm + c = 0$ have:

- Two distinct real roots, λ, μ :

$$y = Ae^{\lambda x} + Be^{\mu x}$$

- Repeated real roots, λ :


$$y = Ae^{\lambda x} + Bxe^{\lambda x}$$


- Purely imaginary roots, $\pm \omega i$:

$$y = A \cos \omega x + B \sin \omega x \text{ or } R \cos(\omega x + \alpha)$$

- Complex roots, $\alpha \pm \beta i$:

$$y = e^{\alpha x} (A \cos \beta x + B \sin \beta x)$$






Practice! (4s.f. where appropriate)

1. Find the general solution to the equation $\frac{d^2x}{dt^2} + 18\frac{dx}{dt} + 81x = 0$.
2. Find the general solution to the equation $\frac{d^2y}{dx^2} + 20\frac{dy}{dx} + 100y = 0$.
3. Find the general solution to the equation $5\ddot{x} + 3x = 0$.
4. Find the general solution to the equation $\ddot{x} + 18\dot{x} + 81x = 0$.
5. Find the general solution to the equation $5\frac{d^2p}{dq^2} + 3\frac{dp}{dq} + 10p = 0$.
6. Find the general solution to the equation $\frac{d^2y}{dx^2} + 6\frac{dy}{dx} + 8y = 0$.

Practice! (4s.f. where appropriate)




1. $x = Ae^{-9t} + Bte^{-9t}$
2. $y = Ae^{-10x} + Bxe^{-10x}$
3. $x = A \sin(0.7746t) + B \cos(0.7746t)$
4. $x = Ae^{-9t} + Bte^{-9t}$
5. $p = e^{-\frac{3}{10}q} (A \sin(1.382q) + B \cos(1.382q))$
6. $y = Ae^{-2x} + Be^{-4x}$

Damping

These second order DEs often describe real-world events. If you think about a spring, for instance, its acceleration is based on the distance from its resting position, but so is its velocity.

Many such systems attempt to reach an equilibrium position.

EG:



Damping


These second order DEs often describe real-world events. If you think about a spring, for instance, its acceleration is based on the distance from its rest position.

Many such

on.

EG:

Moving Examples! 




Damping

When discussing these solutions, we consider **under-damped**, **over-damped** and **critically damped** systems.

The DEs all have particular solutions when these occur:

- **under-damped**: auxiliary equation has complex roots.




Damping


When discussing these solutions, we consider **under-damped**, **over-damped** and **critically damped** systems.

The DEs all have particular solutions when these occur:

- **under-damped**: auxiliary equation has complex roots.
- **over-damped**: auxiliary equation has distinct real roots.



Damping




When discussing these solutions, we consider **under-damped**, **over-damped** and **critically damped** systems.

The DEs all have particular solutions when these occur:

- **under-damped**: auxiliary equation has complex roots.
- **over-damped**: auxiliary equation has distinct real roots.
- **critically damped**: auxiliary equation has repeated negative roots.

Damping



When discussing these solutions, we consider **under-damped**, **over-damped** and **critically damped** systems.

The DEs all have particular solutions when these occur:

- **under-damped**: auxiliary equation has complex roots.
- **over-damped**: auxiliary equation has distinct real roots.
- **critically damped**: auxiliary equation has repeated negative roots.
- **undamped** systems are when the AE roots are purely imaginary (Simple Harmonic Motion).