

Sea detuning sweep report


Global parameters (constant across sweep):

f_Az (sea Larmor)	= 700.000 Hz
f_Rz (rare Larmor)	= 1100.000 Hz
f1A (sea Rabi)	= 100.000 Hz
f1R (rare Rabi)	= 50.000 Hz
gamma_sea	= 4.200
gamma_rare	= 6.600
B0_common	= 1.047e+03
B1_sea	= 1.496e+02
B1_rare	= 4.760e+01
t_final	= 1.000 s
steps	= 2000
n_sea	= 12
phi_sea	= 0.000 rad
phi_rare	= 0.000 rad


Sea detunings ($\Delta_A = f_{Az} - f_{rf,A}$) in Hz:

-300.0, -200.0, -100.0, +0.0, +100.0, +200.0, +300.0


$\Delta_A = -300.0$ Hz




$\Delta_A = -300.0$ Hz (with T-fit overlays)




$\Delta_A = -300.0$ Hz (rare drive OFF)




$\Delta_A = -200.0$ Hz




$\Delta_A = -200.0$ Hz (with T-fit overlays)




$\Delta_A = -200.0$ Hz (rare drive OFF)




$\Delta_A = -100.0$ Hz




$\Delta_A = -100.0$ Hz (with T-fit overlays)




$\Delta_A = -100.0$ Hz (rare drive OFF)




$\Delta_A = +0.0$ Hz




$\Delta_A = +0.0$ Hz (with T-fit overlays)




$\Delta_A = +0.0$ Hz (rare drive OFF)




$\Delta_A = +100.0$ Hz




$\Delta_A = +100.0$ Hz (with T-fit overlays)




$\Delta_A = +100.0$ Hz (rare drive OFF)




$\Delta_A = +200.0$ Hz




$\Delta_A = +200.0$ Hz (with T-fit overlays)




$\Delta_A = +200.0$ Hz (rare drive OFF)




$\Delta_A = +300.0$ Hz



$\Delta_A = +300.0$ Hz (with T-fit overlays)



$\Delta_A = +300.0$ Hz (rare drive OFF)



T-like decay fits (from Ix traces)

delta_Hz	T_Ix_sea_off	T_Ix_R_off	T_Ix_sea_on	T_Ix_R_on
-300.0	NA	NA	NA	5.3
-200.0	3.58	NA	4.19	6.42
-100.0	5.96	16.7	NA	NA
+0.0	8.09	NA	2.11	29.9
+100.0	5.96	16.7	NA	NA
+200.0	3.58	NA	4.17	6.41
+300.0	NA	NA	NA	5.29