

Faraday Wave Experiment

1 Introduction

Bose-Einstein Condensation (BEC) is a state of matter in boson gases that results when atoms, cooled very close to absolute zero, abruptly accumulate in their ground state [9]. This state of matter has been experimentally realized using various atomic vapors [8]. In particular, a laboratory here at Rice led by Professor Randall Hulet has studied Bose-Einstein Condensation in lithium. Professor Hulet's lab is able to tune the interaction strength between atoms by exploiting a shallow zero-crossing of a Feshbach resonance, and, due to nonlinearity, this can result in rich dynamics such as the creation of matter-wave soliton trains [7]. Because it is difficult to study nonlinear systems analytically, computational tools have been important for understanding the behavior of these systems.

1.1 Project Overview

More recently, Professor Hulet's lab has performed experiments on the emergence of Faraday Waves in a Bose-Einstein Condensate by periodically modulating the scattering length for these atoms. In these experiments, they have modulated the scattering length periodically for a fixed period of time. Then after holding the scattering length constant, they observe that perturbations of particular wavenumbers will grow rapidly, resulting in the emergence of faraday waves.

Using a mean field theory, it can be shown that the dynamics of a Bose-Einstein condensate are governed by the time-dependent Gross-Pitaevskii Equation [8]

$$i\hbar \frac{\partial \psi(\mathbf{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{r}, t) + V(\mathbf{r})\psi(\mathbf{r}, t) + g|\psi(\mathbf{r}, t)|^2\psi(\mathbf{r}, t), \quad g \equiv 4\pi\hbar^2 a_s/m \quad (1)$$

where ψ is a field describing the number density of atoms in real space, hence the total particle number N satisfies

$$\int_{\mathbb{R}^3} d\mathbf{r} |\psi(\mathbf{r})|^2 = N \quad (2)$$

The parameter g corresponds to the interaction strength between atoms ($g > 0$ corresponding to repulsive interactions and $g < 0$ corresponding to attractive interactions), and is defined in terms of the scattering length a_s and the atomic mass m .

In this project, I numerically solve this nonlinear, partial differential equation assuming a cigar-shaped trapping potential in three spatial dimensions to computationally confirm recent experimental results from Professor Hulet's lab by showing that the BEC's perturbations in Fourier space grow in the expected way.

2 Methods

2.1 Dimensionless GPE

Though the form given for the Gross-Pitaevskii equation in (1) is physically correct, the units in which the equation has been expressed are not appropriate for the time and length scale of our problem. To remedy this, we will first scale the field

$$\frac{\psi}{\sqrt{N}} \rightarrow \psi \quad (3)$$

which now gives the PDE

$$i\hbar \frac{\partial \psi(\mathbf{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{r}, t) + V(\mathbf{r})\psi(\mathbf{r}, t) + gN|\psi(\mathbf{r}, t)|^2\psi(\mathbf{r}, t), \quad g \equiv 4\pi\hbar^2 a_s/m \quad (4)$$

with a normalization condition that is set to unity

$$\int_{\mathbb{R}^3} d\mathbf{r} |\psi(\mathbf{r})|^2 = 1 \quad (5)$$

To make the GPE dimensionless, we stipulate that the potential is harmonic

$$V(\mathbf{r}) = \frac{m}{2} (\omega_x^2 x^2 + \omega_y^2 y^2 + \omega_z^2 z^2) \quad (6)$$

and then choose a convenient characteristic length a_0 and scale our coordinates as follows

$$\omega_x t \rightarrow t, \quad \mathbf{r}/a_0 \rightarrow \mathbf{r}, \quad a_0^{3/2} \psi \rightarrow \psi \quad \text{where} \quad a_0 \equiv \sqrt{\frac{\hbar}{m\omega_x}} \quad (7)$$

Plugging (7) into (4) and simplifying we find the GPE in dimensionless form can be written as

$$i \frac{\partial \psi(\mathbf{r}, t)}{\partial t} = -\frac{1}{2} \nabla^2 \psi(\mathbf{r}, t) + V(\mathbf{r})\psi(\mathbf{r}, t) + gN|\psi(\mathbf{r}, t)|^2\psi(\mathbf{r}, t), \quad g \equiv 4\pi a_s \quad (8)$$

where the potential has been scaled so that one of the trapping frequencies is unity

$$V(\mathbf{r}) = \frac{1}{2} (x^2 + \gamma_y^2 y^2 + \gamma_z^2 z^2), \quad \gamma_y \equiv \frac{\omega_y}{\omega_x}, \quad \gamma_z \equiv \frac{\omega_z}{\omega_x}$$

For the Faraday Wave experiment that we will be considering, the trapping potential is cylindrically symmetric with $\omega_x = \omega_y$. We thus define ω_ρ as the trapping frequency along the x-axis and y-axis (henceforth called the radial direction), thus $\omega_\rho = \omega_x = \omega_y$. So the trapping potential is really

$$V(\mathbf{r}) = \frac{1}{2} (\rho^2 + \gamma_z^2 z^2), \quad \rho^2 = x^2 + y^2, \quad \gamma_z \equiv \frac{\omega_z}{\omega_\rho}. \quad (9)$$

2.2 Numerical Methods

To solve this nonlinear PDE, I use a time-splitting spectral (TSSP) method while assuming periodic boundary conditions. The general idea behind applying this method is that some operators on the RHS of (8) are easy to apply in momentum space (e.g., the second order derivative) and some are easy to apply in position space (e.g., the potential term). To exploit this fact, this method involves shifting the wave function solution between its real space and momentum space representation via fourier transformation and applying each operator component in the space it is most easily applied.

Before I can start solving the nonlinear PDE, however, I must find the correct initial condition. These experiments in Hulet's lab are performed on BEC that is very close to its ground state. The ground state for the GPE is not known analytically in this trapping potential, so I will use finite difference methods to implement the imaginary-time propagation method [5, 6], which will allow me to numerically calculate the ground state.

The TSSP method is described in detail in [3] for a one dimensional version of the GPE, and the method can be straightforwardly generalized to three spatial dimensions without affecting the method's validity. The advantages of this method are multifold: the TSSP method is unconditionally stable, time reversible, time-transverse invariant, conserves particle number, and is second order accurate in space and time [3]. One noteworthy drawback is that this method is not symplectic, i.e., it does not conserve energy.

Time reversible means that if the solution ψ_{n+1} at t_{n+1} is obtained by applying the TSSP method to the solution ψ_n at t_n , then the past state can be re-obtained from ψ_{n+1} by using TSSP with time step $-\Delta t = -(t_{n+1} - t_n)$. Time-transverse invariance or gauge invariance refers to the property of the GPE that if $V \rightarrow V + \alpha$ where $\alpha \in \mathbb{R}$, then the solution $\psi \rightarrow \psi e^{i\alpha t}$, which implies that number density of atoms $|\psi|^2$ is unchanged under this transformation. This method respects this property by producing the corresponding changes in ψ when V is transformed in this way [2].

2.3 Implementation and Validation

I have implemented this spectral method in Python¹. I used the CuPy package² to implement this method because this package provides a NumPy-like environment that performs computations on a GPU. This allowed my code to achieve considerable speedup (almost an order of magnitude for certain problems) over a serial implementation. I will now consider some of the test cases that I have used to ensure I have correctly implemented the time-splitting spectral method and the imaginary-time propagation method.

2.3.1 Homogeneous Solution

It can be trivially verified that

$$\psi(t) = Ae^{-ig|A|^2 t}, \quad A \in \mathbb{C} \quad (10)$$

is a solution to (8). It should be noted that the method will accurately solve a constant initial condition ($\psi(\mathbf{x}, 0) = A$) only when the time step is small compared to the period of oscillation which is determined by angular frequency $\omega = g|A|^2$. In the file [homogeneous_test.py](#), I solve a homogeneous initial condition with $A = \frac{1}{2}$ and $g = 10$ thus giving $T = \frac{2\pi}{\omega} = 2.51$. Our time step is $\Delta t \approx 0.0179$. For more details on parameters, see the file linked in this document.

After solving the problem with these initial conditions until $t = 2.5$. When the code is run using a GPU, the maximum error at any one gridpoint is on the order of 10^{-14} and the normalization loss ratio (the change in the normalization divided by the norm at $t = 0$) is also on the order of 10^{-14} . Running the code in serial gives a max error and norm loss ratio of the same order of magnitude.

2.3.2 Plane Wave

It is also possible to show that the Gross-Pitaevskii Equation admits a plane wave solution with the form

$$\psi(\mathbf{r}, t) = Ae^{i(\mathbf{k} \cdot \mathbf{r} - \omega t)}, \quad A \in \mathbb{C} \quad (11)$$

provided that ω and \vec{k} satisfy the dispersion relation

$$\omega(\mathbf{k}) = \frac{1}{2}|\mathbf{k}|^2 + g|A|^2 \quad (12)$$

In the file [planewave_test.py](#), I use this analytic solution to again test the accuracy of my solver. I use the same value for A and g but with $\vec{k} = (\frac{2\pi}{L_x}, 2\frac{2\pi}{L_y}, 3\frac{2\pi}{L_z})$ the period is now $T = \frac{2\pi}{\omega} = 1.97$. Solving this plane wave initial condition until $t = 2.5$ gives a max error on the order of 10^{-13} and a norm loss ratio on the order of 10^{-14} . Changing from GPU to serial has no significant effect on the error and loss of normalization.

Because this initial condition is spatially varying, it gives us a little more information on how well the solver is working. The spatial derivative in (8) is handled entirely separately from the potential and nonlinear terms. The TSSP method splits (8) into two equations that each be solved exactly and then combines the

¹My [github](#) repository

²CuPy [website](#)

splitting steps together using Strang splitting to give a second-order accurate solution. The equation with the Laplacian is solved using fourier transforms, and in the code this is handled with FFT. This test and the previous one are useful for testing these different components and isolating which part of the solver has a problem if any.

2.3.3 Thomas-Fermi Ground State

The previous two test cases are meant to ensure that the code is correctly solving equation (8), but it is also necessary to test that I have correctly implemented the imaginary-time propagation method. Though we have no exact ground state to which we can compare our numerical result, it is possible to write down an approximate ground state for certain limiting cases of the GPE [3, 4].

As explained in the references just cited, a stationary solution $\phi(\mathbf{x})$ of the GPE can be written as

$$\psi(\mathbf{r}, t) = e^{-i\mu t} \phi(\mathbf{r}) \quad (13)$$

where μ is the chemical potential. Plugging (13) into (8) gives us the equation $\phi(\mathbf{x})$ must satisfy to be a stationary solution

$$\mu\phi(\mathbf{r}) = -\frac{1}{2}\nabla^2\phi(\mathbf{r}) + V(\mathbf{r})\phi(\mathbf{r}) + g|\phi(\mathbf{r})|^2\phi(\mathbf{r}) \quad (14)$$

with the normalization condition

$$\int_{\mathbb{R}^3} |\phi(\mathbf{r})|^2 d\mathbf{r} = 1. \quad (15)$$

The ground state is the stationary solution satisfying (14) and (15) while also minimizing the energy functional

$$E[\phi] := \int_{\mathbb{R}^3} \left[\frac{1}{2} |\nabla\phi|^2 + V(\mathbf{r})|\phi|^2 + \frac{g}{2} |\phi|^4 \right] d\mathbf{r} \quad (16)$$

This ground state solution is unique if we also require that ϕ be real. In the strongly-interacting regime where $gN \gg a_0$ and $g > 0$, the ground state can be approximated as

$$\mu_g = \frac{1}{2} \left(\frac{15gN\gamma_y\gamma_z}{4\pi} \right)^{2/5}, \quad \phi_g(\mathbf{x}) = \begin{cases} \sqrt{\frac{\mu_g - V(\mathbf{x})}{gN}}, & V(\mathbf{x}) < \mu_g \\ 0, & \text{otherwise} \end{cases} \quad (17)$$

This approximate ground state is known as the Thomas-Fermi ground state. It can be derived from the Thomas-Fermi approximation where the kinetic term in (8) is dropped from the equation.

In the file `TFgs_test.py`, I have used my implementation of the imaginary-time propagation method to find the ground state in a trap with $\omega_x = \omega_y = \omega_z = 1$ and $gN = 2000$. On a grid of points $64 \times 64 \times 64$ points, I have compared my numerically calculated ground state with the analytic approximation given in (17), and I have provided a plot of each in figure 1.

To compare how close the numerical solution is to the analytic approximation, I used equation (16) to calculate the ground state energy for each. The numerical solution has an energy of 8.28 and the analytic approximation has an energy of 8.37. The max error between the wavefunction for each was 0.010.

These figures are not as promising as the tests for the GPE solver were, but there may be good reason for this. The energy of the Thomas-Fermi ground state is actually infinite, namely

$$E[\phi_g] = +\infty \quad (18)$$

due to the low regularity of ϕ_g at the free boundary where $V(x) = \mu_g$. To remedy this issue, an interface layer has to be constructed at this boundary to improve the accuracy of the approximation [4].




Figure 1: These are plots of $|\psi|^2$ for the numerically calculated solution (a) and the analytic approximation (b). The x-dimension has been integrated out to make plotting possible. The two are very similar except at the boundary where the analytic approximation becomes zero.

3 Results

3.1 Experimental Setup

In the experiment, the lithium BEC is confined in a cylindrically symmetric harmonic trap. Let ω_ρ be the trapping frequency in the radial direction and ω_z the trapping frequency in the axial direction (i.e., along the z-axis). The experimental parameters are

$$\left(\frac{\omega_\rho}{2\pi}, \frac{\omega_z}{2\pi}\right) = (476 \text{ Hz}, 7 \text{ Hz}), \quad a_0 = 1.74 \mu\text{m}, \quad N \approx 7 \times 10^5, \quad a_s \approx 4r_{\text{Bohr}}, \quad t_{\text{mod}} = 5 \text{ ms} \quad (19)$$

Because $\omega_\rho \gg \omega_z$, the BEC in its ground state is highly elongated along the axial direction. In the experiment, the scattering length is modulated periodically such that

$$g(t) = \begin{cases} g_0(1 + \epsilon \sin(\omega t)) & \text{if } 0 \leq t \leq t_{\text{mod}} \\ g_0 & \text{else} \end{cases} \quad (20)$$

where g_0 is the value for $g = 4\pi a_s$ before the experiment begins, t_{mod} is the length of time over which the scattering length is modulated, and $\epsilon = 0.2$. The modulation frequency of the scattering length ω varies depending on the experiment being performed.

When the scattering length is modulated in this way, the BEC expands and contracts within the trapping potential, and continues to do so after the scattering length is kept fixed. This expansion and contraction sets up a persistent modulation of the width which in turn drives the growth of periodic spatial perturbations along the axial direction of the BEC. The rate of growth for these spatial perturbations is dependent on the perturbation's wavenumber. There is one mode in particular that will grow faster than all other modes which we call k_{max} , and the timescale on which this dominant growing mode will emerge is dependent on the scattering length modulation frequency ω [1].

In order to reconstruct the experiment in numerical simulation, these parameters must be converted to dimensionless form as described in section 2.1. Doing so we obtain

$$(\omega_\rho, \omega_z) = (1, \frac{7}{476}), \quad g \approx 0.001528, \quad gN \approx 1070, \quad t_{\text{mod}} = 14.952 \approx 5\pi. \quad (21)$$




Figure 2: Energy of condensate over time

We have computationally simulated a condensate with these conditions where $\omega = 2\omega_\rho$, and an animation of its time evolution can be downloaded [here](#). With this simulation, we can test how well energy is conserved by the numerical method. Unfortunately, the time-splitting spectral method is not guaranteed to conserve energy, so checking that the energy is conserved can be an indicator that the time-step is not too large. Figure 2 plots how the energy of the condensate changes with time. As we expect, the energy is increasing before t_{mod} as a result of the changing scattering length. After t_{mod} the scattering length is held constant and the energy remains relatively constant as well.

3.2 Computational Results

Now, we analyze computational results that can be directly compared with experiment. Due to limitations on instrument precision in the laboratory, certain properties of the BEC cannot be measured. The results that can be compared with experimental measurement include: (1) how the time it takes for the dominant mode to reach its maximum amplitude varies with the scattering length modulation frequency ω and (2) how the wavenumber of the dominant mode varies with ω .

First, we find the ground state for the trapping potential and experimental parameters described above and then evolve the system forward in time. After a long enough time has elapsed, the small spatial perturbations along the axial direction will grow until a dominant mode emerges. We will use a Fourier

transformation to analyze the growth and amplitude of each mode for the spatial perturbations. One of the modes grows in amplitude until reaching a maximum. Afterwards, this dominant mode will have a decaying amplitude until it grows in size again. We denote as t_{max} the time it takes for this dominant mode to reach its maximum. In figure 3, I have plotted how t_{max} varies with the modulation frequency. We see that t_{max} increases as ω moves away from the resonant frequency, which is $2\omega_\rho$ or twice the radial trapping frequency [1]. See figure 4 to view this plot with the units used in experiment rather than in dimensionless form.




Figure 3: This plot compares t_{max} with the modulation frequency ω . The maximum emerges the soonest when the modulation frequency is at its resonant frequency, which is twice the radial trapping frequency.

Next, we look at how the wavenumber of the dominant mode varies with ω . Theoretical analysis predicts that the wavenumber of the dominant mode should always be $k_{dom} = 1.1$ or $k_{dom} = 0.632 \mu\text{m}^{-1}$ should be independent of ω [1]. Our computational results are close to this prediction, with the wavenumber of the dominant mode remaining close to this analytic prediction. In figure 5, we have plotted how the wavenumber of the dominant mode varies with the modulation frequency.




Figure 5: This plot compares k_{\max} with the modulation frequency ω .

3.3 Island Formation

One notable phenomenon in the animations that we have produced from these simulations is that modulating the scattering length produces an “island” or isolated accumulation of atoms to the left and right of the central cloud. For simulations with the parameters given in subsection 3.2, the islands tend to form around $t = 285$ as illustrated in figure 7. The full 1D animation of these dynamics can be viewed [here](#). Additionally, we have created a 2D animation of the same simulation, though now displayed as a heat map, which can be viewed [here](#). I speculate the the island formation is the result of the periodic expansion and contraction of the atomic cloud and the growth of spatial patterns on the BEC surface that results from this expansion and contraction, rather than being directly caused by the periodic modulation of the scattering length *per se*.

At the very least, it can be shown that this expansion and contraction can also be produced by modulating the potential for a fixed period of time, which also ultimately produces similar dynamics by causing the atomic cloud to expand and contract. In this 1D animation and in this 2D heat map animation




Figure 7: This plot shows the islands around the time when they first form after the scattering length has been modulated at the resonant frequency. This plot shows the number density of atoms along the z -axis when $t = 285$.

References

- [1] M. Amin. Notes: Resonant emergence of spatio-temporal patterns in elongated bose-einstein condensates.
- [2] X. Antoine, W. Bao, and C. Besse. Computational methods for the dynamics of the nonlinear schrödinger/gross-pitaevskii equations. *Computer Physics Communications*, 184(12):2621–2633, 2013.
- [3] W. Bao, D. Jaksch, and P. A. Markowich. Numerical solution of the gross-pitaevskii equation for bose-einstein condensation. *Journal of Computational Physics*, 187(1):318–342, 2003.
- [4] W. Bao and W. Tang. Ground-state solution of bose-einstein condensate by directly minimizing the energy functional. *Journal of Computational Physics*, 187(1):230–254, 2003.
- [5] M. L. Chiofalo, S. Succi, and M. Tosi. Ground state of trapped interacting bose-einstein condensates by an explicit imaginary-time algorithm. *Physical Review E*, 62(5):7438, 2000.
- [6] P. Muruganandam and S. K. Adhikari. Fortran programs for the time-dependent gross-pitaevskii equation in a fully anisotropic trap. *Computer Physics Communications*, 180(10):1888–1912, 2009.
- [7] J. H. Nguyen, D. Luo, and R. G. Hulet. Formation of matter-wave soliton trains by modulational instability. *Science*, 356(6336):422–426, 2017.

- [8] C. J. Pethick and H. Smith. *Bose-Einstein condensation in dilute gases*. Cambridge university press, 2008.
- [9] D. V. Schroeder. An introduction to thermal physics, 1999.




Figure 4: This plot compares t_{max} with the modulation frequency ω . The maximum emerges the soonest when the modulation frequency is at its resonant frequency, which is twice the radial trapping frequency.




Figure 6: This plot compares k_{\max} with the modulation frequency ω .